This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

Gift of
Estate of
Dr. Herman Knoche

CONSPECTUS

FLORAE GRAECAE

AUCTORE

E. DE HALÁCSY

VOLUMEN II

LIPSIAE

SUMPTIBUS GUILELMI ENGELMANN

1902.

QK315
H3
v. 2

581.9495
H57
v. 2

Jus traductionis reservatum.

Wirceburgi, ex typis H. Stürtz.

LXII. Ordo. **Compositae** Vaill. in act. acad. Paris 1718 p. 143.

Dispositio tribuum generumque:

I. Divisio. *Tubuliflorae* DC. pr. V. p. 8. — Corollae omnes tubulosae vel marginales ligulatae.

1. Subdivisio. *Corymbiferae* Juss. gen. p. 177. — Stylus sub apice neque articulatus nec nodoso-inflatus.

1. Tribus. *Eupatorieae* Less. syn. p. 156. — Capitula homogama vel heterogama, interdum subdioica; receptaculum nudum; antherae ecaudatae; styli crura semicylindrica vel cylindrica, obtusa; achenia cylindrica, costata, pappo piloso coronata.

1. Subtribus. *Adenostyleae* DC. pr. V. p. 103. — Capitula homogama; corollae tubulosae; antherae basi rotundatae.

1. Eupatorium L.

2. Subtribus. *Tussilagineae* Less. syn. p. 158. — Capitula heterogama, plerumque subdioica; corollae florum femineorum limbo oblique truncato vel ligulato, florum hermaphroditorum vel abortu masculorum tubulosae; antherae basi excisae, lobis rotundatis.

a. Capitula heterogama-subdioica, in thyrsum disposita.

2. Petasites Tourn.

b. Capitulum heterogamum-monoicum, unicum, terminale.

3. Tussilago L.

2. Tribus. *Astereae* N. ab E. gen. et sp. ast. p. 158. — Capitula heterogama, rarius homogama; receptaculum nudum; antherae basi rotundatae, ecaudatae; styli crura compressa, apice rotundata, nuda; achenia compressa, ecostata, raro cylindrica, costata, pappo piloso, raro nullo vel scarioso coronata.

1. Subtribus. *Bellieae* DC. pr. V. p. 302, ampl. — Achenia antice compressa, marginata, ecostata, pappo nullo vel scarioso coronaeformi vel e paleis scariosis et setis alternantibus composito.

a. Pappus nullus vel scariosus, brevis, coronaeformis.

4. Bellis L.

b. Pappus e paleis scariosis setisque alternantibus compositus.

5. Bellium L.

2. Subtribus. *Erigeroneae* Gr. et Godr. fl. fr. II. p. 92. — Achenia antice compressa, raro cylindrica, plerumque ecostata, pappo piloso.

a. Achenia compressa, ecostata.

α. Flores marginales feminei, anguste ligulati, disci tubulosi; pappus uniserialis.

6. Erigeron L.

β. Flores marginales feminei vel neutri, ligulati, disci tubulosi; pappus pluriserialis.

7. Aster L.

b. Achenia teretiuscula, costata; flores marginales feminei, ligulati, disci tubulosi; pappus uniserialis.

8. Solidago L.

3. Tribus. *Inuleae* Cass. in ann. sc. nat. 1829 p. 20. — Capitula heterogama vel homogama; receptaculum nudum vel paleaceum; antherae basi caudatae vel setiferae (in gen. Phagnalon interdum ecaudatae); styli crura linearia, obtusa, rarius truncata, non appendiculata, saepius clavata, subcompressa; achenia varia, pappo piloso vel scarioso coronaeformi, rarius setoso vel nullo.

1. Subtribus. *Buphthalmeae* Less. syn. p. 209. — Receptaculum rigide paleaceum; styli crura compressa, apice rotundata pubescentia; achenia conformia vel difformia, marginalia triquetra, cetera oblongo-obconica, facie interna carinata, omnia pappo scarioso coronaeformi lacero munita.

a. Involucri phylla externa inermia; flosculi radii uniseriales, ligulati, 3 dentati, basi biauriculati, disci tubulosi, teretes, 5 dentati, tubo basi incrassato; achenia conformia, subtriquetra, aptera.

9. Asteriscus Moench.

b. Involucri phylla externa spinescentia; flosculi radii biseriales, ligulati, acute trifidi, basi bialati, disci tubulosi, 5 fidi, tubo interne unialato; achenia difformia, radii compresso-bialata, triquetra, disci compresso-triquetra.

10. Pallenis Cass.

2. Subtribus. *Euinuleae* Bois. fl. or. III. p. 184. — Receptaculum nudum; styli crura compressa, apice rotundata, pubescentia; achenia conformia, tetragona vel cylindrica, pappo piloso.

a. Pappus uniserialis, setis conformibus.

11. Inula L.

b. Pappus biserialis, serie externa coronaeformi dentata persistente, interna setis paucis constante.

12. Pulicaria Gaertn.

3. Subtribus. *Gnaphalieae* Less. in Linnaea 1831 p. 221. — Receptaculum nudum aut marginem versus paleis obsitum; styli crura apice truncata vel obtusa; achenia cylindrica vel compressa, pappo plerumque piloso, rarius setoso aut nullo.

a. Receptaculum nudum.

α. Antherae basi sagittata breviter et tenuiter caudatae, interdum ecaudatae; achenia cylindrica, pappo piloso. Suffrutices.

13. Phagnalon Cass.

β. Antherae basi caudatae; achenia oblongo-cylindrica, pappo piloso.

× Flores omnes tubulosi, hermaphroditi, rarius nonnulli exteriores feminei uniseriales angustiores.

14. Helichrysum L.

×× Flores exteriores feminei filiformes pluriseriales, rarius subuniseriales, reliqui tubulosi hermaphroditi.

15. Gnaphalium L.

b. Receptaculum margine vel inter flores femineos paleaceum, ceterum nudum.

α. Achenia pappo piloso fragili caduco munita.

16. Filago L.

β. Achenia epapposa.

× Flores marginales feminei pluriseriales; receptaculum elongato-conicum, summo apice inter flores masculos nudum, inter flores femineos paleaceum.

17. Evax Gaertn.

×× Flores marginales feminei uniseriales; receptaculum angustum, ad peripheriam paleis uniserialibus flores femineos includentibus obsitum

18. Micropus L.

4. Tribus. *Heliantheae* Less. syn. p. 221. — Capitula heterogama vel homogama; receptaculum paleaceum; antherae basi ecaudatae; styli crura truncata vel breviter appendiculata; achenia conformia, pappo aristaeformi vel nullo.

19. Bidens L.

5. Tribus. *Anthemideae* Cass. opusc. phyt. III. p. 61. — Capitula heterogama vel rarius homogama; receptaculum paleaceum vel nudum; antherae basi ecaudatae; styli crura apice rotundata vel truncata, penicillata; achenia varia, pappo coronaeformi vel nullo.

1. Subtribus. *Euanthemideae* DC. pr. VI. p. 1. — Receptaculum paleaceum.

a. Capitula plerumque parva, corymbosa, radio brevi vel nullo; achenia compressa, disco epigyno ecoronato.

α. Capitula discoidea, homogama.

20. Diotis Desf.

β. Capitula radiata, heterogama.

21. Achillea L.

b. Capitula plerumque majuscula, solitaria, radio elongato rarius nullo, achenia varia, disco epigyno latere interno saepius membranaceo-coronato.

α. Limbus corollarum disci aequaliter 5fidus; achenia non alata.

× Tubus corollarum disci compressus, basi demum ampliatus; achenia obconica, circacircum costata, apice nunc calva, nunc margine coronaeformi, nunc auricula laterali superata.

22. Anthemis L.

×× Tubus corollarum disci teres, basi lateraliter in calcar ampliatus, achenia caduca, obovata, subcompressa, apice rotundata, calva, facie exteriori laevia, interiori tenuiter 3costata.

23. Ormenis Gay.

β. Limbus corollarum disci 5fidus, lobis duobus (in nostris) ceteris longioribus; achenia compressa, laevia, saltem exteriora utrinque ala lata in auriculam abeunte obsita.

24. Anacyclus Pers.

2. Subtribus. *Chrysanthemeae* DC. pr. VI. p. 38. — Receptaculum epaleaceum, glabrum, rarius pilosum.

a. Capitula saepissime radiata; styli crura non in conum producta; achenia varia, costata, calva vel saepius pappo scarioso, coronae-, auriculae- vel linguaeformi superata.

α. Achenia conformia.

× Achenia latere ventrali costata, dorso laevia.

○ Receptaculum elongato-conicum, cavum; achenia teretiuscula, ventre 5costata, calva vel coronata, eglandulosa.

25. Matricaria L.

○○ Receptaculum hemisphaericum vel conicum, farctum; achenia obpyramidata vel subcompressa, ventre tricostata, calva vel coronata, apice 1—2glandulosa.

26. Chamaemelum Vis.

XX Achenia circumcirca subregulariter costata.

○ Tubus corollarum disci alato-compressus; achenia calva vel corona scariosa munita.

27. Leucanthemum Tourn.

○○ Tubus corollarum disci teres; achenia corona scariosa munita.

. Capitula radiata, radio longo; achenia eglandulosa, corona scariosa varia munita.

28. Pyrethrum Gaertn.

.. Capitula discoidea vel radio abbreviato munita; achenia pellucide-glandulosa, margine membranacco vel corona abbreviata munita.

29. Tanacetum.

β. Achenia difformia, radii 2—3 alata, disci exalata.

30. Chrysanthemum L.

b. Capitula radiata vel discoidea; styli crura non in conum producta, achenia teretia, ecostata, obsolete striata, pappo coronaeformi brevi fimbriato-lacero superata.

31. Ammanthus Bois. et Heldr.

c. Capitula discoidea; styli crura saepius supra penicillum in conum producta; achenia compressa, ecostata, calva.

32. Artemisia L.

6. Tribus. *Senecioneae* Cass. in dict. sc. nat. XX. p. 377. — Capitula heterogama, rarius homogama; receptaculum nudum; antherae ecaudatae, basi breviter sagittatae; styli crura truncata, penicillata; achenia cylindrica, costata, omnia vel saltem disci pappo piloso coronata.

a. Anthodium explanatum, phyllis 2—3 serialibus; achenia radii calva, disci pappo piloso munita.

33. Doronicum L.

b. Anthodium cylindricum vel conicum, phyllis 1 serialibus; achenia omnia pappo piloso munita.

34. Senecio L.

7. Tribus. *Calenduleae* Less. syn. p. 89. — Capitula heterogama; receptaculum nudum; antherae basi sagittatae; styli crura linearia, compressa, crassa; achenia calva, in eodem capitulo saepius heteromorpha.

35. Calendula L.

2. Subdivisio. *Cynarocephalae* Juss. gen. p. 171. — Stylus florum hermaphroditorum superne nodoso-incrassatus, ad nodum saepe penicillatus et articulatus, cruribus liberis vel concretis. Corollae omnes tubulosae, rarius marginales ampliatae, bilabiatae vel ligulatae; receptaculum saepius dense setosum vel paleis onustum; antherae basi sagittatae.

8. Tribus. *Echinopsideae* Less. in Linnaea VI. p. 88. — Capitula uniflora, in receptaculo communi globoso in glomerulum globosum congesta; achenia sericeo-villosa, areola basi recta; pappus coronaeformis.

36. Echinops L.

9. Tribus. *Carlineae* Less. syn. p. 11. — Capitula pluriflora, discreta vel subaggregata; achenia sericea vel villosa (cf. *Staehelina*), areola basilari recta, rarius subobliqua; pappus uniserialis, paleaceus vel setaceus, setis basi in fasciculos plures rarius in annulum concretis.

a. Herbae.

α. Pappi paleae aristato-acuminatae.

× Herbae spinosae, foliis pinnatipartitis, capitulis numerosissimis, subsessilibus, homogamis, pauci- et aequalifloris, in glomerulos bracteato-spinosissimos terminales, corymbosos congestis; receptaculum fimbrillatum.

37. Cardopatium Juss.

×× Herbae inermes, foliis integris, capitulis solitariis, longepedunculatis, heterogamis (floribus externis paucis neutris, corolla parva bilabiata donatis); receptaculum paleis concavis flores includentibus obsitum.

38. Xeranthemum L.

β. Pappi setae plumosae.

× Herbae spinosae; involucri phylla intima plerumque radiato-patentia; receptaculum paleaceum, paleis fissis; achenia adpresse serica; pappi paleae basi ternatim quaternatim in fasciculos concretae.

39. Carlina L.

×× Herbae spinosae; involucri phylla intima erecta, nec radiantia; receptaculum paleaceum, paleis fissis; achenia dense sericeo-villosa; pappi paleae basi in annulum coalitae.

40. Atractylis L.

b. Suffrutices inermes; capitulis angustis, solitariis vel aggregatis; involucri phylla imbricata; receptaculi paleae augustae, lacero-multifidae; achenia glabra vel sericea; pappo e pilis simplicibus, inaequalibus, sericeis, scabris, basi in fasciculos connatis composito.

41. Staehelina L.

10. Tribus. *Carduineae* Cass. tabl. p. 9. — Capitula pluriflora, saepius discreta; achenia glabra, areola basilari recta; pappus pluriserialis, pilosus vel plumosus, setis basi in annulum concretis.

a. Filamenta libera; pappi setae segregatim caducissimae; receptaculum fimbrillosum. Plantae elatae inermes, foliis amplis subintegris, involucri phyllis uncinatis.

42. Lappa Tourn.

b. Filamenta libera vel in vaginam connata; pappi setae basi in annulum concretae, cum eo deciduae.

α. Plantae spinosae.

× Filamenta libera.

○ Filamenta papilloso-pilosa vel glabriuscula; receptaculum fibrilliferum vel setosum.

. Pappus pilosus, scaber.

43. Carduus L.

.. Pappus plumosus.

; Involucri phylla basi non carnosa.

, Receptaculum fibrilliferum; achenia oblonga, compressa, hilo recto, disco epigyno margine brevi integro cincto.

— Involucri phylla in spinam simplicem attenuata vel inermia; stigmata coalita.

44. Cirsium Tourn.

= Involucri phylla in spinam pinnatam attenuata; stigmata libera.

45. Picnomon Lob.

,, Receptaculum lineari-paleaceum; achenia oblique-obovata, compressa, hilo obliquo, disco epigyno non marginato.

46. Notobasis Cass.

;; Involucri phylla basi carnosa, receptaculum carnosum, alveolatum, fibrilliferum.

47. Cynara L.

○○ Filamenta glabra; receptaculum foveolatum, vix brevissime fibrilliferum; pappus scaber vel plumosus.

48. Onopordon L.

XX Filamenta in vaginam connata. .

○ Pappi pili brevissime ciliolati vel denticulati.

. Capitulum homogamum; receptaculum carnosum; filamenta papillosa; antherae breviter appendiculatae.

49 Silybum Vaill.

.. Capitulum heterogamum; receptaculum non carnosum; filamenta basi hirta; antherae obsolete caudatae.

50. Tyrimnus Cass.

○○ Pappus longe plumosus; capitulum heterogamum, floribus marginalibus radiantibus; filamenta papillosa.

51. Galactites Moench.

β. Plantae inermes.

X Suffrutices, foliis lineari-filiformibus; receptaculum fibrilliferum; filamenta plumosa; achenia obovata; pappus plumosus, seriebus aequilongis.

52. Chamaepeuce Prosp. Alp.

XX Herbae (nostrae), foliis variis; receptaculum setosum; filamenta glabra; achenia obpyramidato-tetragona vel subcompressa; pappus barbellatus vel plumosus, serie intima ceteris longiore.

53. Jurinea Cass.

11. Tribus. *Centaurieae* Less. syn. p. 53. — Capitula pluriflora, discreta; achenia glabra, areola basilari obliqua vel laterali; pappus setosus, paleaceus, rarius nullus.

a. Capitula basi nuda, rarius basi foliis nonnullis inermibus suffulta.

α. Pappus conformis, persistens, plumosus vel barbellatus, pluriserialis, seriebus indistinctis, ab externis ad intimas sensim elongatis.

54. Phaeopappus Bois.

β. Pappus biformis (interdum nullus), externi setae vel paleae pluriseriales, ab externis ad intimas elongatae, interni uniseriati paleae vel et saepissime abbreviatae conniventes, vel et rarius elongatae, pappum externum superantes.

X Pappus persistens.

○ Achenia intra pappum interiorem cupula crenulata non superata.

. Pappus (interdum nullus) externus paleis uniserialibus abbreviatis conniventibus constans.

55. Centaurea L.

.. Pappus internus paleis in squamam unilateralem excavatam acuminatis concretis constans.

56. Microlonchus Cass.

XX Pappus in annulum deciduus.

58. Aegialophila Bois. et Heldr.

a. Capitula bracteis spinoso-dentatis involucrata.

α. Pappi setae pluriseriales.

X Pappus e paleis inaequalibus latis scabridis compositus, interdum nullus.

59. Carthamus L.

XX Pappus e setis ciliatis subplumosis, basi in annulum concretis compositus.

60. Carduncellus Ad.

β. Pappus e setis 10 exterioribus longis et 10 interioribus brevibus, cum illis alternantibus constans.

61. Cnicus L.

II. Divisio. *Liguliflorae* DC. pr. VII. p. 74. — Corollae omnes ligulatae, hermaphroditae; stylus sub apice neque articulatus nec nodoso-inflatus; antherae apice appendiculatae, basi sagittatae.

A. Herbae spinosae (Carthami facie); paleae achenia involventes, pericarpium late alatum simulantes.

12. Tribus. *Scolymeae* Less. syn. p. 126. — Receptaculum paleaceum; pappus coroniformis, brevis.

62. Scolymus L.

B. Herbae inermes; paleae cum adsunt achenia non involventes.

I. Receptaculum paleaceum vel setosum.

13. Tribus. *Catanancheae* Schultz in Bonpl. 1860. p. 367. — Receptaculum setosum vel margine paleaceum; pappus longe paleaceus.

a. Receptaculum setoso-fimbrilliferum.

63. Catananche L.

b. Receptaculum ad peripheriam paleis latis 1—2 serialibus obsitum, centro subfimbriatum.

64. Hymenonema Cass.

14. Tribus. *Hypochaerideae* Less. syn. p. 130. — Receptaculum paleaceum; pappus plumosus.

a. Pappus conformis, uni-vel biserialis, radiis plumosis, basi non dilatatis, externis cum adsunt brevibus setiformibus, aut biformis, acheniorum radii in coronam brevem fimbriatam mutatus, disci plumosus.

65. Hypochaeris L.

b. Pappus acheniorum centralium biserialis, radiis internis plumosis, basi paleaceo-dilatatis, externis brevissimis setiformibus; acheniorum marginalium saepius nullus vel abortivus.

66. Seriola L.

15. Tribus. *Rodigieae* DC. pr. VII. p. 98. — Receptaculum paleaceum vel longe setosum; pappus setosus.

a. Receptaculum paleis lanceolatis, acheniorum rostro vix brevioribus obsitum.

67. Rodigia Spreng.

b. Receptaculum setis achenia aequantibus vel superantibus obsitum.

68. Lagoseris M. B.

II. Receptaculum nec paleaceum nec setosum.

§ Achenia calva, aut pappo paleaceo vel cupuliformi, interiora interdum pappo setoso superata.

16. Tribus. *Hyoserideae* Less. syn. p. 127. — Pappus paleaceus vel cupuliformis, paleae saepe in pilum acuminatae, sed neque plumosae, neque a basi piliformes.

a. Ligulae coeruleae (raro roseae vel albae); achenia pappo brevi, 2—3 seriali, paleolis vel setulis constante coronata.

69. Cichorium L.

b. Ligulae flavae.

α. Achenia difformia, externa teretiuscula, breviter apice setulosa, intermedia compressa, late alata, pappo setis brevibus paleisque dilatatis, intima angulato-cylindrica, pappo e paleis longe cuspidatis constante superata, interdum deficientia.

70. Hyoseris L.

β. Achenia omnia teretiuscula, externa pappo brevi cupuliformi vel pelviformi, laciniato-denticulato, interna paleis lanceolatis vel subulatis, setis brevioribus stipatis, superata.

71. Hedypnois Tourn.

17. Tribus. *Lapsaneae* Less. syn. p. 126. — Achenia saltem marginalia calva vel coronulata, interiora interdum pappo setoso superata.

a. Achenia subtetragona, peripherica pappo brevissimo, scarioso, fimbriato, reliqua pappo eodem setisque longis asperis superata.

72. Tolpis Gaertn.

b. Achenia plus minus compressa, calva.

α. Involucri phylla externa minima, interna 8—10, fructifera immutata, erecto-conniventia; achenia compressa, striata, margine obsoleto terminata, decidua.

73. Lapsana L.

β. Involucri phylla externa minima, interna 5—8, fructifera accreta, indurata, patentia; achenia a dorso subcompressa, ecostata, omnino epapposa, externa persistentia.

74. Rhagadiolus Tourn.

§§. Pappus acheniorum interiorum saltem plumosus.

18. Tribus. *Leontodonteae* Schultz in Koch. syn. p. 417. — Pappus acheniorum omnium aut saltem interiorum plumosus, pilis setarum non intertextis, aut pappus acheniorum marginalium coroniformis.

a. Involucrum imbricatum.

α. Pappus difformis, acheniorum marginalium brevis, coroniformis, dentatus, acheniorum disci plumosus, radiis basi dilatatis.

75. Thrincia Roth.

β. Pappus omnium acheniorum conformis, radiis 10 vel pluribus, basi interdum dilatatis, plumosis, additis interdum setis nonnullis externis brevibus piliformibus vel plumosis.

× Caulis aphyllus; pappus persistens, radiis liberis, interdum basi dilatatis.

76. Leontodon L.

×× Caulis foliatus; pappus (in nostris) deciduus, radiis basi in annulum concretis, interioribus basi dilatatis.

77. Picris L.

b. Involucrum duplex, externum phyllis 3—5 foliaceis latis, interius subbiseriale, phyllis angustis; achenia rugulosa, in rostrum capillare ruptile et cum pappo uniseriali secedens, abrupte abeuntia.

78. Helminthia Juss.

c. Involucrum simplex, 8 phyllum, phyllis basi connatis; achenia muricato-lamellata, rostro cum achenio articulato, basi fistuloso et e semine septo separato, superata.

79. Urospermum Juss.

19. Tribus. *Scorzoneereae* Schultz in Koch syn. p. 422. — Pappus acheniorum omnium aut saltem interiorum plumosus, pilis setarum intertextis.

a. Receptaculum ad peripheriam alveolato-fibrillosum; involucrum simplex, 8 phyllum, uniseriale; achenia in rostrum longum sensim attenuata, marginalia paleis 5 scabridis superata, disci multiradiata, radiis plumosis.

80. Geropogon L.

b. Receptaculum nudum.

α. Involucrum simplex, phyllis 5—16 subaequalibus, basi subconnatis; achenia in rostrum attenuata, raro erostria, pappo conformi, multiradiato, plumoso.

81. Tragopogon L.

β. Involucrum imbricatum.

× Achenia apicem versus attenuata, erostria, basi callo brevi nunquam inflato suffulta; pappus plumosus, radiis numerosis saepius inaequalibus, longioribus tum apice nudis; folia integerrima.

82. Scorzonera L.

×× Achenia apicem versus non attenuata, erostria, basi callo elongato subinflato, achenio ipso crassiore suffulta; pappus plumosus, radiis numerosis aequalibus; folia divisa.

83. Podospermum DC.

§§§ Pappus pilosus.

20. Tribus. *Chondrilleae* Koch syn. p. 427. — Achenia teretiuscula, saepius rostrata et (in nostris) superne muricata vel coronula squamularum cincta.

a. Involucrum imbricatum, subcalyculatum; flores multiseriales; achenia subcompresso-tetragona, undique vel superne squamuloso-muricata, in rostrum filiforme abeuntia; herbae acaules.

84. Taraxacum Hall.

b. Involucrum calyculatum, oligophyllum; flores 7—12, biseriales; achenia pentagona, superne squamuloso-muricata et coronula basin rostri cingente terminata; herbae caulescentes, ramosissinae.

85. Chondrilla L.

21. Tribus. *Crepideae* Bois. fl. or. III. p. 712. — Achenia compressa vel teretia, nuda.

a. Achenia plus minus a dorso compressa.

α. Achenia erostria.

× Involucrum imbricatum; flores multiseriales, flavi.

86. Sonchus L.

×× Involucrum subcalyculatum; flores 5, uniseriales, purpurei.

87. Prenanthes L.

β. Achenia rostrata; involucrum imbricatum, interdum subbiseriale; flores 2—3 seriales.

88. Lactuca L.

b. Achenia prismatica vel teretia, interdum a latere compressa.

α. Achenia biformia.

× Involucrum calyculatum, fructiferum angulato-torulosum, phyllis interioribus calloso-incrassatis, achenia marginalia foventibus; achenia marginalia a latere compressa, dorso valde gibba, in rostrum brevissimum attenuata, disci teretiuscula, erostria.

89. Zacintha Gaertn.

×× Involucrum imbricatum, fructiferum immutatum.

○ Capitulum pauciflorum; achenia externa 4—5 gona, 4—5 sulcata, transverse profunde crenato-rugosa, interna a basi sensim attenuata, subconica, saepe sublaevia.

90. Picridium Desf.

2. Tribus. **ASTEREAE** N. a. E. gen. et sp. ast. p. 158.

4. Bellis L. gen. n. 962.

a. Perennes, scapigerae.

α. Folia obovato-spathulata, in petiolum abrupte contracta.

1. **B. perennis** L. sp. p. 886; S. et S. pr. II. p. 184; Pieri corr. fl. p. 119; Ch. et B. exp. p. 251. Fl. pelop. p. 58; Friedr. Reise p. 267; Fraas fl. class. p. 213; Raul. cret. p. 748; Bois. fl. or. III. p. 173, cum β. *microcephala* (f. capitulis minoribus, non *B. microcephala* Lange): Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 43; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1896 p. 42, 1897 p. 32; Hal. Beitr. fl. Epir. p. 28, Beitr. fl. Thessal. p. 14, Beitr. fl. Achaia p. 24, f. *microcephala* Bois.; Haussk. symb. p. 114. — *B. longifolia* Bois. fl. or. suppl. p. 290. — *B. amoena* Heldr. it. gr. septentr. a. 1879. — Icon: Fl. dan. t. 503. — Exsicc.: Hal. it. gr. II. a. 1803.

Laete virens, pilis papillosis plus minus obsita vel subglabra; foliis rosulatis, uninerviis, obsolete crenatis vel serratis, scapis apice incrassatis, involucri phyllis oblongis, obtusis; ligulis albis vel subtus purpurascentibus.

In herbidis, pascuis ut videtur praesertim montanis et subalpinis, in regionem alpinam adscendens. Epirus: mt. Tsumerka, Strungula, Peristeri (Hal.); Thessalia: mt. Oxya (Hal.), Ghavellu, in oropedio Neuropolis, ad monasterium Korona (Haussk.), mt. Said Pascha, Karava, Baba inter Krania et Klinovo in Pindo, pr. Rapsani in Olympo (Form.); Eurytania: mt. Chelidoni (Heldr.); Attica: ad Cephissum pr. Athenas (Friedr.); Elis: pr. Lechaena, mt. Olenos (Heldr.); Achaia: mt. Panachaicon, Chelmos, Kyllene (Hal.); Argolis: pr. Tirynthum (Bois.); Laconia: mt. Taygetos (Chaub.); Creta: pr. Canea (Heldr.); Cephalonia: pr. Argostoli (Heldr.); Corcyra (Pieri); loca nonnulla ad sequentem pertinere videntur. — Mart. Aug. ♃.

2. **B. hybrida** Ten. syll. p. 436. — *B. perennis* × *silvestris* Haussk. symb. p. 14. — Exsicc.: Heldr. herb. norm. n. 944; Sint. it. thessal. n. 85; Baen. herb. europ. n. 9215. —

Differt a praecedente, quacum intermediis conjuncta esse videtur, indumento saepius copiosiori, foliis majoribus trinerviis, evidentius et argutius dentatis, scapis elatioribus, basi saepe 1—2 foliatis, capitulis saepe majoribus.

In herbidis, humidiusculis regionis inferioris et montanae, praecedenti ut videtur multo vulgatior. Thessalia: mt. Karava (Heldr.), ad Lechonia pr. Volo (Sint.); Attica: mt. Parnes, in valle Cephissi pr. Chelidonu (Heldr.); Achaia: mt. Omplo pr. Patras (Heldr.); Corcyra: pr. urbem (Sagburg), Kastrades (Baenitz.); — huc certe loca plura ad speciem praecedentem indicata. — Febr. Jun. ♃.

β. **Folia** oblongo-spathulata, sensim in petiolum attenuata

3. **B. silvestris** Cyr. pl. rar. II. p. 42, t. 4; Ch. et B. exp. p. 250, Fl. pelop. p. 58; Marg. et R. fl. Zante p. 56; Friedr. Reise p. 274; Ung. Reise p. 123; Raul. cret. p. 778; Bois. fl. or. III. p. 74; Heldr. fl. cephal. p. 43, Chlor. mykon. p. 245 *v. glabrescens*; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 115; Bald. riv. coll. bot Alb. 1896 p. 65. — *Doronicum bellidiastrum* S. et S. pr. II. p. 183, non L. — Exsicc.: Orph. fl. gr. n. 476; Heldr. herb. norm. n. 1350.

Obscure viridis, pubescens, foliis rosulatis, trinerviis, obsolete denticulatis vel serratis, scapis apice non vel vix incrassatis; involucri phyllis oblongo-lanceolatis, acutiusculis; ligulis albis vel roseis.

In herbidis regionis inferioris et submontanae. Epirus: mt. Micikeli (Bald.); Thessalia: pr. Litochoron (Krüper); Attica: pr. Cephissia, Tatoi (Heldr.); Corinthia: pr. Corinthum (Orph.), Kalamaki (Haussk.); Achaia: pr. Patras (Heldr.); Argolis: pr. Nauplia (Sart.), Poros (Friedr.); Messenia: pr. Phigalea, Messene, Methone, Navarin (Chaub.), Kalamata (Zahn); Cycladum insula Melos (Leon.), Mykonos (Heldr.); Creta: pr. Platania (Reverch.), Malaxa (Raul.); Zante (Sibth.); Cephalonia (Schimp.); Corcyra (Ung.). — Oct. Mart. ♃.

4. **B. longifolia** Bois. et Heldr. diagn. XI. p. 1, Fl. or. III. p. 174; Raul. cret. p. 778. — Exsicc.: Rev. pl. cret. n. 253. —

Differt a praecedente foliis uninerviis, capitulis minoribus, involucri phyllis obtusissimis.

β. **glabrata** Raul. cret. p. 778. — Subglabra. — Exsicc.: Bald. it. cret. alt. n. 173. —

In regione subalpina Cretae: ad Amalos in mt. Sphacioticis (Rev.), mt. Aphendi-Kavutsi (Heldr.); — *β.* mt. Aphendi-Lassiti (Heldr.). — Apr. Jun. ♃.

b. Annuae, caulescentes.

5. **B. annua** L. sp. p. 887; S. et S. pr. II. p. 184; Sieb. avis p. 5; Ten. annot. p. 19; Ch. et B. exp. p. 251, Fl. pelop. p. 58; Marg. et R. fl. Zante p. 56, Friedr. Reise p. 267; Ung. Reise p. 123; Raul. cret. p. 778; Bois. fl. or. III. p. 175; Heldr. fl. cephal. p. 43; Haussk. symb. p. 114. — *B. dentata* DC. pr. V. p. 304; Weiss in z. b. G. 1869 p. 42. — *B. annua v. dentata* Clem. p. 59. — Icon: Fl. gr. t. 876. — Exsicc.: Orph. fl. gr. n. 34; Heldr. herb. fl. hellen. n. 54 et 55; Baen. herb. europ. n. 9214.

Puberula; caulibus pumilis, simplicibus vel basi ramosis, inferne foliatis, rarius scapiformibus; foliis obovato-spathulatis, dentatis crenatisve, in petiolum abrupte attenuatis; scapis apice vix incrassatis; involucri phyllis ellipticis, obtusis; ligulis albis, subtus purpurascentibus.

β. **minuta** DC. pr. IV. p. 304. — Minuta, capitulis minimis. — Habitu *Bellio minuto* similis, ab eo pappo coronaeformi discedens. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889.

In arenosis, pascuis, collibus regionis inferioris. Attica: pr. Athenas, ad Piraeum, Phaleram (Heldr.); Peloponnesus (Sibth.): pr. Arcadia Messeniae (Chaub.); Creta: pr. Canea (Raul.); Zante, ad Maltri (Marg.); Cephalonia (Heldr.); Corcyra: pr. Potamo (Ung.). — ♂ Cycladum insula Melos (Leon.). — Jan. Maio. ⊙.

5. Bellium L. mant. p. 157.

1. **B. minutum** L. mant. p. 286; Raul. cret. p. 778; Bois. fl. or. III. p. 173; Heldr. chlor. Thera p. 14. — *B. bellidioides* S. et S. pr. II. p. 184; Urv. enum. p. 112; non L., quod ex Bois. l. c. phyllis involucri 12—14 oblongo-lanceolatis, ligulis multo numerosioribus, linearibus, involucro plus duplo longioribus differt. — Exsicc.: Forsyth Major n. 802 (ex insula Kassos).

Nanum, papilloso-hirtulum; foliis omnibus radicalibus, ovato-spathulatis, obtusissimis, in petiolum eis sublongiorem attenuatis; scapis folio sublongioribus; capitulis minimis; involucri phyllis 7—8, subbiseriatis, ovato-oblongis, cucullatis, obtusissimis, margine scarioso-rubellis; ligulis tubo aequilongis, involucro sublongioribus; acheniis hirtis; pappi squamis 4, ovato-orbiculatis, truncatis, setis 4, 3—4plo longioribus, tubum corollae aequantibus.

In rupestribus. Euboea (Sibth.); Cycladum insula Therasia pr. Thera (Urv.); Creta (Tourn. cor. 37). — Apr. Jun. ⊙. N. v.

6. Erigeron L. gen. n. 951.

a. Flores femimei marginales discum aequantes.

α. Flores feminei marginales involucro parum longiores.

1. **E. linifolium** Willd. sp. pl. III. p. 1955; Bois. fl. or. III. p. 169; Heldr. fl. cephal. p. 43; Bald. viagg. cret. p. 61. — *Conyza ambigua* DC. fl. fr. V. p. 468; Haussk. symb. p. 119. — *E. crispum* Pourr. chl. narb. n. 470. — Exsicc.: Heldr. herb. norm. n. 332.

Cinereo-virens, scabridum; caule erecto, simplici vel apicem versus ramoso, folioso, ramis axim primariam superantibus; foliis undique breviter hirtis, inferioribus lanceolatis, in petiolum attenuatis, ceteris linearibus, sessilibus; capitulis laxe corymbosis, mediocribus; involucri phyllis subuniseriatis, linearibus, acuminatis, pappo brevioribus; floribus marginalibus pluriserialibus, filiformibus, flavescentibus, truncato 2—3 dentatis; acheniis parce hirtulis, pappo rufescenti 15—20seto triplo brevioribus.

In ruderatis, locis aridis, ad vias in cultis regionis inferioris. Thessalia: pr. Volo (Haussk.); Aetolia: pr. Mesolongion (Nieder); Attica: pr. Athenas (Heldr.); Achaia: pr. Aegion (Heldr.); Argolis: pr. Vromolimni in peninsula Methana (Heldr.); pr. Methone Messeniae, ad radices mt. Taygetos (Chaub.); Creta: pr. Canea (Bald.); Cephalonia: pr. Pessada. [illegible] Kutavo (Heldr.). — Maio, Sept. ⊙.

[illegible]dense L. sp. p. 863; Ch. et B. exp. p. 248. Fl. pelop. [illegible] fl. Zante p. 56; Bois. fl. or. III. p. 163; Form.

in D. bot. Mon. 1890 p. 16, in Ver. Brünn 1895 p. 25, 1896 p. 42, 1897 p. 32. — Icon: Fl. dan. t. 1274. — Exsicc.: Tunt. herb. a. 1900.

Virens, scabriusculum; caule erecto, simplici vel a medio pyramidatim ramosissimo, folioso; foliis lineari-lanceolatis, glabris vel breviter hirtis, margine rigido-ciliatis, inferioribus in petiolum attenuatis, superioribus sessilibus; capitulis minutis, numerosissimis, saepe paniculam pyramidatam amplam formantibus; involucri phyllis laxis, lineari-lanceolatis, acuminatis, pappo subbrevioribus; floribus marginalibus pluriserialibus, ligulatis, sordide albis; acheniis glabriusculis, pappo albo pauciseto triplo brevioribus. — Differt a praecedente indumento valde scabro, capitulis duplo minoribus, floribus femineis ligulatis et pappo albido.

Ex America boreali ortum, nunc inquilinum. Frequens in ruderatis regionis inferioris in Epiro et Thessalia a Kalabaka ad Volo (Form.); Euboea: pr. Kurbatzi (Wild), Politika (Holzm.); Attica: pr. Athenas (Heldr.); Achaia: pr. Patras (Heldr.); Laconia: in distr. Alagonia (Zahn); Cycladum insula: Andros (Sart.), Tenos (Chaub.); Zante (Marg.); Corcyra (Form.); et probabiliter alibi. — Maio, Sept. ⊙.

β. Flores feminei marginales involucro subduplo longiores.

3. **E. acre** L. sp. p. 863; Friedr. Reise p. 263; Bois. fl. or. III. p. 166; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1895 p. 25, 1897 p. 32; Haussk. symb. p. 114. — Icon: Rchb. germ. t. 26. —

Scabrido-pubescens; caule erecto, simplici vel apicem versus ramoso, remotiuscule foliato; foliis inferioribus oblongo-spathulatis, in petiolum attenuatis, superioribus oblongo-vel lineari-lanceolatis, sessilibus; capitulis mediocribus, racemosis vel laxe corymbosis; involucri phyllis lineari-lanceolatis, acuminatis, pappo brevioribus; floribus marginalibus incarnatis. exterioribus ligulatis, interioribus filiformibus; acheniis hirtulis, pappo rufescenti triplo brevioribus.

In saxosis, sterilibus regionis praesertim montanae, in alpinam usque adscendens. Thessalia: mt. Zygos (Haussk.), pr. Vendista et Ochsa Despot, mt. Oxya, Chassia (Form.), mt. Olympus (Heldr); Achaia: pr, Patras (Friedr.); Corcyra: pr. urbem (Friedr.), Stratia (Form.). — Jun. Aug. ⊙ et ♃. N. v.

b. Flores feminei marginales disco subduplo longiores.

4. **E. alpinum** L. sp. p. 864; Bois. fl. or. III. p. 165, cum *β. majus* (f. pleiocephala); Hal. in z. b. G. 1888 p. 758, Beitr. fl. Epir. p. 28, in ö. b. Z. 1897 p. 285; Heldr. chlor. Parn. p. 20, f. *majus* Bois.; Form. in Ver. Brünn 1896 p. 42. — *E. olympicum* Schott in ö. b. Wochenbl. VII. p. 230. — *E. uniflorum* Bald. riv. coll. bot. Alb. 1895 p. 45, 1896 p. 65, non L. — Icon: Rchb. t. 23. — Exsicc.: Hal. it. gr. II. a. 1893.

Pluricaule, pilis patulis plus minus obsitum; caulibus erectis vel adscendentibus, simplicibus vel pauciramosis, remote foliatis; foliis inferioribus-oblongo-spathulatis, in petiolum attenuatis, superioribus lineari-lanceolatis sessilibus; capitulis majusculis, solitariis vel 3—5; involucri

hirsuti phyllis lineari-lanceolatis, acuminatis, pappo subaequilongis; floribus marginalibus purpureis, exterioribus ligulatis, interioribus filiformibus; acheniis hirtulis, pappo rufescenti duplo brevioribus.

In saxosis regionis alpinae. Epirus: mt. Papignon et Gamila distr. Zagorion, mt. Tsumerka (Bald.), mt. Peristeri (Hal.); Thessalia: mt. Olympus (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), mt. Parnassus (Heldr.). — Jul. Sept. ♃.

7. **Aster** L. gen. n. 954.

a. Caulis monocephalus.

1. **A. cylleneus** Bois. et Orph. diagn. ser. 2, III. p. 7 pro var. *A. alpini*. — *A. alpinus* Bois. fl. or. III. p. 157 quoad pl. graec., non L. — Exsicc.: Orph. fl. gr. n. 285.

Cinereo-pubescens, caespitosum; caulibus pumilis, foliatis; foliis integris, radicalibus oblongo-spathulatis, in petiolum brevem attenuatis, caulinis diminutis sessilibus; involucri phyllis lanceolatis, obtusis; ligulis pallide violaceis, disco duplo longioribus; acheniis hirtis. — Subspecies *A. alpini* L., a quo statura minori, indumento denso cano, foliis multo minoribus, involucri phyllis obtusis et ligulis pallidis discedit.

In rupestribus regionis alpinae, rarissime. Achaia: mt. Kyllene (Orph.). — Jul. Aug. ♃.

b. Caulis polycephalus.

2. **A. amellus** L. sp. p. 873; S. et S. pr. II. p. 179; Ch. et B. fl. pelop. p. 58; Fraas fl. class. p. 210; Bois. fl. or. III. p. 157. — Icon: Jacq. fl. austr. V. t. 425.

Pubescenti-scaber; caulibus erectis, firmis, foliatis, laxe corymbosis; foliis integris vel subserratis, inferioribus ellipticis, in petiolum attenuatis, superioribus lanceolatis, sessilibus; involucri phyllis oblongis, obtusis; ligulis coeruleis, disco subduplo longioribus; acheniis hirtis.

Attica: pr. Athenas (Sibth.), ad Phalerum (Fraas); a recentioribus non lectum. — Graeciae civis maxime dubia. — Aug. Sept. ♃. N. v.

3. **A. tripolium** L. sp. p. 872; S. et S. pr. II. p. 178; Pieri corc. fl. p. 118; Friedr. Reise p. 274; Bois. fl. or. III. p. 158; Haussk. symb. p. 114. — *Tripolium vulgare* N. ab E. gen. et sp. aster. p. 153. — Icon: Fl. dan. t. 615. — Exsicc.: Heldr. herb. norm. n. 943; Dörfl. fl. gr. n. 159. —

Glaber; caulibus erectis vel adscendentibus, foliatis, inferne saepe fistulosis, a basi interdum ramosis; foliis glaucescentibus, carnosis, subintegris, inferioribus oblongo-ellipticis, petiolatis, reliquis lineari-lanceolatis, sessilibus; involucri phyllis oblongis, obtusis; ligulis violaceis, disco subduplo longioribus; acheniis scabridulis.

In maritimis, humidis salsuginosis. — Aetolia: pr. Mesolongion (Nied.), insula Hagios Sostis (Reinhold); Attica: ad Phalerum, Eleusis (Heldr.); Euboea: pr. Kurbatzi (Wild); Argolis: pr. Nauplia (Haussk.), Poros (Friedr.); Messenia: pr. Pylos (Zahn); Archipelagus (Sibth.); Leucas (Mazz.), Corcyra (Pieri). — Oct. Apr. ♃.

8. Solidago L. gen. n. 955.

1. **S. virgaurea** L. sp. p. 880; Bois. fl. or. III. p. 156; Heldr. chlor. Parn. p. 20; Hal. in ö. b. Z. 1892 p. 371. — Icon: Fl. dan. t. 663. — Exsicc.: Heldr. herb. norm. n. 286; Sint. et Bornm. it. turc. n. 1294.

Caule elato, glabro vel superne puberulo, simplici vel superne ramoso; foliis ovatis oblongo-lanceolatisve, glabris vel puberulis, inferioribus in petiolum attenuatis, superioribus sessilibus; involucri phyllis inaequalibus, lineari-lanceolatis, glabriusculis; ligulis flavis, involucro sesquilongioribus.

β. **vestita.** — Caule folioso, superne dense puberulo, apice in paniculam brevem confertam abeunti; foliis minutis, ovatis, acuminatis, utrinque, subtus densius, aspero-tomentellis; involucri phyllis adpresse puberulis. — Exsicc.: Heldr. herb. a. 1895.

In rupestribus regionis abietinae et alpinae, rare. Thessalia: in valle Megarema pr. Litochori (Sint.); mt. Parnassus (Heldr.); — *β.* mt. Dirphys Euboeae (Heldr.). — Aug. Sept. ♃.

3. Tribus. INULEAE Cass. in ann. sc. nat. 1829 p. 20.

9. Asteriscus Moench meth. p. 592.

1. **A. maritimus** L. sp. p. 903; S. et S. pr. II. p. 196; Pieri corc. fl. p. 121; *(Buphthalmum)*; Less. syn. comp. p. 210; Friedr. Reise p. 274; Bois. fl. or. III. p. 179. — Icon: Rchb. t. 48. —

Perennis, papilloso-hirtus; caulibus e rhizomate lignoso ramoso adscendentibus, simplicibus vel ramosis; foliis obovato vel oblongo-spathulatis, in petiolum attenuatis; capitulis terminalibus, 1—2 foliis floralibus suffultis; involucri phyllis basi demum induratis, externis spathulatis, radio subaequilongis; ligulis aureis vel citrinis, disco longioribus; acheniis sericeis; pappi paleis denticulatis.

In rupestribus maritimis Atticae et Boeotiae (Sibth.); insula Poros (Friedr.); Zante (Bois.); Corcyra (Pieri). — Mart. Jun. ♃. N. v.

2. **A. aquaticus** L. sp. p. 903; S. et S. pr. II. p. 196; Pieri corc. fl. p. 121; Ch. et B. exp. p. 254, Fl. pelop. p. 59; *(Buphthalmum)*; Less. syn. comp. p. 210; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 278; Clem. sert. p. 59; Raul cret. p. 780; Weiss in z. b. G. 1869 p. 42, cum f. *discoidea*; Bois. fl. or. III. p. 178, cum f. *nana*, forma macra caule brevissimo, 1—2 cephalo; Heldr. fl. cephal. p. 43, Fl. Aegina p. 304; Spreitz. in z. b. G. 1887 p. 662, 1890 p. 296; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1895 p. 25, 1896 p. 42, 1897 p. 32; Haussk. symb. p. 115, cum f. *nana* Bois.; Hal. in ö. b. Z. 1895 p. 411. — Icon: Fl. gr. t. 899. — Exsicc.: Orph. fl. gr. n. 1145; Sint. et Bornm. it. turc. n. 1285; Sint. it. thessal. n. 407.

Annuus, villosulo-pubescens; caule erecto, simpliciter vel repetite bi-trichotomo; foliis oblongo-lanceolatis, inferioribus in petiolum attenuatis, superioribus sessilibus, semiamplexicaulibus; capitulis sessilibus, foliis summis cinctis, primario semper inter ramos sessili, axim primariam terminante; involucri phyllis basi demum induratis, externis radio multo longioribus; ligulis brevibus, flavis; acheniis sericeis; pappi paleis lacerodentatis. — Differt a praecedente radice annua, foliorum forma, caule dichotomo nec alterne ramoso, involucri phyllis radium multo excedentibus et ligulis brevibus.

In fossis, locis humidis, arenosis regionis inferioris, frequens in Epiro, Thessalia, Attica, Peloponneso, Creta, Cythera, in insulis Jonicis, indicatur deinde in Sporadum insula Psathura, in Cycladum insula Melos, Syra, in insula Aegina et in Pharmacusis. — Apr. Jun. ○.

3. **A. citriodorus** Heldr. et Hal. fl. aeg. exs. a. 1889, in ö. b. Z. 1895 p. 411. —

Differt a praecedente pubescentia glanduloso-viscosa citriodora, caule divaricato ramosissimo, ramis tenuioribus albidis, internodiis elongatis, foliis omnibus sessilibus, superioribus angustioribus acutiusculis, capitulis minoribus pallidioribus, praecipue alaribus valde diminutis, saepe vix pisum aequantibus, pappi paleis brevioribus.

In saxosis Cycladum insulae Jos, copiose circa urbem et inter urbem et portum (Heldr.). Jun. Jul. ⊙.

10. Pallenis Cass. dict. sc. nat. 37 p. 275.

1. **P. spinosa** L. sp. p. 903; S. et S. pr. II. p. 195, Pieri corc. fl. p. 121; Sieb. avis p. 5, rem. p. 6; Ch. et B. exp. p. 254, Fl. pelop. p. 59; *(Buphthalmum);* Cass. l. c.; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 263; Ung. Reise p. 123; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 780; Bois. fl. or. III. p. 180; Heldr. fl. cephal. p. 43, Fl. Aegina p. 304; Haussk. symb. p. 115; Hal. in ö. b. Z. 1899 p. 25. — Icon: Fl. gr. t. 898. — Exsicc.: Sint. et Bornm. it. turc. n. 1284; Baen. herb. europ. n. 9210; Dörfl. fl. aeg. n. 4, pl. cret. n. 129.

Annua, molliter subvillosa, interdum sublanata, rarius puberula; caule erecto, superne dichotome ramoso, rarius simplici; foliis inferioribus oblongo-spathulatis, in petiolum attenuatis, superioribus lanceolatis, sessilibus, subcordato-semiamplexicaulibus; capitulis pedunculatis, basi nudis; involucri phyllis externis elevatim nervosis, spinosis, stellatim patentibus, radio multo longioribus; ligulis flavis, brevibus; acheniis subglabris; pappo breviter coronaeformi. — *Asterisco aquatico* similis, ab eo praeter alias notas, capitulis foliis non cinctis et involucri phyllis spinosis statim diagnoscenda. — Specimina graeca nonnulla, praesertim e Creta et Cycladibus indumento sparso, pedunculis elongatis subnudis et capitulis diminutis a typo discedunt *(f. microcephala).* —

In herbidis, olivetis, collibus saxosis regionis inferioris per totam **Graeciam.** — **Apr. Jul.** ⊙.

11. Inula L. gen. n. 956.

1. Sectio. *Corvisartia* Mer. fl. par. p. 328 pro gen. — Involucri phylla exteriora late foliacea, recurvata; receptaculum fimbriatum; achenia 5—4 gona, truncata, obsolete costata, glabra; pappi setae liberae. Capitula speciosa.

1. **I. helenium** L. sp. p. 881; S. et S. pr. II. p. 180; Bois. fl. or. III. p. 186; Heldr. in Sitzungsb. acad. Berlin 1883 p. 7. — Icon: Fl. gr. t. 873. — Exsicc.: Orph. herb. a. 1857.

Caule robusto, erecto, ramoso; foliis amplis, dentato-serratis, subtus plus minus molliter tomentosis, inferioribus elliptico-oblongis, in petiolum attenuatis, caulinis cordato-amplexicaulibus; involucri phyllis exterioribus ovatis, tomentosis, interioribus lineari-spathulatis, subcoriaceis, erectis; ligulis flavis, involucro plus duplo longioribus.

In dumosis regionis montanae. Thessalia (Sibth.): inter *Pterides* ad radices Olympi pr. Caterina (Orph.), pr. Laspochori ad mt. Ossa (Heldr.), ad Zagora in mt. Pelion (Aphentulis); Achaia: mt. Kyllene (Heldr.). — Jul. Sept. ♃.

2. Sectio. *Enula* Duby bot. gall. I. p. 267. — Involucri phylla exteriora apice appendiculo parvo foliaceo aucta et recurvata; receptaculum glabrum, achenia subcylindrica, costata, glabra vel hirsuta, apice truncata vel breviter attenuata; pappi setae liberae. Capitula mediocria, rarius parva.

a. Achenia glabra; pappus multisetus.

α. Folia caulina basi cordato-semiamplexicaulia.

2. **I. germanica** L. sp. p. 883; Bois. fl. or. III. p. 189; Haussk. symb. p. 115; Form. in Ver. Brünn 1895 p. 25, 1896 p. 42, 1897 p. 32. — Icon: Jacq. fl. austr. t. 134. — Exsicc.: Orph. fl. gr. n. 787; Sint. et Bornm. it. turc. n. 1299 b.

Caule erecto, hirto vel villosulo, apice corymboso; foliis oblongo-lanceolatis, acutis, integris vel denticulatis, reticulato-venosis, supra adpresse pilosulis, subtus praesertim nervo medio villosulis; capitulis cylindricis; involucri phyllis adpresse hirtulis; ligulis flavis, involucrum parum excedentibus.

In collibus dumosis, apricis regionis inferioris et montanae. Epirus: pr. Han Kamberga, Han Balduma, in valle Dipotami (Form.); Thessalia: pr. Malakasi, Kalabaka, ad monasterium Korona, Pharsalus (Haussk.), pr. Asproklisia, Kucuro, Kerasia Sina, mt. Cuka, mt. Chassia, pr. Vlacharo, Konisko, Velestinos (Form.), pr. Litochoron ad radices Olympi (Orph.). — Jun. Jul. ♃.

3. **I. cordata** Bois. diagn. IV. p. 3; Hal. in ö. b. Z. 1892 p. 371. — *I. salicina β. aspera* Beck. imul. europ. p. 22; Haussk. symb. p. 115, vix *I. aspera* Poir. enc. suppl. III. p. 154, quae sec. Haussk. l. c. probabiliter hybrida. — Icon: Fl. gr. t. 875. — Exsicc.: Sint. et Bornm. it. turc. n. 1299.

Caule erecto, setulis e tuberculo ortis plus minus scabrido, apice oligo rarius monocephalo; foliis coriaceis, oblongo-lanceolatis, acuminatis, reticulato-venosis, copiose scabro-denticulatis, supra glabris, subtus scabro-pilosis; capitulis hemisphaericis; involucri phyllis glabris, ciliatulis; ligulis flavis, involucro duplo longioribus.

In dumosis regionis montanae. Thessalia: ad monasterium Korona in Pindo (Haussk.), ad monasterium Metoji pr. Litochoron in Olympo (Sint.). — Jun. Jul. ♃.

β. Folia caulina basi rotundata vel attenuata.

4. **I. hirta** L. sp. p. 883; S. et S. pr. II. p. 182; Bois. fl. or. III. p. 187. — Icon: Jacq. fl. austr. t. 358.

Caule erecto, hirto, mono rarius oligocephalo; foliis oblongis vel oblongo-lanceolatis, acutis, reticulato-venosis, integris vel subdenticulatis, utrinque hirsutis, basi rotundata sessilibus; capitulis mediocribus; involucri phyllis reticulato-venosis, strigulosis, setuloso-ciliatis; ligulis flavis, involucro duplo longioribus.

In Archipelagi insulis (Sibth.); — a recentioribus non lecta. — Maio. Jun. ♃. N. v.

5. **I. ensifolia** L. sp. p. 883; Clem. sert. p. 60. — Icon: Jacq. fl. austr. t. 162.

Caule erecto, glabro vel superne albo-lanuginoso, mono vel 2—5 cephalo; foliis anguste lineari-lanceolatis, acutis, parallele-nerviis, margine scabridulis et parce lanuginosis, ceterum glabris, basi attenuata sessilibus; capitulis mediocribus; involucri phyllis parallele-nerviis, lanatulis; ligulis flavis, involucro duplo longioribus.

In Corcyra et Attica (Clem.), sed a recentioribus ut videtur non lecta. — Jul. Aug. ♃. N. v.

b. Achenia hirsuta.

α. Herbaceae; pappus multisetus.

× Ligulae brevissimae, involucrum vix superantes.

6. **I. vulgaris** Lam. fl. fr. II. p. 73 (*Conyza*); Trevis. fl. eugan p. 29. — *Conyza squarrosa* L. sp. p. 861. — *I. conyza* DC. pr. V. p. 464; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1895 p. 26, 1896 p. 42, 1897 p. 32. — Icon: Fl. dan. t. 622. —

Pubescenti-subtomentosa; caule erecto, superne paniculato-corymboso, polycephalo; foliis elliptico-lanceolatis, integris vel denticulatis, inferioribus in petiolum attenuatis, superioribus basi angustata sessilibus; capitulis parvis; involucri phyllis puberulis; ligulis subtubulosis, flavis.

In rupestribus dumosis regionis inferioris et montanae. Corcyra: pr. Evropuli, Afra (Form.); Epirus: pr. Han Balduma, in valle Dipotami (Form.); Thessalia: pr. Han Kuraneos in valle Penei, pr. Kalabaka, mt. Mitrica in mt. Chassia, pr. Rapsani in Olympo (Form.); mt. Pelion pr. Zagora (Aphentulis). — Jul. Aug. ♃. N. v.

×× Ligulae involucro subduplo longiores.

7. **I. oculus Christi** L. sp. p. 881; S. et S. pr. II. p. 181; Ch. et B. exp. p. 250, Fl. pelop. p. 58; Bois. fl. or. III. p. 193; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Thessal. p. 14, in ö. b. Z. 1895 p. 412; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1895 p. 26, 1896 p. 42, 1897 p. 32; Haussk. symb. p. 115; Bald. riv. coll. bot. Alb. 1895 p. 46. — ? *I. bubonium* Pieri corc. fl. p. 119, non L. — Icon: Jacq. fl. austr. t. 223. — Exsicc.: Orph. fl. gr. n. 89; Sint. it. thessal. n. 1128.

Molliter et adpresse plus minus villosa; rhizomate repente; caule erecto, apice corymbose-oligocephalo; foliis oblongis vel oblongo-lanceolatis, integris vel subdenticulatis, inferioribus in petiolum attenuatis, superioribus basi cordata amplexicaulibus; capitulis mediocribus; involucri phyllis adpressis, dense sericeo-villosis, exterioribus sensim brevioribus; ligulis aureis; acheniis villosis.

In siccis, silvaticis regionis montanae et subalpinae. Epirus: pr. Janina (Bald.), mt. Peristeri (Form.); Thessalia: frequens in Pindo, mt. Chassia, Othrys (Form.), mt. Olympus (Orph.), Pelion (Heldr.); Aetolia: mt. Tymphrestus (Samar.), Korax (Heldr.); mt. Kiona (Hal.); Peloponnesus: mt. Kyllene, Malevo, pr. Zatuna (Orph.) et ad lacum Stymphalidem (Heldr.) Arcadiae, mt. Taygetos (Psarid.), in agro Messeniaco et Eliensi (Sibth.); Corcyra: pr. Kephalomanduchio, Potamos, Evropuli, Afra (Form.). — Jun. Jul. ♃.

8. **I. britannica** L. sp. p. 882; S. et S. pr. II. p. 181; Bois. fl. or. III. p. 193, cum f. *rupestris* Griseb. et Schenk it. hung. p. 337, forma foliis angustioribus, superioribus basi subamplexicaulibus fere exauriculatis, capitulis saepe minoribus, phyllis exterioribus a medio squarrosis; intermediis ad typum transiens; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 6, Chlor. Parn. p. 20; Form. in Ver. Brünn 1896 p. 42, 1897 p. 32. — Icon: Fl. dan. t. 412. — Exsicc.: Heldr. it. thessal. n. 73; Sint. it. thessal. n. 1166.

Adpresse villosula; radice cylindracea, nodosa; caule erecto, apice 2—3 cephalo vel saepe a medio ramoso et corymbose-polycephalo; foliis oblongo-lanceolatis, integris vel subdenticulatis, inferioribus in petiolum attenuatis, superioribus basi cordata vel auriculata amplexicaulibus; capitulis mediocribus; involucri phyllis laxis, sericeis et subtiliter glandulosis, exterioribus interiora aequantibus vel superantibus; ligulis flavis; acheniis hirtis, apice glandulis paucissimis obsitis.

In humidiusculis montanis. Thessalia: pr. Sermeniko (Sint.), pr. Phlambures, Vlachava, Verenci, Koprana, Povelci, Larissa, Pharsalus (Form.), pr. Ampelakia in mt. Ossa (Heldr.); mt. Dirphys Euboeae, mt. Parnassus, ad lacum Stymphalidem Arcadiae (Heldr.). — Jul. Aug. ⊙.

β. Basi suffrutescentes, pappus pauci (5—10)-setus.

⁂ Involucri phylla adpressa.

9. **I. verbascifolia** Willd. sp. III. p. 1924 (*Conyza*); Haussk. symb. p. 115, ? Form. in Ver. Brünn 1895 p. 43, 1897 p. 32; Hal.

in z. b. G. 1899 p. 714. — *Conyza candida* Clem. sert. p. 59; non L. — *I. candida* Cass. dict. XXIII. p. 554; Spreitz. in z. b. G. 1887 p. 662. — *I. candida v. radiata* Bois. fl. or. suppl. p. 291. — Icon: Ten. fl. nap. t. 76. — Exsicc.: Hal. it. gr. a. 1888.

Tomento candido densissime pannosa; caulibus crassiusculis, adscendentibus vel erectis, simplicibus; foliis radicalibus oblongo-spathulatis, obtusis, in petiolum attenuatis, integris crenatisve, caulinis ovatis, sensim brevius petiolatis; capitulis magnis, diametro 20—25 mm, breviter pedunculatis subsessilibusve, solitariis, saepius approximatis, bracteis magnis foliis similibus suffultis; involucri phyllis exterioribus obtusissimis, ad interiora obtusa sensim auctis; flosculis flavis, marginalibus ligulatis, tridentatis; acheniis apice setoso-barbatis.

Ad rupes, muros regionis inferioris et submontanae. Corcyra (Clem.): in urbe (Hal.), pr. Palaeokastrizza (Spreitz.), Garuna (Letourn.); Thessalia: pr. Tyrnovo (Haussk.) et pr. Miluna, Patsios, Rapsani, in valle Tempe, ad Limogardi pr. Lamia, si *I. verbascifolia* Form. huc pertineat. — Jun. Jul. ♃.

10. **I. parnassica** Bois. et Heldr. diagn. ser. 2 III. p. 11; Hal. in z. b. G. 1888 p. 759, 1899 p. 714, cum var. *euboea*; Heldr. chlor. Parn p. 20. — *I. candida v. radiata* Bois. fl. or. III. p. 196. — *I. oxylepis* Schultz herb. sec. Nym. — *I. attica v. macrophylla* Heldr. herb. norm. n. 1553. — Exsicc.: Heldr. herb. norm. n. 49; Orph. fl. gr. n. 343.

Cinereo-tomentosa; caulibus gracilibus, adscendentibus vel declinatis, simplicibus; foliis oblongo-spathulatis, obtusis, in petiolum attenuatis, integris crenatisve, subtus reticulato-venosis, superioribus decrescentibus, brevius petiolatis; capitulis 1—3, solitariis, remotis, 15—20 mm. diametro, saepius longiuscule pedunculatis, basi non vel bractea oblonga vel oblongo-lineari suffultis; involucri phyllis exterioribus obtusiusculis, ad interiora tenuiora acuta sensim auctis; flosculis flavis, marginalibus ligulatis, tridentatis; acheniis glabrescentibus. — Differt a praecedente indumento tenuiore, caulibus gracilibus, remote oligocephalis, capitulis minoribus, saepe longiuscule pedunculatis.

Ad rupes regionis montanae et subalpinae. Aetolia: pr. Mesolongion (Nieder); Doris: mt. Kiona pr. Segditza (Hal.); mt. Parnassus pr. Rachova, mt. Telethrion Euboeae (Heldr.); Achaia: in faucibus Diakophto, mt. Kyllene pr. Trikala (Heldr.); Laconia: mt. Selitza pr. Kalamata (Zahn). — Jun. Jul. ♃.

11. **I. attica** Hal. in Heldr. herb. norm. n. 1040, in z. b. G. 1899 p. 713; Heldr. fl. Aegina p. 304. — *Conyza candida* S. et S. pr. II. p. 173; Ch. et B. exp. p. 247, Fl. pelop. p. 58, non L. — *I. verbascifolia v. limonifolia* DC. pr. V. p. 464 p. p. — *C. candida v. verbascifolia* Clem. sert. p. 59. — *I. candida* Bois. fl. or. III. p. 196 p. p.; ? Raul. cret. p. 779 *v. integrifolia*; Heldr. fl. cephal. p. 44; Form. in D. bot. Monat 1890 p. 16; Bald. riv. coll. bot. Alb. 1895 p. 46. — *I. oxylepis* Schultz herb. sec. Haussk. symb.

p. 115. — *I. Aschersoniana* Hal. in ö. b. Z. 1892 p. 371, non Janka — *I. methanea* Haussk. symb. p. 116. — *I. verbascifolia* Form. in D. bot. Monat. 1898 p. 78. — Icon: Fl. gr. t. 864. — Exsicc.: Heldr. herb. norm. n. 465, 1040, 1445, in Magn. fl. sel. n. 3796; Orph. fl. gr. n. 788; Sint. et Bornm. it. turc. n. 1305; Bald. it. alb. epir. III. n. 70.

Praecedenti maxime affinis et ad eam ut videtur intermediis transiens, differt tamen indumento saepissime densiore, foliis saepius non reticulatis, capitulis minoribus, discoideis, saepius longe pedunculatis.

β. **limonella** Heldr. ap. Hal. in z. b. G. 1899 p. 713. — Caulibus humilioribus, foliis oblongo-lanceolatis, dense adpresse albo-pannosis, subnitentibus, non reticulato-venosis, capitulis breviter pedunculatis. — Exsicc.: Heldr. herb. norm. n. 775.

In rupibus regionis submontanae, ad subalpinam adscendens. Epirus: mt. Zalongo pr. Prevesa (Bald.); Thessalia: pr. Velestinos (Form.), in fauce Megarema pr. Litochori (Sint.); Attica: mt. Lycabettus (Orph.), Parnes, Pentelicon, Hymettus (Heldr.), pr. Sunium (Gerold); insula Aegina ad Paleochora (Heldr.); Argolis (Sprun.): pr. Vromolimni (Haussk.); Achaia: pr. Kalavryta, mt. Chelmos (Orph.), mt. Kyllene (Heldr.); Laconia: pr. Megali Anastasova (Zahn), mt. Taygetos (Heldr.), pr. Lutro, Thuria, in Maina, pr. Kardamyle, Vitilos (Chaub.); Creta (Sibth.); Cephalonia: pr. Asso, Charakas (Heldr.); loca nonnulla probabiliter ad speciem praecedentem spectant; — *β.* mt. Dirphys Euboeae (Heldr.). — Jun. Jul. ♃.

×× Involucri phylla squarroso-patentia.

12. **I. candida** L. sp. p. 862, saltem pro max. p.; Willd. sp. III. p. 1923; Sieb. avis p. 5, rem. p. 6 (*Conyza*). — *C. limonifolia* S. et S. pr. II. p. 174, Fl. gr. IX. p. 50, t. 865; Sieb. avis p. 5; Ch. et B. exp. p. 247, Fl. pelop. p. 58. — *I. candida v. limonifolia* DC. pr. V. p. 464 p. p.; Bald. viagg. cret. p. 61. — *I. limonifolia* Bois. fl. or. III. p. 198; Raul. cret. p. 778; Heldr. in ö. b. Z. 1880 p. 346; Spreitz. in z. b. G. 1887 p. 662, 1890 p. 296; Fors. in bull. herb. Bois. V. p. 399; Hal. in z. b. G. 1899 p. 714. — Exsicc.: Heldr. pl. cret. n. 1579; Bald. it. cret. n. 22. —

Albo-pannosa; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis integris, subnitentibus, radicalibus ovato- vel oblongo-spathulatis, in petiolum attenuatis, superioribus sensim diminutis, breviter petiolatis; capitulis majusculis, discoideis, subcorymboso-paniculatis, breviter pedunculatis subsessilibusque, basi saepe bracteatis; involucri phyllis obtusis; exterioribus subbrevioribus, intimis acutis; flosculis flavis; acheniis glabrescentibus. — Tomento adpressissimo subnitente et involucro squarroso insignis.

β. **rotundifolia.** — Foliis infimis rotundatis, ceteris ellipticis oblongisve. — Exsicc.: Chaub. pl. exs. Morée; Spreitz. it. ion. a. 1880. —

In rupibus maritimis, collibus saxosis regionis inferioris. Creta (Sibth.): pr. Kissamos Kastelli (Heldr.), ad promontorium Grabusa, pr.

Palaeokastro (Raul.), Aradhena (Spreitz.), Gonia (Bald.); Cythera: pr. Mylopotamo (Fers.), in fauce Kalamo, insula Prassonizza (Spreitz.); Laconia: promontorium Ierea et Malea (Chaub.). — Jun. Jul. ♃.

3. Sectio. *Limbarda* Adans. fam. II. p. 125 pro gen. — Involucri phylla exteriora membranacea, adpressa, omnino exappendiculata; receptaculum glabrum; achenia subcylindrica, obsolete costata, hirsuta apice truncata, pappi setae liberae. Capitula mediocria, folia carnosa.

13. **I. crithmoides** L. sp. p. 883; S. et S. pr. II. p. 182; Ch. et B. exp. p. 250. Fl. pelop. p. 58; Marg. et R. fl. Zante p. 56; Raul. cret. p. 779; Bois. fl. or. III. p. 195; Heldr. fl. cephal. p. 44; Form. in Ver. Brünn 1895 p. 26. — *I. crithmifolia* Willd. sp. III. p. 2101; Pieri corc. fl. p. 119; Sieb. avis p. 5, rem. p. 6; Friedr. Reise p. 274; Ung. Reise p. 123. — Icon: Engl. bot. t. 68.

Fruticulosa, glaberrima; caulibus erectis vel adscendentibus, apice mono vel oligocephalis; foliis linearibus, obtusis, inferioribus saepe apice tridentatis, axillis fasciculiferis; pedunculis elongatis, bracteolatis, apice incrassatis; involucri phyllis lineari-lanceolatis, acuminatis, exterioribus subbrevioribus; ligulis flavis, involucro subduplo longioribus.

Ad oras et rupes maritimas. Corcyra (Pieri); pr. Canone, Hagios Joannes (Form.); Cephalonia: pr. Drapano, Lixuri (Heldr.); Zante (Sibth.); Aetolia: pr. Mesolongion (Nied.); Thessalia: pr. Volo (Orph.); Peloponnesus: insula Poros (Friedr.), ad fl. Eurotas, pr. Navarin (Chaub.); Creta: ad Mandromolo pr. Kissamos, pr. Grabusa (Raul.); Archipelagus (Bois.). — Aug. Oct. — ♃. N. v.

4. Sectio. *Cupularia* Gr. et Godr. fl. fr. II. p. 180. pro gen. — Involucri phylla exteriora subfoliacea, adpressa, omnino exappendiculata; receptaculum nudum; achenia cylindrica, obtusangula, ecostata, hirsuta, apice constricta et glandulis pellucidis obsita, pappi setae basi in cupulam breviter coalitae. — Capitula parva.

14. **I. viscosa** L. sp. p. 863; S. et S. pr. II. p. 174; Sieb. avis p. 5, rem. p. 6; Ch. et B. exp. p. 247. Fl. pelop. p. 57; Fraas fl. class. p. 209; (*Erigeron*); Ait. hort. Kew. III. p. 233; Marg. et R. fl. Zante p. 56; Friedr. Reise p. 264; Clem. sert. p. 60; Raul. cret. p. 779; Bois. fl. or. III. p. 198; Heldr. fl. cephal. p. 44, chlor. Thera p. 14, chlor. Mykonos p. 245; Hal. in z. b. G. 1888 p. 759; Haussk. symb. p. 115; Form. in Ver. Brünn 1895 p. 26, 1897 p. 32. — *Solidago viscosa* Lam. fl. fr. II. p. 144. — *Pulicaria viscosa* Cass. ex. Koch syn. p. 361. — *Cupularia viscosa* Gr. et Godr. fl. fr. II. p. 181; Heldr. fl. Aegina p. 304. — Icon: Jacq. hort. vind. II. t. 165. — Exsicc.: Orph. fl. gr. n. 90; Sint. et Bornm. it. turc. n. 1300.

Glandulo[illegible] viscosa, basi suffrutescens; caulibus virgatis, racemo[illegible] lanceolatis, sessilibus, remote denticulatis, su[illegible]is; capitulis breviter pedunculatis, approxi-[illegible] lanceolatis, valde inaequalibus, margine [illegible]volucro sesquilongioribus.

In arenosis, olivetis, saxosis regionis inferioris et montanae. Epirus (Form.); Thessalia: ad Plaka pr. Litochori (Sint.), pr. Lechonia, Agria, Stylida, Lamia (Form.); Euboea: pr. Kurbatzi (Wild.), Kumi (Wurlisch); Doris: pr. Amphissa (Hal.); Attica: pr. Athenas (Orph.), ad Ilissum, insula Aegina (Heldr.); Argolis: pr. Nauplia (Sart.); Cyclades (Sibth.): insula Keos, Mykonos, Thera (Heldr.); Creta: pr. Canea, Nerokuru (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli, Pessada, Piscardo (Heldr.); Corcyra (Friedr.). — Aug. Oct. ♃.

15. **I. graveolens** L. sp. ed. 2 p. 1210; S. et S. pr. II, p. 174; Sieb. avis p. 5, Ch. et B. fl. pelop. p. 58; Friedr. Reise p. 264; Fraas fl. class. p. 209 (*Erigeron*); Desf. fl. atl. II. p. 275; Marg. et R. fl. Zante p. 56; Clem. sert. p. 60; Raul. cret. p. 779; Bois. fl. or. III. p. 199; Heldr. fl. cephal. p. 44, chlor. Thera p. 14, chlor. Mykonos p. 246. — *Solidago graveolens* Lam. fl. fr. II. p. 145. — *Cupularia graveoleus* Gr. et Godr. fl. pr. II. p. 180; Heldr. fl. Aegina p. 304. — Icon: Fl. gr. t. 866. — Exsicc.: Heldr. herb. fl. hellen. a. 1882.

Glanduloso-pubescens, viscosa, annua; caule erecto, a basi pyramidatim ramoso; foliis lanceolatis, sessilibus, integris vel obsolete denticulatis, basi interdum subauriculatis; capitulis breviter pedunculatis, approximatis; involucri phyllis lineari-lanceolatis, exterioribus herbaceis, interioribus scariosis, vitta viridi notatis; ligulis paucis, luteis vel violascentibus, involucrum vix excedentibus.

In arenosis humidis, ruderatis, rupestribus regionis inferioris. Boeotia (Fraas); Attica: pr. Athenas, Kephissia, Tatoi (Heldr.), Megara (Clem.), insula Aegina (Heldr.); Euboea: pr. Kurbatzi (Wild); Achaia: pr. Patras (Friedr.); Creta: mt. Psiloriti (Raul.); Cyclades (Sibth.): insula Mykonos, Thera (Heldr.); Zante (Marg.); Cephalonia: pr. Argostoli, Pessada (Heldr.); Corcyra (Pichler). — Aug. Oct. ⊙

12. Pulicaria Gaertn. fr. II. p. 461.

a. Perennes; flosculi radii discum subduplo superantes,

1. **P. odora** L. sp. p. 881; S. et S. pr. II. p. 180; Sieb. avis p. 5, rem. p. 6; Ch. et B. exp. p. 249, Fl. pelop. p. 58; Fraas fl. class. p. 211; (*Inula*); Rchb. fl. germ. exc. p. 239; Marg. et R. fl. Zante p. 57; Weiss in z. b. G. 1869 p. 42; Raul cret. p. 779; Bois. fl. or. III. p. 201; Heldr. fl. cephal. p. 44; Spreitz. in z. b. G. 1887 p. 662; Hal. Beitr. fl. Epir. p. 28; Haussk. symb. p. 118. — Icon: Desf. cor. Tourn. t. 38. — Exsicc.: Baen. herb. europ. n. 9316.

Rhizomate fibroso; caule erecto, fastigiato-ramoso, rarius simplici, villoso, foliis infimis ovalibus vel ovali-lanceolatis, in petiolum attenuatis, subtus lanuginosis, caulinis multo minoribus, cordato-semiamplexi caulibus; capitulis majusculis, longiuscule pedunculatis; involucri villosi phyllis inaequalibus, lineari-acuminatis, externis subherbaceis; ligulis aureis; pappi coronula laciniata.

In collibus, dumosis regionis inferioris et montanae. Epirus: pr. Kalentini, Mazuki (Hal.); Thessalia: mt. Ghavellu, in oropedio Neuropolis in Pindo (Haussk.); Aetolia: pr. Mesolongion (Nied.); Attica: mt. Parnes pr. Dekeleia (Hal.); Euboea: pr. Kurbatzi (Wild), mt. Dirphys (Sprun.); Elis: pr. Lintzi (Heldr.); Argolis (Sibth.): pr. Poros (Heldr.); Messenia: pr. Methone, Corone, Nisi, Kalamata, Zarnate, in Maina, mt. Taygetos Laconiae (Chaub.); Cyclades (Sibth.): insula Cythnos (Tunt.), Tenos (Sart.); Creta: pr. Canea (Weiss), Khalepa, Nerokuru, Perivolia, Spili, Preveli, Voriza (Raul.); Zante (Marg.); Cephalonia: pr. Pessada, Lixuri (Heldr.); Corcyra: pr. Manducchio (Spreitz.), Palaeokastrizza (Baen.). — Maio, Jul. ♃

2. **P. dysenterica** L. sp. p. 882; Pieri corc. fl. p. 118; Ch. et B. exp. p. 249, Fl. pelop. p, 59; (*Inula*); Gaertn. fruct. II. p. 462; Marg. et R. fl. Zante p. 57; Clem. sert. p. 60; Hal. in ö. b. Z. 1890 p. 39; Form. in D. bot. Monat 1890 p. 16, in Ver. Brünn 1895 p. 26. — Icon: Fl. dan. t. 410. — Exsicc.: Sint. it or. a. 1889 n. 1830.

Rhizomate crassiusculo, turiones squamatos edente; caule erecto, ramoso, lanuginoso; foliis oblongis, undulatis, denticulatis, subtus canescenti-tomentosis, infimis in petiolum attenuatis, ceteris cordato-semiamlexicaulibus, late auriculatis; capitulis mediocribus, laxe cymoso-corymbosis; involucri phyllis inaequalibus, lineari-cuspidatis, ciliatis, dorso glandulosis, externis foliaceis; ligulis flavis; pappi coronula denticulata.

β. **dentata** S. et S. pr. II. p. 181 pro sp. (*Inula*). — *P. uliginosa* Stev. in DC. pr. V. p. 478; Haussk. symb. p. 118; Heldr. in ö. b. Z. 1898 p. 184. — *P. dentata* DC. l. c. p. 480; Raul. cret. p. 779. — *P. dysenterica v. microcephala* Bois. fl. or. III. p. 202; Heldr. fl. cephal. p. 44; Form. in Ver. Brünn 1896 p. 43, 1897 p. 32. — A basi saepe ramosissima, viscidula, capitulis fere dimidio minoribus, ligulis brevioribus. Formis intermediis ad typum transit. — Icon: Fl. gr. t. 874. — Exsicc.: Heldr. herb. norm. n. 675 et 1151.

In humidis regionis inferioris et montanae per totam Graeciam β. typo multo vulgatior. Jul. Oct. ♃

b. Annuae flosculi radii discum non vel vix superantes.

3. **P. vulgaris** Gaertn. fruct. II. p. 461, t. 173; Marg. et R. fl. Zante p. 56; Bois. fl. or. III. p. 204; Form in D. bot. Monat. 1890 p, 16, in Ver. Brünn 1896 p. 43, 1897 p. 32; Haussk. symb. p. 118. — *Inula pulicaria* L. sp. p. 882; Pieri corc. fl. p. 119; Ch. et B. exp. p. 250. — *I. arabica* Sieb. avis p. 5, non L. — *P. crispa* Link in Linnaea IX. p. 583, non *Inula crispa* Pers. — *I. dentata* Ch. et B. fl. pelop. p. 58, non S. et S. — *I. pulegium* Friedr. Reise p. 264. — *I. undulata* Friedr. Reise p. 274, non L. — Icon: Fl. dan. t. 613. — Exsicc.: Orph. fl. gr. n. 748; Rev. pl. cret. n. 69; Sint. et. Bornm. it. turc. n. 1304; Sint. it. thessal. n. 1064.

Crispule tomentosa; caule erecto, fere a basi ramoso; foliis oblongis vel oblongo-lanceolatis, undulatis, obsolete denticulatis, inferioribus in petiolum attenuatis, ceteris basi rotundata semiamplexicaulibus; capitulis parvis, laxe cymoso-paniculatis; involucri villosi phyllis lineari-lanceolatis, acuminatis; ligulis flavis, discum vix superantibus, pappi coronula laciniata, setis 8—10, achenio subaequilongis.

In humidis regionis inferioris. Frequens in Thessalia; dein indicatur in Aetolia: pr. Vrachori (Samar.); Argolis: pr. Nauplia, Argos (Chaub.), insula Poros (Friedr.); Cyclades (Chaub.); Creta (Sieb.): pr. Canea (Rev.); Zante: in palude Macri (Marg.); Corcyra (Pieri): Lago di valle (Friedr.). — Jun. Aug. ⊙

4. **P. sicula** L. sp. p. 864; Sieb. avis p. 7, rem. p. 6; (*Erigeron*); Moris fl. sard. II. p. 363; Bois. fl. or. III. p. 205; Bald. riv. coll. bot. Alb. 1892 p. 46, in nuovo giorn. bot. ital. 1894 p. 97. — *Jasonia sicula* DC. in ann. sc. nat. 1834 p. 261; Marg. et R. fl. Zante p. 56; Raul. cret. p. 779. – Icon: Bocc. t. 31, f. 4. — Exsicc: Heldr. herb. norm. n. 540; Bald. it. alb. a. 1892 p. 249.

Puberula; caule erecto, saepe a basi ramoso; foliis rigidis, oblongo-lanceolatis vel linearibus, obsolete denticulatis, inferioribus in petiolum attenuatis, ceteris aurito-semiamplexicaulibus; capitulis parvis, laxe cymoso-racemosis; involucri puberuli phyllis linearibus, acutis; ligulis flavis, discum subaequantibus; pappi coronula ad basin partita, setis 18—25, achenio duplo longioribus.

In herbidis, humidis maritimis. Epirus: pr. Skafidaki et Camarina distr. Prevesa (Bald.); Aetolia: pr. Mesolongion (Nieder); Phocis: pr. Itaea (Heldr.); Argolis: pr. Tirynthum (Sart.); Zante: ad Agria (Marg.); Creta: pr. Canea (Sieb.). — Jul. Oct. ⊙

13. Phagnalon Cass.

1. **P. methanaeum** Haussk. in Mitth. thür. bot. Ver. 1887 V. p. 85, symb. p. 113. — *P. saxatile* Heldr. fl. Aegina p. 303, non (L.) Cass. — Exsicc.: Heldr. herb. norm. n. 941.

Suffruticosum; caulibus erectis vel adscendentibus, albo-tomentosis; foliis linearibus vel lineari-lanceolatis, integris, basi adnato-semiamplexicaulibus, supra viridibus subfloccosis, subtus adpresse albo-tomentosis; pedunculis elongatis, monocephalis, bracteis paucis squamaeformibus obsitis; involucri phyllis pallidis, glabris, membranaceo-scariosis, ad marginem subundulatis, infimis subpatulis brevibus, ovato-oblongis, obtusis, intermediis longioribus, oblongis, obtusis, intimis lineari-oblongis, abrupte mucronatis. — Ab affini *P. saxatili* (L.) squamis infimis non squarrosis, rodundatis, nec acutato-mucronatis facile diagnoscitur.

In rupibus calcareis regionis inferioris. Argolis: pr. Vromolimni in peninsula Methana (Haussk.), in urbe Poros, insula Hydra, Aegina Jos Cycladum (Heldr.). — Apr. Maio. ♄

2. **P. graecum** Bois. et Heldr. diagn. XI. p. 6, Fl. or. III. p. 221; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 778; Heldr. fl.

cephal. p. 44, Fl. Aegina p. 304, Chlor. Thera. p. 15, chlor. Mykonos p. 245; Spreitz. in z. b. G. 1887 p. 662; Hal. Beitr. fl. Aetol. p. 7; Haussk. symb. p. 113; Form. in Ver. Brünn 1895 p. 26. — *Conyza saxatilis* S. et. S. pr. II. p. 173, Fl. gr. t. 862 ex loco, sed icon dubium; Sieb. avis p. 5; Ch. et B. exp. p. 246, Fl. pelop. p. 58; non L. — *C. rupestris* Sieb. avis. rem. p. 6, non L. — *P. saxatile* DC. pr. V. p. 396, quoad loca graeca; Friedr. Reise p. 265 et 278; Marg. et R. fl. Zante p. 56; non (L.) Cass. — *P. rupestre* Clem. sert. p. 59; Ung. Reise p. 123; Raul. cret. p. 778; Gelmi in bull. soc. bot. ital. 1889 p. 450; Bald. viagg. Creta p. 63; Hal. in ö. b. Z. 1899 p. 25; non (L.) DC. — *P. Tenorii* Bald. in nuovo giorn. bot. ital. 1894 p. 97, non Presl, quod ad *P. rupestrem* spectat. — Exsicc.: Heldr. herb. norm. n. 483, 942; Orph. fl. gr. n. 478; Rev. pl. cret. n. 68, in Baen. herb. europ. n. 4790; Bald. it. cret. n. 20; Baen. herb. europ. n. 9300; Sint. it. thessal. n. 76; Dörfl. pl. cret. n. 131.

Suffruticosum; caulibus erectis vel adscendentibus, albo-tomentosis; foliis oblongo-vel lineari-lanceolatis, undulato-sinuatis, basi adnato-semiamplexicaulibus, supra viridibus subfloccosis, subtus adpresse albotomentosis; pedunculis elongatis, monocephalis, bracteis paucis squamaeformibus obsitis; involucri phyllis adpressis, fuscescentibus, glabris, membranaceo-scariosis, obtusis, ad marginem non subundulatis, infimis brevibus triangulari-lanceolatis, intermediis longioribus, obtusiusculis, linearilanceolatis, sensim acutatis. — Differt a praecedente foliis undulato-sinuatis, capitulis subminoribus, involucri phyllis adpressis, angustis, acutis; — a *P. rupestri* (L.) involucri phyllis inferioribus conspicue angustioribus, longioribus, minusque scariosis, interioribus acutis. Planta dalmatica ob phylla interiora acutata medium tenet inter nostram et plantam occidentalem.

In rupestribus regionis inferioris, frequens. Thessalia: pr. Volo (Heldr.); Aetolia: mt. Chalkis et Taphiassos ad sinum Patranum (Hal.); Attica: pr. Athenas, in Acropoli, mt. Lycabettus, Kerata, pr. Laurion, Pharmacusarum insula Lero et Mikra Kyra, insula Aegina (Heldr.); Achaia: pr. Patras (Bois.), Psatopyrgos (Heldr.); Acrocorinthus (Haussk.); Argolis: pr. Nauplia (Friedr.), mt. Palamidi, Tolon, in peninsula Methana (Haussk.), insula Hydra (Heldr.); Laconia: mt. Taygetos (Chaub.); Messenia: pr. Pylos (Heldr.), Methone, Arcadia (Chaub.); Cycladum insula: Mykonos, Keos, Thera (Heldr.), Syra (Weiss), Delos (Tunt.); Creta: pr. Topolia (Bald.), Kissamos, Malaxa (Rev.), Akroteri, Canea, Lutro, Candia (Raul.); Strophades (Reiser); Zante (Marg.); Cephalonia: pr. Argostoli, Steno (Heldr.); Corcyra: in urbe (Gelmi), pr. Palaeokastrizza (Ung.), Manducchio (Form.). — Mart. Jun. ♄.

3. **P. pumilum** S. et S. pr. II. p. 173, Fl. gr. IX. p. 49, t. 863. (*Conyza*); DC. pr. V. p. 397; Raul. cret. p. 778; Bois. fl. or. III. p. 222; Spreitz. in z. b. G. 1890 p. 297; Bald. viagg. Creta p. 64. — *Conyza pygmaea* Sieb. avis p. 5, Reise II. p. 322, t. 10.

Suffruticosum, nanum; rhizomate crasso, lignoso; caulibus brevissimis, ad rosulam saepe reductis; foliis oblongo-spathulatis, integris; pedunculis

monocephalis, bracteis 1—2 squamaeformibus obsitis; involucri phyllis laxiusculis, pallidis, pauciseriatis, scariosis, non undulatis, lanceolatis, acuminatis, externis brevioribus. — Species insignis.

α. **tomentosum** Raul. cret. p. 778; Spreitz. in z. b. G. 1890 p. 297. — Totum cano-tomentosum. — Exsicc.: Heldr. pl. cret. n. 1620; Spreitz. it. cret. a. 1882. —

β. **glabrum** Bois. fl. or. III. p. 222; Bald. viagg. Creta p. 64. — Viride, glabrum, glandulis sessilibus obsitum. — Exsicc.: Heldr. pl. cret. a. 1846; Bald. it. cret. n. 71.

In rupium fissuris regionis subalpinae et superioris Cretae: ad Hellinoseli, Stravopodia (Raul.), Mavrus Lakus, Hagios Theodoros (Spreitz.), Hagion Pneuma (Heldr.) in mt. Sphacioticis (Sibth.); — β. mt. Hagion Pneuma, Volakia, Stravopodia (Heldr.), mt. Gigilos (Bald.). — Jun. Jul. ♄.

14. Helichrysum DC. pr. VI. p. 169.

a. Involucrum niveum, phyllis intermediis discum valde superantibus, patulis, demum radiantibus.

1. **H. amorginum** Bois. et Orph. diagn. ser 2, V. p. 110. — Exsicc.: Orph. fl. gr. n. 471; Dörfl. fl. aeg. n. 52, herb. norm. n. 3528.

Cano-tomentosum, basi suffruticosum; caulibus erectis, apice corymbosis; foliis radicalibus congestis, caulinisque inferioribus oblongo-spathulatis, in petiolum attenuatis, superioribus multo minoribus, linearibus, sessilibus; capitulis breviter pedicellatis, majusculis; involucri phyllis ellipticis, obtusis, glabris, nitidis, intermediis sublongioribus, infimis et intimis brevioribus, hisce lineari-spathulatis. — Species pulcherrima.

In rupibus insulae Amorgos Cycladum supra urbem (Griparis). — Apr. Maio. ♄.

b. Involucrum flavum, phyllis discum non vel vix superantibus, erectis, non radiantibus.

α. Folia latiuscula, plana, saltem infima oblongo-spathulata; capitula majuscula.

2. **H. orientale** L. sp. p. 853; S. et S. pr. II. p. 169, Fl. gr. IX. p. 45 t. 858; (*Gnaphalium*); DC. pr. VI. p. 183; Ung. Reise p. 124; Raul. cret. p. 784; Bois. fl. or. III. p. 230; Heldr. fl. cephal. p. 44; Spreitz. in z. b. G. 1887 p. 662; Bald. viagg. Creta p. 65. — Exsicc.: Orph. fl. gr. n. 1143; Spreitz. it. ion. a. 1880.

Dense et adpresse albo-lanatum, basi suffruticosum, lignosum, tortuosum; caulibus erectis, apice densiuscule corymbosis; foliis radicalibus congestis, caulinisque inferioribus oblongo-spathulatis, obtusis, basi longe attenuatis, superioribus sensim minoribus, linearibus, acutiusculis, sessilibus; capitulis ovato-hemisphaericis, 1 cm latis, breviter pedicellatis; involucri citrini, nitidi, phyllis laxis, concavis, glabris, ab externis ovatis triplo brevioribus ad intima lineari-spathulata sublanata sensim auctis.

In rupestribus regionis calidae. Creta: pr. Katholiko, Hagios Joannes pr. Cydonia (Heldr.), ad promontorium Maleka (Raul.); Cythera: inter Hagios Nicolaos et Kapsali (Spreitz.); Cycladum insula: Pholegandros, Amorgos (Orph.); Cephalonia: in rupe Platyvalo pr. Lasi, pr. Same (Philippides); indicatur quoque ad lacum Hylike Boeotiae (Ung.). — Maio, Jul. ♄.

3. **H. plicatum** DC. pr. VI. p. 183; Bald. riv. coll. bot. Alb. 1896 p. 65. — *H. anatolicum* Bois. diagn. IV. p. 11, ex ipso in Fl. or. III. p. 231. — *H. plicatum v. anatolicum* Haussk. symb. p. 112; Form. in Ver. Brünn 1896 p. 46, 1897 p. 34. — Exsicc.: Orph. fl. gr. n. 1067 (Bitolia); Bald. it. alb. epir. IV. p. 176.

Parce areneosum et dense glandulosum, virens, basi suffruticosum; caulibus erectis, apice corymbosis; foliis radicalibus congestis, caulinisque inferioribus oblongo-spathulatis, obtusiusculis, in petiolum attenuatis, ceteris vix decrescentibus, lanceolatis, acutis vel acuminatis, sessilibus; capitulis globosis, 5 mm latis, breviter pedicellatis; involucri aurei, nitidi, phyllis laxiusculis, subconcavis, glabris, ab externis ovatis triplo brevioribus, ad intima oblongo-linearia sensim auctis. —

In herbidis regionis subalpinae. Epirus: mt. Kuruna supra Diovisda distr. Ljaskovik (Bald.); Thessalia: pr. Chaliki, Mandra Hodza (Form.), mt. Zygos, Oxya, Ghavellu et Karava (Haussk.) in Pindo. — Jul. Aug. ♄.

β. Folia angusta, linearia, margine revoluta; capitula parva.
× Involucri phylla inferiora intimis 2—3 plo tantum breviora.

4. **H. Heldreichii** Bois. fl. or. III. p. 229. — *H. panormitanum* Bois. diagn. XI. p. 31, non Tiff — *H. panormitanum v. angustifolium* Raul. cret. p. 784. — Exsicc.: Heldr. pl. cret. n. 1629.

Basi suffruticosum; caulibus adscendentibus, inferne crebre et longe foliatis; foliis linearibus, obtusis, utrinque dense cano-tomentosis; corymbo composito, denso; capitulis minutis, campanulatis, 4—5 mm. longis; involucri phyllis basi lanatis, superne glabris, stramineis, nitidis, externis oblongo-triangularibus acutis, internis duplo longioribus, linearibus, acuminatis. — Capitulis minutis et involucri phyllis acutis insigne. — Ab affinibus *H. rupestre* DC., *H. panormitano* Tin et *H. nebrodensi* Heldr., capitulis duplo minoribus, phyllisque involucri duplo angustioribus et acutioribus, basi lanatis eximie differt.

In rupibus faucis Hagia Rumeli Cretae, rare (Heldr.). — Jun. ♄.

5. **H. siculum** Spreng. syst. III. p. 476 (*Gnaphalium*); Bois. fl. or. III. p. 229; Spreitz. in z. b. G. 1877 p. 713, 1887 p. 662; Heldr. fl. cephal. p. 44, Fl. Aegina p. 303; Haussk. symb. p. 112; Form. in D. bot. Monat. 1898 p. 78. — *G. scandens* Sieb. avis p. 5, rem. p. 6. — *G. stoechas* Ch. et B. exp. p. 246, Fl. pelop. p. 57; Fraas fl. class. p. 208; non L., quod differt odore aromatico, foliis tenuioribus, capitulis minus dense conglobatis, involucri phyllis numerosioribus, magis

inaequalibus, infimis acutiusculis. — *H. rupestre v. Cambessedesii* DC. pr. VI. p. 182 quoad pl. graec.; Marg. et R. fl. Zante p. 58. — *H. stoechas* Marg. et R. l. c. err. et corr. p. 96, non (L.) DC. — *G. conglobatum* Friedr. Reise p. 271 et 277, non Viv., quod = *H. Fontanesii* Camb. — *H. conglobatum* Ung. Reise p. 124; Weiss in z. b. G. 1869 p. 41, non (Viv.) Steud. — *H. scandens* Raul. cret. p. 784. — ? *H. stoechas v. incanum* Bald. viagg. cret. p. 65. — Exsicc.: Orph. fl. gr. n. 470; Heldr. herb. norm. n. 462, herb. fl. hellen. n. 93, in Baen. herb. europ. n. 3600, in Magn. fl. sel. n. 2498; Reverch. pl. cret. n. 77, in Baen. herb. europ. n. 4880; Dörfl. pl. cret. n. 31.

Basi suffruticosum, multicaule; caulibus tenuibus, erectis vel adscendenti-decumbentibus, inferne crebre foliatis; foliis anguste-linearibus, obtusiusculis, supra arachnoideo-virentibus, subtus dense cano-tomentosis; corymbo oligocephalo, dense capitato; capitulis parvis, campanulato-subglobosis, 4—6 mm. longis; involucri phyllis basi glabris, stramineis, nitidis, ovatis, obtusis, internis duplo-longioribus, lineari-spathulatis, acutiusculis, dorso tomentellis. — Differt a praecedente foliis angustioribus brevioribusque, supra virentibus, capitulis majoribus, phyllis latioribus, obtusis, basi non lanatis.

β. **laureoticum** Haussk. symb. p. 112. — Caules elongati, folia angustiora, subtus leviter griseo-tomentosa, capitula minora, involucri phylla angustiora, acuta, interiora mucronulata. — Exsicc.: N. v.

γ. **brachyphyllum** Bois. fl. or. III. p. 230; Spreitz. in z. b. G. 1890 p. 296; Haussk. symb. p. 112. — *H. decumbens v. spathulata* Raul. cret. p. 784. — Caules abbreviati, folia multo breviora, inferiora 3—5 mm. tantum longa, oblongo-spathulata, superiora lineari-subspathulata, 5—10 mm. longa. — Exsicc.: Heldr. pl. cret. n. 1452.

In collibus saxosis et rupestribus regionis inferioris et montanae, freqnens. Boetia: mt. Helicon (Fraas); Attica: pr. Athenas, Phalerum, mt. Lycabettus, in valle Cephissi pr. Podoniphti, mt. Parnes, Pentelicon, Hymettus, pr. Raphina, Sunium, Laurion, Kamariza, Pharmacusarum insula Mikra Kyra, insula Aegina et adjacente Metopi (Heldr.); Petalium insula Xeronisi, (Holzm.); isthmus Corinthiacus pr. Lutraki (Heldr.); Argolis: pr. Poros (Friedr.), Epidaurus (Chaub.); Messenia: pr. Pylos, Arcadia, Methone, Corone (Chaub.), Kalamata (Zahn); Cycladum insula: Cythnos (Orph.), Syra, Tenos (Weiss), Naxos (Bois.); Creta: pr. Vukolies (Bald.), Kissamos (Reverch), Canea, Suda (Weiss), Sitia, Khalepa, Akroteri, Francocastello, Guves (Raul.), Males distr. Hierapetra (Leon.); Cythera: pr. Hagios Nicolaos (Spreitz.); Zante et adjac. insula Peluso (Marg.); Cephalonia: pr. coenobium Sisia (Heldr.); Corcyra: pr. Benizze (Spreitz.); — *β.* pr. Kamariza distr. Laurion Atticae (Haussk.); — *γ.* Creta: pr. Askyphos (Spreitz.), ad promontorium Sidero (Heldr.); indicatur quoque ad Phalerum Atticae et Neocorinthum (Haussk.). — Ap.r Jun. ♄.

XX Involucri phylla ab infimis brevissimis ad intima multo longiora regulariter imbricata.

6. **H. italicum** Roth cat. bot. II. p. 115 (*Gnaphalium*); Guss. fl. sic. syn. II. p. 469; Weiss in z. b. G. 1869 p. 41; Bois. fl. or. III. p. 234; Hal. in ö. b. Z. 1895 p. 411; Heldr. chlor. Thera p. 14. — *G. stoechas* S. et S. pr. II. p. 169, Fl. gr. IX. p. 44, t. 857; Urv. enum. p. 108; non L. — *H. angustifolium* DC. fl. fr. V. p. 467, p. p. — *H. serotinum v. orientale* Bois. voy. esp. p. 328. — Exsicc.: Orph fl. gr. n. 1144; Dörfl. fl. aeg. n. 168.

Basi suffruticosum, multicaule, adpresse canescens; caulibus tenuibus, virgatis; foliis lineari-filiformibus, supra demum virentibus, inferioribus confertis brevioribus, superioribus saepe subincurvis; corymbo composito, capitulis parvis, oblongo-cylindricis; involucri phyllis adpresse imbricatis, scariosis, concavis, obtusis, ab externis ovatis, tomentellis, ad intima lineari-oblonga, dorso glandulis adspersa sensim auctis; acheniis minute glandulosis.

β. **microphyllum** Willd. sp. III. p. 1863; Sieb. avis p. 5, rem. p. 6, in Flora I. p. 275; pro sp.; (*Gnaphalium*); Bois. fl. or. III. p. 234; Heldr. in ö. b. Z. 1898 p. 184. — *H. microphyllum* Camb. in mem. mus. hist. nat. XIV. p. 272; Raul. cret. p. 784; Bald. viagg. Creta p. 65. — Caules abbreviati, folia multo breviora. — Exsicc.: Heldr. pl. cret. a. 1846; Bald. viagg. Creta n. 73.

In saxosis regionis inferioris. Cycladum insula: Syra (Weiss.), Evreo Kastron pr. Paros (Reis.), Kythnos, Seriphos, Tenos (Sart.), Siphnos, Ios, Naxos, Thera, Therasia (Heldr.); — β. Creta: pr. Akroteri, Drakona, Askyphos, Niato, mt. Volokia (Raul.), mt. Omalo (Bald.), mt. Ida (Bois.); Cycladum insula Naxos (Heldr.). — Maio, Jul. ♄.

15. Gnaphalium L. gen. n. 949.

a. Annua; capitula glomerata.

1. **G. luteoalbum** L. sp. p. 851; S. et S. pr. II, p. 170; Ch et B. exp. p. 246, Fl. pelop. p. 57; Friedr. Reise p. 265 et 277; Raul. cret. p. 785; Bois. fl. or. III. 224; Haussk. symb. p. 112; Form. in Ver. Brünn 1897 p. 34. — Icon: Fl. dan. t. 1763. — Exsicc.: Orph. herb. n. 158; Heldr. herb. gr. dimorph. n. 34.

Albo-lanatum; caulibus erectis vel adscendentibus, simplicibus vel apice ramosis; foliis oblongo-lanceolatis, lanceolatisve, inferioribus obtusis, superioribus acutis; capitulis glomerato-cymosis, aphyllis; involucri phyllis glabris, pallide stramineis, nitidis, obtusis; floribus lutescentibus; acheniis tenuiter tuberculatis.

In arenosis praesertim humidiusculis regionis inferioris et montanae. Thessalia: pr. Limogardi (Form.); Attica: ad lacusculum Thalas in mt. Pentelicon (Heldr.); Corinthus, insula Poros (Friedr.); Argolis: pr. Astros (Sart.); Laconia: mt. Malevo pr. Castanitza et Platanos (Orph.), ad fluvium Eurotas (Chaub.); Creta (Sibth.): pr. Rumata, Platania (Raul.). — Apr. Oct. ⊙.

2. **G. uliginosum** L. sp. 856; S. et S. pr. II. p. 171; Ch. et B. exp. p. 246, Fl. pelop. p. 57; Bois. fl. or. III. p. 225. — Icon: Fl. dan. t. 859. —

Molliter albo-tomentosum; caule a basi ramoso, ramis patulis vel diffusis; foliis lanceolato-linearibus, acutis; capitulis glomerato-aggregatis, foliosis; involucri phyllis apice glabriusculis, fuscis, nitidis, obtusiusculis; floribus lutescentibus; acheniis puberulis.

In Peloponneso (Sibth.), postea ut videtur a nemini lectum. — Jun. Sept. ⊙. N. v.

b. Perennia; capitula racemosa.

α. Phylla involucri extima capitulo triplo breviora.

3. **G. silvaticum** L. sp. p. 856; Bois. fl. or. III. p. 226; Heldr. chlor. Parn. p. 20; Haussk. symb. p. 112. — *G. Sophiae* Heldr. ex ipso l. c. — Icon: Fl. dan. t. 1229. — Exsicc.: Heldr. pl. fl. hellen. a. 1887.

Caulibus firmis, erectis, adpresse lanatis, foliosis, 20—50 cm. altis; foliis cano-tomentosis, supra demum glabrescentibus, acutis, basin versus attenuatis, uninerviis, infimis lanceolatis, caeteris lineari-lanceolatis; capitulis in axillis foliorum superiorum glomeratis, racemum spiciformem, elongatum, foliosum, interruptum formantibus; involucri phyllis adpressis, a basi supra medium usque herbaceis, lanatulis, caeterum scariosis, obtusis, exterioribus ovatis, caeteris oblongis linearibusque, parte scariosa brunneis vel fuscis; acheniis puberulis.

In rupestribus regionis mediae. Thessalia: mt. Zygos in Pindo (Haussk.), mt. Pelion pr. Zagora (Aphentulis); mt. Parnassus (Guicc.). — Jul. Aug. ♃.

4. **G. Roeseri** Bois. et Heldr. diagn. ser. 2. VI. p. 102; Hal in z. b. G. 1899 p. 713. — *G. Hoppeanum* Bois. fl. or. III. p. 227 non Koch. — Exsicc.: Heldr. fl. gr. a. 1855. —

Caespitosum, nanum; caulibus subfiliformibus, erectiusculis, adpresse lanatis, oligophyllis, 2—7 cm. altis; foliis utrinque adpresse sericeo-lanatis, spathulato-lanceolatis, acutis, basi longe attenuatis, trinerviis; capitulis paucis, racemum spiciformem, ovatum, subinterruptum formantibus; involucri phyllis adpressis, a basi supra medium usque herbaceis, lanatis, caeterum scariosis, obtusis, exterioribus ovatis, caeteris oblongis, parte scariosa brunneis; acheniis adpresse sericeis. — *G. Hoppeano* Koch, capitulis numerosioribus, magis confertis, involucri phyllis acutiusculis fuscis, acheniis puberulis diverso, maxime affine.

In rupestribus regionis alpinae. Phocis mt. Parnassus (Guicc.); Laconia: mt. Taygetos (Zahn). — Jul. Aug. ♃.

β. Phylla involucri extima capitulum dimidium superantia.

5. **G. supinum** L. syst. ed. 12. III. p. 234. — *Omalotheca supina* Cass. dict. 56. p. 218. — Icon: Schk. t. 267. — Exsicc.: Bald. it. alb. epir. III. n. 161.

Caespitosum, nanum; caulibus subfiliformibus, erectiusculis, adpresse lanatis, oligophyllis, 2—10 cm. altis; foliis utrinque albo-lanatis, linearibus vel lineari-lanceolatis, acutis, basi attenuatis, uninerviis; capitulis paucis, racemum spiciformem, brevem, subinterruptum formantibus; involucri phyllis adpressis, a basi supra medium usque herbaceis, lanatis, caeterum scariosis, obtusis, exterioribus ovatis, caeteris oblongis, parte scariosa fuscis; acheniis sparsim puberulis.

β. **subacaule** Wahlenb. fl. lapp. p. 402 pro sp.; DC. pr. VI. p. 245 pro var. *Omalothecae supinae*; Bald. riv. coll. bot. Alb. 1896 p. 65. — *G. pusillum* Haenke Riesengeb. p. 93. — Caules brevissimi vel subnulli, capitula inter folia radicalia subsessilia. — Exsicc.: Bald. it. alb. epir. IV. n. 175. —

In pascuis alpinis. Epirus: mt. Tsumerka (Bald.); — β. Epirus: sub jugo Gamila distr. Zagorion (Bald.). — Jun. Aug. ♃.

Obs. *G. supracanum* S. et S. pr. II. p. 171, Fl. gr. IX. p. 47, t. 861 = *Ifloga spicata* Schultz, in Graecia erronee indicatum.

16. Filago L. gen. n. 995.

a. Receptaculum filiforme, elongatum; involucri phylla 5 seriata, cuspidata, in capitulo fructifero convergentia; achenia libera.

1. **F. germanica** L. sp. p. 857; S. et S. pr. II. p. 71; Sieb. avis rem. p. 6; Ch. et B. exp. p. 246, Fl. pelop. p. 57; Friedr. Reise p. 275; (*Gnaphalium*); L. sp. ed. 2 p. 1311; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 785.

Molliter lanato-tomentosa, simplex vel a collo pluricaulis; caulibus erectis vel adscendentibus, apice breviter et stricte ramoso-subcorymbosis; foliis crebris, sessilibus, lineari vel oblongo-lanceolatis; glomerulis densis, capitulis 20—30 constantibus, in dichotomiis atque ad ramorum apicem sessilibus, foliis floralibus eis brevioribus involucratis; involucro ad medium usque lana immerso, obconico, obtuse pentagono, phyllis laxis, longitudinaliter plicatis, longe subulato-acuminatis, acuminibus erectis.

α. **canescens** Jord. obs. III. p. 202 t. 7, f. a; Haussk. symb. p. 112; pro sp. — ? *F. pyramidata* L. sp. p. 1199. — *F. pyramidata v. canescens* Form. in D. bot. Mon. 1890 p. 18, 1898 p. 78, in Ver. Brünn 1895 p. 26, 1896 p. 46, 1897 p. 34. — Tota planta albo-tomentosa; phylla involucri apice pallida. — Exsicc.: Sint. it. thessal. n. 183.

β. **lutescens** Jord. obs. III. p. 201, t. 7, f. b; Raul. cret. p. 785; pro sp. — Tota planta lutescenti-tomentosa; phylla involucri fuscidula, apice saepe purpurascentia. — Exsicc.: N. v.

γ. **eriocephala** Guss. pl. rar. p. 344 t. 69; Hal. Beitr. fl. Epir. p. 27, in ö. b. Z. 1897 p. 95, 1899 p. 25; Haussk. symb. p. 112: pro sp.; Bois. fl. or. III. p. 245; Bald. riv. coll. bot. Alb. 1896 p. 65. — *F. germanica v. lanuginosa* DC. pr. VI. p. 247; Marg.

et R. fl. Zante p. 58. — Albo-tomentosa; involucrum usque ad $^3/_4$ longitudinis lana immersum, phyllis brevius acuminatis, saepe tantum acutis. — Intermediis ad var. α transit. — Exsicc.: Heldr. herb. norm. n. 850 a et b, 1444; Sint. it. thessal. n. 1132.

In arenosis, arvis, campis, olivetis, quericetis regionis inferioris et montanae. — α. Epirus: pr. Prevesa, Mitikas. Han Balduma (Form.); frequens in Thessalia tota (Form.); β. indicatur pr. Canea Cretae (Raul.); — γ. ut videtur multo vulgatior: Epirus: mt. Profetae Elias distr. Liaskovik (Bald.), pr. Kalentini (Hal.); Thessalia: pr. Sermenikon (Sint.), ad monasterium Korona, pr. Pharsalus (Heldr.); Euboea; pr. Kastaniotissa (Heldr.); Sporadum insula Skopelos (Leon.); Cycladum insula: Cythnos, Keos, Jos, Tenos (Heldr.), Amorgos (Orph.); Argolis: pr. Vromolimni in peninsula Methana (Heldr.); Messenia: pr. Kalamata, Diabolitzi (Zahn); Elis: in colle Chlemutzi (Heldr.); in Strophadum insula majore (Reiser); Zante (Marg.) — Typus dein indicatur in Peloponneso: iusula Poros (Friedr.), ad fluvium Eurotas, in Zarnate, pr. Methone, Corone, Messene (Chaub.); Creta: pr. Canea, Malaxa, Istovai ad promontorium Sidero (Raul.). — Maio, Jul. ⊙

2. **F. spathulata** Presl. del prag. p. 93; Weiss in z. b. G. 1869 pag. 41; Bois. fl. or. III. p. 246; Hal. Beitr. fl. Aetol. p. 7, Beitr. fl. Achaia p. 24 (*f. decalvans*), in ö. b. Z. 1897 p. 95; Haussk. symb. p. 112, cum *f. affinis*, caulibus pumilis erectis, dense foliatis, foliis abbreviatis, cauli adpressis; Form. in Ver. Brünn 1896 p. 46, 1897 p. 34. — *F. Jussiaei* Coss. et Germ. in ann. sc. nat. ser. 2 XX. p. 284, t. 13; Clem. sert. p. 60.

Differt a praecedente foliis oblongo-spathulatis, basi attenuatis; glomerulis e capitulis 12—15 constantibus, foliis floralibus eis longioribus involucratis; involucri basi vel ultra medium lanati, ovoidei, acutiuscule pentagoni, phyllis adpressis, carinatis, longe subulato-acuminatis, acuminibus subpatulis.

α. **typica.** — Simplex vel a collo pluricaulis; caulibus erectis vel basi adscendentibus, 30 cm. usque altis, apice vel a medio ramosis. — Exsicc.: Heldr. herb. norm. n. 849, mixta cum var. β.; Dörfl. pl. cret. n. 143.

β. **prostrata** Parl. pl. nov. p. 11; Raul. cret. p. 785; pro sp. onn DC. pr. VI. p. 249; Bois. fl. or. III. p. 246; Heldr. fl. cephal. p. 44, Fl. Aegina p. 303; Haussk. symb. p. 112. — Prostrata, ramosissima, glomeruli numerosi, saepe contigui, phylla praeter cuspidem lanata. — Intermediis ad typumtransit. Exsicc.: Heldr. herb. norm. n. 531.

γ. **siria** Heldr. herb. norm. n. 1443. — Pusilla, a collo ramosa, glomeruli parvi, involucri basi lanati phyllis glabriusculis, nitidis, brevius acuminatis vel tantum acutis. — An species propria.

In herbidis, campis, dumosis regionis inferioris et montanae. Epirus: mt. Peristeri (Form.); Thessalia: pr. Orman Magula, Pharsalum (Heldr.),

pr. Neraida, mt. Pelion (Form.); Sporadum insula Scopelos (Leon.); Aetolia: pr. Mesolongion (Nied.), mt. Taphiassos (Hal.); Boeotia: pr. Mulki (Heldr.); Attica: pr. Athenas, Eleusis (Haussk.), mt. Pentelicon (Clem.), in valle Cephissi (Heldr.); Acrocorinthus (Haussk.); Achaia: mt. Olenos (Hal.); Cycladum insula Syra (Weiss); Creta: pr. Canea (Weiss), Males distr. Hierapetra (Leon.); — *β.* typo vulgatior, Thessalia: pr. Trikala, Sophates, Aivali, Pharsalus (Haussk.); Attica: pr. Athenas, Eleusis, mt. Pentelicon, insula Aegina (Heldr.); Argolis: peninsula Methana (Haussk.); Arcadia: pr. Carithena (Prinopulos); Messenia: pr. Kalamata (Zahn); Cycladum insula Keos, Cythnos, Tenos (Tuntas); Creta: pr. Francocastello, Rhodakino, mt. Aphendi Kavutsi (Raul.); Cephalonia: in oropedio Omala (Heldr.); — *γ.* insula Keos, in rimis humentibus pr. urbem (Heldr.). — Apr. Jul. ⊙

Obs. Formas hybridas: *F. canescens* × *spathulata* (*F. intermedia*) *et F. eriocephala* × *spathulata* (*F. similata*) Haussk. symb. p. 113, observavit auctor pr. Pharsalum Thessaliae et pr. Vromolimni in peninsula Methana Argolidis.

b. Receptaculum breve, applanatum; involucri phylla 3 rarius 4-seriata, non cuspidata, in capitulo fructifero stellatim patentia.

3. **F. arvensis** L. sp. p. 856 (*Gnaphalium*); L. sp. in add. p. ult. post indicem n. 5; Form. in D. bot. Mon. 1890 p. 18, in Ver. Brünn 1896 p. 46, 1897 p. 34. — Icon: Fl. dan. t. 1275. — Exsicc.: N. v.

Molliter lanato-tomentosa; caulibus erectis, stricte paniculatis; foliis lineari-lanceolatis, acutis, floralibus glomerulos aequantibus; glomerulis oligocephalis, lateralibus terminalibusque; involucri ovati phyllis subbiseriatis, externis subfoliaceis, internis longioribus, scariosis, obtusiusculis, lanatis, apice glabrescentibus; acheniis liberis.

β. **lagopus** Steph. in Willd. sp. III. p. 1897 pro sp. (*Gnaphalium*); DC. pr. VI. p. 249; Bois. fl. or. III. p. 247; Haussk. symb. p. 113; Bald. riv. coll. bot. Alb. 1895 p. 47; Form. in Ver. Brünn 1897 p. 34. — *F. lagopus* Parl. pl. nov. p. 13; Raul. cret. p. 785, *v. alpina*. — Folia latiora, capitula minus aggregata, saepe solitaria, dense floccoso-lanata. — Exsicc.: Heldr. herb. norm. n. 347, it. thessal. n. 8; Orph. fl. gr. n. 776; Bald. it. alb. epir. III. n. 162; Sint. it. thessal. n. 590. —

In aridis, vineis, nemorosis siccis regionis inferioris, in alpinam usque adscendens. Thessalia: pr. Velitsena, Malakasi, Makrichori, Kalabaka, Vlachava, Kritsotades, mt. Cuka in mt. Chassia, Larissa, Volo (Form.); — *β.* Epirus: mt. Tsumerka (Bald.); Thessalia: mt. Zygos, pr. Malakasi, monasterium Korona (Haussk.), Phlambures et Mitrica in mt. Chassia (Form.), in cacumine mt. Pelion (Heldr.); mt. Parnassus, Pateras, Hymettus (Heldr.); Laconia: mt. Malevo pr. Castanitza (Orph.); Creta: mt. Ida (Raul.). — Jun. Aug. ⊙

4. **F. gallica** L. sp. p. 857; S. et S pr. II. p. 171; Ch. et B. exp. p. 246, Fl. pelop. p. 57; (*Gnaphalium*); L. sp. in add. p. ult. post indicem n. 6; Urv. enum. p. 109; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 785; Bois. fl. or. III. p. 248; Form. in D. bot. Mon. 1890 p. 18, in Ver. Brünn 1896 p. 46; Haussk. symb. p. 113; Heldr. fl. Aegina p. 393, Chlor. Mykonos p. 245. · *F. tenuifolia* Presl del. prag. p. 101; Raul. cret. p. 785. — ? *F. minima* Form. in D. bot. Monat. 1890 p. 18, vix Fr. — Icon: Rchb. germ. t. 947. — Exsicc.: Orph. fl. gr. n. 775.

Adpresse albo-tomentosa, subsericea; caulibus erectis, saepe jam a basi ramosis, ramis gracilibus; foliis lineari-subulatis, floralibus glomerulos superantibus; glomerulis oligocephalis, lateralibus terminalibusque; involucri obtuse pentagoni phyllis triseriatis, externis herbaceis, internis longioribus, scariosis, obtusis, glabrescentibus; acheniis marginalibus phyllorum involucri omnino cinctis, ceteris liberis.

In campis, collibus aridis regionis inferioris et montanae. Thessalia: pr. Karditza, Palaeokastro, Korona, Kalabaka, Meteora (Haussk.), Han Kukleus in valle Penei, pr. Larissa, Makrichori, Patsios in Olympo (Form.), Caterina (Orph.); mt. Parnassus pr. Krisso (Guicc.); Attica: mt. Pentelicon, Hymettus (Haussk.); pr. Kurbatzi Euboeae (Wild.); insula Aegina, peninsula Methana Argolidis, pr. Patras Achaiae, pr. Chlemutzi Elidis (Heldr.); mt. Diaforti, mt. Kupe pr. Pylos (Chaub.), pr. Kalamata (Zahn); Cycladum insula Keos, Jos, Tenos (Heldr.), Cythnos (Tunt.), Melos (Urv.); Creta (Sibth.); pr. Enneachoria, Canea, Akroteri, Malaxa, Pilialimata (Raul.). — Apr. Aug. ⊙.

17. Evax Gaertn. fr. II. p. 393.

a. Folia floralia glomerulis longiora.

1. **E. pygmaea** L. sp. ed. 2 p. 1311; S. et S. pr. II. p. 107; (*Filago*); Pers. syn. II. p. 422; Marg. et R. fl. Zante p. 56; Friedr. Reise p. 270; Fraas fl. class. p. 209; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 778; Bois. fl. or. III. p. 242; Spreitz. in z. b. G. 1877 p. 713, 1887 p. 662; Heldr. fl. cephal. p. 44, Fl. Aegina p. 304; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Achaia p. 24, in ö. b. Z. 1895 p. 412, 1897 p. 95, 1899 p. 25; Haussk. symb. p. 118. — *Micropus pygmaeus* Desf. atl. II. p. 307; Urv. enum. p. 109; Ch. et B. exp. p. 246, Fl. pelop. p. 58. — Icon: Fl. gr. t. 921. — Exsicc.: Orph. fl. gr. n. 188; Baen. herb. europ. n. 9256; Dörfl. pl. cret. n. 17. —

Cano-tomentosa, simplex vel a collo ramosa; caule 1—5 cm. alto; foliis oblongo-spathulatis, oblongis, obtusis; capitulis in glomerulos terminales conglobatis, rarius solitariis; involucri phyllis paleisque dorso glabriusculis, ovatis, acuminatis, apice abrupte patulis; receptaculo elongato; acheniis obovatis, margine scabriusculis.

In arenosis, rupestribus, pascuis siccis regionis inferioris. Aetolia: pr. Mesolongion (Nied.), Krioneri ad sinum Patranum (Reis.); Attica:

ad Phalerum (Orph.), Piraeum (Heldr.), Geraka in planitie Mesogia; ad Acrocorinthum (Haussk.), Neocorinthum (Heldr.); insula Aegina: cap Perdicca (Friedr.); Achaia: pr Patras (Hal.), mt. Corphi (Orph.); Elis (Sibth.); Messenia: pr. Diabolitzi (Zahn), Arcadia, Methone, Messene, Zarnate (Chaub.); mt. Parnon (Fraas); Argolis (Sprun.); Sporadum insula Skiathos (Leon.); Cycladum insula: Keos (Heldr.), Cythnos, Mykonos (Tunt.), Syra (Weiss), Melos (Urv.); Creta: pr. Canea (Weiss), Akroteri (Raul.), Anatoli distr. Hierapetra (Leon.); Strophadum insula major (Reis.); Zante (Marg.); Cephalonia: pr. Argostoli, Pessada, Piscardo (Heldr.); Corcyra: pr. Benizze, cap Bianco (Spreitz.), Palaeokastrizza (Baen.). — Mart. Jun. ⊙

b. Folia floralia glomerulis aequilonga vel breviora.

2. **E. exigua** S. et S. pr. II. p. 207 (*Filago*); DC. pr. V. p. 458; Parl. in ann. sc. nat. Paris 1841 p. 300; Bois. fl. or. III. p. 244; Heldr. chlor. Thera p. 14. — *Micropus exiguus* Urv. enum. p. 109. — *Filago congesta* Guss. in DC. pr. VI. p. 248; ex ipso in Fl. sic. syn. II. p. 460. — Icon: Fl. gr. t. 922.

Cano-tomentosa, a collo breviter ramosa; caule vix bipollicari; foliis oblongo-spathulatis, obtusis; capitulis in glomerulos terminales dense congestis, paucifloris; involucri phyllis externis lanatis, paleisque externis concavis, subcarinatis, acuminatis, internis obtusis, glabris; receptaculo brevi; acheniis oblongis scabriusculis.

In Archipelagi insulis (Sibth.); in summo vulcano novae Camini pr. Thera (Urv.). — Maio. ⊙ N. v.

3. **E. perpusilla** Bois. et Heldr. diagn. ser. 2, III. p. 18, Fl. or. III. p. 245; Heldr. chlor. Parn. p. 20. — Exsicc.: Heldr. herb. norm. n. 244; Orph. fl. gr. n. 329.

Molliter lanato-tomentosa, simplex vel a collo pauciramosa; caule 5—20 mm. alto; foliis oblongis, obtusis; capitulis in glomerulos parvos dense congestis, paucifloris; involucri phyllis paleisque oblongis, concavis, obtusis, praeter basin glabram albo-lanatis; receptaculo brevi; acheniis ovato-oblongis, papilloso-hirtis. — A praecedentibus phyllis et paleis obtusis egregie differt.

In pascuis siccis regionis abietinae mt. Parnassus pr. Hagios Nicolaos, nec non in regione superiori pr. Messino nero (Orph.). — Jun, Jul. ⊙

18. Micropus L. gen. n. 996.

1. **M. erectus** L. sp. in add. p. ult. post indicem n. 2; S. et S. pr. II. p. 208; Ung. Reise p. 123; Bois. fl. or. III. p. 241. — Icon: Lam. ill. III. t. 694. — Exsicc.: Orph. fl. gr. n. 773 (Macedonia).

Molliter lanato-tomentosus; caule erecto, strictiuscule ramoso vel a collo ramoso, ramis diffusis; foliis oblongis, margine undulatis, floralibus glomerulos superantibus; glomerulis globosis, lateralibus terminalibusque; involucri phyllis oblongis, obtusis, lanatis. — *Filagini arvensi* similis, sed acheniis epapposis ab ea statim diagnoscenda.

In variis Archipelagi insulis (Sibth.) et in Cephalonia (Ung.). — Maio, Jul. ⊙. N. v.

2. **M. bombycinus** Lag. gen. et sp. p. 32; Bois. fl. or. III. p. 241; Spreitz. in z. b. G. 1887 p. 662. — Icon: Rchb. germ. t. 943. — Differt a praecedente lana copiosiori et longiori; foliis non undulatis, floralibus glomerulos non superantibus.

In aridis. Attica: mt. Kerata pr. Eleusis (Heldr.); Achaia: pr. Trikala (Heldr.); Leucas: pr. San Giovanni (Spreitz.). — Maio, Jul. ⊙. N. v.

4. Tribus. **HELIANTHEAE** Less. syn. p. 221.

19. Bidens L. gen. n. 932.

1. **B. tripartita** L. sp. p. 831; Form. in D. bot. Monat. 1890 p. 16. — Icon: Fl. dan. t. 2178. — Exsicc.: Heldr. pl. fl. hellen. a. 1898.

Glabriuscula; caule erecto, ramoso; foliis petiolatis, tripartitis, segmentis lanceolatis, dentatis; capitulis erectis, solitariis, discoideis; acheniis biaristatis.

In paludosis, ad fossas, rare. Thessalia: pr. Trikala (Form.); Messenia: inter Kalamata et Nisi (Zahn). — Jul. Nov. ⊙.

2 **B. cernua** L. sp. p. 832; Hal. in ö. b. Z. 1890 p. 39; Bald. riv. coll. bot. Alb. 1896 p. 65. — Icon: Fl. dan. t. 841. — Exsicc.: Sint. et Bornm. it. turc. n. 1289.

Glabriuscula; caule erecto, ramoso; foliis sessilibus, indivisis, lanceolatis, dentatis; capitulis cernuis, solitariis, radiatis vel discoideis; acheniis quadriaristatis.

In paludosis, ad fossas, rare. Epirus: ad lacum Janina (Bald.); Thessalia: ad Braiabalti pr. Litochori (Sint.). — Jul. Nov. ⊙.

Obs. *Helianthus tuberosus* L. sp. p. 904; Heldr. Nutzpfl. p. 26. — Colitur pr. Athenas (Heldr.). —

5. Tribus. **ANTHEMIDEAE** Cass. opusc. phyt. III. p. 61.

20. Diotis Desf. fl. atl. II. p. 261.

1. **D. maritima** L. sp. p. 927 (*Filago*); Sm. engl. fl. III. p. 403; Bois. fl. or. III. p. 253; Hal. in ö. b. Z. 1895 p. 411; Heldr. in ö. b. Z. 1898 p. 184, chlor. Mykonos p. 245. — *Athanasia maritima* L. sp. ed. 2 p. 1182. — *Santolina maritima* Sm. fl. brit. II. p. 860; S. et S. pr. II. p. 165; Sieb. avis p. 5; Ch. et B. exp. p. 245, Fl. pelop. p. 57; Fraas fl. class. p. 208. — *D. candidissima* Desf. l. c.; Sieb. avis rem. p. 6; Raul. cret. p. 782; Spreitz. in z. b. G. 1887 p. 662; Bald. viagg. Creta p. 62. — *Otanthus maritimus* Lk.

fl. port. II. p. 365. — Icon: Fl. gr. t. 850. — Exsicc.: Sint. et Bornm. it. turc. n. 1282; Dörfl. fl. aeg. n. 131.

Albo-pannosa, basi suffruticosa; caulibus erectis vel adscendentibus simplicibus vel parce ramosis; foliis oblongis, semiamplexicaulibus, obtusis, integris; capitulis globosis, breviter pedunculatis, in corymbum dispositis; floribus luteis.

In arenosis maritimis. Thessalia: pr. Litochori (Sint.), pr. Zagora ad mt. Pelion (Aphent.); Sporadum insula Psathura (Reis.); Aetolia: pr. Mesolongion (Nied.); Attica: ad Phalerum (Fraas), Raphina, Laurion (Heldr.), Sunium (Chaub.), Salamis (Orph.); Argolis: peninsula Methana (Chaub.); Messenia: pr. Gialova (Zahn); Cycladum insula: Mykonos (Sart.), Tenos (Chaub.), Naxos (Reis.); Creta: pr. Kissamos, Platania, Rethymo, insula Dia, promontorium Sidero (Raul.); Zante (Mazz.); Corcyra: pr. Pelleca (Spreitz.). — Maio, Aug. ♃.

21. Achillea L. gen. n. 646.

Dispositio specierum:

1. Sectio. *Ptarmica* Tourn. — Ligulae 8—10, albae, saepius magnae et involucro aequilongae. Foliorum segmenta non transverse imbricata.

a. Folia subintegerrima vel minute crenata vel subpinnatifida; capitula majuscula plerumque solitaria, rarissime 2—3 in corymbo laxiusculo.

1. A. ageratifolia S. et S. **2. A. aizoon** Griseb.

b. Folia pinnatim partita vel secta, rarissime integra; capitula parva, fere semper numerosa, corymbosa, rarissime pauca vel solitaria.

α. Caulis et folia plus minus tomentosa vel sericea, cana.

× Ligulae adultae involucro aequilongae.

3. A. Clavennae L. **4. A. umbellata** S. et S.

×× Ligulae involucrum dimidium aequantes.

○ Folia inferiora pectinatim pinnatipartita segmentis integris, superiora integra.

5. A. Barbeyana Heldr. et Heim.

○○ Folia inferiora bipinnatisecta, superiora pinnatisecta.

6. A. ambrosiaca Bois. et Heldr. **7. A. Fraasii** Schultz.

β. Caulis et folia glabra vel hirtula, viridia.

8. A. olympica Heim. **9. A. abrotanoides** Vis.

2. Sectio. *Millefolium* Tourn. — Ligulae 4—5, rarissime numerosiores, albae vel flavae, involucro saepissime dimidio breviores, interdum brevissimae.

a. Ligulae 4—5, albae vel dilute ochroleucae; foliorum segmenta non transverse imbricata.

α. Foliorum segmenta utrinque 3—8, lata.

10. A. grandifolia Friv. **11. A. ligustica** All.

β. Foliorum saltem inferiorum segmenta numerosa, angusta.

× Folia ambitu ovalia vel oblongo-ovalia.

12. A. nobilis L. **13. A. odorata** L.

×× Folia ambitu oblongo- vel lineari-lanceolata, rhachide edentula.

14. A. crithmifolia W. et K. **15. A. fililoba** Freyn. **16. A. setacea** W. et K.

b. Ligulae 4—5, flavae.

α. Foliorum segmenta non transverse imbricata.

× Ligulae involucro duplo vel parum breviores.

17. A. absinthoides Hal. **18. A. holosericea** S. et S.

×× Ligulae minimae, involucro multo breviores.

◯ Folia oblonga, indivisa, serrata.

19. A. ageratum L.

◯◯ Folia pinnatisecta.

. Caules herbacei; corymbus majusculus.

20. A. clypeolata S. et S. **21. A. coarctata** Poir.

.. Caules basi suffruticosi; corymbus parvus.

22. A. aegyptiaca L.

β. Foliorum segmenta transverse imbricata.

23. A. chrysocoma Friv.

c. Ligulae 6—10, albae; foliorum segmenta transverse imbricata.

24. A. cretica L.

1. Sectio. *Ptarmica* Tourn. inst. t. 283. — Ligulae 8—10, albae, saepius magnae et involucro aequilongae. Foliorum segmenta non transverse imbricata.

a. Folia subintegerrima vel minute crenata vel subpinnatifida; capitula majuscula, plerumque solitaria, rarissime 2—3 in corymbo laxiusculo.

1. **A. ageratifolia** S. et S. pr. II. p. 191, Fl. gr. IX. p. 68, t. 888; Raul. cret. p. 780 (*Anthemis*); Bois. fl. or. III. p. 275. — *Ptarmica ageratifolia* Nym. consp. p. 364. — *A. ageratifolia subsp. euageratifolia* Heim. monogr. ptarm. p. 21. — Exsicc.: Orph. fl. gr. n. 781; Sint. it. or. a. 1889 n. 1826.

Adpresse cano-tomentosa, caespitosa; caulibus adscendentibus, simplicibus, remote foliatis; foliis anguste linearibus, basi apiceque attenuatis, pectinatim et crebre pinnatifidis, segmentis brevissimis, rhachide brevioribus, saepe ad crenas reductis, foliis summis integris subintegrisve; capitulo majusculo, hemisphaerico, 15—25 mm. diametro; involucri phyllis pallidis, villosulis, oblongis, obtusiusculis, margine anguste fuscescenti-scariosis; ligulis oblongis, tricrenatis, involucro sublongioribus.

In rupestribus regionis mediae mt. Olympus Thessaliae supra Hagios Dionysios (Orph.); indicatur a Sibthorp in Creta, sed ut videtur per errorem. — Jul. Aug. ♃.

2. **A. aizoon** Griseb. Reise Rumel. II. p. 165 (*Ptarmica*). — *Anthemis aizoon* Griseb. sp. II. p. 210. — *Anth. aizoides* Bois. et Orph. in Orph. fl. gr. exs. n. 244. — *Ach. ageratifolia β. aizoon* Bois. fl. or. III. p. 276. — *Ach. ageratifolia subsp. aizoon* Heim. monogr. ptarm. p. 22. — Exsicc.: Orph. l. c.; Heldr. herb. norm. n. 611. —

Differt a praecedente foliis innovationum lineari-lingulatis, apicem versus latioribus, nec attenuatis, minutissime crenulatis vel subintegris.

In rupestribus regionis alpinae mt. Parnassus loco Dakalia (Heldr.) et Mavralitharia (Orph.) dicto. — Jun. Jul. ♃.

b. Folia pinntim partita vel secta, rarissime integra; capitula parva, fere semper numerosa, corymbosa, rarissime pauca vel solitaria.

α. Caulis et folia plus minus tomentosa vel sericea, cana.

× Ligulae adultae involucro aequilongae.

3. **A. Clavennae** L. sp. p. 897; Bald. riv. coll. bot. Alb. 1895 p. 48. — *A. capitata* Willd. tract. achill. p. 12, t. 1. — *A. Clavennae v. capitata* Heim. monogr. ptarm. p. 41; Hal. Beitr. fl. Epir. p. 27; Form. in Ver. Brünn 1896 p. 45. — *Ptarmica Clavennae* DC. pr. VI. p. 21. — Exsicc.: Hal. it. gr. II. a. 1893; Bald. it. alb. epir. III. n. 160.

Adpresse subsericeo-tomentosa, caespitosa; caulibus erectis vel adscendentibus, simplicibus, superne subnudis; foliis inferioribus oblongo-cuneatis, basi attenuatis, pinnatifidis, segmentis oblongis, obtusis, integris, foliis superioribus dentatis vel integris, sessilibus; corymbo simplici, rarius composito; involucri glabriusculi phyllis nigro-marginatis; ligulis oblongis vel obovatis, integris vel obsolete crenatis.

β. **pindicola** Haussk. symb. p. 109. — Caules glabrati, folia juniora tantum subcinerea, adulta glabrata, involucri phylla glaberrima. — Exsicc.: N. v.

γ. **integrifolia** Hal. Beitr. fl. Epir. p. 27; Bald. riv. coll. bot. Alb. 1895 p. 48; Form. in Ver. Brünn 1896 p. 45. — Folia omnia integerrima. — Varietas insignis; occurrit quoque (f. *depauperata*), caule monocephalo. — Exsicc.: Hal. it. gr. II. a. 1893.

In rupestribus regionis alpinae. Epirus: mt. Tsumerka et Peristeri (Hal.); — β. mt. Karava in Pindo (Haussk.); — γ. mt. Peristeri (Hal.). — Jul. Aug. ♃.

3 × 7. **A. Clavennae × Fraasii (A. Kerneri)** Hal. Beitr. fl. Epir. p. 27, t. 2. — Ab *A. Clavennae* tomento copiosiori longiori, foliis pinnaticectis, segmentis linearibus integris vel dentatis; — ab *A. Frasii* foliis minus dissectis corymbo laxo, capitulis majoribus et ligulis longioribus, involucro aequilongis, distincta. — In regione alpina mt. Peristeri in Epiro, ubi caespitem unicum tantum legi (Hal.).

4. **A. umbellata** S. et S. pr. II. p. 192; Schultz in Flora XXV. Beibl. p. 159: Bois. fl. or. II. p. 276: Hal. in z. b. G. 1888 p. 758, Beitr. fl. Achaia p. 24: Heldr. chlor. Parn. p. 20. — Exsicc.: Heldr. herb norm. n. 401. 1348 et 1550: Orph. Fl. gr. n. 44: Dörfl. fl. gr. n. 418.

Molliter albo-tomentosa, opaca, caespitosa: caulibus erectis vel adscendentibus, simplicibus, superne subnudis: foliis inferioribus ovato-spathulatis, petiolatis, pinnatifidis vel partitis, segmentis subspathulato-ovatis oblongisve, obtusis, saepissime integris, foliis superioribus brevius dentatis, sessilibus: corymbo simplici vel composito: involucri hirtuli phyllis fusco-marginatis: ligulis ovatis, integris vel obsolete crenatis. —

A praecedente indumento molliori, non sericeo, foliis inferioribus latioribus brevioribusque, basi non attenuatis, segmentis plerumque subspathulatis, relative multo longioribus specifice distincta. Occurrit quoque interdum caule monocephalo (*A. umbellata v. monocephala* Heldr. in Heim. monogr. ptarm. p. 44). —

β **pauciloba** Heldr. in Heim. l. c. p. 45. — *A. Millii* Heldr. in sched. — Folia calvescentia, segmentis utrinque perpaucis munita. — Exsicc,: Heldr. it. gr. septentr. a. 1879.

γ. **major** Bois. fl. or. III. p. 276. — *A. major* (*grandifolia* × *umbellata*) Heim. l. c. p. 45, t. 3. — Elatior, foliis majoribus, minus tomentosis, virescentibus, segmentis varie incisis integrisve, corymbo composito. — Meo sensu varietas luxurians, umbrosa, subalpina, intermediis cum typo conjuncta, nec hybrida. — Exsicc.: Heldr. reliqu. Orph. a. 1887.

Ad rupes regionis alpinae et subalpinae. Phthiotis: mt. Oeta (Heldr.); Doris: mt. Kiona (Hal.), mt. Parnassus (Heldr.); Euboea: mt. Xerovuni (Fraas), mt. Dirphys (Pichl.); Achaia: mt. Kyllene (Heldr.), Chelmos (Hal.), pr. Megaspilaeon f. *monocephala* (Mill.); Elis: mt. Olenos (Heldr.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.); — β. mt. Korax Aetoliae (Heldr.); mt. Chelmos Achaiae (Orph.); — γ. pr. Hagios Vlasios in mt. Kyllene (Orph.). — Jun. Aug. ♃.

×× Ligulae involucrum dimidium aequantes.

○ Folia inferiora pectinatim-pinnatipartita segmentis integris, superiora integra.

5. **A. Barbeyana** Heldr. et Heim. in Heim. monogr. ptarm. p. 49, t. 2. — *A. Boissieriana* Heldr. it. gr. septentr. a, 1879, non *A. Boissieri* Haussk. in Bois. fl. or. III. p. 272. —

Adpresse sericeo-villosa, nitens, caespitosa; caulibus humilibus, 5—15 cm. altis, erectiusculis simplicibus, superne subnudis; foliis inferioribus anguste-lanceolatis, in petiolum attenuatis, pectinatim pinnatipartitis, segmentis angusto-lanceolatis, acutis, integris, foliis superioribus diminutis, linearibus, integris, sessilibus; corymbo simplici, denso; involucri villosi phyllis nigro-marginatis; ligulis transverse latioribus, 2—3 crenatis. — Species elegans, indumento sericeo-nitido, foliis inferioribus simpliciter pinnatipartitis, superioribus integris insignis.

Ad summum cacumen mt. Korax Aetoliae rarissime (Heldr.). — Jul. Aug. ♃.

○○ Folia inferiora bipinnatisecta, superiora pinnatisecta.

6. **A. ambrosiaca** Bois. et Heldr. diagn. ser. 2, III. p. 19 (*Ptarmica*); Bois. fl. or. III. p. 276. — Icon: Rouy ill. p. 53 t. 59. — Exsicc.: Heldr. herb. n. 2469; Orph. fl. gr. n. 780.

Cano-tomentosa, opaca, caespitosa; caulibus 10—20 cm. altis, adscendentibus, simplicibus vel ramulo auctis, superne subnudis; foliis inferioribus oblongo-lanceolatis, petiolatis, in lacinias breviter oblongo-lineares, obtusas bipinnatisectis, superioribus pinnatisectis, sessilibus;

corymbo simplici, denso; involucri campanulati, crispule hirtuli phyllis fusco-marginatis; ligulis suborbicularibus, tricrenatis.

In rupestribus regionis alpinae mt. Olympus Thessaliae (Heldr.). — Jul. Aug. ♃.

7. **A. Fraasii** Schultz in Flora 1842 I. p. 159; Bois. fl. or. III. p. 277; Hal. in z. b. G. 1888 p. 758, Beitr. fl. Epir. p. 26; Heldr. chlor. Parn. p. 20; Haussk. symb. p. 109; Bald. riv. coll. bot. Alb. 1895 p. 48, 1896 p. 66. — *A. nivea* Sprun. in Flora 1842 II. p. 638. — *Ptarmica Fraasii* Nym. consp. p. 364. — *A. Neumayeri* Heldr. in sched. ex Heim. monogr. ptarm. p. 51. — Exsicc.: Orph. fl. gr. n. 250; Heldr. herb. norm. e. 603 ot 1349; Sint. it. thessal. n. 727; Dörfl. fl. gr. n. 317.

Sericeo-tomentosa, opaca vel subnitens, caespitosa; caulibus 10—40 cm. altis, erectis vel adscendentibus, simplicibus, superne subnudis; foliis inferioribus oblongo- vel lineari-lanceolatis, petiolatis, in lacinias 2—3 fidas et integras obtusas bipinnatisectis, superioribus, pinnatipartitis, sessilibus; corymbo composito, denso; involucri subcylindrici, hirti, phyllis pallide fusco-marginatis; ligulis transverse latioribus, tricrenatis. — Differt a praecedente caulibus robustioribus, indumento subsericeo, foliis angustioribus, capitulis numerosioribus, subcylindricis, in corymbum densiorem compositum congestis.

In saxosis regionis subalpinae et alpinae. Epirus: mt. Micikeli, Olycika (Bald.), Tsumerka, Strungula, Peristeri (Hal.), mt. Plaka (Sint.); Thessalia: mt. Ghavellu, Karava (Haussk.), Phteris (Neumayer), mt. Oeta (Heldr.); Aetolia: mt. Tymphrestus (Fraas), Korax (Leon.); Doris: mt. Kiona (Hal.); mt. Parnassus (Heldr.); Achaia: mt. Kyllene, Chelmos (Orph.); Laconia: mt. Malevo (Leon.); Zante: mt. Daphni (Mazz.); Leucas (Vrioni); Corcyra: ad Fano (Mazz.). — Jun. Jul. ♃.

β. Caulis et folia glabra vel hirtula, viridia.

8. **A. olympica** Heim. monogr. ptarm. p. 33; Bois. fl. or. suppl. p. 296; pro subspec. *A. moschatae* Wulf. — ? *A. moschata* S. et S. pr. II. p. 193. —

Caulibus erectis, simplicibus superne subnudis, cum pedunculis dense glanduliferis; foliis glabris, caulinis petiolatis, oblongo-ellipticis, pectinato-pinnatipartitis, segmentis lineari-oblongis, obtusiusculis, antice breviter attenuatis, inferioribus integris, mediis saepe lacinulis paucis auctis; corymbo dense conferto, oligocephalo; involucri glaberrimi phyllis rufo-brunneo marginatis, ligulis rotundatis, truncatis, involucro duplo brevioribus. — Ex binis speciminibus incompletis tantum nota.

In saxosis, ad nives aeternas mt. Olympus Thessaliae (Heldr.). — Jul. Aug. ♃. N. v.

9. **A. abrotanoides** Vis. fl. dalm. II. p. 81; Bald. riv. coll. bot. Alb. 1896 p. 66. — *Ptarmica abrotanoides* Vis. fl. dalm. t. 10. — Bald. it. alb. a. 4892 n. 142, it. alb. epir. IV. n. 174.

Caulibus erectis, simplicibus, pubescentibus, ad apicem usque decrescenti-foliatis; foliis breviter pubescentibus, oblongis, in lacinias angusto-lineares acutas divergentes bipinnatisectis, inferioribus petiolatis, superioribus sessilibus, pinnatifidis; corymbo composito, laxo; involucri phyllis puberulis, subconcoloribus; ligulis suborbicularibus, tricrenatis, involucrum aequantibus.

In rupestribus regionis subalpinae et alpinae. Epirus: mt. Papignon et Gamila distr. Zagorion ad confines ditionis nostrae, nec non mt. Kiore Acroceraunia (Bald.). — Jun. Aug. ♃.

2. Sectio. *Millefolium* Tourn. inst. t. 283. — Ligulae 4—5, (rarissime numerosiores), albae vel flavae, involucro saepissime dimidio breviores, interdum brevissimae.

a. Ligulae 4—5, albae vel dilute ochroleucae; foliorum segmenta non transverse imbricata.

a. Foliorum segmenta utrinque 3—8, lata.

10. **A. grandifolia** Friv. in Flora 1836 p. 433; Bois. fl. or. II. p. 258, Heldr. in Sitzungsber. acad. Wiss. Berlin 1883 p. 4, chlor. Parn. p. 20; Haussk. symb. p. 110. — *A. pallescens* DC. pr. VII. p. 296 et *A. silvatica v. subvelutina* DC. l. c. p. 295, sec. Bois. l. c. — Cf. Celak. in Sitzungsb. böhm. Ge.s Wiss. 1887 p. 441. — Exsicc.: Orph. fl. gr. n. 339; Heldr. herb. norm. n. 703; Sint. it, thessal. n. 723.

Adpresse pubescens; caule elato, simplici, ad apicem usque foliato; foliis amplis, ambitu late ovato-oblongis, pinnatisectis, segmentis utrinque 4—8, secus rhachidem late alatam integram decurrentibus, cuneato-lanceolatis, dentato-lobatis, dentibus utrinque 2—4 acutis vel obtusiusculis; corymbo composito; involucri ovati hirtuli phyllis pallidis, obtusis, externis ovatis, caeteris oblongis; ligulis albis, involucrum dimidium aequantibus. — Planta graeca a balkanica capitulis majoribus differt.

In silvaticis regionis montanae et subalpinae. Thessalia: in valle Negerli pr. Chaliki (Sint.), ad monasterium Korona in Pindo (Haussk.), pr. Kissos et Makrinitza in mt. Pelion (Heldr.); mt. Parnassus pr. Rachova (Orph.); Achaia: mt Kyllene, Olenos. — Maio, Jul. ♃.

11. **A. ligustica** All. ped. I. p. 181, t. 53; S. et S. pr. II. p. 195, Fl. gr. t. 897; Raul. cret. p. 782; Bois. fl. or. III. p. 258; Heldr. fl. cephal. p. 44; Hal. in z. b. G. 1888 p. 758, Beitr. fl. Achaia p. 24; Haussk. symb. p. 110; Bald. in bull. herb. Bois. 1896 p. 205. — *A. magna* S. et S. pr. II. p. 194, Fl. gr. t. 896; Fraas. fl. class. p. 215; non L., quae probabiliter ad formas *A. millefolii* spectat. — *A. silvatica* Ten. syll. p. 442. — Exsicc.: Orph. fl. gr. n. 750 et 1141; Heldr. herb. norm. n. 751 et 1551; Bald. it. alb. epir. III. n. 68.

Adpresse hirta, caule stricto, folioso, simplici vel apice ramoso; foliis abbreviatis, ambitu ovatis, pinnatisectis, segmentis utrinque 3—4, secus rhachidem late alatam subintegram decurrentibus, lineari-lanceolatis, pinnatifidis vel paucidentatis, dentibus acutis; corymbo composito; in-

volucri oblongi, hirtuli phyllis pallidis, obtusis; ligulis albis, involucro dimidio brevioribus. — Caule rigido, saepius valde folioso, foliis multo minoribus, capitulis oblongis minoribus et toto habitu a praecedente egregie diversa.

In dumosis regionis montanae et subalpinae. Thessalia: ad monasterium Korona (Heldr.), in oropedio Neuropolis (Haussk.); Aetolia: mt. Tymphrestus, Korax (Fraas); mt. Kiona (Hal.), Parnassus (Sibth.); Attica: mt. Parnes (Heldr.), mt. Dirphys Euboeae (Heldr.); Achaia: pr. Saradi, Megaspilaeon, Planideri (Hal.), ad lacum Pheneon, mt. Kyllene (Heldr.); Arcadia: pr. Zatuna (Orph.); Argolis: pr. Poros (Wiedem.); Laconia: mt. Malevo pr. Castanitza (Orph.), in Alagonia (Zahn), mt. Taygetos pr. Hagios Petros (Sart.), Gaitzies (Psarides); Cyclades: mt. Pittrophoro in insula Andros (Sart.); Creta (Sibth.): ad Enneachoria (Raul.); Cephalonia: pr. Balsamata, mt. Aenos, Gerolaki supra Akoli, mt. Rhudi (Heldr.); Leucas: pr. Eukluvi (Bald.). — Jun. Aug. ♃.

β. Foliorum saltem inferiorum segmenta numerosa, angusta, × Folia ambitu ovalia vel oblongo-ovalia.

12. **A. nobilis** L. sp. p. 899; Ch. et B. exp. p. 253, Fl. pelop. p. 59; Bois. fl. or. III. p. 257; Form. in D. bot. Monat. 1890 p. 17. — Icon: Rchb. germ. t. 133. — Exsicc.: N. v.

Adpresse hirta; caule erecto, folioso, simplici vel superne ramoso; foliis ambitu ovalibus, bipinnatisectis, segmentis remotiusculis, laciniis dentatis, rachide angusta a medio ad apicem folii lobulifera; corymbo composito, denso; involucri ovati, hirtuli phyllis pallidis obtusis; ligulis albis, involucro dimidio brevioribus.

β. **Neilreichii** Kern. in ö. b. Z. 1871 p. 141; Hal. Beitr. fl. thessal. p. 13; pr. sp.; Form. in Ver. Brünn 1897 p. 33. — *A. punctata* Ten. pr. fl. neap. p. 50, non Moench. — Flores ochroleuci. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Hal. it. gr. II. a. 1893.

In dumosis regionis montanae et subalpinae. Thessalia: pr. Velitsena, Kastania (Form.); Eurytania: mt. Tymphrestus (Heldr.); pr. Carithena Arcadiae, mt. Taygetos (Chaub.); — β. Thessalia: pr. Kastania (Hal.), monasterium Korona (Heldr.) et pr. Neochorion in mt. Agrapha (Form.); Aetolia: pr. Musinitza in mt. Korax (Heldr.). — Jun. Jul. ♃.

13. **A. odorata** L. sp. ed. 2 p. 1268; S. et S. pr. II. p. 195; Haussk. symb. p. 110 *v. virescens* Fenzl. in Tchih. as. min. II. p. 264 et *v. lacmonica* Haussk. l. c. (f. patule dense villosa). — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Differt a praecedente foliorum segmentorum laciniis integris vel unidentatis, rachide edentula vel subedentula; species ambae formis intermediis conjunctae esse videntur.

In dumosis regionis montanae et subalpinae. In montibus Graeciae (Sibth.); Thessalia: pr. Chaliki, Said Pascha (Form.), mt. Zygos supra Metzovo (Heldr.), pr. Malakasi, Uranaeos, Tsungeri, Meteora, Neuropolis, Korona, Karditza (Haussk.). — Jun. Jul. ♃.

XX Folia ambitu oblongo-vel lineari-lanceolata, rhachide edentula.

14. **A. crithmifolia** W. et K. pl. rar. hung. I. p. 68 t. 66; Haussk. symb. p. 110; Form. in Ver. Brünn 1896 p. 44, 1897 p. 33. — Exsicc.: Orph. fl. gr. n. 749 (Macedonia); Heldr. it. thessal. n. 7; Sint. et Bornm. it. turc. n. 1296, 1297 et 1825; Sint. it. thessal. n. 919.

Villoso-pubescens; caule erecto, folioso, simplici vel ramoso; foliis difformibus, rosularum late oblongo-lanceolatis, bipinnatisectis, segmentis approximatis laciniis fere setaceo-linearibus, caulinis lanceolatis saepe semel pinnatisectis, segmentis abbreviatis, in lobos breviter lineari-oblongos partitis; corymbo composito, fastigiato; involucri ovato-oblongi phyllis pallidis, obtusis; ligulis albidis, involucro dimidio brevioribus.

In dumosis regionis montanae et subalpinae. Thessalia: pr. Malakasi (Sint.), Kalabaka (Haussk.), Vlachava, Phlamburo in mt. Oxya (Form.), pr. Hagios Dionysios in Olympo (Sint.), mt. Pelion (Heldr.). — Jun. Aug. ♃.

15. **A. fililoba** Freyn. in Flora 1881 p. 210; Form. in D. bot. Monat. 1896 p. 17, in Ver. Brünn 1897 p. 33; Haussk. symb. p. 110. — *A. setacea v. filifolia* Bois. fl. or. suppl. p. 295. — Exsicc.: Heldr. it. gr. septentr. a. 1879, it. thessal. a. 1885.

Villosa; caule adscendente, ad apicem usque foliato, simplici vel ramoso; foliis conformibus, oblongo-lanceolatis, bipinnatisectis, segmentis tenuissimis, elongatis, remotiusculis, laciniis fere setaceo-linearibus; corymbo composito, fastigiato; involucri ovato-oblongi phyllis pallidis, obtusis; ligulis albidis, involucro dimidio brevioribus. — A praecedente foliis, omnibus in lacinias tenuissimas bipinnatisectis, capitulis minoribus; a sequente foliis ambitu latioribus, segmentis paucioribus laxioribus discedit.

In dumosis, silvaticis regionis montanae et abietinae. Thessalia: pr. Chaliki, Klinovo in Pindo (Haussk.), pr. Mavrika et Katasara in mt. Othrys (Form.), mt. Pelion, mt. Korax Aetoliae (Heldr.). — Jul. Aug. ♃.

16. **A. setacea** W. et K. pl. rar. hung. I. p. 82, t. 80; Bois. fl. or. III. p. 256; Hal. in z. b. G. 1888 p. 758, Beitr. fl. Epir. p. 27; Heldr. chlor. Parn. p. 20; Form. in D. bot. Monat. 1890 p. 17, in Ver. Brünn 1897 p. 33; Haussk. symb. p. 110. — *A. millefolium v. setacea* Koch. syn. p. 373; Bald. riv. coll. bot. Alb. 1896 p. 65. — *A. dolopica* Freyn. et Sint. in bull. herb. Bois. V. p. 625 (f. magis lanata). — Huc spectant probabiliter: *A. millefolium* Form. in D. bot. Monat. 1890 p. 17 et *A. pannonica* Form. l. c. et in Ver. Brünn 1896 p. 44. — Exsicc.: Orph. fl. gr. n. 43 et 1142; Sint. it. thessal. n. 1061 et 1062; Heldr. herb. norm. n. 1552 a et b. —

Plus minus lanato-hirsuta; caule erecto, simplici, foliato; foliis conformibus anguste lineari-lanceolatis, bipinnatisectis, segmentis tenuibus, abbreviatis, approximatis, numerosissimis, laciniis confertis, saepe imbri-

catis; corymbo composito, fastigiato; involucri ovato-oblongi phyllis pallidis, obtusis; ligulis albidis, involucro dimidio brevioribus.

In collibus apricis, dumosis regionis inferioris in subalpinam adscendens. Epirus: pr. Dobra (Bald.), Vulgarelion, Mazuki (Hal.); frequens in Thessalia tota; mt. Tymphrestus (Nied.), Kiona (Hal.), Parnassus (Heldr.); Achaia: mt. Kyllene (Heldr.); pr. Zatuna Arcadiae, mt. Malevo (Orph.), mt. Xirovuni et in faucibus Langada mt. Taygetos (Zahn) Laconiae. — Maio, Aug. ♃.

b. Ligulae 4—5, flavae.

α. Foliorum segmenta non transverse imbricata.

× Ligulae involucro duplo vel parum breviores.

17. **A. absinthoides** Hal. Beitr. fl. Epir. p. 27, t. 2; Bald. riv. coll. bot. Alb. 1895 p. 47. — Exsicc.: Hal. it. gr. II. a. 1893; Bald. it. alb. epir. III. n. 163.

Breviter et adpresse sericeo-hirta; caule erecto, simplici; foliis turionum radicalibusque longe petiolatis, ambitu ovatis, pinnatipartitis, segmentis infimis integris, reliquis plerumque trifidis, laciniis anguste linearibus, foliis caulinis sursum decrescentibus, superioribus sessilibus; corymbo composito, denso; involucri ovati phyllis oblongis obtusis, margine ciliatis; ligulis late obovatis, trilobis, involucrum subaequantibus. — Species valde aromatica, distinctissima, cum nulla alia comparanda.

In rupestribus regionis mediae et superioris mt. Tsumerka supra Vulgarelion loco Spathes dicto (Hal.), nec non in ejusdem montis jugo Strungula et Bresani (Bald.). — Jun. Jul. ♃.

18. **A. holosericea** S. et S. pr. II. p. 194; Bois. fl. or. III. p. 259; Hal. in z. b. G. 1888 p. 758; Beitr. fl. Epir. p. 27, in ö. b. Z. 1895 p. 409; Heldr. chlor. Parn. p. 20; Haussk. symb. p. 111; Bald. riv. coll. bot. Alb. 1895 p. 47, 1896 p. 66; Form. in Ver. Brünn 1897 p. 33. — *A. flabelliformis* S. et S. fl. gr. IX. p. 72, t. 894. — ? *A. tomentosa* Fraas. fl. class. p. 215, non L. — Exsicc.: Orph. fl. gr. n. 251, herb. n. 3715; Heldr. herb. norm. n. 302; Sint. et Bornm. it. turc. n. 1298; Sint. it. thessal. n. 724; Bald. it. alb. 1892 p. 143, it. alb. epir. IV. n. 342; Dörfl. fl. gr. n. 194.

Sericeo-villosa; caule erecto, simplici; foliis turionum radicalibusque petiolatis, ambitu obovatis oblongisve, lyratis, segmentis ovatis, inciso-dentatis, ab inferioribus parvis distinctis ad superiora confluentia auctis, foliis caulinis sursum descrescentibus, lanceolatis pinnatipartitis; corymbo simplici vel composito, denso; involucri ovati phyllis oblongis, obtusis, villosulis; ligulis rotundatis, trilobis, involucro duplo subtriplo brevioribus. — Species pulcherrima, nulla cum alia comparanda.

In rupestribus regionis mediae et superioris. Epirus: in mt. Acroceraunii, mt. Gamila, Nimercka, Kuruna, Micikeli, Murga (Bald.), Peristeri, Strungula, Tsumerka (Hal.); Thessalia: mt. Plaka pr. Chaliki (Sint.), mt. Zygos (Heldr.), Karava, Ghavellu, Gionscala (Haussk.), mt. Olympus (Orph.), mt. Pirgo, Mavrika et Katasara in mt. Othrys (Form.); Aetolia: mt. Tymphrestus (Sprun.), Korax (Reiser); Doris: mt. Kiona

(Hal.), mt. Parnassus (Sibth.); Euboea: mt. Dirphys et Xerovuni (Heldr.); Attica: mt. Pateras, Parnes (Heldr.), Pentelicon (Sprun.); Elis: mt. Olenos (Heldr.); Achaia: mt. Kyllene (Orph.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.). — Maio, Aug. ♃

×× Ligulae minimae, involucro multo breviores.

○ Folia oblonga, indivisa, serrata.

19. **A. ageratum** L. sp. p. 897; Ch. et B. fl. pelop. p. 59. — Icon: Rchb. germ. t. 122.

Glabra vel puberula, graveolens; caule erecto, folioso, simplici vel ramoso; foliis oblongis, radicalibus petiolatis, caulinis sessilibus, fasciculatis; corymbo composito, denso; involucri ovati phyllis oblongis, obtusiusculis.

Messenia: pr. Kalamata (Chaub.). — Species pro Graecia dubia, a recentioribus non lecta. — Jun. Sept. ♃

○○ Folia pinnatisecta.

. Caules herbacei; corymbus majusculus.

20. **A. clypeolata** S. et S. pr. II. p. 193, Fl. gr. IX. p. 71, t. 863; Haussk. symb. 111; Bald. riv. col. bot. Alb. 1895 p. 47, 1896 p. 66. — Exsicc.: Bald. it. alb. epir. III. n. 66, IV. n. 173.

Adpresse cano-tomentosa; caule erecto, simplici; foliis ambitu lineari-lanceolatis, pinnatisectis, inferioribus petiolatis, caulinis decrescentibus, subsessilibus, segmentis ellipticis, acute inciso-pauciserratis, secus rhachidem integram cuneato-decurrentibus, foliorum superiorum subintegerrimis; corymbo composito, denso; involucri ovati, lanati phyllis oblongis, obtusiusculis; ligulis orbiculatis, involucri 5 plo brevioribus.

In rupestribus regionis montanae et subalpinae. Epirus: mt. Kuruna distr. Lyaskovik, mt. Micikeli et ad Syraku distr. Janina (Bald.); Thessalia: mt. Ghavellu pr. Sermeniko (Heldr.). — Jun. Jul. ♃

21. **A. coarctata** Poir. suppl. I. p. 94; Form. in D. bot. Monat. 1890 p. 17, in Ver. Brünn 1896 p. 45, 1897 p. 33; Hal. Beitr. fl. Thessal. p. 13. — *A. compacta* Willd. sp. III. p. 2206; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 4; Haussk. symb. p. 111; non Lam. dict. I. p. 27, quae ad formas *A. magnae* L. spectat. — *A. glomerata* M. B. taur. cauc. III. p. 585. — *A. velutina* Desf. cat. h. par. ed. 3 p. 180. — *A. sericea* Janka in Linnaea 1860 p. 579; Form. in D. bot. Monat. 1890 p. 18. — Exsicc.: Orph. fl. gr. n. 751 (Macedonia); Heldr. it. thessal. n. 46; Sint. it. thessal. n. 477.

Dense villosa; caule erecto, simplici, rarius ramoso; foliis ambitu lineari-lanceolatis, pinnatisectis, inferioribus petiolatis, caulinis sessilibus, segmentis numerosissimis, oblongis, in lacinias lineares obtusas pinnatipartitis, rhachide tenui, hinc inde dentifera; corymbo composito, compacto; involucri turbinati, tomentosi phyllis oblongis, obtusis; ligulis minimis. — Differt a praecedente indumento villoso, foliis angustioribus, segmentis numerosioribus minoribus pinnatipartitis et ligulis exiguis.

In collibus apricis, dumosis regionis inferioris et montanae.

Thessalia: pr. Malakasi, monasterium Korona, Kalabaka (Haussk.), Han Kriavris, Trikala, Asproklisia, Vlachava, Konisko, Kastri, Kucuro, Agia Paraskevi, Phlambures, mt. Mtrica in mt. Chassia, pr. Tyrnovo, Patsios, Tafilvris, Baraktari, Miluna et Rapsani in Olympo, pr. Kapurna, Portaria (Form.) et Volo in mt. Pelion (Heldr.). — Jun. Aug. ♃

13 × 21. **A. odorata × coarctata.** — *A. compacta × odorata v. virescens* (*A. tymphaea*) Haussk. symb. p. 111. — Ab *A. odorata* caule subvilloso, foliis angustioribus, oblongis, cinereo-hirtis, segmentis abbreviatis, corymbo compacto, involucro dense et molliter hirto et ligulis sulfureis; ab *A. coarctata* foliis brevioribus latioribusque, cinereo-hirtis, segmentis angustioribus, longius dentatis, corymbo minus compacto et ligulis pallidioribus, discedens.

Inter parentes in herbidis inter Kalabaka et Tsungeri Thessaliae (Haussk.). — N. v.

.. Caules basi suffruticosi; corymbus parvus.

22. **A. aegyptiaca** L. sp. p. 900; S. et S. pr. II. p. 193 p. p.; Hal. in ö. b. Z. 1895 p. 410. — *A. Tournefortii* DC. pr. VI. p. 28; excl. pl. Laconiae; Bois. fl. or. III. p. 260. — Icon: Tourn. voy. p. 87. — Exsicc.: Dörfl. fl. aeg. n. 119.

Dense cano-tomentosa, saepe pannosa; caulibus adscendentibus, simplicibus, superne subnudis; foliis ambitu linearibus, pinnatisectis, inferioribus petiolatis, caulinis decrescentibus, subsessilibus, segmentis ovato-cuneatis, lobato-crenatis vel inciso-serratis; corymbo subcomposito, denso; involucri ovati, lanati, phyllis oblongis, obtusiusculis; ligulis minimis. — *A. clypeolatae* affinis, sed omnibus partibus minor.

β. **taygetea** Bois. et Heldr. diagn. XI. p. 16, Fl. or. III. p. 261; Hal. in ö. b. Z. 1895 p. 411; pro sp. — *A. aegyptiaca* S. et S. pr. II. p. 193 p. p.; Ch. et B. exp. p. 253, Fl. pelop. p. 59. — *A. Tournefortii* DC. pr. VI. p. 28, quoad pl. laconicam. — Caulibus saepius elatioribus, foliorum indumento saepius tenuiore, segmentis saepius magis incisis. — Aliquam differentiam graviorem inter plantam insularem et peloponnesiacam detegere nequeo. — Exsicc.: Orph. fl. gr. n. 1082; Heldr. herb. norm. n. 1442.

In rupestribus regionis inferioris. Cycladum insula: Stenosa (Tourn.), Makaria (Reiser), Caloyero (Sibth.); — β. Laconia (Sibth.): in regione alpina mt. Malevo, mt. Taygetos l. d. Kakochioni et Megala Zonaria (Heldr.). — Jun. Jul. ♄

β. Foliorum segmenta transverse imbricata.

23. **A. chrysocoma** Friv. in Flora 1835 p. 336; Bois. fl. or. III. p. 265; Haussk. symb. p. 111; Form. in Ver. Brünn 1896 p. 45. — ? *A. tomentosa* Ch. et B. exp. p. 253, Fl pelop. p. 59. — Exsicc.: Heldr. it. thessal, IV. a. 1885; Sint. it. thessal. n. 917.

Sericeo-villosa; caulibus erectis vel adscendentibus, simplicibus, foliosis; foliis ambitu lanceolato-linearibus, pinnatisectis, segmentis per-

multis, confertis, in lacinias minimas, lineari-setaceas, mucronatas partitis, in foliis superioribus subintegris; corymbo composito, denso; involucri ovato-campanulati, hirsuti phyllis ovatis, obtusis, margine fusco-scariosis; ligulis trilobis, involucro duplo brevioribus. — Subspecies austro-orientalis *A. tomentosae* L., a qua differt indumento densiori, foliorum laciniis tenuioribus, involucri phyllis fuscomarginatis, saepe fimbriatis, capitulis submajoribus et ligulis intensius coloratis.

In herbidis regionis subalpinae. Thessalia: mt. Zygos (Heldr.). Dokimi (Form.); nec non in Arcadia: mt. Diaforti, Gortys, si *A. tomentosa* Ch. et B. re vera huc spectet. — Jun. Aug. ♃

c. Ligulae 6—10, albae; foliorum segmenta transverse imbricata.

24. **A. cretica** L. sp. p. 899: Sieb. avis p. 5, rem. p. 6; Weiss in z. b. G. 1869 p. 40; Raul. cret. p. 782; Bois. fl. or. III. p. 269. — *A. santolina* S. et S. fl. gr. IX. p. 70, t. 891, non L., quae ligulis minutissimis luteis a nostra discedit. — Exsicc.: Heldr. pl. cret. n. 1343.

Basi suffruticosa; caulibus erectis, simplicibus, foliosis, adpresse tomentosis; foliis linearibus, ad rhachidem canescentibus, pinnatisectis, segmentis minutis, plus minus arcte imbricatis, glabrescentibus, in lobos ovatos, concavos, margine callosos tripartitis; corymbo composito, laxiusculo; involucri lanatuli phyllis ovatis, obtusis; ligulis truncato-trilobis, involucro subbrevioribus. — Species ligulis numerosis magnis transitum ad *Ptarmicas* praebet.

In rupibus regionis inferioris Cretae: ad promontorium Maleka (Sieb.), in peninsula Akroteri pr. Canea (Weiss), Suda, promontorium Spada in fauce Katholiko (Raul.). — Mart. Jun. ♄

Obs. 1. Species incerta: *A. pubescens* S. et S. pr. II. p. 194, Fl. gr. t. 895 (e mt. Parnasso), non L., quae ex Bois. fl. or. III. p. 260 *Pyrethrum achilleaefolium* vel speciem affinem sistit. — Secundum auctores nonnulus = *A. micrantha* M. B., qua specie autem icon Sibthorpii minime quadrat, quae insuper in mt. Parnasso vix crescit.

Obs. 2. *Santolina chamaecyparissus* L. sp. p. 842, a Dallap. prosp. p. 111 in Cephalonia et *S. rosmarinifolia* L. l. c., a Raulin cret. p. 782 in Creta, probabiliter ex confusione indicantur.

22. Anthemis L. gen. n. 645.

Dispositio specierum:

1. Sectio. *Cota* Gay. — Achenia tetragono-compressa, ad angulos marginales acuta; paleae complicatae, flores et achenia amplexantes. Receptaculum in nostris hemisphaericum.

a. Perennes.

1. A. tinctoria L.

b. Annuae.

α. Pedunculi fructiferi valde incrassati.

2. A cota L.

β. Pedunculi fructiferi vix incrassati.

3. A. brachycentros Gay. **4. A. Brachmanni** Bois. et Heldr.

2. Sectio. *Euanthemis* Bois. — Achenia obconica vel turbinata, saepe angulata, nunquam compressa; paleae planiusculae vel carinatae, flores et achenia non amplexantes. Ligulae (cum adsunt) in nostris albae.

a. Perennes.

α. Foliorum laciniae obtusae vel acutae, nunquam mucronatae.

× Receptaculum convexum.

5. A. montana L. **6. A. pindicola** Heldr.

×× Receptaculum elongato-conicum.

7. A. panachaica Hal.

β. Foliorum laciniae mucronatae vel setaceo-apiculatae.

8. A. meteorica Haussk. **9. A. pectinata** (Ch. et B.).

b. Annuae.

α. Achenia calva vel margine brevi obsita.

× Pedunculi fructiferi incurvi, incrassati; capitula discoidea.

10. A. cretica (L.)

×× Pedunculi fructiferi recti; capitula radiata, rarissime (*A. Muenteriana*) discoidea.

○ Ligulae fertiles.

. Virentes vel plus minus pubescentes; foliorum laciniae lineares, acutae, mucronatae.

11. A. arvensis L. **12. A. ruthenica** M. a. B. **13. A. parnesia** Bois. et Heldr.

.. Lanato-tomentosa; foliorum laciniae ovatae, obtusae.

14. A. Muenteriana Heldr.

○○ Ligulae steriles.

15. A. tomentosa L. **16. A. peregrina** L. **17. A. Guicciardii** Heldr. et Sart.

β. Achenia saltem exteriora auricula eis aequilonga vel subaequilonga superata.

× Receptaculum conico-cylindricum.

18. A. auriculata Bois.

×× Receptaculum hemisphaericum vel ovatum.

19 .A. chia L.

3. Sectio. *Maruta* Cass. — Achenia tuberculata, calva (in nostra teretiusculo-turbinata); paleae subulatae. Ligulae albae.

20. A. cotula L.

1. Sectio. *Cota* Gay in Guss. fl. sic. syn. II. p. 866. — Achenia tetragono-compressa, ad angulos marginales acuta; paleae complicatae, flores et achenia amplexantes. Receptaculum in nostris hemisphaericum.

a. Perennes.

1. **A. tinctoria** L. sp. p. 896; S. et S. pr. II. p. 192; Ch. et B. fl. pelop. p. 59; Clem. sert. p. 62; Form. in D. bot Monat. 1890 p. 16, in Ver. Brünn 1896 p. 43; 1897 p. 32; Haussk. symb. p. 106 f. *brachyglossa*; Hal. Beitr. fl. Epir. p. 26, Beitr. fl. Aetol. p. 7,

Beitr. fl. Achaia p. 24. — *Cota tinctoria* Gay l. c. p. 867. — *A. brachyglossa* C. Koch in Linn. XXIV. p. 318, f. ligulis abbreviatis. — Icon: Fl. dan. t. 741. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Caulibus erectis vel adscendentibus, simplicibus vel ramosis, superne longe nudis; foliis adpresse hirtis, pinnatisectis, segmentis oblongis inciso-serratis, dentibus breviter cuspidatis; involucri glabriusculi phyllis pallidis, externis lanceolatis, acutiusculis, internis oblongo-linearibus, apice obtuso scariosis; ligulis luteis, discum subaequantibus; paleis lanceolatis, in mucronem discum aequantem abrupte attenuatis; acheniis utraque facie pluristriatis, corona brevi scariosa subintegra.

β. **discoidea** Vahl. symb. I. p. 74; Bois. fl. or. III. p. 281; Heldr. chlor. Parn. p. 21; Haussk. symb. p. 106; Hal. Beitr. fl. Thessal. p. 13. — *A. discoidea* Willd. sp. III. p. 2188; S. et S. pr. II. p. 191, Fl. gr. IX. p. 69 t. 890. — Ligulae nullae. — Exsicc.: Orph. fl. gr. n. 785; Sint. it. thessal. n. 914; Dörfl. fl. gr. n. 407.

γ. **pallida** DC. pr. VI. p. 11; Bois. fl. or. III. p. 281; Haussk. symb. p. 106; Hal. Beitr. fl. Achaia p) 24; Form. in Ver. Brünn 1896 p. 43, 1897 p. 32, in D. bot. Mon. 1898 p. 78. — *A. Triumfetti* Bald. riv. coll. bot Alb. 1895 p. 48, 1896 p. 66, an *A. tinctoria v. Triumfetti* L. sp. p. 896, quae sec. DC. pr. VI. p. 11 varietatem *A. austriacae* Jacq. sistit. — Huc forsan: *A. pyrethrum* Fraas class. p. 215. — Ligulae albae. — Exsicc.: Orph. fl. gr. n. 782; Heldr. it. Thessal. n. 6; Sint. it. thessal. n. 915; Dörfl. fl. gr. n. 277.

δ. **parnassica** Bois. et Heldr. diagn. ser. 2, III. p. 20 pro sp. (*Cota*); Bois. fl. or. III. p. 281; Haussk. symb. p. 106; Form. in Ver. Brünn 1897 p. 32. — Huc refert sec. Bois. l. c.: A. *maritima* S. et S. pr. II. p. 188, Fl. gr. IX. p. 64, t. 882; Marg. et R. fl. Zante p. 57; an etiam Pieri corc. fl. p. 121 et Ung. Reise p. 123. — Adpresse canescens; involucri phylla fusco-marginata; ligulae albae, elongatae. — Intermediis ad *γ.* manifeste transit. — Exsicc.: Heldr. herb. norm. n. 210; Orph. fl. gr. n. 332.

In herbidis, rupestribus regionis inferioris in alpinam usque adscendens. Epirus: pr. Kalentini (Hal.); Thessalia: mt. Said Pascha, pr. Malakasi, Kastania, Han Kuraneos, mt. Agrapha in Pindo, mt. Chassia, pr. Spilia et Ambelakia in mt. Ossa, mt. Pelion, pr. Volo (Form.), mt. Olympus (Heldr.); Acarnania: pr. Kravassaras (Hal.); Achaia: pr. Patras (Hal.); Messenia: inter Pyrgos et Katakolo (Chaub.); Corcyra (Clem.); — *β.* Thessalia: pr. Kastania (Hal.), Malakasi, Kalabaka, Sermeniko, Korona (Haussk.); Eurytania: pr. Laspi (Samar.); mt. Parnassus (Sibth.); Attica: mt. Pateras (Heldr.); Achaia: pr. Kalavryta (Leon.), mt. Kyllene (Heldr.); Arcadia: pr. Tripolis (Sart.); Laconia: mt. Malevo pr. Kastanitza, mt. Taygetos (Heldr.); — *γ.* Epirus: mt. Murga, Micikeli (Bald.), pr. Chaliki (Sint.), Said Pascha, Malakasi (Form.), mt. Zygos, mt. Olympus, Pelion, pr. Aivali (Heldr.), Velestinos, Konisko, Stylida (Form.); Aetolia: mt. Korax (Leon.); Attica: mt. Citheron, Parnes, Pentelicon, mt. Dirphys Euboeae (Heldr.); Achaia: mt. Omplo

pr. Patras (Heldr.), pr. Lopesi (Hal.); — δ. cum formis intermediis: Thessalia: mt. Zygos, Karava, Ghavellu, Korona, Orman Magula, Aivali, Pharsalus (Haussk.), mt. Olympus (Orph.); mt. Parnassus (Heldr.) f. typica *Cotae parnassicae*; Aetolia: mt. Arapocephalo pr. Prusso (Samar.); Zante (Sibth.); Corcyra (Pieri): pr. Palaeokastro (Ung.). — Maio, Aug. ♃

b. Annuae.

α. Pedunculi fructiferi valde incrassati.

2. **A. cota** L. sp. p. 893; Ch. et B. exp. p. 252, Fl. pelop. p. 59; Haussk. symb. p. 106; Form. in Ver. Brünn 1897 p. 33. — *A. altissima* L. sp. p. 893; S. et S. pr. II. p. 188; Ch. et B. l. c.; Weiss in z. b. G. 1869 p. 39; Raul. cret. p. 780; Bois. fl. or. III. p. 282; Bald. viagg. Creta p. 62. — Nomen *A. cota* jure anterioritatis *A. altissimae* anteferendum est. — *Cota altissima* Gay in Guss. fl. sic. syn. II. p. 867. — Icon: Rchb. germ. t. 117. — Exsiss.: Orph. fl. gr. n. 783; Reverch. pl. cret. n. 71; Sint. it. thessal. n. 469.

Glabriuscula vel parce puberula, caule erecto, patule ramoso; foliis 2—3 pinnatipartitis, lacinulis lanceolatis, mucronato-spinulosis; capitulis magnis; involucri phyllis ab externis lanceolatis ad intima oblonga magis scariosa sensim auctis; paleis subspathulatis, abrupte in cuspidem rigidam aequilongam, flosculos disci superantem contractis; ligulis albis, disco aequilongis; acheniis utraque facie tenuiter 10-costatis, ad angulos marginales anguste marginatis, coronula brevissima superatis.

In cultis, vinetis, ruderatis regionis inferioris. Thessalia: ad Guwelzi pr. Kalabaka (Sint.), pr. Trikala, Kerasia Sina (Form.), Pharsalus, Volo (Haussk.); Euboea: pr. Kastaniotissa (Heldr.); pr. Athenas (Sart.); Achaia: pr. Trikala (Heldr.); Argolis: pr. Nauplia (Haussk.), Tirynthum (Orph.); Messenia: pr. Arcadia (Chaub.); Creta: pr. Rumata distr. Kissamos (Bald.), Canea (Weiss), Messara, Koenuriokorio, Sabres, mt. Aphendi Kavutsi (Raul.). — Maio, Jul. ⊙.

β. Pedunculi fructiferi vix incrassati.

3. **A. brachycentros** Gay in Koch syn. ed. 2. p. 414; Bois. fl. or. III. p. 284; Haussk. symb. p. 106; Bald. riv. coll. bot. Alb. 1896 p. 66; Form. in Ver. Brünn 1897 p. 33. — *A. cota* Koch syn. p. 376, non L. — *A. pseudocota* Vis. fl. dalm. II. p. 78, t. 50. — *Cota brachycentros* Gay in Guss. fl. sic. syn. II. p. 867. — Exsicc.: Baen. pl. corc. a. 1896.

Glabriuscula vel parce puberula; caule erecto, patule ramoso; foliis bipinnatipartitis, lacinulis tenuiter linearibus, mucronato-spinulosis; capitulis mediocribus; involucri phyllis ab externis lanceolatis ad intima oblonga magis scariosa sensim auctis; paleis oblongo-lanceolatis, in cuspidem rigidam, brevem, flosculos disci subaequantem sensim attenuatis; ligulis albis, disco aequilongis; acheniis utraque facie tenuiter 5—7 costatis, ad angulos marginales argute marginatis, coronula brevissima superatis. — Praecedenti similis, differt pedunculis vix incrassatis, capitulis minoribus et praesertim paleis sensim, nec abrupte, in cuspidem breviorem attenuatis, nec non acheniis facie 5—7 costatis.

In arvis, herbidis regionis inferioris. Epirus: mt. Cuka distr. Janina (Bald.); Thessalia: Mavrika et Katasara in mt. Othrys (Form.); Arcadia: pr. Andrizena (Heldr.); Messenia: pr. Methone (Bois.); Corcyra (Haussk.): pr. Palaeokastrizza (Baen.). — Maio, Jul. ⊙.

4. **A. Brachmanni** Bois. et Heldr. diagn. VI. p. 84, Fl. or. III. p. 285; Hal. Beitr. fl. Achaia p. 24. — *Cota Brachmanni* Bois. et Heldr. diagn. ser. 2, III. p. 21. — Exsicc: Heldr. pl. gr. n. 179.

Differt a praecedente paleis abrupte in cuspidem disco breviorem contractis, acheniis laevibus, parum compressis, tetragonis. Ultima nota transitum ad sectionem sequentem praebet. — *A. cota* differt ab ea pedunculis fructiferis incrassatis, capitulis majoribus, paleis in cuspidem elongatam contractis et acheniis evidenter compressis, 10 costatis.

In rupestribus regionis montanae et subalpinae. Achaia: mt. Panachaicon (Hal.), pr. Megaspilaeon (Heldr.); Arcadia: pr. Karithena (Sart.); mt. Ithome Messeniae, pr. Androvista in mt. Taygetos Laconiae (Heldr.), — Maio, Jun. ⊙.

2. Sectio. *Euanthemis* Bois. fl. or. III. p. 278. — Achenia obconica vel turbinata, saepe angulata, nunquam compressa; paleae planiusculae vel carinatae, flores et achenia non amplexantes. Ligulae (cum adsunt) in nostris albae.

a. Perennes.

α. Foliorum laciniae obtusae vel acutae, nunquam mucronatae.

× Receptaculum convexum.

5. **A. montana** L. sp. ed. 2. p. 1261; Bois. fl. or. III. p. 291. — *A. saxatilis* DC. syn. fl. fr. p. 291. — *A. pontica* Willd. sp. III. p. 2184; ? Sieb. avis p. 5 et Raul. cret. p. 780. — *A. montana v. Linnaeana* Gr. et Godr. fl. fr. II. p. 155; Haussk. symb. p. 106. cum f. *thracica* Griseb. spic. II. p. 209 (foliis demum glabratis, segmentis angustioribus, involucri phyllis pallidis et f. *macedonica* Griseb. l. c. (foliis incanis, segmentis latioribus, involucri phyllis fusco-marginatis); Hal. Beitr. fl. Achaia p. 24 p. p.; Bald. riv. coll. bot. Alb. 1895 p. 48, 1896 p. 67; Form. in Ver. Brünn 1895 p. 33. — *A. Gerardiana* Jord. obs. VII. p. 31. — *A. montana v. saxatilis* Heldr. fl. Aegina p. 303. — Huc pertinere videtur: *A. carpathica* Haussk. symb. p. 106, non W. et K., quae foliis glabriusculis, radicalium segmentis linearibus obtusis, dente obtuso uno alterove donatis, caulinorum segmentis integerrimis obtusis, capitulis majoribus et involucri phyllis late nigro-marginatis ab *A. montana* discedit et in Graecia vix crescit. —

Adpresse-pubescens vel sericea; caulibus adscendentibus, simplicibus, raro parce ramosis, superne longe nudis; foliis pinnatisectis, segmentis integris vel 2—3 fidis vel pinnatisectis, laciniisque linearibus; capitulis mediocribus; involucri phyllis exterioribus ovato-lanceolatis acutis, interioribus lanceolatis obtusis; paleis oblongo-lanceolatis, in mucronem brevem attenuatis, exterioribus saepe tridentatis vel sub mucrone excisis;

acheniis obtuse tetragonis, corona brevi superatis. — Stirps valde polymorpha.

a. Capitula radiata.

α. **typica.** — Adpresse pubescens, virescens; caulibus 10—30 cm., altis, saepissime simplicibus; foliis pinnatisectis, segmentis integris vel 2—3 fidis, latiusculis, obtusis. — Exsicc.: Heldr. it. thessal. IV. a. 1885. —

β. **olenaea.** — Adpresse pubescens, virescens; caulibus elatis, usque 40 cm. altis, infra medium in 2—3 ramos elongatos partitis; foliis saltem inferioribus bipinnatisectis, laciniis valde angustis, acutis. — Formis intermediis ad var. sequentem transire videtur. — Exsicc.: Heldr. herb. n. 2082.

γ. **cronia** Bois. et Heldr. diagn. ser. 2 III. p. 22 pro sp.; Bois fl. or. III. p. 291. — Adpresse sericea, canescens; caulibus saepius humilibus, sed interdum usque 30 cm. quoque altis, saepissime simplicibus; foliis pinnatisectis, segmentis 2—3fidis vel integris, laciniisque nunc angustatis acutiusculis, nunc brevibus, latiusculis, interdum (*f. breviloba*) brevissimis 2—3 mm. tantum longis (Heldr. herb. norm. n. 103, specimina a. 1876 lecta). — Exsicc: Heldr. herb. norm. n. 1250; Bald. it. alb. epir. III. n. 63, IV. n. 374; Dörfl. fl. gr. n. 305 (*f. angustisecta*).

b. Capitula discoidea.

δ. **pentelica** Bois. fl. or. III. p. 292; Hal. in z. b. G. 1888 p. 758; Haussk. symb. p. 106. — *A. chrysocephala* Heldr. herb. norm. n. 405 et Orph. fl. gr. n. 280, non Bois. — Adpresse-sericea, canescens; caulibus 10—30 cm. altis, saepius simplicibus; foliorum laciniis nunc angustis (*f. gracilior*, Heldr.), nunc latioribus, ut in var. γ., cujus formam discoideam probabiliter sistit; paleae apice erosulae; acheniorum corona saepius longior, dimidiata. — Exsicc.: Heldr. herb. norm. n. 405 et 1037, herb. dimorph. n. 32; Orph. fl. gr. n. 280.

ε. **tenuiloba** DC. pr. VI. p. 15 pro sp. (*Lyonnetia*); Bois. fl. or. III. p. 291. — *Anacyclus orientalis* L. sp. p. 892, sec. Bois. l. c. — Adpresse-sericea, canescens, gracilior; foliorum laciniis tenuibus, acutiusculis; capitulis minoribus; paleis eximie truncato-retusis; acheniorum margo brevissimus. — Exsicc.: Heldr. it. thessal. n. 5; Bald. it. cret. alt. n. 80.

ζ. **incana** Bois. diagn. ser. 2 III. p. 26; Raul. cret. p. 780; pro sp.; Bois. fl. or. III. p. 292 p. p. — *Santolina alpina* S. et S. pr. II. p. 165, Fl. gr. IX. p. 39 t. 851; Sieb. avis p. 5; non L. — *S. anthemoides* Sieb. in Flora V. 2 p. 639, non L. — ? *S. incana* Ch. et B. fl. pelop. p. 57, non Lam. — Sericeo-tomentosa, cana; caulibus 10—30 cm. altis, simplicibus vel in duos ramos elongatos, superne longe nudos abeuntibus; foliis inferioribus bipinnatisectis, laciniis brevibus, anguste linearibus, obtusiusculis; capitulis mediocribus, phyllis

sericeo-tomentosis; paleis apice erosis; acheniis corona longiuscula aequali superatis. — Intermediis formis conjuncta esse videtur cum var. δ. — Exsicc.: Heldr. herb. norm. n. 1440.

η. **Spruneri** Bois. et Heldr. diagn. ser. 2 III. p. 24; Hal. in ö. b. Z. 1895 p. 409; pro sp. — *A. absinthifolia* Bois. et Sprun. diagn. IV. p. 8, p. p. — *A. montana v. incana* Bois. fl. or. III. p. 292 p. p.; Heldr. chlor. Parn. p. 21; Haussk. symb. p. 106. — Elatior, foliis multo majoribus, laciniis latiuscule lineari-spathulatis, obtusis, caeteris uti ζ. — Exsicc.: Orph. fl. gr. n. 333; Heldr. herb. norm. n. 5 (forma foliis minoribus, laciniis angustioribus transitum ad ζ. praebet).

In rupestribus regionis montanae et alpinae. Thessalia: mt. Ghavellu et Karava in Pindo dolopico (Heldr.); indicatur ceterum: mt. Pelion et Othrys (Form.); Aetolia: mt. Tymphrestus (Samar.); Boeotia; mt. Cithaeron pr. Platea (Bois.); Attica: mt. Parnes (Bois.); Euboea: mt. Dirphys (Sart.); insula Aegina: mt. Oros (Heldr.), sed loca probabiliter ad unam vel alteram varietatem pertinent; — β. mt. Olenos (Heldr.); γ. Epirus: mt. Kuruna, Murga, Micikeli (Bald.); Aetolia: mt. Korax (Leon.); Attica: mt. Parnes (Heldr.); Achaia: mt. Chelmos (Hal.); Laconia: mt. Malevo (Orph.); — δ. Thessalia: mt. Pelion (Heldr.); Attica: mt. Pateras (Bois.), Parnes (Sprun.), Pentelicon (Heldr.); Achaia: mt. Kyllene (Orph.); Creta: mt. Ida (Heldr.); — ε. Thessalia: mt. Pelion (Heldr.); Creta: mt. Ida (Bald.); — ζ. Laconia: mt. Malevo, Taygetos (Heldr.); Creta: mt. Sphaciotici (Sibth.), mt. Ida (Heldr.), indicatur quoque in mt. Pelion (Haussk.), unde tantum var. δ. et ε. vidi, dein pr. Mykenas Argolidis, si *Santolina incana* Ch. et B. huc spectet; — η. Doris: mt. Kiona (Hal.), mt. Parnassus (Orph.). — Apr. Jul. ♃

6. **A. pindicola** Heldr. it. Thessal. IV. a. 1885. — Exsicc.: Heldr. l. c.; ? Sint. it thessal. n. 1167.

Adpresse-pubescens, virescens; caulibus adscendentibus vel erectis, elatis, usque 40 cm. altis, a medio corymbose-ramosis, ramis nudis vel subnudis; foliis bipinnatisectis, laciniis anguste-linearibus, elongatis, acutiusculis; involucri glabriusculi phyllis acutis, exterioribus ovato-lanceolatis, interioribus lanceolatis; paleis oblongo-lanceolatis, in mucronem brevem attenuatis; acheniis obtuse-tetragonis, corona brevi superatis. — A praecedente caulibus corymbose-ramosis, 4—6 cephalis discedit.

In rupestribus regionis superioris Pindi summi mt. Karava (Heldr.) in mt. Gionscala pr. Sermeniko (Sint.). — Jun. Jul. ♃

×× Receptaculum elongato-conicum.

7. **A. panachaica.** *A. montana v. Linnaeana* Hal. Beitr. fl. Achaia p. 24 p. p. — Exsicc.: Hal. it. gr. II. a. 1893.

Adpresse-pubescens, virescens; caulibus adscendentibus, 15—20 cm. altis, simplicibus vel parce ramosis, superne longe nudis; foliis pinnatisectis, segmentis brevibus, breviter 2—3 fidis vel integris; involucri hirtuli phyllis exterioribus ovato-lanceolatis, acutis, interioribus lanceo-

latis, obtusis; paleis oblongo-lanceolatis, in mucronem brevem abruptiuscule attenuatis; ligulis disco aequilongis; acheniis obtuse-tetragonis, corona brevissima superatis. — A praecedentibus receptaculo conico egregie distincta, *A. graveolenti* Bois. foliis tripinnatisectis, involucri phyllis glabris et acheniis exterioribus grosse costato-sulcatis diversae, affinis.

In rupestribus regionis mediae mt. Panachaicon Achaiae (Hal.). — Maio, Jun. ♃

β. Foliorum laciniae mucronatae vel setaceo-apiculatae.

8. **A. meteorica** Haussk. symb. p. 106. — Exsicc.: Haussk. it. gr. a. 1885.

Caulibus adscendentibus, 10—20 cm. altis, a basi ad apicem squarroso-ramosis, denique glabris, ramis elongatis, corymbosis, inferne foliatis; foliis indumento brevissimo griseo-viridibus, rosularum petiolatis, in lacinias divaricato-patentes lineares mucronatas pinnatisectis, caulinis inferioribus sessilibus, sensim diminutis, pinnatis, segmentis simplicibus, valde divaricatis, summis linearibus sensim indivisis; involucri glabri phyllis ab inferioribus lanceolatis acutis, ad suprema ovato-oblonga, late marginata, apice mucronulata sensim auctis; receptaculi conico-hemisphaerici paleis ovato-oblongis, apice abrupte et longiuscule mucronatis; ligulis disco aequilongis; acheniis cylindrico-turbinatis, in marginem brevissimum abeuntibus. — Sequenti affinis, quae foliis rosularum sessilibus, laciniis angustissimis subulato-setaceis et caulibus parce ramosis vel simplicibus discrepat.

In apricis saxosis supra Kalabaka Thessaliae ad monasteria meteora praecipue ad monasterium Hagios Stephanos (Haussk.). — Jun. Jul. ♃

9. **A. pectinata** Ch. et B. exp. p. 251, t. 28, Fl. pelop. p. 59, t. 30 (*Anacyclus*); Bois. et Reut. diagn. IV. p. 6 p. p.; non Bois. flor. III. p. 297, quae = *Cotula complanata* S. et S. pr. II. p. 187, Fl. gr. IX. p. 62 t. 879 et *Anthemis complanata* nominanda. — *A. taygetea* Bois. et Heldr. diagn. VI. p. 83, Fl. or. III. p. 296. — *Lyonnetia pectinata* DC. pr. VII. p. 295 p. p. — Exsicc.: Orph. fl. gr. n. 93 et 786; Heldr. herb. norm. 1548 et 1549.

Ad basin et rhachidem foliorum inferiorum adpresse-pubescens, ceterum glabra; caulibus adscendentibus, simplicibus vel pauciramosis, ramis divaricatis, inferne foliatis; foliis viridibus, sessilibus, pinnatis, segmentis subulato-setaceis mucronatis, simplicibus vel 2—3 fidis, foliis superioribus sensim diminutis, summis setaceis indivisis; involucri glabri phyllis ab inferioribus lanceolatis acutis ad suprema oblongo-linearia, apice obtuso scariosa sensim auctis; receptaculi conico-hemisphaerici paleis carinatis, apice breviter attenuato mucronatis; ligulis disco aequilongis vel nullis; acheniis cylindrico-turbinatis, in marginem brevissimum abeuntibus, exterioribus subquadrangulis, obsolete tuberculatis. — Variat: *α. ligulata* et *β. eligulata.*

In vinetis, pinetis regionis montanae, in alpinam adscendens. Achaia: mt. Kyllene (Orph.); Laconia: mt. Malevo pr. Kastanitza

(Orph.), pr. Hagios Petros ad Sarantapotamos (Sart.); Laconia: pr. Megali Anastasova (Zahn), Anavryti (Heldr.); Messenia: mt. Diaforti (Chaub.). — Maio, Jul. ♃

b. Annuae.

α. Achenia calva vel margine brevi obsita.

× Pedunculi fructiferi incurvi, incrassati; capitula discoidea.

10. **A. cretica** L. sp. p. 892; S. et S. pr. II. p. 188; Urv. enum. p. 113; (*Anacyclus*); Nym. syll. p. 7; Bois. fl. or. III. p. 299; Bald. viagg. Creta p. 62; Forsyth in bull. herb. Bois. III. p. 87; Heldr. fl. Aegina p. 303, in ö. b. Z. 1898 p. 184, chlor. Thera p. 14, chlor. Mykon. p. 245. — *Santolina anthemoides* L. sp. ed. 2, p. 1180; S. et S. pr. II. p. 166, Fl. gr. t. 854. — *S. rigida* S. et S. pr. II. p. 166, Fl. gr. t. 853. — *Cotula abrotanifolia* Willd. sp. III. p. 2167; Sieb. avis p. 5. — *Lyonnetia pusilla* Cass. dict. XXXIV. p. 106. — *L. rigida* DC. pr. VI. p. 14; Raul. cret. p. 781; Weiss in z. b. G. 1869 p. 40. — *L. abrotanifolia* Less. syn. p. 259; Raul. cret. p. 781. — *A. abrotanifolia* Guss. fl. sic. syn. II. p. 490. — *A. Urvillei* Nym. syll. p. 7. — Exsicc.: Orph. fl. gr. n. 291; Heldr. herb. norm. n. 503 et 1039; Reverch. pl. cret. n. 73; Bald. it. cret. n. 72; Dörfl. fl. aeg. n. 24 et 126.

Villoso-pubescens vel glabriuscula; caulibus a collo saepissime pluribus, diffusis, simplicibus vel ramosis; foliis petiolatis, pinnatisectis, segmentis indivisis vel saepius 2—6 fidis, laciniis oblongis, brevibus, acutis vel obtusis; involucri phyllis adpressis, lanceolatis, apice praesertim pilosis; receptaculi ovati paleis apice erosulis, breviter et abrupte mucronatis; acheniis turbinatis, striatis, margine brevissimo superatis.

In arenosis maritimis, locis aridis, herbidis regionis inferioris et montanae usque 1500 m. Boeotia (Sibth.); Euboea: pr. Chalkis (Sart.); insula Angistri pr. Aegina (Heldr.); Cyclades: Syra, Tenos (Weiss), Keos (Heldr.), Mykonos (Sart.), Cythnos, Delos, Rhenia (Tunt.), Naxos (Leon.), Melos, Kimolos, Thera, Mikra Kameni (Urv.), Amorgos (Sibth.); Creta: pr. Kissamos (Rev.), Canea (Weiss), Khalepa, Suda, Lutro, Francocastello, Askyphos, Anopolis (Raul.), Omalos (Bald.). — Mart. Jul. ☉.

×× Pedunculi fructiferi recti; capitula radiata, rarissime (*A. Muenteriana*) discoidea.

○ Ligulae fertiles.

. Virentes vel plus minus pubescentes; foliorum laciniae lineares, acutae, mucronatae.

11. **A. arvensis** L. sp. p. 894; S. et S. pr. II. p. 189; Urv. enum. p. 114; Ch. et B. exp. p. 253, Fl. pelop. p. 59; Bois. fl. or. III. p. 301; Form. in D. bot. Monat. 1890 p. 16, in Ver. Brünn 1897 p. 33; Haussk. symb. p. 107; Hal. Beitr. fl. Epir. p. 26; Bald. riv. coll. bot. Alb. 1895 p. 49. — Icon: Fl. dan. t. 1178. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Bald. it. alb. epir. III. n. 62.

Adpresse-pubescens; caulibus erectis, adscendentibus vel diffusis, ramosis; foliis tripinnatisectis, laciniis lanceolatis, mucronatis; pedunculis elongatis, demum vix aut parum incrassatis; involucri pallidi phyllis ovato-lanceolatis, exterioribus acutis, ceteris obtusis, late scariosis; receptaculi conici paleis lanceolatis, integris, in mucronem discum superantem subabrupte attenuatis; ligulis ellipticis, disco subaequilongis; acheniis turbinato-quadrangulis, sulcatis, exterioribus annulo tumido, saepe radiatim rugoso-plicato, interioribus margine brevi acuto terminatis.

β. **incrassata** Lois. nat. p. 129 pro sp.; Bois. voy. bot. esp. p. 894; Haussk. symb. p. 107. — *A. multicaulis* Bois. et Heldr. in sched. — Pedunculi post anthesin valde incrassati, saepe fere tubiformes, fistulosi. — Exsicc.: Orph. fl. gr. n. 784; Sint. it thessal. n. 405 et 536; Heldr. herb. norm. n. 1441 f. *microcephala*.

In herbidis, cultis regionis inferioris in subalpinam adscendens, varietas typo multo frequentior. Epirus: pr. Philippiada (Bald.), pr. Kalentini, mt. Tsumerka, Strungula, Peristeri (Hal.); Thessalia: mt. Zygos, Karava, Ghavellu, ad monasterium Korona (Haussk.) et Meteora, pr. Kalabaka (Sint.), Trikala, Vlachava, Kastri (Form.); Aetolia: mt. Tymphrestus (Samarit.), Mesolongion (Heldr.); Attica: pr. Athenas, mt. Hymettus (Heldr.); Euboea: pr. Kurbatzi (Wild.); Achaia: pr. Patras (Hal.), mt. Kyllene (Orph.); Argolis: pr. Tirynthum, Nauplia, Astros, pr. Castanitza in mt. Malevo (Orph.); Messenia: pr. Kalamata (Zahn); Cycladum insula: Melos (Urv.). — Apr. Jul. ⊙.

12. **A. ruthenica** M. a B. taur. cauc. II. p. 330; Haussk. symb. p. 107. — *A. Neilreichii* Ortm. in z. b. G. 1852 II. p. 138. — Huc forsan: *A. austriaca* Form. in D. bot. Monat. 1890 p. 16, vix Jacq. —

Adpressiuscule-lanuginosa; caulibus adscendentibus vel diffusis, ramosis; foliis pipinnatisectis, laciniis lanceolatis, mucronatis; pedunculis elongatis, non incrassatis; involucri pallidi phyllis ovato-lanceolatis, exterioribus acutis, ceteris obtusis, late scariosis; receptaculi conici paleis oblongo-linearibus, antice eroso-denticulatis, in mucronem discum aequantem subabrupte attenuatis; ligulis ellipticis, disco subaequilongis; acheniis turbinato-quadrangulis, sulcatis, apice angustissime acute et saepe obsolete marginatis. — A praecedente odore aromatico, paleis eroso-denticulatis et acheniis discedit.

In collibus apricis supra Tyrnovo Thessaliae (Haussk.) nec non pr. Kapurna et Volo, si *A. austriaca* Form. huc pertineat. — Maio, Jul. ⊙. N. v.

13. **A. parnesia** Bois. et Heldr. fl. or. III. p. 305; Haussk. symb. p. 107. — *A. misella* Heldr. et Sart. in Heldr. herb. norm. n. 6.

Adpresse-pubescens; caulibus adscendentibus vel diffusis, simplicibus vel parce ramosis; foliis 1—2 pinnatisectis, laciniis breviter-lanceolatis, mucronatis; pedunculis elongatis, non incrassatis; involucri phyllis margine late fusco-scariosis, exterioribus ovato-lanceolatis acutis, ad interioribus ellipticis sensim auctis; receptaculi conici paleis lineari-lanceolatis,

in mucronem discum subaequantem attenuatis; ligulis ellipticis, disco subaequilongis; acheniis turbinato-quadrangulis, costato-striatis, auricula brevi subfungosa superatis. — A duabus antecedentibus phyllis eximie fusco-marginatis et acheniis breviter subfungoso-auriculatis discedit.

In herbidis, collibus, lapidosis regionis inferioris et abietinae. Aetolia: mt. Aropokephalo (Samarit.), mt. Vardusia (Heldr.); Attica: mt. Parnes (Heldr.); Corinthus, pr. Vromolimni in peninsula Methana et mt. Palamidi pr. Nauplia Argolidis (Haussk.). — Maio, Jul. ⊙.

.. Lanato-tomentosa; foliorum laciniae ovatae, obtusae.

14. **A. Muenteriana** Heldr. in Bois. fl. or. suppl. p. 298. — Exsicc.: Heldr. herb. norm. n. 1038.

Caulibus adscendentibus vel diffusis, simplicibus vel ramosis; foliis pinnatisectis, segmentis breviter cuneatis, pinnati-lobatis vel individis, foliis superioribus sensim diminutis, summis indivisis, cuneatis, apicem versus paucilobatis vel lanceolatis, integris; pedunculis demum incrassatis; involucri pallidi phyllis externis triangularibus acutis, internis apice scariosis obtusis; receptaculi conici paleis lineari-lanceolatis, in mucronem discum subaequantem plus minus abrupte attenuatis; ligulis disco subbrevioribus vel nullis; acheniis subquadrangulis, sulcatis, margine acuto brevissimo coronatis. — Sequentibus habitu valde affinis. — Variat: *α. radiata* et *β discoidea.*

In arenosis maritim,s. Attica: pr. Porto Rhaphti, nec non in Petalium insulis Phuntas, Pontikonisi, Xeonisi et Prasios (Heldr.). — Apr. Jun. ⊙.

○○ Ligulae steriles.

15. **A. tomentosa** L. sp. p. 893; S. et S. pr. II. p. 189; Ch. et B exp. p. 252, Fl. pelop. p. 59; Bois. fl. or. III. p. 309; Spreitz. in z. b. G. 1887 p. 662; Haussk. symb. p. 107; Bald. in nuov. giorn. bot. ital. 1894 p. 97. — Exsicc.: Orph. herb. n. 3652.

Lanato-tomentosa; caulibus decumbentibus vel adscendentibus, crassiusculis, divaricatim-ramosis; foliis parvis, in segmenta pauca obovata vel oblonga, obtusa, integra vel pauciloba pinnatipartitis, summis saepe lanceolatis, integris; pedunculis demum incrassatis; involucri phyllis externis acutis, internis obtusis, apice scariosis; receptaculi ovato-conici paleis oblongis, apice plus minus abrupte attenuatis, mucronatis, praeter nervum fuscum albohyalinis; ligulis disco brevioribus; acheniis subquadrangulis striatis, apice oblique truncatis vel margine angusto unilaterali brevi coronatis.

In arenosis maritimis. Thessalia: pr. Caterina (Orph.); Euboea: in Petalium insula Platera (Holzm.); Attica: pr. Ergastiria, Sunium (Haussk.); ad Corinthum et Lechaeum (Orph.); Argolis (Sibth.); Laconia: pr. Kranidi, Scutari, Marathonisi, Corone, Methone Messeniae (Chaub.); Elis: pr. Lintzi (Heldr.); Cycladum insula: Delos, Naxos (Chaub.); Leucas: pr. Amaxichi (Spreitz.); Corcyra: pr. Pyrgi, Ipso (Bald.); sed loca nonnulla probabiliter ad sequentem spectant, quae enim vix specifice diversa videtur. — Apr. Jul. ⊙

16. **A. peregrina** L. syst. nat. ed. 10 p. 1223; **Marg.** et R. fl. Zante p. 57; Friedr. Reise p. 268; Bois. fl. or. III. p. 308; Heldr. fl. cephal. p. 45; Fl. Aegina p. 303, prosth. chlor. Thera p. 2; Haussk. symb. p. 107. — *A. tomentosa* Hal. Beitr. Fl. Achaia p. 24; Form. in D. bot. Monat. 1898 p. 78. — *A. Muenteriana* Hal. in ö. b. Z. 1897 p. 95, non Heldr. — Exsicc.: Orph. fl. gr. n. 472; Heldr. herb. norm. n. 104; Dörfl. fl. gr. n. 208.

Lanato-tomentosa; caulibus decumbentibus vel adscendentibus, crassiusculis, saepe divaricatim-ramosis; foliis mediocribus, in segmenta lineari-vel oblongo-cuneata, obtusiuscule pinnatilobata vel partita vel in foliis summis indivisa pinnatisectis; pedunculis demum incrassatis; involucri phyllis externis acutis, internis obtusis, apice scariosis; receptaculi breviter conici paleis oblongis, albohyalinis, apice erosulo plus minus abrupte mucronatis; ligulis disco aequilongis vel longioribus; acheniis subquadrangulis, sulcatis, auricula rotundata brevi coronatis. — Valde affinis praecedenti, a qua foliis magis dissectis laciniis minus obtusis, ligulis sublongioribus et acheniis valde sulcatis auriculatis discedit.

β. **heracleotica** Bois. et Heldr. diagn. ser. 2, III. p. 21, Fl. or. III. p. 309; Haussk. symb. p. 107. — Caules abbreviati; pedunculi valde incrassati; foliorum laciniae pauciores, abbreviatae; capitula minora, ligulis abbreviatis. — Exsicc.: Heldr. herb. norm. n. 502.

γ. **platyloba** Haussk. symb. p. 107. — Robustior, caulibus crassioribus, magis ramosis; foliis majoribus, in segmenta oblonga vel ovato-oblonga, integra vel apice 2—3 dentata, obtuse pinnatiloba vel partita, pinnatisectis; capitulis ligulisque majoribus. — Exsicc.: N. v.

In arenosis maritimis, in collibus et campis siccis. Thessalia: ad Lechonia pr. Volo (Heldr.); Euboea: pr. Kurbatzi (Wild.); Attica: pr. Athenas, ad Piraeum, Phalerum, mt. Hymettus (Heldr.), Laurion (Haussk.); insula Aegina (Friedr.), isthmus Corinthiacus (Heldr.); Argolis: peninsula Methana (Haussk.), pr. Astros (Bois.); Achaia: pr. Rhion (Hal.); Zante (Marg.); Cephalonia: pr. Drapano (Heldr.); Cycladum insula Thera (Heldr.); Sporadum insula Skiathos (Leon.). — β. Attica: mt. Lycabettus, pr. Heracleon (Heldr.), Kamariza Laurii (Haussk.). — γ. inter scorias pr. Kamariza Laurii (Haussk.). — Mart. Jun. ⊙

17. **A. Guicciardii** Heldr. et Sart. in Heldr. pl. exs. a. 1855 n. 2238, herb. norm. n. 406, herb. fl. hellen. n. 56, Fl. Aegina p. 303. *A. peregrina v. Guicciardii* Bois. fl. or. III. p. 308; Haussk. symb. p. 107.

Differt a praecedente indumento adpresse-sericeo, caulibus gracilibus saepius simplicibus, foliorum laciniis tenuioribus acutiusculis, pedunculis parum incrassatis et acheniorum exteriorum auricula dimidio achenio saepe longiore. — Notis indicatis medium tenet inter praecedentem et sequentem.

In arvis, campis. Attica: pr. Athenas, Eleusis, Daphni, Heracleon, mt. Hymettus, Bari, insula Lero, Salamis, Aegina (Heldr.); isthmus

Corinthiacus: pr. Kalamoki, Corinthus (Haussk.); Argolis (Orph.); pr. Trikala Achaiae, Tripolis Arcadiae (Heldr.); Cynuria; pr. Hagios Lukas (Orph.). — Apr. Maio. ⊙

β. Achenia saltem exteriora auricula eis aequilonga vel subaequilonga superata.

× Receptaculum conico-cylindricum.

18. **A. auriculata** Bois. diagn. IV. p. 5; Fl. or. III. p. 310; Haussk. symb. p. 109; Heldr. fl. Aegina p. 303. — *A. metallorum* Heldr. Haussk. et Holzm. in Heldr. herb. norm. n. 940; cf. Haussk. symb. p. 109. — Huc probabiliter: *A. melampodina* Ung. Reise p. 123, non Del.

Adpresse griseo-canescens; caulibus erectis vel adscendentibus, ramosis; foliis in lacinias lineares, acutas, integras vel lobatas bipinnatisectis; pedunculis elongatis, non incrassatis; involucri phyllis pallidis, praeter infima brevia acutata, oblongis, obtusis, late scariosis; paleis lineari-lanceolatis, in mucronem discum aequantem attenuatis; ligulis discum aequantibus; acheniis exterioribus brevioribus, turbinatis, costato-sulcatis, in auriculam subfungosam eis vix breviorem abeuntibus, interioribus cylindricis, striatis, auricula breviore hyalina superatis.

In arvis, pinetis regionis inferioris. Euboea: pr. Kastaniotissa (Heldr.); Attica: pr. Rheitro (Ung.), Athenas (Sart.), pr. Kamariza Laurii (Haussk.); insula Aegina: mt. Oros; insula Keos Cycladum (Heldr.). — Apr. Jun. ⊙

×× Receptaculum hemisphaericum vel ovatum.

19. **A. chia** L. sp. p. 894; S. et S. pr. II. p. 189; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 266; Fraas fl. class. p. 215; Clem. sert. p. 62; Ung. Reise p. 123; Weiss in z. b. G. 1869 p. 39; Raul. cret. p. 780; Bois. fl. or. III. p. 311; Spreitz. in z. b. G. 1877 p. 712; Heldr. fl. cephal. p. 45; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. in ö. b. Z. 1895 p. 409, 1897 p. 95; Haussk. symb. p. 109; Bald. riv. coll. bot. alb. 1895 p. 49 *v. conica* (? f. receptaculo conico); Boissieu in bull. soc. bot. Fr. 1896 p. 286. — *A. Visianii* Weiss in z. b. G. 1869 p. 40; Haussk. symb. p. 108; quae ab *A. chia* receptaculo hemisphaerico, paleis medio herbaceis, acheniis pauci (7) costatis, omnibus coronatis, differe dicitur. — Huc pertinet probabiliter ex descriptione: *A. fuscata* Ch. et B. exp. p. 253, Fl. pelop. p. 59; non Brot., quae in Graecia non crescit. — Icon: Fl. Gr. t. 884. — Exsicc.: Orph. herb. n. 3458 (Chios), Fl. gr. n. 473; Heldr. herb. norm. n. 404 et 1251, herb. hellen. n. 92; Reverch. pl. cret. n. 72, in Baen. herb. europ. n. 5658; Sint. it. thessal. n. 86; Baen. herb. europ. n. 9206; Dörfl. pl. cret. n. 107.

Glabriuscula; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis in lacinias breves, oblongas, acutas, saepe 3—5 fidas bipinnatisectis; pedunculis elongatis, non incrassatis; involucri phyllis fusco-marginatis, ab externis triangularibus acutiusculis, ad interna

oblongo-linearia obtusa sensim auctis; paleis oblongo-linearibus, acutiusculis discum subaequantibus; ligulis disco longioribus; acheniis cylindricis, grosse costatis, exterioribus subcurvatis, auricula hyalina saepe aequilonga, interioribus auricula brevi vel corona aequali brevi acuta, superatis.

In cultis, vineis, rupestribus regionis inferioris et montanae frequens. — Mart. Maio. ⊙

3. Sectio. *Maruta* Cass. dict. XXIX. p. 174. — Achenia tuberculata, calvà (in nostra teretiusculo-turbinata); paleae subulatae. Ligulae albae.

20. **A. cotula** L. sp. p. 894; Ch. et B. fl. pelop. p. 59; Bois. fl. or. III. p. 315; Heldr. fl. cephal. p. 45; Form. in D. bot. Mon. 1890 p. 16, in Ver. Brünn 1897 p. 33; Hal. Beitr. fl. Epir. p. 26; Haussk. symb. p. 108. — *Maruta cotula* DC. pr. VI. p. 13; Friedr. Reise p. 273. — *A. psorosperma* Ten. viagg. Abruzz. p. 128 (f. acheniis glandulosis). — *Maruta cotula v. psorosperma* Raul. cret. p. 781. — Icon: Fl. dan. t. 1179. — Exsicc.: Orph. fl. gr. n. 33; Reverch. pl. cret. n. 200, in Baen. herb. europ. n. 5659; Sint. et Bornm. it. turc. n. 1290; Sint. it. thessal. n. 497.

Glabriuscula, foetida; caulibus erectis, corymbose ramosis; foliis in lacinulas anguste lineares 2—3 pinnatisectis, pedunculis non incrassatis; involucri phyllis obtusis, anguste scariosis; receptaculo elongato-conico, parte inferiori paleis destituto; ligulis sterilibus; acheniis pallidis.

In cultis, arvis, ruderatis regionis inferioris et montanae. Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.); Thessalia: frequens a Pindo, per planitiem usque ad mare Aegaeum; Acarnania: pr. Voniza (Lakon.); Aetolia: pr. Mesolongion (Nied.), Karpenisi (Samar.); Attica: pr. Tatoi (Holzm.); Elis: pr. Lintzi (Heldr.), Olympia (Nied.); Achaia: pr. Trikala in mt. Kyllene (Orph.); Messenia: pr. Kalamata (Chaub.); Creta: pr. Canea, Askyphos, mt. Kedros, pr. Voriza ad pedem mt. Ida (Raul.); Cycladum insula: Keos (Heldr.), Andros (Sart.); insula Jonicae (Mazz.). — Mai, Sept. ⊙

Obs. *A. nobilis* L. sp. p. 894. — Erronee indicatur in Corcyra (Pieri corc. fl. p. 122) et in Cephalonia (Dallap. prosp. p. 117).

23. Ormenis Cass. dict. XXXVI. p. 355.

1. **O. mixta** L. sp. p. 894; Ch. et B. exp. p. 252, Fl. pelop. p. 59 (*Anthemis*); DC. pr. VI. p. 18; Raul. cret. p. 781; Bois. fl. or. III. p. 320; Bald. riv. coll. bot. Alb. 1895 p. 49. — *Anthemis cota* S. et S. pr. II. p. 188, Fl. gr. t. 880, non L. — Exsicc.: Heldr. pl. fl. hellen. a. 1899.

Pubescens; caule erecto saepe a basi ramoso, ramis interdum diffusis; foliis oblongis vel linearibus, inferioribus petiolatis, 1—2 pinnatipartitis, superioribus pectinatim pinnatifidis, laciniis brevibus mucronatis; pedunculis demum incrassatis; involucri phyllis oblongis, obtusis, margine scariosis; receptaculi elongato-conici paleis lineari-lanceolatis, complicatis,

achenium involventibus; ligulis albis, basi flavescentibus; acheniis cuneato-obovatis, subcompressis, disco epigyno saepe obliquo. — Facie *Anthemidis*, a quo corollae et achenii structura differt.

In arenosis maritimis, herbidis, ad vias, rare. Epirus: ad pagum Kanali distr. Prevesa (Bald.); Elis: pr. Manolada, Lechaena, Kunupeli (Heldr.); Messenia: pr. Methone (Chaub.); Creta: pr. Suda, Acroteri, Malaxa, Askyphos, Anopolis (Raul.). — Maio, Oct. ⊙

24. **Anacyclus** Pers. syn. II. p. 464.

1. **A. clavatus** Desf. fl. atl. II. p. 287 (*Anthemis*); Pers. syn II. p. 465; Bois. fl. or. III. p. 321; Haussk. symb. p. 108. — *A tomentosus* DC. fl. fr. V. p. 481. — Huc probabiliter: *Anthemis pedunculata* Ch. et B. exp. p. 252, Fl. pelop. p. 59 (= *Anacyclus Candollei* Nym. syll. p. 8), non Desf. — Icon: Rchb. t. 108, fr. 2. — Exsicc.: Heldr. herb. norm. n. 207 et 1252.

Villosulus; caule erecto, divaricatim ramoso; foliis bipinnatisectis, laciniis linearibus mucronulatis; pedunculis demum incrassatis; involucri phyllis ovato-lanceolatis, margine anguste scariosis; receptaculi breviter conici paleis cuneato-trapezoidiis, apice brevissime mucronatis, piliferis glabrisve; ligulis oblongis, albis; acheniis cuneiformibus, exterioribus alatis, alis latis, in auriculam acutam productis. — Facie *Anthemidis*, a qua lobis corollarum disci duobus ceteris longioribus statim diagnoscendus.

In aridis, arenosis maritimis, submontosis. Boeotia (Sprun.); Attica: pr. Villia ad radices mt. Cithaeron, pr. Marathon, in oliveto Athenarum, pr. Phalerum (Heldr.); Corinthus, Nauplia (Haussk.); Achaia: pr. Pellene in mt. Kyllene (Heldr.), Patras (Chaub.). — Apr. Jul. ⊙

2. **A. radiatus** Lois. fl. gall. p. 583; Raul. cret. p. 780; Bois. fl. or. III. p. 321. — *Anthemis valentina* L. sp. p. 895, non *Anacyclus valentinus* L. sp. p. 892.; Ch. et B. fl. pelop. p. 59. — Icon: Rchb. t. 108, f. 3.

Differt a praecedente involucri phyllis apice appendice scariosa lacera munitis; paleis glabris; ligulis ellipticis, luteis; acheniis exterioribus latissime alatis, alis in auriculam longam erectam eroso-denticulatam productis.

In arenosis, herbidis regionis inferioris. Messenia: in maritimis pr. Nisi (Chaub.); Creta (Raul.); a recentioribus non lectum. — Apr. Maio. ⊙ N. v.

Obs. *A. pyrethrum* L. sp. p. 895 (*Anthemis*); DC. fl. fr. suppl. p. 480; Raul. cret. p. 781. — A praecedentibus radice perenni, ligulis subtus purpureis, corollarum lobis disci aequalibus discedens, indicatur a Linnaeo in Creta, sed a recentioribus non lectum.

25. Matricaria L. gen. n. 967.

1. **M. chamomilla** L. sp. p. 891; S. et S. pr. II. p. 187; Dallap. prosp. p. 116; Ch. et B. exp. p. 251, Fl. pelop. p. 59; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 265; Fraas fl. class. p. 215; Clem. sert. p. 63; Weiss in z. b. G. 1869 p. 41; Raul. cret. p. 782; Bois. fl. or. III. p. 323; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 45; Form. in D. bot. Monat. 1891 p. 17, in Ver. Brünn 1895 p. 26; Hal. Beitr. fl. Achaia p. 24. — *M. suaveolens* L. fl. suec. p. 297; Ch. et B. exp. p. 251, Fl. pelop. p. 59. — *Chamomilla officinalis* C. Koch in Linnaeaa XVII. p. 45. — *M. chamaemelon* Heldr. fl. Aegina p. 303, chlor. Thera p. 14. — Icon: Fl. dan. t. 1764. — Exsicc.: Reverch. pl. cret. n. 74; Sint. it. thessal. n. 406.

Glabra, aromatica; caule erecto vel diffuso, ramoso; foliis in lacinias setaceas bipinnatisectis; involucri phyllis oblongis, late scariosis; receptaculo elongato-conico, ligulis albis.

α. **typica.** Acheniis margine minimo prominulo coronatis.

β. **pappulosa** Marg. et Reut. fl. Zante errat. et addend. p. 96. — *M. coronata* Gay ex Koch. syn. ed. 2 p. 416. — *M. chamomilla v. coronata* Bois. fl. or. III. p. 324; Haussk. symb. p. 112. — *Chamomilla meridionalis* C. Koch in Linnaea XVII. p. 45. — Acheniis omnibus vel ligularum corona longiuscula lobata instructis.

γ. **eradiata** Rupr. fl. ingr. p. 594; Baen. herb. europ. n. 9288. Ligulis nullis vel rudimentariis.

In ruderatis, cultis, inter segetes regionis inferioris, in abietinam adscendens, per totam Graeciam, *β.* typo ut videtur vulgatior, *γ.* in Corcyra: pr. Kastrades, Scripero (Baen.). — Non indicatur in insulis maris Aegaei. — Mart. Jul. ⊙

Obs. Quid sit *M. montana* Pieri corc. fl. p. 120, nescio.

26. Chamaemelum Vis. fl. dalm. II. p. 84.

1. **C. tenuifolium** Kit. in Schult. oest. Flora II. p. 498 (*Chrysanthemum*). — *Chrysanthemum trichophyllum* Bois. diagn. IV. p. 10. — *Pyrethrum trichophyllum* Griseb. spic. II. p. 201. — *Matricaria trichophylla* Bois. diagn. VI. p. 88; Haussk. symb. p. 111 p. p. — *Chamaemelum uniglandulosum* Vis. fl. dalm. II. p. 85, t. 51. — *Chamaemelum trichophyllum* Bois. diagn. XI. p. 21, Fl. or. III. p. 330. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Glabrum; caule erecto, elato, in corymbum simplicem vel compositum abeunti; foliis in lacinias capillaceas, elongatas, 2—3 pinnatisectis; capitulis radiatis; involucri pallidi phyllis oblongis, obtusis; scarioso-marginatis, receptaculo hemisphaerico; ligulis albis, diametro disci longioribus; acheniis obconicis, crasse albo-costatis, ad dorsi subrugulosi apicem plus minus distincte 1—2 glandulosis, calvis.

In dumosis regionis montanae. Thessalia: ad monasterium Korona (Heldr.), in oropedio Neuropolis, mt. Ghavella, Karava (Haussk.) — Jun. Jul. ⊙ vel ♃

2. **C. Tempskyaum** Freyn et Sint. in bull. herb. Bois. V. p. 791. — *Matricaria trichophylla v. discoidea* Form. in D. bot. Monat. 1891 p. 17; Haussk. symb. p. 111. — *Chrysanthemum tenuifolium v. discoideum* Hal. Beitr. fl. Thessal. p. 13. — *Chamaemelum trichophyllum v. discoideum* Form. in Ver. Brünn 1896 p. 44. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Sint. it. thessal. n. 744.

Glabrum; caule erecto, elato, in corymbum simplicem abeunti; foliis in lacinias lineari-setaceas, breves, 2—3 pinnatisectis; capitulis saepissime discoideis; involucri pallidi phyllis oblongis, obtusis, scarioso-marginatis; receptaculo hemisphaerico; ligulis albis, disci dimidium aequantibus; acheniis obconicis, crasse albo-costatis, ad dorsi evidenter rugulosi apicem plus minus distincte 1—2 glandulosis, corona brevissima marginiformi superatis. — Praecedenti maxime affine, differt ab eo foliorum laciniis brevioribus, ligulis (cum adsunt) brevibus et praesertim acheniis dorso dense rugulosis, apice corona brevi munitis. — Formam discoideam tantum vidi.

In dumosis montanis. Thessalia: mt. Zygos (Heldr.), Dokimi, Said Pascha (Sint.), Oxya (Hal.) — Jun. Jul. ⊙ vel ♃

3. **C. rosellum** Bois. et Orph. in Bois. fl. or. III. p. 334. — Exsicc.: Orph. fl. gr. n. 778.

Glabriusculum, caule adscendenti, 1—2 cephalo; foliis in lacinias lineari-setaceas longiusculas 1—2 pinnatisectis; capitulis radiatis; involucri phyllis oblongis, obtusis, fuscescenti-marginatis; receptaculo convexo; ligulis pallide roseis, disco subbrevioribus; acheniis (immaturis) apice biglandulosis, corona lobato-partita superatis.

In regione media mt. Malevo Laconiae, rarissime (Orph.). — Maio, Jun. ♃.

Obs. *Plagius ageratifolius* L'Her. in DC. pr. VI. p. 135, — In Creta (tantum sec. P. Alp.) probabiliter erronee indicatur.

27. Leucanthemum Tourn. inst. p. 492.

a. Achenia omnia calva.

1. **L. vulgare** Lam. fl. or. II. p. 137, Bois. fl. or. III. p. 335. — *Chrysanthemum leucanthemum* L. sp. p. 888; S. et S. pr. II. p. 185. — Icon: Fl. dan. t. 994.

Glabrum vel pubescens, caule erecto, 1-oligocephalo; foliis inferioribus petiolatis, obovato-cuneatis, crenatis, ceteris sessilibus, oblongo-lanceolatis vel lanceolato-linearibus, serratis vel pinnatifidis; involucri phyllis externis lanceolatis, margine fuscescenti-scariosis, internis apice dilatatis, late lacero-scariosis; receptaculo convexo; ligulis albis.

In Archipelagi pascuis montosis (Sibth.), a recentioribus non lectum. — Maio, Aug. ♃ N. v.

b. Achenia disci calva, radii corona superata.

2. **L. pallens** Gay in Guill. ann. bot. 1833 II. p. 545 (*Chrysanthemum*); DC. pr. VI. p. 47; Boissieu in bull. soc. bot. Fr. 1896 p. 286. — Icon: Rchb. t. 98.

Plus minus hispidum, caule erecto, 1-oligocephalo; foliis inferioribus petiolatis, obovato-cuneatis, crenatis, ceteris sessilibus, ligulatis, dentatis, summis integris; involucri phyllis ovali-oblongis, margine pallide-scariosis vel apicem versus dilute fuscis; receptaculo convexo; ligulis albis; acheniis radii corona scariosa bipartita munitis.

Creta: pr. Canea, forsan introducta (Boissieu). — Jun. Jul. ♃ N. v.

3. **L. montanum** L. sp. p. 888; S. et S. pr. II. p. 185 (*Chrysanthemum*); DC. pr. VI. p. 48. — Icon: Jacq. obs. IV. t. 91.

Glabrum; caule erecto vel adscendenti, monocephalo; foliis crassiusculis, dentatis, infimis obovato-lanceolatis, in petiolum attenuatis, ceteris sessilibus, oblongo-lanceolatis, summis oblongo-linearibus; involucri phyllis externis ovato-lanceolatis, anguste hyalino-scariosis, internis lanceolatis, apice late lacero-scariosis; acheniis radii corona scariosa, completa, cupuliformi, lacera superatis.

In Archipelagi pascuis montosis (Sibth.); a recentioribus non lectum. — Jun. Sept. ♃ N. v.

28. Pyrethrum Gaertn. fruct. p. 430.

a. Foliorum lacinulae acutiusculae.

P. parthenium L. sp. p. 890; Dallap. prosp. p. 116; Fraas fl. class. p. 214 (*Matricaria*); Sm. fl. brit. II. p. 900; S. et S. pr. II. p. 186; Raul. cret. p. 783; Bois. fl. or. III. p. 344; Heldr. fl. cephal. p. 45; Haussk. symb. p. 111; Form. in Ver. Brünn 1895 p. 26, 1896 p. 44, 1897 p. 33. — *Chrysanthemum parthenium* Pers. syn. II. p. 462; Form. in D. bot. Monat. 1891 p. 17. — Icon: Fl. dan. 674. — Exsicc.: Orph. fl. gr. n. 772; Sint. it. thessal. n. 914.

Puberulum; caule erecto, corymboso-ramoso; foliis petiolatis, pinnatisectis, segmentis elliptico-oblongis, pinnatifidis vel crenatis, superioribus confluentibus; involucri demum umbilicati phyllis oblongis, obtusis, apice scariosis; receptaculo convexo; ligulis obovatis, albis, discum aequantibus; acheniis corona brevissima crenata munitis.

In ruderatis, silvestribus regionis montanae. Epirus: pr. Govrica, in valle Dipotami (Form.); Thessalia: pr. Chaliki (Sint.), Koturi, Vendista, Kastania, Bisula, Sermeniko, mt. Ghavellu (Form.), Zygos, ad monasterium Korona (Haussk.) in Pindo, mt. Pelion pr. Zagora (Aphentulis); Euboea: pr. Steni (Fraas); Achaia: pr. Trikala in mt. Kyllene (Heldr.); Arcadia: ad Platano pr. Tripolis (Sart.), pr. Zatuna (Orph.); Laconia: mt. Malevo pr. Kastanitza (Orph.); Creta: pr. Canea (Raul.); Cephalonia (Dallap.). — Jun, Aug. ♃

b. Foliorum lacinulae mucronatae.

2. **P. corymbosum** L. sp. p. 890 (*Chrysanthemum*); Willd. sp. III. p. 2155; Haussk. symb. p. 111; Form. in Ver. Brünn 1897 p. 33; Bald. riv. coll. bot. Alb. 1896 p. 67. — Icon: Jacq. fl. austr. t. 379. — Exsicc.: Bald. it. alb. epir. IV. n. 123.

Caule erecto, apice corymboso, foliis pinnatisectis, adpressiuscule hirsutis, segmentis oblongo-lanceolatis, pinnatifidis et argute serratis, summis confluentibus, foliis superioribus sessilibus, segmentis lanceolatis, serratis; involucri non umbilicati, hirti, phyllis pallidis, obtusis, interioribus apice scariosis; ligulis oblongo-ellipticis, albis, discum aequantibus; acheniorum corona elongata, apice fissa.

In silvaticis regionis montanae. Epirus: mt. Maria distr. Ljaskovik (Bald.); Thessalia in oropedio Neuropolis (Haussk.), mt. Kerasia Sina in mt. Chassia (Form.); Phocis: mt. Parnassus (Kruper); Laconia: mt. Malevo pr. Kastanitza (Orph.). — Maio, Aug. ♃

3. **P. cinereum** Griseb. spic. II. p. 202. — Exsicc.: Sint. it. thessal. n. 719.

Differt a praecedente, cui proximum, foliis supra canescentibus subtus sericeo-canis, involucri phyllis longioribus acutis. Secundum Velenovsky fl. bulg. p. 269 varietas tantum praecedentis, nam occurrunt formae, quae transitum manifestum ad illum praebent.

Thessalia: in valle Negerli pr. Chaliki (Sint.). — Maio, Aug. ♃

Obs. *P. fuscatum* Willd. sp. III. p. 2156, species Tunetana, erronee indicatur in Graecia a S. et S. pr. II. p. 196.

29. Tanacetum L. gen. n. 944.

1. **T. vulgare** L. sp. p. 844; Dallap. prosp. p. 112; Form. in Ver. Brünn 1896 p. 44, 1897 p. 33. — Icon: Fl. dan. t. 871.

Glabriusculum; caule erecto, apice composite-corymboso; foliis pinnatisectis, segmentis lineari-lanceolatis, elongatis, inciso-serratis vel pinnatifidis; capitulis hemisphaericis; involucri phyllis oblongis, obtusis, apice lacero-scariosis; corollis flavis, radii tubulosis, limbo brevissimo unilaterali.

In dumetis, nemorosis. Thessalia: pr. Mavreli, in valle fluvii Longi, pr. Jerakari, Vlachava, mt. Hagia Paraskevi et Cuka in mt. Chassia, mt. Godaman in Olympo (Form.); Cephalonia (Dallap.), sed sec. Heldr. fl. cephal. p. 45 ex confusione. — Jul. Aug. ♃

Obs. *T. annuum* L. sp. p. 844, a Ch. et B. fl. pelop. p. 57 pr. Philiatra in Peloponneso probabiliter ex confusione indicatur. — *T. balsamita* L. sp. p. 845; Dallap. prosp. p. 112; = *Pyrethrum tanacetum* DC. pr. VI. p. 63; Heldr. Nutzpfl. p. 26. Colitur in hortis.

30. Chrysanthemum L. gen. n. 966.

a. Achenia sensim in coronam amplam vaginaeformem abeuntia.

1. **C. Myconis** L. sp. ad 2 p. 1254; Weiss in z. b. G. 1896 p. 40; Bois. fl. or. III. p. 335; Spreitz. in z. b. G. 1877 p. 713,

1887 p. 662; Heldr. fl. cephal. p. 45; Bald. riv. coll. bot. alb. 1795 p. 49. — *Pyrethrum Myconis* Moench. suppl. p. 287. — *Coleostephus Myconis* Cass. dict. XLI. p. 43. — *Chrysanthemum hybridum* Guss. cat. hort. Bocc. 1821 p. 16 et 70; Gelmi in bull. soc. bot. ital. 1889 p. 450. — *Pyrethrum hybridum* Guss. fl. sic. syn. p. 483. — Huc sec. Guss. fl. sic. syn. p. 483: *Chrysanthemum paludosum* Sieb. avis p. 5, non Desf. = *Leucanthemum setabense* Raul. cret. p. 782, non DC.; nec non sec. Heldr. fl. cephal. p. 45: *Chrysanthemum leucanthemum* Dallap. prosp. p. 115, non L.; et *Chrysanthemum viscidum* Friedr. Reise p. 277. — Icon: Jacq. obs. IV. t. 94. — Exsicc.: Baen. herb. europ. n. 9242.

Glabrum vel pubescens; caulibus erectis vel adscendentibus, simplicibus vel saepius stricte ramosis; foliis serratis, inferioribus petiolatis, obovato-cuneatis, superioribus obovato-oblongis, oblongis linearibusve, basi subaurita semiamplexicaulibus; involucri phyllis lineari-oblongis, apice scariosis; ligulis luteis; acheniis incurvis, radii sterilibus, corona vaginaeformi eis sublongiore, apice lacera superatis, disci corona tubulosa, oblique truncata instructis.

β. **discolor** Guss. fl. sic. syn. II. p. 483 pro var. *Pyrethri hybridi*. — *Pyrethrum Myconis v. albidum* DC. pr. VI. p. 61; Raul. cret. p. 783; Weiss in z. b. G. 1869 p. 40. — Radio albo, basi luteolo. — Exsicc.: Reverch. pl. cret. p. 75.

In pratis, cultis, vineis, ad agrorum margines regionis inferioris. Epirus: pr. Mitikas (Bald.); Corcyra: pr. Potamo (Spreitz.); Cephalonia; pr. Lixuri (Heldr.); insula Poros (Friedr.); Creta: pr. Suda (Weiss); — β. Creta: pr. Kissamos (Rev.), ad fluvium Jordanus pr. Canea (Weiss), pr. Nerokuru, Suda (Raul.). — Apr. Jun. ⊙

b. Achenia ecoronata.

2. **C. segetum** L. sp. p. 889; S. et S. pr. II. p. 185; Ch. et B. fl. pelop. p. 59; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 272; Weiss in z. b. G. 1869 p. 41; Raul. cret. p. 783; Bois. fl. or. III. p. 336; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 45; Fl. Aegina p. 303, prosth. chor. Thera p. 3; Gelmi in bull. soc. bot. ital. 1889 p. 450; Form. in D. bot. Monat. 1890 p. 17; Hal. Beitr. fl. Epir. p. 27; Haussk. symb. p. 111. — Icon: Rchb. germ. t. 95. — Exsicc.: Heldr. pl. fl. hellen. a. 1878 et 1890.

Glabrum; caule erecto, simplici vel ramoso; foliis inferioribus petiolatis, subcuneiformibus, inciso dentatis, caeteris subspathulatis, amplexicauli-auriculatis, apice trifidis, basi dentato-serratis, summis lanceolatis integris; involucri phyllis ovatis, obtusis, apice scariosis; ligulis luteis; acheniis radii compresso-trigonis, dorso plano 3 costatis, facie ventrali obtusangulo 4 costatis, disci subteretibus, circacircum 10 costatis.

In cultis, inter segetes. Epirus: pr. Kalentini (Hal.); Thessalia: pr. Orman Magula (Haussk.), Kapurna, Baba (Form.), Volo (Heldr.); Euboea: pr. Kurbatzi (Wild.), Kastaniotissa (Heldr.), Kumi (Wurlisch); Attica: pr. Stephani, Pikermi, Thoricos (Heldr.), Phaleron, Mesogia,

Laurion (Haussk.); insula Aegina: pr. Janitzio (Heldr.); Argolis: pr. Vromolimni (Friedr.); Elis: ubique in planitie maritima (Heldr.); Messenia: pr. Kalamata, Armyros (Chaub.); Cycladum insula: Mykonos (Sart.), Tenos (Weiss), Naxos (Oertzen), Keos, Thera (Heldr.); Creta: pr. Canea, Suda (Weiss); Zante (Marg.); Cephalonia: pr. Lixuri (Spreitz.), Argostoli, Omala (Heldr.); Corcyra: pr. Gasturi, Benizze (Gelmi). — Apr. Jun. ⊙

3. **C. coronarium** L. sp. p. 890; S. et S. pr. II. p. 186; Ch. et B. exp. p. 251, Fl. pelop. p. 59; Marg. et R. fl. Zante p. 57; Friedr. Reise p. 271; Fraas fl. class. p. 213; Raul. cret. p. 783; Bois. fl. or. III. p. 336; Heldr. fl. cephal. p. 45; Bald. in nuov. giorn. bot. ital. 1894 p. 97. — *Matricaria coronaria* Desr. in Lam. dict. III. p. 737; Pieri corc. fl. p. 120. — *Pinardia coronaria* Less. syn. p. 255; Weiss in z. b. G. 1869 p. 40; Spreitz. in z. b. G. 1887 p. 662; Haussk. symb. p. 111; Form. in Ver. Brünn 1895 p. 26; Heldr. fl. Aegina p. 303; chlor. Thera p. 14; Hal. in ö. b. Z. 1899 p. 25. — Icon: Fl. gr. t. 877. — Exsicc.: Reverch. pl. cret. n. 76; Baen. herb. europ. n. 9305.

Glabrum; caule erecto, ramoso; foliis bipinnatisectis, rhachide lobato-dentata, laciniis lanceolatis; involucri phyllis ovato-oblongis, late scariosis; ligulis luteis vel ochroleucis; acheniis striatis, inter strias glandulosis, radii compresso-triquetris, disci lateraliter compressis.

β. **discolor** Urv. enum. p. 112. — Ligulae albae. — Exsicc.: N. v.

Inter segetes, in cultis frequens. Attica: pr. Athenas, in Acropoli, ad Phalerum (Heldr.), Laurion (Haussk.); insula Xeronisi et Pontikonisi Petalium (Holzm.), Aegina (Friedr.); Corinthus (Haussk.); Elis: ubique in planitie maritima (Heldr.); Arcadia: pr. Zulatika Gortyniae (Nied.); Messenia: pr. Kalamata (Zahn); Cycladum insula: Kythnos (Tunt.), Jos, Keos, Thera (Heldr.), Syra, Tenos (Weiss); Creta: pr. Canea (Tourn. voy. p. 10), Kissamos (Reverch.), Retymo, Candia (Raul.); Strophades (Reiser); Zante (Marg.); Cephalonia: pr. Argostoli, Lixuri, Kontagionata (Heldr.); Corcyra (Pieri): pr. urbem (Baen.), Canali (Form.), Melichia (Spreitz.); — *β.* copiose in insulis Archipelagi (Urv.). — Mart. Jun. ⊙

31. Ammanthus Bois. et Heldr. diagn. XI. p. 18.

1. **A. filicaulis** Bois. et Heldr. l. c. p. 19, Fl. or. III. p. 357; Raul. cret. p. 781. — Icon: Rouy ill. t. 10. — Exsicc.: Heldr. pl. cret. n. 1459.

Adpresse hirtus; caulibus filiformibus, adscendentibus, subsimplicibus, monocephalis; foliis inferioribus petiolatis, in segmenta parva, pauca, oblonga, integra vel lobata pinnatisectis, foliis superioribus subsessilibus diminutis, summis linearibus integris; capitulis radiatis; involucri phyllis oblongis, obtusis, latiuscule scariosis; ligulis albidis, ovatis,

disco brevioribus; acheniis parce glandulosis, corona dimidiata, triplo breviore superatis. — Herba pusilla, facie *Anthemidis.*

In saxosis inter frutices prope promontorium Sidero Cretae orientalis (Heldr.). — Apr. Maio. ⊙

2. **A. maritimus** Bois. et Heldr. l. c. p. 19, Fl. or. III. p. 358; Raul. cret. a. 1846.

Differt a praecedente foliis obtuse 5—7 lobatis; capitulis discoideis; involucri phyllis angustius scariosis, acheniis pellucide glanduloso-tuberculatis. — An varietas discoidea praecedentis?

In arenosis maritimis promontorii Sidero Cretae orientalis (Heldr.). — Apr. Maio. ⊙

32. Artemisia L. gen. n. 945.

1. Sectio. *Absinthum* DC. fl. fr. IV. p. 189. — Capitula heterogama, floribus radii femineis, disci hermaphroditis, omnibus fertilibus; receptaculum pilosum; involucri phylla exteriora tomentella.

a. Petioli exauriculati.

1. **A. arborescens** L. sp. ed 2 p. 1188; S. et S. pr. II. p. 168; Urv. enum. p. 108; Sieb. in Flora I. p. 270; Ch. et B. exp. p. 245, Fl. pelop. p. 57; Marg et R. fl. Zante p. 58; Fraas fl. class. p. 207; Heldr. Nutzpfl. p. 26, Fl. cephal. p. 45, chlor. Thera p. 14; Raul. cret. p. 783; Bois. fl. or. III. p. 372; Spreitz. in z. b. G. 1887 p. 662; Bald. in nuov. giorn. bot. ital. 1894 p. 97; Form. in Ver. Brünn 1895 p. 26. — *A. absinthium* Dallap. prosp. p. 113, non L. — Icon: Fl. gr. t. 856. — Exsicc.: Baen. herb. europ. n. 9209; Dörfl. fl. aeg. n. 45.

Sericeo-argentea, frutescens; caulibus erectis, dense foliosis; foliis inferioribus petiolatis, tripinnatisectis, ceteris 2—1 pinnatisectis, brevius petiolatis sessilibusve, laciniis linearibus, obtusis; capitulis cernuis, demum erectis, pedicellatis, in racemos secundos, laterales, erectos, paniculam amplam angustam formantes, dispositis; involucri extus sericei phyllis oblongis, apice scariosis; acheniis flavo-glandulosis.

In rupestribus maritimis. Epirus: pr. Govrica (Form.); Corcyra: ad Fortozza vecchia in urbe (Baen.), pr. Palaeokastrizza (Spreitz.); Cephalonia: pr. Kalligata et in Sarlata (Heldr.); Zante: pr. urbem (Marg.); frequens in regione maritima Messeniae et Laconiae (Chaub.): pr. Gythion (Pichl.); Creta: pr. Kalyves, Canea (Raul.), in fauce Serisso (Sieb.); Cycladum insula: Neokameni pr. Thera (Urv.), Anaphi (Leon.). — Maio, Aug. ♄

2. **A. absinthium** L. sp. p. 848; Haussk. symb. p. 112; Hal. Beitr. fl. Epir. p. 27, Beitr. fl. Thessal. p. 14; Form. in Ver. Brünn 1895 p. 26, 1896 p. 45, *v. montana,* 1897 p. 34. — Icon: Fl. dan. t. 1654. — Exsicc.: Sint. it. thessal. n. 721.

Sericeo-cana, herbacea; caulibus erectis, foliosis; foliis inferioribus petiolatis, tripinnatisectis, ceteris 2—1 pinnatisectis, brevius petiolatis

sessilibusve, laciniis lineari-oblongis vel linearibus, capitulis cernuis, pedicellatis, in racemos secundos laterales, erecto-patulos, paniculam pyramidato-conicam formantes, dispositis; involucri extus sericei phyllis oblongis, apice scariosis; acheniis glabris.

In dumosis, silvaticis regionis montanae et subalpinae. Epirus: pr. Janina, mt. Micikeli, pr. Govrica (Form.), Vulgarelion (Hal.); Thessalia: pr. Chaliki (Sint.), mt. Oxya (Hal.), Zygos, pr. Sermeniko, Bezula, mt. Karamanoli, ad monasterium Korona (Haussk.), mt. Dokimi, Ghavellu, pr. Neochorion, in valle fluvii Longi in Pindo, ad Godaman in Olympo (Form.). — Jun. Aug. ♃

b. Petioli ad basin auriculati.

3. **A. camphorata** Vill. fl. dauph. III. p. 242; Form. in D. bot. Monat. 1896 p. 18. — Icon: Rchb. t. 142. — Exsicc.: Sint. it. thessal. n. 1278.

Plus minus canescens, suffruticosa; caulibus, adscendentibus erectisve; foliis petiolatis, impresso-punctatis, inferioribus, bi-, superioribus pinnatisectis, summis tripartitis; capitulis cernuis, pedicellatis, in racemos secundos, laterales, erectos, paniculam saepius angustam formantes, dispositis; involucri extus tomentelli phyllis oblongis, apice scariosis; acheniis glabris.

In saxosis regionis inferioris et montanae. Thessalia: pr. Koturi Lepeniza, mt. Dokimi in Pindo, pr. Karditza, Portaria (Form.), Volo (Sint.). — Aug. Oct. ♄

2. Sectio. *Abrotanum* Bess. in nouv. mem. soc. Mosc. II. p. 105. — Capitula heterogama, floribus radii femineis, disci hermaphroditis, omnibus fertilibus; receptaculum glabrum; involucri phylla exteriora tomentella.

4. **A. vulgaris** L. sp. p. 848; S. et S. pr. II. p. 169; Dallap. prosp. p. 113; Bois. fl. or. III. p. 371; Heldr. fl. cephal. p. 45; Form. in D. bot. Mon. 1890 p. 18, in Ver. Brünn 1895 p. 26, 1896 p 46, 1897 p. 34. — Icon: Fl. dan. t. 1176. — Exsicc.: Sint. it. thessal. n. 1202.

Herbacea; caulibus erectis, elatis 60—130 cm. altis, glabris vel superne lanatulis; foliis supra glabris, obscure viridibus, subtus canotomentosis, ambitu ovatis, inferioribus petiolatis, pinnatipartitis, segmentis lanceolatis, integris vel inciso-serratis, superioribus sessilibus, subpalmatopartitis, basi auriculatis, summis lanceolatis, integris; capitulis sessilibus, secus ramulos glomeratis, paniculam longam pyramidatam formantibus; involucri extus tomentelli phyllis oblongis, late scariosis; acheniis glabris.

Ad sepes, in dumosis regionis inferioris et montanae. Epirus: pr. Luros, Han Levka et Hagios Nicolaos pr. Janina, Han Kanberga (Form.); Thessalia: pr. Neochorion, Pezula, Velitsena, Trikala, ad monasterium Korona (Form.), pr. Megala Kalyvia (Heldr.), mt. Olympus (Orph.); Eurytania: pr. Mikrochorio (Heldr.); Cephalonia (Dallap.). — Jul. Sept. ♃

5. **A. petrosa** Baumg. en. Trans. III. p. 90 (*Absinthium*); Fristch in Fl. exsicc, austro-hung. n. 2262. — *A. eriantha* Ten. sem. cat. hort. neap. 1730 p. 14. — *A. Baumgartenii* Bess. tent. abrot. p. 73. — *A. Villarsii* Gr. et Godr. fl. fr. II. p. 130. — *A. spicata β. eriantha* DC. pr. VI. p. 118; Bald. riv. coll. bot. Alb. 1896 p. 67. — Huc probabiliter: *A. spicata* S. et S. pr. II. p. 168, vix Wulf., quae foliis caulinis praesertim pinnatifidis discedit.

Herbacea, caulibus adscendentibus, 5—15 cm. altis, foliisque sericeo-tomentosis; foliis inferioribus longe petiolatis, ambitu oblongo-cuneatis, palmato-multifidis, basi non auriculatis, caulinis mediis digitato-partitis, breviter petiolatis, segmentis linearibus, summis simplicibus indivisis; capitulis breviter pedunculatis, in racemum simplicem dispositis; involucri tomentosi phyllis oblongis, margine spadiceo-scariosis; acheniis glabris.

In rupestribus alpinis summis mt. Smolika supra Kerasovo distr. Konitza in Epiro (Bald.); in montibus Graeciae (Sibth.). — Jul. Aug. ♃ N. v.

Obs. *A. abrotanum* L. sp. p. 845; Dallap. prosp. p. 113; Fraas fl. class. p. 207; Heldr. fl. cephal. p. 45. — Colitur in hortis.

3. Sectio *Dracunculus* Bess. in nouv. mem. soc. Mosc. VIII. p. 16. — Capitula heterogama, floribus radii femineis, fertilibus, disci hermaphroditis, sterilibus; receptaculum et involucri phylla glabra.

6. **A. campestris** L. sp. p. 846; S. et S. pr. II. p. 168; Ch. et B. exp. p. 245, Fl. pelop. p. 57; Bois. fl. or. III. p. 363; Heldr. in Sitzungsb. acad. Berlin 1883 p. 4 et 8; Form. in D. bot. Monat. 1890 p. 18; Haussk. symb. p. 112. — Icon: Fl. dan. 1175. — Exsicc.: Sint. et Bornm. it. turc. n. 1291.

Glabrescens, suffruticosa; caulibus adscendentibus, elatis; foliis inferioribus bipinnatisectis, ceteris pinnatifido-dentatis, laciniis linearibus, mucronatis; capitulis minutis, breviter pedicellatis, in racemos subsecundos, paniculam amplam formantes dispositis; involucri phyllis oblongis, scarioso-marginatis, acheniis glabris.

In saxosis, sterilibus regionis inferioris et submontanae. Thessalia: ad monasterium Meteora (Haussk.), pr. Kalabaka (Form.), Litochoron (Sint.); Peloponnesus (Chaub.), sine loci specialis indicatione. — Jun. Sept. ♄

7. **A. scoparia** W. et K. pl. rar. hung. I. p. 66, t. 65; Haussk. symb. p. 112. — Exsicc.: Heldr. herb. norm. n. 1149 (Thessalonica); it. thessal. n. 64; Sint. et Bornm. it. turc. n. 1292.

Herbacea, praeter folia radicalia, adpresse canescentia glabra; caule erecto, elato; foliis radicalibus petiolatis, in lacinias oblongo-spathulatas, obtusas bipinnatisectis, caulinis subsessilibus, in lacinias setaceas sectis; capitulis minimis, breviter pedicellatis, in racemos subsecundos, paniculam amplam ramosissimam, effusam formantes dispositis; involucri phyllis oblongis, scarioso-marginatis; acheniis glabris. —

Praecedenti interdum persimile, ab ea radice annua, foliis radicalibus rosulatis, capitulis minoribus discedens.

In arenosis, glareosis regionis inferioris. Thessalia: pr. Aivali, Orman-Magula (Haussk.), mt. Pelion pr. Portaria (Heldr.), pr. Litochoron (Sint.). — Aug. Sept. ⊙ et ⊙.

Obs. *A. pontica* L. sp. p. 847 a Sieber in Flora I. p. 271 in fauce Perivoglia Cretae certissime ex confusione indicatur.

6. Tribus. **SENECIONEAE** Cass. in dict. sc. nat. XX. p. 377.

33. Doronicum L. gen. n. 959.

a. Caulis ample paniculatus

1. **D. Orphanidis** Bois. fl. or. III. p. 378; Haussk. symb. p. 105; Form. in Ver. Brünn 1896 p. 47, *v. oligocephalum.* — Exsicc.: Charrel fl. emp. ottom. a. 1888 (Macedonia).

Elatum, inferne subhirtum, superne glandulosum; foliis inferioribus e basi cordata ovato-oblongis, acutiusculis, crenatis, intermediis auriculato-petiolatis, superioribus amplexicaulibus, sessilibus, floralibus lineari-lanceolatis; involucri glandulosi phyllis lanceolatis, acuminatis, radio duplo brevioribus; receptaculo villoso; ligulis flavis basi hirtulis, disco $2^1/_2$plo longioribus; acheniis omnibus hirsutis.

Ad silvarum margines inter jugum Zygos et Oxya (Haussk.) et mt. Dokimi (Form.) in Pindo tymphaeo. — Jul. ♃.

b. Caulis saepissime monocephalus.

2. **D. caucasicum** M. a. B. fl. taur. cauc. II. p. 322; Bois. fl. or. III. p. 380; Spreitz in z. b. G. 1877 p. 712; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 105; Boissieu in bull. bot. Fr. 1896 p. 286. — *D. orientale* Willd. en. hort. berol. p. 898. — *D. eriorrhizon* Guss. in litt. ad DC. a. 1841, ex ipso in Fl. sic. syn. II. p. 482. — *D. pardalianches* S. et S. pr. II. p. 183; Ch. et B. exp. p. 250, Fl. pelop. p. 58; Friedr. Reise p. 280; Fraas fl. class. p. 211; non L. — ? *D. Columnae* Link. in Linnaea IX. p. 583; Ung. Reise p. 124. — Exsicc.: Heldr. herb. norm. n. 122 et 1249; Orph. fl. gr. n. 196; Sint. it. thessal. n. 80.

Puberulum; radice horizontali, nodosa, stolonifera, squamosa, ad collum et squamas lanata; foliis radicalibus, petiolatis, orbiculari-cordatis, repando-dentatis vel obtuse crenatis, caulinis 1—2, cordato-amplexicaulibus, ovatis vel oblongis; involucri parce setulosi phyllis lineari-lanceolatis, acuminatis, radium subaequantibus; receptaculo pubescente; ligulis flavis, basi crispule hirsutis, disco $2^1/_2$plo longioribus; acheniis radii glabris, disci hirsutis.

In umbrosis, silvaticis regionis montanae et subalpinae. Thessalia: in oropedio Neuropolis (Haussk.), pr. Kalabaka (Boissieu), mt. Bulgara in mt. Agrapha, mt. Itamos, in mt. Othrys (Mazaraki), mt. Pelion (Heldr.), ad Thermopylas (Kupassari); mt. Parnassus (Sibth.); Boeotia:

mt. Helicon (Orph.); Euboea: mt. Kandyli (Wild), mt. Dirphys (Orph.); Attica: mt. Cithaeron, Parnes, Pentelicon, Hymettus (Heldr.); Achaia: mt. Olenos (Hal.), pr. Patras (Link), Kyllene, Chelmos (Orph.); Arcadia: pr. Alonistena (Heldr.), Zulatika (Nied.), mt. Diaforti (Friedr.); Messenia: mt. Kupe, pr. Phigalea, Methone (Chaub.); Laconia (Sibth.): mt. Taygetos (Heldr.); Corcyra: mt. Pantocrator (Spreitz.). — Apr. Jun. ♃

3. **D. cordatum** Wulf. in Roem. arch. III. p. 408 (*Arnica*); Schultz in öst. bot. Wochenbl. 1854 p. 411; Hal. in z. b. G. 1888 p. 758; Beitr. fl. Epir. p. 26, in ö. b. Z. 1897 p. 284; Form. in Ver. Brünn 1896 p. 47; Bald. riv. coll. bot. Alb. 1896 p. 67. — *D. Columnae* Ten. fl. nap. pr. p. 49; Haussk. symb. p. 105. — *D. cordifolium* Sternb. in Denkschr. bayer. Ges. Regensb. II. p. 147; Bois. fl. or. III. p. 381; Heldr. chlor. Parn. p. 21. — *Arnica scorpioides* S. et S. pr. II. p. 183, non L. — Icon: Ten. fl. nap. t. 79. — Exsicc.: Heldr. herb. norm. n. 38; Orph. fl. gr. n. 340; Sint. it. thessal. n. 733; Dörfl. fl. gr. n. 184 et 373.

Glabrum; radice obliqua, praemorsa, glaberrima; foliis acutiuscule dentato-lobatis, radicalibus petiolatis, cordato-reniformibus, caulinis saepius pluribus, inferioribus petiolo brevi basi auriculato, superioribus cordato-amplexicaulibus, ovatis; involucri glanduloso-puberuli phyllis lanceolatis, acuminatis, radio subdimidio brevioribus; receptaculo pubescente; ligulis flavis, basi hirtis, disco 2plo longioribus; acheniis radii glabriusculis, disci pubescentibus.

In rupestribus regionis subalpinae et alpinae. Epirus: mt. Olycika (Bald.), Peristeri, Strungula, Tsumerka (Hal.); Thessalia: mt. Karava (Haussk.), Olympus (Heldr.); Aetolia: mt. Korax (Heldr.); Doris: mt Kiona (Hal.); mt. Parnassus (Sibth.). — Jun. Aug. ♃

34. Senecio L.

Dispositio specierum:

a. Frutescentes, albo-tomentosae.
  α. Folia lyrata vel pinnatipartita.
    1. **S. bicolor** (Willd.). 2. **S. taygeteus** Bois. et Heldr.
  β. Folia anguste linearia, longa, integra vel hinc inde lobulo minutissimo aucta.
    3. **S. gnaphalodes** Sieb.
b. Perennes vel biennes.
  α. Involucrum ecalyculatum.
    × Achenia glabra.
      4. **S. Aucheri** DC.
    ×× Achenia pubescentia.
      5. **S. Parnassi** Bois. et Heldr. 6. **S. Heldreichii** Bois.
  β. Involucrum bracteolis (interdum perpaucis) calyculatum.
    × Capitula magna, 2—3 cm. diametro.
      ○ Caulis corymbosus.
      7. **S. euboeus** Bois. et Heldr. 8. **S. macedonicus** Griseb.

○○ Caulis monocephalus.

9. S. arachnoideus Sieb.

×× Capitula parva, ad 1 cm. diametro vel minora.

○ Folia subtus adpresse niveo-tomentosa.

10. S. thapsoides DC.

○○ Folia utrinque viridia.

. Folia indivisa; caulis elatus, corymboso-paniculatus, polycephalus.

11. S. nemorensis L.

.. Folia incisa, pinnatisecta vel lyrata, rarissime indivisa et tunc caulis humilis, oligocephalus.

; Folia oblongo-vel lineari-lanceolata, inferiora runcinata.

12. S. barckhausiaefolius Bois. et Heldr.

;; Folia varie partita vel interdum indivisa, sed nunquam runcinata.

, Folia obovato-vel lanceolato-oblonga, indivisa, acutiuscule dentata.

13. S. fruticulosus S. et S.

,, Folia obovato-oblonga vel oblonga, pinnatifida, segmentis rhachique angulato-dentatis.

14. S. rupestris W. et K.

,,, Folia varie partita, rhachi integra.

— Rhizoma repens; folia omnia pinnatipartita, segmentis lineari-lanceolatis, integris vel dentatis; achenia omnia hirta.

15. S. erucifolius L.

= Radix obliqua, praemorsa; folia lyrata vel lyrato-pinnatifida, infima interdum indivisa, summa pinnatipartita, segmenta oblonga vel oblongo-linearia, latiora; achenia marginis glabra, disci puberula.

16. S. Jacobaea L. **17. S. barbaraefolius** Krock.

c. Annui.

α. Ligularum lamina involucrum valde superans.

18. S. coronopifolius Desf. **19. S. vernalis** W. et K.

β. Ligulae nullae vel brevissimae, involucrum vix superantes, revolutae.

× Achenia glabra.

20. S. viscosus L.

×× Achenia pubescentia.

21. S. lividus L. **22. S. vulgaris** L.

a. Frutescentes, albo-tomentosae.

α. Folia lyrata vel pinnatipartita.

1. **S. bicolor** Willd. sp. III. p. 2085 (*Cineraria*); Bois. fl. or. III. p. 395; Heldr. fl. cephal. p. 46; Spreitz. in z. b. G. 1867 p. 662; Hal. in ö. b. Z. 1897 p. 284. — Huc probabiliter: *Cineraria maritima* Ch. et B. exp. p. 249, Fl. pelop. p. 58, vix L., quae foliis magis dissectis et capitulis subminoribus discedit. — Exsicc.: Heldr. herb. norm. n. 1036.

Caulibus adpresse-tomentosis, simplicibus vel superne patule ramosis, inferne densiuscule, superne remote foliatis; foliis supra glabrescen-

tibus, subtus dense niveo-tomentosis, ambitu oblongis, lyratis, vel pinnatifidis, segmentis integris vel obtuse 2—3 lobis; capitulis hemisphaericis, circa 1 cm. diametro latis, in corymbum simplicem vel compositum dispositis; involucri tomentosi, squamulis paucis brevissimis fulti, phyllis breviter acuminatis; ligulis luteis, disco parum longioribus; acheniis glabris.

In saxosis, vinetis regionis inferioris. Insula Leucas: ad promontorium Zuana (Spreitz.); Cephalonia: pr. Asso, Poros, Kipuria (Heldr.); insula Oxya Echinadum (Reis.); Peloponnesus: pr. Navarin (Sart.), ad promontorium Malea (Chaub.); Cycladum insula: Jos (Heldr.), Melos (Chaub.). — Jun. Aug. ♄

2. **S. taygeteus** Bois. et Heldr. diagn. VI. p. 95, Fl. or. III. p. 394. — *Cineraria taygetea* Nym. syll. p. 2. — Exsicc.: Heldr. pl. fl. hellen. a. 1844 et 1880.

Caulibus adpresse-tomentosis, simplicibus, fere ad apicem usque aequaliter foliatis; foliis supra canis, subtus niveo-tomentosis, ambitu oblongis, lyratis, segmentis integris vel obtuse 2—3 lobatis, terminali multo majore; capitulis hemisphaericis, circa 5 mm. diametro, in corymbum compositum dispositis; involucri glabrescentis vel farinosi phyllis breviter acuminatis, basi elevatim binerviis, squamulis subnullis; ligulis luteis, disco vix longioribus; acheniis glabris. — Indumento tenuiore, caulibus simplicibus, aequaliter foliatis, foliis minus profunde dissectis, capitulis duplo minoribus a praecedente specifice differt.

In regione abietina, rarissime. Euboea: mt. Xerovuni et Dirphys (Heldr.); Laconia: mt. Taygetos (Heldr.). — Jul. Aug. ♄

β. Folia anguste-linearia, longa, integra vel hinc inde lobulo minutissimo aucta.

4. **S. gnaphalodes** Sieb. Reise II. p. 322, t. 10 (*Conyza*); Sieb. avis. p. 5, rem. p. 6; Spreng. syst. III. p. 554; Raul. cret. p. 786; Bois. fl. or. III. p. 397. — *Cineraria gnaphalodes* Nym. consp. p. 351. — Exsicc.: Sieb. pl. cret. a. 1825.

Caulibus lanato-tomentosis, simplicibus; foliis supra glabrescentibus, subtus niveo-tomentosis; capitulis hemisphaeris, circa 5 mm. diametro, in corymbum compositum dispositis; involucri glabri phyllis acutis, squamulis paucis ovatis vel ovato-lanceolatis fultis; ligulis luteis, discum aequantibus, acheniis glabris.

In rupestribus regionis inferioris Cretae: ad Magula in mt. Lassiti, pr. Sitia, in promontorio Sidero (Sieb.); — nuper etiam in insula Karpathos a Pichler lectum. — Jun. Aug. ♄

b. Perennes vel biennes.

α. Involucrum ecalyculatum.

× Achenia glabra.

4. **S. Aucheri** DC. pr. VI. p. 361; Bois. fl. or. III. p. 413. — Nym. consp. p. 351. — Exsicc.: Heldr. herb. n.

Araneoso-lanatus; caule erecto, simplici; foliis radicalibus oblongis, repando-denticulatis, in petiolum eis breviorem attenuatis, caulinis remotis, inferioribus oblongo-lanceolatis, basi sessili attenuatis, superioribus diminutis linearibus; corymbo simplici, oligocephalo; involucro tandem glabrescente; ligulis flavis; pappo corollae tubum aequante. — Mihi ex unico specimine florifero tantum nota; habitu *Cinerariae ovirensi* Koch simillima. Ulteriu observandum, au re vera speciem propriam sistet.

In pascuis alpinis ad nives mt. Olympus Thessaliae (Heldr.). — Jul. Aug. ♃

×× Achenia pubescentia.

5. **S. Parnassi** Bois. et Heldr. fl. or. III. p. 412; Heldr. chlor. Parn. p. 21. — Exsicc.: Heldr. reliqu. Orph. a. 1887; Heldr. herb. norm. n. 1546 potius huc, nec ad *S. Heldreichii* spectare videtur.

Arachnoideo-canus; caule erecto, simplici; foliis integris vel repando-denticulatis, radicalibus oblongis, in petiolum eis breviorem attenuatis, caulinis remotis, inferioribus oblongo-lanceolatis, sessilibus, summis linearisubulatis; corymbo simplici vel parce composito; involucro hirto; ligulis luteis; pappo tubum flosculorum superante. — Facies praecedentis, a quo achenis dense puberulis praesertim differt.

In pascuis regionis alpinae. — Phocis: mt. Parnassus (Heldr.); Euboea: mt. Kandyli (Orph.); Laconia: mt. Taygetos (Heldr.). — Jul. Aug. ♃

6. **S. Heldreichii** Bois. diagn. VI. p. 95, Fl. or. III. p. 396; Heldr. chlor. Parn. p. 21. — Exsicc.: Heldr. pl. gr. n. 385.

Arachnoideo-canus; caule erecto, simplici; foliis integris vel repando-denticulatis, inferioribus oblongis, in petiolum eis aequilongum vel subaequilongum attenuatis, mediis oblongo-lanceolatis, sessilibus, superioribus a basi sessili obtuse auriculata attenuatis, lanceolato-linearibus; corymbo subsimplici; involucro hirto; ligulis luteis; pappo flosculorum tubo superante. — Praecedenti robustior, folia ampliora, superiora auriculata.

In herbidis regionis mediae et superioris. Phocis: mt. Parnassus ad Cagna (Guicc.); Peloponnesus: mt. Olenos (Hal.), mt. Taygetos l. d. Chupata et Megali Zonaria (Heldr.). — Jul. Aug. ♃

β. Involucrum bracteolis (interdum perpaucis) calyculatum.

× Capitula magna, 2—3 cm. diametro.

○ Caulis corymbosus.

7. **S. euboeus** Bois. et Heldr. diagn. ser. 2, III. p. 36, Fl. or. III. p. 405. — Exsicc.: Heldr. pl. gr. a. 1848 et 1895.

Caule erecto, rigido, adpresse araneoso-cano, superne vel a medio sursum corymboso, corymbi ramis saepius simplicibus, elongatis, erecto-patulis; foliis rigidis, adpresse araneosis, demum glabrescentibus, integris vel obsolete repandis, inferioribus oblongo-spathulatis, in petiolum attenuatis, ceteris sessilibus, subsemiamplexicaulibus, oblongis vel lanceolatis; involucri campanulati, adpresse nivei phyllis coriaceis, lanceolatis, acu-

minatis; squamulis 6—8, lanceolatis, involucro tertia parte brevioribus; ligulis croceis, disco duplo longioribus; acheniis exterioribus hirtis, ceteris glabris. — Planta speciosa, sesquipedalis, capitulis magnis adpresse niveis insignis. — Achenia ab auctoribus glaberrima dicuntur, dum in speciminibus meis achenia peripherica semper hirta video.

In rupestribus regionis mt. Dirphys, Xerovuni (Heldr.) et Kandyli (Orph.) Euboeae. — Aug. ♃

8. **S. macedonicus** Griseb. spic. II. p. 221; Bois. fl. or. III. p. 403; Heldr. chlor. Parn. p. 21. — Exsicc.: Heldr. herb. norm. n. 792 et 939.

Caule erecto, glabriusculo, apice subsimpliciter corymboso; foliis subcoriaceis, demum glabratis, inferioribus argute denticulatis, oblongo-lanceolatis, in petiolum attenuatis, ceteris sessilibus, integris, intermediis lanceolatis, summis oblongo-subrhombeis; involucri hemisphaerici, glabriusculi phyllis linearibus, acuminatis; squamulis 10—12, lanceolatis, involucro duplo brevioribus; ligulis intense flavis, disco duplo longioribus; acheniis hirtis.

β. **bracteatus** Bois. et Orph. diagn. ser. 2, III. p. 36 pro sp. — *S. macedonicus β. brachyphyllus* Bois. fl. or. III. p. 404. — Folia minora, etiam inferiora integra, floralia confertiora. — An species propria. — Exsicc.: Orph. fl. gr. n. 100.

In regione abietina. Thessalia: mt. Olympus pr. Hagios Dionysius (Orph.); mt. Parnassus (Guicc.); Peloponnesus: mt. Kyllene, Malevo (Orph.), Taygetos (Heldr.); — *β.* mt. Kyllene (Orph.). — Jun. Jul. ♃

○○ Caulis monocephalus.

9. **S. arachnoideus** Sieb. pl. rar. n. 117 (1812); DC. pr. VI. p. 357. — *S. lanatus* Scop. fl. carn. ed. 2, II. p. 165, non L. — *S. Scopolii* Hoppe et Hornsch. Tageb. Reise Küstenl. p. 34. — *Cineraria arachnoidea* Rchb. fl. exc. p. 242. — Exsicc.: Sint. it. thessal. n. 742.

Floccoso-lanuginosus; caule erecto, monocephalo; foliis inaequaliter serratis denticulatisve, inferioribus ovalibus lanceolatisque, in petiolum attenuatis, superioribus lanceolato-acuminatis, semiamplexicaulibus; involucri hemisphaerici phyllis lineari-acuminatis, squamulis lineari-filiformibus, involucro subbrevioribus; ligulis citrinis; acheniis glabris.

In pratis subalpinis. Epirus: mt. Tsumerka (Topali); Thessalia: pr. Mandra Hodza in Pindo tymphaeo (Sint.). — Maio, Jun. ♃

×× Capitula parva, ad 1 cm. diametro vel minora.

○ Folia subtus adpresse niveo-tomentosa.

10. **S. thapsoides** DC. pr. VII. p. 301; Bois. fl. or. III. p. 396; Hal. in z. b. G. 1888 p. 758, Beitr. fl. Epir. p. 26, Beitr. fl. thessal. p. 13, in ö. b. Z. 1897 p. 284; Heldr. chlor. Parn. p. 21; Form. in D. bot. Monat. 1890 p. 18, Bald. riv. coll. bot. Alb. 1895 p. 49. — *Cacalia verbascifolia* S. et S. pr. II. p. 164; Ch. et B. exp. p. 245; Fl. pelop. p. 57; Fraas fl. class. p. 210; non *S. verbascifolius* L. — *Cineraria thapsoides* Nym. sylt. p. 2

Icon: Fl. gr. t. 848. — Exsicc.: Heldr. herb. norm. n. 281 et 1439; Orph. fl. gr. n. 348; Bald. it. alb. a. 1892 n. 141; Sint. it. thessal. n. 740; Dörfl. fl. gr. n. 217 et 413.

Caulibus erectis, adpresse tomentosis, in racemum thyrsoideum compositum abeuntibus; foliis subtus niveo-tomentosis, supra glabrescentibus, inferioribus oblongo-spathulatis, crenulatis, in petiolum attenuatis, caulinis late auriculato-amplexicaulibus; capitulis hemisphaericis, discoideis, pedicellis eis aequilongis suffultis; involucri tomentosi phyllis linearibus, acuminatis, squamulis brevissimis; acheniis glabris.

In saxosis regionis subalpinae et alpinae, fere in omnibus montibus elatioribus frequens. Epirus: mt. Tsumerka, Peristeri (Hal.); Thessalia: pr. Chaliki, mt. Oxya, Zygos (Haussk.), Dokimi, Ochsa Despot, Baba, pr. Krania, Mandra Hodza (Form.), mt. Oeta (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Sibth.); Euboea: mt. Dirphys (Fraas); Peloponnesus: mt. Olenos (Heldr.), Chelmos (Orph.), pr. Carithena, mt. Diaforti, Taygetos (Chaub.), Malevo (Orph.). — Jun. Aug. ♃

○○ Folia utrinque viridia.

. Folia indivisa; caulis elatus, corymboso-paniculatus, polycephalus.

11. **S. nemorensis** L. sp. p. 870; Bois. fl. or. III. p. 408; Hal. in ö. b. Z. 1895 p. 409. — *S. Jacquinianus* Rchb. ic. crit. III. p. 80. — *S. nemorensis v. Gmelini* Bald. riv. coll. bot. Alb. 1896 p. 68, vix *S. cacaliaster γ. Gmelini* Vis. fl. dalm. II. p. 71.

Caule erecto, glabro vel puberulo, superne corymboso-paniculato; foliis ovatis vel ovato-lanceolatis, acuminatis, serrulato-dentatis, glabris vel subtus papilloso-puberulis, inferioribus breviter petiolatis, ceteris basi attenuata sessilibus; capitulis obconico-cylindricis; involucri glabri vel parce hirtuti phyllis linearibus, abrupte acutatis, squamulis subulatis, involucro parum brevioribus; ligulis 5, flavis, involucro duplo longioribus; acheniis glabris.

β. **expansus** Bois. et Heldr. diagn. ser. 2, III. p. 34 pro var. *S. Jacquiniani.* — *S. nemorensis v. Gmelini* Bald. riv. coll. bot. Alb. 1896 p. 68, vix *S. cacaliaster γ. Gmelini* Vis. fl. dalm. II. p. 71 = *S. nemorensis v. dalmaticus* Grsb. ap. Pant. adnot. fl. Herceg. Crnag. p. 43. — Caulibus, foliis subtus praesertim ad nervum medium, pedicellis et involucris glanduloso-puberulis. — Glandulositate aliquo modo ad varietatem citatam dalmaticam propius accedit, quae autem ut videtur pedicellis tantum glandulosis gaudet. — Exsicc.: Heldr. it. gr. septentr. a. 1879, pl. fl. hellen. a. 1899; Reis. fl. gr. a. 1894.

In silvaticis montanis et subalpinis, rare. Epirus: mt. Gamila distr. Vratedon (Bald.); Thessalia: mt. Olympus, Oeta (Heldr.); Aetolia: mt. Vardusia pr. Granitza (Reiser); Achaia: mt. Chelmos (Orph.); Laconia: mt. Taygetos pr. Sitzova Alagoniae (Zahn); probabiliter loca

omnia ad varietatem spectant. Typum e Graecia non vidi. — Jul. Aug. ♃

.. Folia incisa, pinnatisecta vel lyrata, rarissime indivisa et tunc caulis humilis oligocephalus.

; Folia oblongo-vel lineari-lanceolata, inferiora runcinata.

12. **S. barckhausiaefolius** Bois. et Heldr. diagn. ser. 2. VI. p. 101, Fl. or. III. p. 402. — Exsicc.: Heldr. pl. hellen. n. 3197.

Caule crassiusculo, erecto, glabrescenti, apice in corymbum 4—7 cephalum abeunti; foliis glabriusculis, inferioribus oblongo-lanceolatis, runcinatis, in petiolum attenuatis, lobis utrinque 2—3, triangularibus, 1—3 denticulatis, foliis superioribus sessilibus, lineari-lanceolatis, subdentatis, summis integris; capitulis mediocribus, hemisphaericis; involucri phyllis glabris, duris, linearibus, carinatis, squamulis numerosis, involucro tertia parte brevioribus; ligulis croceis, disco duplo longioribus; acheniis hirtis. — Species distinctissima.

In regione abietina mt. Pateras Atticae rarissime (Guicc.); postea a nemini lectus. — Maio. ♃

; Folia varie partita vel interdum indivisa, sed nunquam runcinata.

, Folia obovato-vel lanceolato-oblonga, indivisa, acutiuscule dentata.

13. **S. frutiulosus** S. et S. pr. II. p. 178, Fl. gr. IX. p. 54, t. 870; Sieb. avis p. 5, rem. p. 6, in Flora I. p. 275; Raul. cret. p. 785; Bois. fl. or. III. p. 391; Spreitz. in z. b. G. 1890 p. 296; Bald. viagg. Creta p. 63. — Exsicc.: Spreitz. it. cret. a. 1882 n. 48.

Glaber; caulibus erectis, 5—15 cm. altis, apice mono-vel laxe oligocephalis; foliis inferioribus in petiolum attenuatis, intermediis basi sessili minute auriculatis, summis anguste linearibus; involucri phyllis linearibus, bicarinatis, squamulis paucis, involucro 3plo brevioribus; ligulis flavis, radiantibus; acheniis aspero-puberulis, — Sequenti valde affinis, a quo statura humili et foliis indivisis imprimis differt. — Nomen improprium, nam caules basi tantum indurati.

In saxosis regionis mediae et superioris Cretae: ad Krioneriti pr. Palaeolutra, mt. Stravopodia, Hagion Pneuma et Theodori (Raul.) in mt. Sphacioticis (Sibth.), mt. Ida (Heldr.). — Jul. Aug. ♃

,, Folia obovato-oblonga vel oblonga, pinnatifida, segmentis rhachique angulato-dentatis.

14. **S. rupestris** W. et K. pl. rar. hung. II. p. 136, t. 128; Friedr. Reise p. 279; Fraas fl. class. p. 210; Raul. cret. p. 785; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Epir. p. 26, Beitr. fl. Thessal. p. 13, Beitr. fl. Achaia p. 24, in ö. b. Z. 1897 p. 284; Form. in Ver. Brünn 1896 p. 46, 1897 p. 34. — *S nebrodensis* Raul. cret. p. 785; Bois. fl. or. III. p. 391; Hal. in ö. b. Z. 1888 p. 758; Heldr. chlor. Parn. p. 21; Haussk. symb. p. 106; Bald.

riv. coll. bot. Alb. 1896 p. 67; non L., qui foliis saepius obovatis, segmentis obtusioribus, brevius et minus acute dentatis (an semper) dis cedit; cf. Strobl Flora der Nebrod. p. 277. — Exsicc.: Heldr. herb. norm. n. 384 et 1547; Orph. fl. gr. n. 108; Sint. it. thessal. n. 741; Dörfl. fl. gr. n. 208 et 414.

Glaber vel parce arachnoideus; caulibus erectis, 10—40 cm. altis; laxe corymbosis; foliis inferioribus in petiolum attenuatis, ceteris auricula dentata amplexicaulibus; involucri phyllis linearibus, bicarinatis; squamulis involucro 4plo brevioribus; ligulis flavis, radiantibus; acheniis pubescentibus. — Variat caulibus humilioribus vel elatioribus, poly-vel oligocephalis, foliis amplioribus vel angustis, valde dissectis vel tantum inciso-dentatis.

In rupestribus regionis subalpinae et alpinae. Epirus: mt. Smolika (Bald.), Tsumerka, Peristeri, pr. Chaliki (Hal.); Thessalia: pr. Krania, mt. Baba (Form.), Zygos, Karava, Ghavellu (Haussk.) et ad monasterium Korona in Pindo, mt. Olympus, Oeta (Heldr.); Euboea: mt. Kandyli (Orph.), Dirphys (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Guicc.); Achaia: mt. Olenos (Heldr.), Panachaicon, Chelmos (Hal.); Arcadia: pr. Carithena (Friedr.); Laconia: mt. Malevo (Orph.), Taygetos (Zahn); Creta: ad Hagios Joannes Kaimenos (Raul.); Corcyra: pr. Signes (Gelmi). — Jun. Aug. ♃

,,, Folia varie partita, rhachi integra.

— Rhizoma repens; folia omnia pinnatipartita, segmentis lineari-lanceolatis, integris vel dentatis; achenia omnia hirta.

15. **S. erucifolius** L. sp. p. 869 n. 15, cf. errata; Bois. fl. or. III. p. 391; Form. in D. bot. Monat. 1890 p. 18, in Ver. Brünn 1896 p. 46. — *S. tenuifolius* Jacq. fl. austr. III. p. 42, t. 278; Ch. et B. exp. p. 249, Fl. pelop. p. 58; forma segmentis angustioribus, iterum saepe pinnatipartitis.

Araneoso-pubescens vel glabriusculus; caulibus erectis, elatis, laxe et composite-corymbosis; foliis inferioribus in petiolum attenuatis, ceteris sessilibus; involucri phyllis oblongis, squamulis involucrum dimidim aequantibus; ligulis flavis, radiantibus.

In fruticetis, ad vias regionis inferioris et montanae. Thessalia: pr. Velestinos, pr. Lokatamburia in mt. Olympo (Form.); Cyclades (Chaub.). — Jul. Aug. ♃. — N. v.

= Radix obliqua, praemorsa; folia lyrata vel lyrato-pinnatifida, infima interdum indivisa, summa pinnatipartita, segmenta oblonga vel oblongo-linearia, latiora; achenia marginis glabra, disci puberula.

16. **S. Jacobaea** L. sp. p. 870; Form. in D. bot. Monat. 1890 p. 18; Bald. riv. coll. bot. Alb. 1895 p. 49. — *S. erucifolius v. macrophyllus* Form. in Ver. Brünn 1897 p. 34 ex descriptione potius

huc pertinere videtur. — Icon: Fl. dan. t. 944. — Exsicc.: Bald. it. alb. epir. III. n. 158; Sint. it. thessal. n. 743.

Glaber vel araneoso-puberulus; caulibus erectis, elatis, composite et stricte corymbosis; foliis infimis petiolatis, oblongo-obovatis, lyrato-pinnatipartitis, interdum indivisis, caulinis auricula multipartita amplexicaulibus, in lacinias dentatas pinnatipartitis; pedunculis erectis; involucri phyllis lineari-lanceolatis; squamulis involucro multo brevioribus; ligulis flavis, patentibus, disco aequilongis.

In herbidis montanis, rarissime. Epirus: pr. Syraku ad radices mt. Peristeri (Bald.); Thessalia: in valle Negerli pr. Chaliki (Sint.), pr. Kerasia Sina in mt. Chassia (Form.). — Jul. Aug. ⊙

17. **S. barbaraefolius** Krock. fl. siles. II. p. 421; Form. in Ver. Brünn 1895 p. 26. — *S. erraticus* Bert. amoen. p. 92; Marg. et R. fl. Zante p. 58; Friedr. Reise p. 264; Bois. fl. or. III. p. 392; Heldr. fl. cephal. p. 46; Form. in Ver. Brünn 1897 p. 34. — *S. aquaticus* S. et S. pr. II. p. 178; Marg. et R. fl. Zante p. 58; non L. — Cf. Pollak in ö. b. Z. 1896 p. 168 et Scholz in ö. b. Z. 1899 p. 284. — Icon: Rchb. pl. crit. IV. p. 538. — Exsicc.: Heldr. it. thessal. n. 82; Sint. it. thessal. n. 1229.

Differt a praecedente caule divaricatissime corymboso, foliorum segmentis patentissimis, terminali maximo, ovato, saepe subcordato, pedunculis patentibus, ligulis saepe brevioribus.

In pratis, salicetis, ad fossas regionis inferioris. Epirus: pr. Philippiades, in valle Dipotami (Form.); Thessalia: pr. Kalabaka (Sint.), ad ostia fluvii Peneios (Heldr.), ad Megalivris et Mavromandila pr. Lamia (Form.); Aetolia: pr. Mesolongion (Nied.); Boeotia: pr. Lebadia (Fraas); Euboea: ad Oreus (Orph.); Achaia: pr. Patras (Friedr.); Messenia: inter Nisi et Petalidi (Zahn); Zante (Marg.); Cephalonia: pr. Same (Heldr.); Corcyra: pr. urbem, Kipuria, Kanoni, Europuli, Afra (Form.). — Jul. Sept. ⊙

c. Annui.

α. Ligularum lamina involucrum valde superans.

18. **S. coronopifolius** Desf. fl. alt. II. p. 273; Bois. fl. or. III. p. 390; Heldr. chlor. Thera p. 14, chlor. Mykon. p. 245. — *S. squalidus* Urv. enum. p. 110, non L. — Exsicc.: Sart. herb. n. 3370; Heldr. et Hal. fl. aeg. a. 1889.

Glabriusculus; caule erecto, saepe a basi ramoso; foliis pinnatipartitis, segmentis obovatis, oblongis vel linearibus, inciso-dentatis subintegrisve, rhachi subintegra, infimis petiolatis, ceteris auriculato-amplexicaulibus; capitulis laxe corymbosis, longe pedunculatis; involucri phyllis linearibus, acuminatis; squamulis lineari-lanceolatis, involucro multo brevioribus; ligulis flavis, subrevolutis, involucro subbrevioribus; acheniis costatis, adpresse hirtis, pappo deciduo, discum aequante.

Inter segetes, ad muros regionis inferioris, rare. Attica: in scopulo Rhaphti (Heldr.); Cycladum insula: Mykonos (Sart.), Melos (Urv.), Cythnos, Thera (Heldr.); Creta: mt. Cavutsi (Heldr.). — Mart. Maio. ⊙

19. **S. vernalis** W. et K. pl. rar. hung. I. p. 23, t. 24; Marg. et R. fl. Zante p. 58; Friedr. Reise p. 266, 268 et 277; Clem. sert. p. 63, *v. nigropunctatus* (f. phyllis involucri exterioribus nigropunctatis), Ung. Reise p. 124; Bois. fl. or. III. p. 389; Spreitz. in z. b. G. 1877 p. 712; Heldr. fl. cephal. p. 46; Haussk. symb. p. 106; Hal. in ö. b. Z. 1897 p. 284. — Exsicc.: Heldr. herb. norm. n. 686; Orph. fl. gr. n. 1140; Sint. it. thessal. n. 77.

Araneoso-lanatus, tandem glabrescens; caule erecto, simplici vel ramoso, laxe corymboso; foliis infimis oblongis, breviter petiolatis, sinuatis vel dentatis, ceteris auriculato-amplexicaulibus, oblongo-linearibus, varie pinnatifidis, lobis oblongis, rhachique crispato-dentatis; capitulis longe pedunculatis; involucri phyllis linearibus, acuminatis; squamulis setaceis, involucro multo brevioribus; ligulis flavis, patentibus, involucro aequilongis; acheniis striatis, dense pubescentibus, pappo persistente, discum aequante. — Praecedenti affinis, ab eo distinctus indumento, foliis minus dissectis, lobis rhachique crispato-dentatis et ligulis longioribus.

In arenosis maritimis, nec non in saxosis montanis et subalpinis. Epirus: pr. Kestoration (Chodses), jam extra ditionem nostram; Thessalia: ad monasterium Hagios Stephanos pr. Kalabaka (Sint.); Acarnania: pr. Astakos (Nied.), insula Petala Echinadum (Reiser); Aetolia: pr. Mesolongion, mt. Chalkis (Nied.), mt. Tymphrestus (Samar.); Boeotia: pr. Lebadia (Orph.); Attica: pr. Eleusis (Haussk.), Marussi, Athenas (Friedr.), mt. Parnes, Pentelicon, Hymettus, ad Phalerum, insula Lero (Heldr.); Acrocorinthus (Haussk.); Argolis: pr. Nauplia (Orph.), Poros (Friedr.); Arcadia: pr. Zatuna (Orph.); Laconia: pr. Selitza (Zahn); Zante: ad colles Acroteri (Marg.); Cephalonia: pr. Argostoli, Pessada, Spartia (Heldr.); Corcyra: mt. Deca (Ung.), Pantocrator (Spreitz.). — Febr. Maio. ⊙

β. Ligulae nullae vel brevissimae, involucrum vix superantes revolutae.

× Achenia glabra.

20. **S. viscosus** L. sp. p. 868; S. et S. pr. II. p. 177; Friedr. Reise p. 278; Ung. Reise p. 123; Bois. fl. or. III. p. 386; Haussk. symb. p. 106; Form. in Ver. Brünn 1896 p. 46. — *S. tymphresteus* Bois. et Heldr. diagn. ser. 2, VI. p. 99. — Icon: Fl. dan. t. 1230. — Exsicc.: Heldr. it thessal. IV. a. 1885.

Glanduloso-hirtus, viscosus; caule erecto, ramoso; foliis oblongo-spathulatis oblongisve, inaequaliter lobatis vel sinuato-dentatis, inferioribus in petiolum attenuatis, superioribus sessilibus; involucri phyllis linearibus, acutis; squamulis involucro 2—3plo brevioribus; ligulis flavis, minimis, revolutis; pappo disco longiore.

In saxosis montanis, in regionem alpinam adscendens. Thessalia: mt. Baba inter Krania et Klinovo in Pindo (Haussk.); Eurytania: mt. Tymphrestus (Samarit.); Argolis: ad lacum Lerna (Friedr.); in agro Laconico (Sibth.); Corcyra (Ung.). — Jun. Aug. ⊙

×× Achenia pubescentia.

21. **S. lividus** L. sp. p. 867; S. et S. pr. II. p. 177; Urv. enum. p. 110; Ch. et B. exp. p. 249, Fl. pelop. p. 58; Bois. fl. or. III. p. 388. — *S. trilobus* S. et S. pr. II. p. 177, Fl. gr. IX. t. 869, non L., sec. Bois. l. c. — *S. foeniculaceus* Ten. fl. nap. II. p. 216, t. 78.

Pube papillari articulata plus minus obsitus, superne glanduloso-puberulus, viscidus; caule erecto, simplici vel ramoso, laxe corymboso; foliis sinuato-lobatis vel pinnatifidis, inferioribus obovatis, in petiolum attenuatis, ceteris oblongo-lanceolatis, basi auriculato-amplexicaulibus, laciniis angulato-dentatis; involucri phyllis linearibus, acuminatis; squamulis involucro multo brevioribus; ligulis flavis, revolutis, parte exserta nvolucro 3—4plo brevioribus; acheniis pappo brevioribus.

In arenosis regionis inferioris et montanae. Peloponnesus (Sibth.): pr. Carithena, Pylos, Messene, mt. Taygetos (Chaub.); Cycladum insula Skinosa (Sibth.), Camini pr. Thera (Urv.). — Mart. Jul. ⊙ N. v.

22. **S. vulgaris** L. sp. p. 867; S. et S. pr. II. p. 176; Pieri corc. fl. p. 117; Dallap. prosp. p. 115; Ch. et B. exp. p. 248, Fl. pelop. p. 58; Marg. et R. fl. Zante p. 58; Friedr. Reise p. 273; Fraas fl. class. p. 210; Weiss in z. b. G. 1869 p. 39; Raul. cret. p. 785; Bois. fl. or. III. p. 386; Heldr. fl. cephal. p. 46; Haussk. symb. ip. 106; Hal. Beitr. fl. Epir. p. 26. — *S. delubrorum* Heldr. herb. norm. n. 1347. — Icon: Fl. dan. t. 513. — Exsicc.: Heldr. pl. fl. hellen. a. 1878.

Glaber vel arachnoideo-lanatus; caule erecto, simplici vel ramoso, apice coarctato-corymboso; foliis oblongis vel oblongo-spathulatis, sinuato-lobatis vel pinnatifidis, inferioribus in petiolum attenuatis, ceteris basi auriculato-amplexicaulibus, laciniis angulato-dentatis; involucri phyllis linearibus, acuminatis; squamulis involucro multo brevioribus; acheniis pappo brevioribus, — Differt a praecedente indumento eglanduloso, capitulis discoideis.

In herbidis, ruderatis, cultis, per omnem Graeciam. — Floret toto anno. — ⊙

19 × 22. **S. vernalis × vulgaris.** — *S. vulgaris × vernalis f. pseudo-vernalis* Zabel in Verh. bot. Ver. Brand. 1869 p. 139. — *S. Weylii* Vatke in Verh. bot. Ver. Brandenb. 1872 p. 48. — Habitu *S. vernalis*, sed folia minus crispato-dentata, ligulae breviores vel nullae, capitula eis *S. vulgaris* majora. — Inter parentes ad monasterium Hagios Stephanos pr. Kalabaka Thessaliae (Sint. it. thessal. n. 87).

7 Tribus. CALENDULEAE Less, syn. p. 89.

35. Calendula L. gen. n. 990.

1. **C. arvensis** L. sp. ed 2 p. 1303; S. et S. pr. II. p. 207; Pieri corc. fl. p. 122; Ch. et B. exp. p. 257, Fl. pelop. p. 60; Friedr. Reise p. 266 et 272; Ung. Reise p. 124; Weiss in z. b. G. 1869

p. 45, *f. rugosa* Vis. fl. dalm. II. p. 26 (acheniis fere omnibus dorso transverse rugosis plicatisve, plicis granulatis vel breviter muricatis); Raul. cret. p. 786; Bois. fl. or. III. p. 418; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 46, Fl. Aegina p. 304; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 118. — *C. micrantha* Tin. et Guss. in Guss. fl. sic. syn. II p. 874; Heldr. chlor. Thera. p. 15; (f. ligulis brevioribus, involucro parum longioribus). — Forsan ad hanc formam pertinet: *C. arvensis pygmaea* Clem. sert. p. 54. — *C. sublanata* Rchb. ic. XV. p. 99, t. 160 (f. caule foliisque tomentellis). — Huc quoque: ? *C. parviflora* Friedr. Reise p. 284, an Raf.; nec non *C. sicula* Friedr. l. c., non Cyr. — Icon: Fl. gr. t. 920. — Exsicc.: Heldr. herb. norm. n. 1041; Sint. it. thessal. n. 400.

Pubescens; caulibus erectis vel decumbentibus, ramosis; foliis oblongo-lanceolatis, subdentatis, infimis in petiolum attenuatis, ceteris sessilibus semiamplexicaulibus; capitulis mediocribus, radio explanato circa 2 cm. latis; ligulis sulfureis vel aurantiacis, involucro subduplo longioribus, disco concolore, acheniis exterioribus dorso plus minus longe aculeatis, rostratis, involucro sesqui vel duplo longioribus, intermediis cymbiformi inflatis intimisque annulatis dorso rugosis.

β. **bicolor** Raf. caract. p. 82; Bois. fl. or. III. p. 418; Haussk. symb. 118; Hal. in ö. b. Z. 1897 p. 95; Heldr. fl. Aegina p. 304; pro sp.; DC. pr. VI. p. 453; Marg. et R. fl. Zante p. 58. — Huc probabiliter: *C. officinalis* Pieri corc. fl. p. 122; Dallap. prosp. p. 117; Marg. et R. l. c.; Heldr. fl. cephal. p. 46; non L., quae foliis omnibus basi paulo angustioribus, floribus duplo majoribus, aureis atque acheniis plerisque cymbaeformibus discedit et in Graecia non indigena. — Involucri phylla apice crocea, corollae disci croceo-nigricantes. — Exsicc.: Orph. fl. gr. n. 474; Heldr. herb. norm. n. 315 et 1254.

In cultis, inter segetes, herbidis, vinetis, olivetis, collibus, aridis, regionis inferioris per totam Graeciam frequens. — Febr. Maio. ⊙

C. aegyptiaca Desf. cat. hort. Par. 1804 p. 100; Bois. fl. or. III. p. 419; Heldr. Fl. Aegina p. 304, chlor. Thera p. 15, chlor. Mykon. p. 246. — Exsicc.: Orph. fl. gr. n. 475; Heldr. herb. norm. n. 514 et 1042; Dörfl. fl. aeg. n. 435, pl. cret. n. 136. —

Papilloso-hirta; caulibus decumbentibus vel adscendentibus, ramosis; foliis oblongo-lanceolatis lanceolatisve, remote denticulatis, infimis in petiolum attenuatis, ceteris sessilibus, semiamplexicaulibus; capitulis parvis, radio explanato circa 1 cm. latis; ligulis flavis, involucrum vix superantibus, disco concolore; acheniis exterioribus dorso longe aculeatis, in rostrum tenue aequilongum vel longius, involucrum superans, abeuntibus, intermediis alato-cristatis vel cymbaeformibus, dorso echinatis vel laevibus, intimis annularibus rugosis.

In arenosis, cultis, collibus aridis regionis inferioris. Attica: mt. Lycabettus, pr. Trakones ad radices mt. Hymetti, insula Aegina (Heldr.); isthmus Corinthiacus: pr. Lechaeum (Orph.); Cycladum insula: Syra Sart.), Melos (Armenis), Jos, Mykonos, Thera (Heldr.), Cythnos (Tunt.),

Tenos, Naxos (Leon.); Cretae: pr. Males distr. Hierapetra (Leon.). — Sept. Jun. ⊙.

2. Subdivisio. **CYNAROCEPHALAE** Juss. gen. p. 171.

8. Tribus. **ECHINOPSIDEAE** Less. in Linnaea VI. p. 88.

36. Echinops L.

a. Involucri phylla quina intima inferne in tubum concreta.

1. **E. viscosus** DC. pr. VI. p. 525; Bois. fl. or. III. p. 429; Heldr. fl. cephal. p. 46, in ö. b. Z. 1898 p. 184, Chlor. Thera p. 15, Chlor. Mykon. p. 246; Bald. viagg. Creta p. 66; Hal. in z. b. G. 1899 p. 185. — *E. sphaerocephalus* S. et S. pr. II. p. 208, non L. — *E. spinosus* Sieb. avis p. 5, rem. p. 6; Raul. cret. p. 786; Spreitz. in z. b. G. 1890 p. 296; an etiam S. et S. pr. II. p. 209 quoad plantam ex Archipelago, nec non Ch. et B. exp. p. 257, Fl. pelop. p. 60; non L. mant. p. 119, qui indumento eglanduloso, foliis parenchymate fere destitutis in spinas longas acerosas ramosas mutatis et penicilli (id est phyllorum involucri series extima) setis involucrum aequantibus discedit. — *E. creticus* Bois. et Heldr. diagn. X. p. 87; Raul. cret. p. 786. — *E. glandulosus* Weiss in z. b. G. 1868 p. 433, 1869 p. 42. — Icon: Fl. gr. t. 923. — Exsicc.: Rev. pl. cret. n. 78, in Baen. herb. europ. n. 4872; Bald. it. cret. n. 74; Dörfl. fl. aeg. n. 267.

Glanduloso-viscosus; caule elato, ramoso; foliis in lacinias lanceolatas lobatas, spinosas, pinnatim vel subbipinnatim partitis vel sectis, superne virentibus, glandulosis, subtus cano-tomentosis; penicilli involucri tertiam vel dimidiam partem aequantis setis scabridulis; involucri phyllis infimis a basi tenui abrupte triangulari-deltoideis, saepe transverse latioribus, glabris vel glandulis sessilibus obsitis, intermediis versus medium denticulatis et attenuato-subulatis, intimis a parte media in tubum cylindricum concretis, omnibus vel tribus brevioribus obtusis fimbriatis; corolla pallide coerulea; pappi setis paleaceis barbellatis, basi vel ad tertiam partem concretis.

In aridis, dumetosis regionis inferioris. **Attica**: mt. Hymettus (Heldr.); ? Corinthus, **Argolis et Laconia** (Chaub.); **Cycladum insula** Keos, Andros, Naxos, **Siphnos, Thera (Heldr.), Mykonos (Sart.)**, Syra (Weiss); Creta: pr. **Khalamia (Bald.), Kissamos, Khalepa, Candia**, mt. Lassiti (Raul.), Anopolis, Lutro (Spreitz.); Cephalonia: inter Drapano et Pharaelata, pr. Hagios Soter (Heldr.). — Jun. Aug. ♃.

2. **E. graecus** Mill. dict. n. 3; Fraas fl. class. p. 203 et 216; Clem. sert. p. 54; Bois. fl. or. III. p. 432; Hal. Beitr. fl. Thessal. p. 14; Haussk. symb. p. 119; Form. in Ver. Brünn 1896 p. 55, 1897 p. 41, 1898 p. 78. — *E. lanuginosus* Lam. dict. III. p. 317; S. et S. pr. II. p. 209; Urv. enum. p. 118; Ch. et B. fl. pelop. p.

60. — *E. tenuissimus* S. et S. fl. gr. X. p. 19 t. 926. — Exsicc.: Orph. fl. gr. n. 269; Sint. it. thessal. n. 1275.

Eglandulosus, araneosus; caule humili, ramoso; foliis in lacinias angustas, lineari-lanceolatas, spinosas, pinnatisectis, supra nitidis, subtus cano-tomentosis; penicilli involucri tertiam partem aequantis setis laeviusculis; involucri glabri phyllis externis a basi tenui apice triangulari ovatis, 5—7 dentatis, intermediis supra medium subulato-attenuatis et adscendenti-dentatis, intimis longioribus, medium versus puberulis, ad $^2/_3$ in tubum oblongo-pentagonum concretis, apice subulatis; corolla, coerulea; pappi brevissimi paleis barbellatis, ultra medium concretis. — Statura humili, indumento eglanduloso, foliis creberrimis tenuissime sectis, capitulis minoribus coeruleis, involucri structura et toto habitu a praecedente diversissimus.

In arvis, ruderatis. Attica: frequens ubique circa Athenas; pr. Vatika (Fraas); Thessalia: pr. Volo (Hal.), Lalomeion, Koryza (Form.); Euboea: pr. Pisona (Orph.); Cycladum insula: Keos (Heldr.), Kythnos (Tunt.), Melos (Urv.). — Maio, Aug. ♃.

b. Involucri phylla omnia libera.

α. Capitula albo-virentia, floribus albis vel pallide coerulescentibus.

× Caulis totus valde glandulosus; involucri phylla saltem inferiora glandulosa.

3. **E. sphaerocephalus** L. sp. p. 814; Bois. fl. or. III. p. 438. — Icon: Fl. dan. t. 2179.

Caule elato, glanduloso-tomentoso, simplici vel ramoso; foliis mollibus, in segmenta oblongo-triangularia, lobata, spinulosa pinnatifidis, utrinque glandulosis, supra viridibus, subtus cano-tomentosis; penicilli setis dimidio involucro sublongioribus; involucri phyllis inferioribus glanduloso-puberulis, apice triangularibus, ceteris glabris, subulato-acuminatis, a medio sursum ciliato-pectinatis; pappi setis ad medium usque concretis.

In fruticetis Peloponnesi circa Megaspilaeon Achaiae (Heldr. ex Bois. l. c.). — Jul. Aug. ♃. N. v.

4. **E. taygeteus** Bois. et Heldr. diagn. VI. p. 98. — *E. sphaerocephalus β. taygeteus* Bois. fl. or. III. p. 439; Heldr. chlor. Parn. p. 21. — Exsicc.: Orph. herb. a. 1851; Hal. it. gr. a. 1888.

Differt a praecedente foliis coriaceis, in lacinias angustas, parenchymate saepe fere omnino destitutas, in spinas longas validasque terminales lateralesque abeuntes pinnatipartitis, involucri phyllis omnibus subulatis, saepius omnibus glanduloso-puberulis.

In regione abietina mt. Parnassus pr. Gurna (Heldr.), mt. Chelmos Achaiae (Orph.), mt. Taygetos supra Varvari (Heldr.). — Jul. Aug. ♃.

×× Caulis eglandulosus vel basi tantum glandulosus; involucri phylla omnia eglandulosa, glabra.

5. **E. albidus** Bois. et Sprun. diag. VI. p. 97; Clem. sert. p. 54; Haussk. symb. p. 118; Form. in D. bot. Monat. 1898 p. 78, *v. longifolius*. — *E. sphaerocephalus γ. albidus* Bois. fl. or. III. p. 439; Form. in D. bot. Monat. 1895 p. 29, 1897 p. 40. — Huc probabiliter: *E. sphaerocephalus* Ch. et B. exp. p. 257, Fl. pelop. p. 60; Form. in D. bot. Monat. 1890 p. 21. — Exsicc.: Heldr. herb. norm. n. 338 et 1255; Orph. fl. gr. n. 351.

Caule mediocri, araneoso-tomentoso, monocephalo vel apice oligocephalo, foliis coriaceis, in segmenta breviter triangularia, lobulata, crebre et longe spinosa pinnatifidis, supra virescentibus, dense breviterque glandulosis, subtus cano-tomentosis; penicilli setis involucro dimidio brevioribus; involucri phyllis inferioribus apice triangularibus, ceteris subulato-acuminatis, a medio sursum ciliato-pectinatis; pappi setis ad medium usque concretis. — Capitula magnitudinis *E. sphaerocephali*, qui autem longe differt caule ad apicem usque dense glanduloso, foliis majoribus mollibus, tenuius spinosis, supra longius glandulosis et involucri phyllis glandulosis; *E. taygeteus* foliis magis dissectis et indumento glanduloso discedit.

In collibus saxosis. Epirus: pr. Panegeia, Ephemiades, Han Mescho, Janina, Sadovica, mt. Prosgoli, in valle Dipotami (Form.); Thessalia: pr. Tyrnavos (Haussk.), Kapurna, Volo, Lalomeion, Velestinos, Limogardi, Mégalivris (Form.); Euboea: ad radices mt. Dirphys (Heldr.); Attica: in colle Lycabetto, Ardetto, mt. Corydalus, Hymettus, ad Ilyssum, pr. Munychia, Phanari, Daphne (Heldr.); Boeotia: pr. Thebas (Sprun.); Acrocorinthus (Hal.); Argolis (Sart.); Laconia: pr. Vitilos, Chimova, ad fluvium Eurotas (Chaub.). — Jun. Aug. ♃.

β. Capitula cyanea.

× Penicillum involucro 3—4 plo brevius.

6. **E. banaticus** Roch. cat. hort. vindob. a. 1823, in Schrad. diss. blum. p. 48; Form. in Ver. Brünn 1895 p. 29, 1897 p. 41. — *E. ruthenicus* Roch. pl. ban. rar. t. 37, non M. a B. — *E. Rochelianus* Griseb. spic. II. p. 229. —

Eglandulosus; caule elato, arachnoideo-tomentoso, simplici vel ramoso; foliis membranaceis, teneris, supra viridibus, breviter papillosis, subtus cano-tomentosis, inferioribus petiolatis, lyratis vel in segmenta oblonga, grosse et parce lobata pinnatifidis, margine ciliato-spinulosis, lobis in spinam minutam debilem abeuntibus, foliis superioribus diminutis, amplexicauli-auriculatis; penicilli paleis involucro triplo brevioribus; involucri glabri phyllis extimis apice minute triangulari-spathulatis, denticulatis, ceteris lanceolato-attenuatis, a medio longe ciliatis; pappi setis inferne connatis. — Foliis teneris, grosse et parce lobatis, lobis minutissime spinuloso-ciliatis insignis.

In valle Dipotami in Epiro et in mt. Cuka in mt. Chassia Thessaliae (Form.). — Jul. Aug. ♃. N. v.

7. **E. ritro** L. sp. p. 815; Bois. fl. or. III. p. 439; Hal. in z. b. G. 1888 p. 759, in ö. b. Z. 1892 p. 41; Form. in D. bot. Monat. 1890 p. 22, in Ver. Brünn 1897 p. 55, 1897 p. 41; Haussk. symb. p. 119; Bald. riv. coll. bot. Alb. 1896 p. 68. — Exsicc.: Orph. herb. n. 2957.

Eglandulosus; caule mediocri, tomentoso; foliis coriaceis, supra viridibus, glabrescentibus vel sublanatis, subtus cano-tomentosis, inferioribus petiolatis, pinnatisectis, segmentis lanceolatis linearibusve, pinnatipartitis, breviter spinosis, foliis superioribus diminutis, sessilibus; penicilli paleis involucro 3—4plo brevioribus; involucri glabri phyllis extimis apice minute triangulari-spathulatis, denticulatis, ceteris lanceolato-attenuatis, a medio longe ciliatis; pappi setis fere ad medium irregulariter concretis.

β. **ruthenicus** M. a. B. fl. taur. cauc. III. p. 597 pro sp. — *E. ritro β. elegans* Bert. in Vis. stirp. dolm. sp. p. 34. — *E. ritro β. tenuifolius* DC. pr. VI. p. 524. — Foliorum segmenta angusta, longiora, linearia. — Exsicc.: Sint. et Bornm. it thessal. n. 1321.

γ. **Sartorianus** Bois. diagn. VI. p. 97; Heldr. chlor. Parn. p. 21; pro sp.; Bois. fl. or. III. p. 440. — Pappi setae longius et interdum prope apicem usque coalitae. — Exsicc.: Heldr. herb. n. 2478.

In saxosis regionis montanae, var. β. ut videtur aliis frequentior. Epirus: mt. Smolika (Bald.); Thessalia: pr. Malakasi (Haussk.), Said Pascha, Krania, Klinovo, Chaliki in Pindo (Form.), pr. Leptokaryes in Olympo (Sint.), pr. Selicany et Spilia in mt. Ossa, mt. Othrys (Form.); Aetolia: mt. Tymphrestus (Samar.), mt. Korax (Heldr.); mt. Kiona (Hal.), Parnassus (Guicc.); Peloponnesus: mt. Chelmos, Malevo (Orph.), Taygetos (Heldr.). — Jul. Aug. ♃.

×× Penicillum subnullum.

8. **E. microcephalus** S. et S. pr. II. p. 209; Bois. fl. or. III. p. 440; Heldr. in Sitzungsb. Acad. Wiss. Berl. 1883 p. 8; Form. in D. bot. Monat. 1890 p. 22, in Ver. Brünn 1896 p. 55, 1897 p. 41; Hal. Beitr. fl. Thessal. p. 14; Haussk. symb. p. 119. — Icon: Fl. gr. t. 925. — Exsicc.: Heldr. herb. norm. n. 40 et 1256; Orph. fl. gr. n. 350.

Araneoso-canescens, eglandulosus; caulibus mediocribus, fere a basi ramosis, dumosis; foliis rigidis, in segmenta lanceolata vel ovato-lanceolata, lobata, breviter spinosa, pinnatipartitis vel sectis; capitulis parvis; penicilli brevissimi setis paucis; involucri glabri phyllis pectinato-ciliatis; ab externis apice triangularibus ad superiora lanceolato-subulata sensim auctis; pappi paleis ad medium coalitis.

In campis, ad vias. Copiose in planitie Thessaliae; Aetolia (Sprun.); Boeotia: pr. Lebadea (Orph.); Achaia: pr. Xylocastron (Heldr.); Arcadia: pr. Tripolis (Sart.). — Jul. Aug. ♃.

9. Tribus. **CARLINEAE** Less. syn. p. 11.

37. **Cardopatium** Juss. in ann. mus. Par. VI. p. 324.

1. **C. corymbosum** L. sp. p. 815 *(Echinops)*; Pers. syn. II. p. 500; Bois. fl. or. III. p. 442; Heldr. fl. cephal. p. 46, in Sitzungsb. acad. Wiss. Berl. 1883 p. 8, Fl. Aegina p. 304; Form. in D. bot. Monat. 1890 p. 21, 1898 p. 78, in Ver. Brünn 1896 p. 53, 1897 p. 38; Haussk. symb. p. 119. — *Carthamus corymbosus* L. sp. ed. 2. p. 1164; S. et S. pr. II. p. 161; Ch. et B. exp. p. 244, Fl. pelop. p. 57; Fraas fl. class. p. 206. — *Brotera corymbosa* Willd. sp. III. p. 2399. — *Onobroma corymbosum* Spreng. syst. III. p. 393. — *C. orientale* Spach. in ann. sc. nat. 3, ser. V. p. 237; Raul. cret. p. 786. — Icon: Fl. gr. t. 844; Jaub. et Sp. ill. or. t. 426. — Exsicc.: Orph. fl. gr. n. 317; Heldr. herb. norm. n. 241 et 851.

Glabrum vel parce arachnoideum; caule humili, ramoso dense corymboso; foliis radicalibus petiolatis, oblongo-lanceolatis, pinnatipartitis, segmentis oblongis, inciso-pinnatifidis, spinosis, caulinis sursum diminutis sessilibus, secus caulem decurrentibus; capitulis numerosissimis, ovato-conicis, 8—10 floris, in glomerulos bracteatos spinosissimos congestis; involucri squarrosi phyllis spinulosis, in spinam recurvam abeuntibus, intimis subscariosis, subintegris; corolla cyanea, rarissime alba; pappi uniseriati paleis lanceolatis, fimbriatis, acuminatis, corollae tubo aequilongis.

In campis, collibus aridis regionis inferioris. Thessalia: frequens in planitie a Trikala, Pharsala, Karditza usque Larissa et Volo (Heldr.); Attica: pr. Athenas, Phalerum, insula Aegina (Heldr.); Peloponnesus: ad fluvium Eurotas, pr. Epidaurus Limera, Monembasia, Damala (Chaub.); Cycladum insula Naxos (Chaub.); Creta: in planitie Messara (Raul.); Cephalonia: pr. Argostoli (Heldr.). — Jun. Jul. ♃.

2. **C. Vrionis** Heldr. in Bois. fl. or. III. p. 443. — Icon: Rouy. ill. t. 160. — Exsicc.: Heldr. herb. fl. hellen. n. 112.

Praecedenti maxime affine, ab eo parum differt foliis minus spinosis, caulinis brevius decurrentibus, capitulis submajoribus, ovatis et pappi paleis corollae tubum dimidium aequantibus.

Aetolia: in halipedo pr. Mesolongion et ad ripas Eueni fluminis (Nieder). — Jun. Jul. ♃.

3. **C. Boryi** Spach in ann. sc. nat. 3 ser. V. p. 243; Bois. fl. or. III. p. 443. —

Ex descriptione praecedentibus simile, sed foliorum segmenta paulo latiora et capitula submajora, spinis validioribus; ab iis praesertim differt pappo duplici, externo paleis 10—13 liberis, ovario et tubo corollae duplo brevioribus constante, interno brevissime coroniformi 5-denticulato.

Laconia (Bory), sine loci specialis indicatione. Jun. Jul. ♃. N. v.

38. Xeranthemum L. gen. n. 947.

a. Involucri phylla glabra, abrupte mucronata; pappi paleae 5.

1. **X. annuum** L. sp. p. 857; S. et S. pr. II. p. 172; Bois. fl. or. III. p. 444; Hal. in z. b. G. 1888 p. 759; Form. in D. bot. Monat. 1890 p. 21, in Ver. Brünn 1896 p. 15, 1897 p. 40; Haussk. symb. p. 121; Bald. riv. coll. bot. Alb. 1896 p. 68. — Icon: Jacq. fl. austr. t. 388. — Exsicc.: Bald. it. alb. epir. IV. n. 126.

Lanato-tomentosum; caule erecto, stricte ramoso, ramis virgatis, apice nudis; foliis lanceolatis, sessilibus; capitulis hemisphaericis; involucri phyllis scariosis, externis pallidis, ovatis, intimis lanceolatis, disco duplo longioribus, radiantibus, roseis; antherarum caudis fimbriatis; pappi paleis e basi triangulari subulatis, achenio brevioribus.

In lapidosis, collibus apricis regionis inferioris et montanae. Epirus: pr. Vromonero distr. Ljaskovik (Bald.); Thessalia: pr. Aivali (Haussk.), pr. Pharsalum (Heldr.), Trikala, Drepani, mt. Chassia, Kapurna, Kalyvia (Form.); Doris: pr. Segditza ad radices mt. Kiona, Amphissa (Hal.). — Jun. Jul. ⊙

2. **X. inapertum** L. sp. p. 858 pro var. *X. annui;* Willd. sp. III. p. 1902; Bois. fl. or. III. p. 445; Heldr. chlor. Parn. p. 21; Hal. Beitr. fl. Achaia p. 24; Bald. riv. coll. bot. Alb. 1896 p. 68. — *X. erectum* Presl del. Prag. p. 106; Haussk. symb. p. 121. — Icon: Rchb. t. 6. — Exsicc.: Heldr. herb. norm. n. 700; Orph. fl. gr. n. 753.

Differt a praecedente capitulis ovatis; involucri phyllis brevioribus, vix radiantibus, erectis; antherarum caudis integris; pappi paleis setaceo-acuminatis, achenio sublongioribus.

In locis similibus, quibus praecedens. Epirus: mt. Olycika (Bald.); Thessalia: mt. Ghavellu, ad monasterium Korona (Haussk.); Phocis: mt. Parnassus pr. Rachova (Samarit.); Attica: mt. Cithaeron, Pateras, Parnes, Hymettus (Heldr.); Achaia: mt. Kastro pr. Kalavryta (Hal.); Laconia: mt. Malevo pr. Hagios Joannes (Orph.); Creta: mt. Volakia (Reverch.). — Maio, Jul. ⊙.

b. Involucri phylla exteriora obtusa, dorso tomentella; pappi paleae 8—10.

3. **X. cylindraceum** S. et S. pr. II. p. 172; Bois. fl. or. III. p. 446; Heldr. chlor. Parn. p. 21; Haussk. symb. p. 121; Bald. riv. coll. bot. Alb. 1896 p. 68; Form. in Ver. Brünn 1897 p. 40. — Icon: Rchb. t. 7. — Exsicc.: Orph. fl. gr. n. 754; Sint. et Bornm. it. turc. n. 1310; Sint. it. thessal. n. 916.

Lanato-tomentosum; caule erecto, stricte ramoso, ramis virgatis, apice nudis; foliis lanceolatis, sessilibus; capitulis oblongo-cylindricis, involucri phyllis scarioso-marginatis, externis pallidis, ellipticis, intimis lanceolatis acutis, vix radiantibus, roseis; antherarum caudis minute mucronatis; pappi paleis inaequalibus, e basi triangulari-lanceolata, subulatis, achenio brevioribus.

In aridis, ruderatis, vineis regionis inferioris. Epirus: mt. Maria distr. Ljaskovik (Bald.); Thessalia: pr. Malakasi (Sint.), Kastania (Form.), monasterium Korona (Haussk.), pr. Palaeokastron (Heldr.), Litochori (Sint.), Volo (Orph.); Euboea: pr. Kastaniotissa (Heldr.); mt. Parnassus pr. Pania (Orph.); Boeotia (Sprun.). — Jun. Jul. ⊙

39. Carlina L. gen. n. 929.

a. Involucri phylla intima non radiantia; pappus biserialis.

1. **C. gummifera** L. sp. p. 829; Sieb. avis p. 5, rem. p. 6, in Flora I. p. 275; Dallap. prosp. p. 110; Urv. enum. p. 104; Ch. et B. exp. p. 244, Fl. pelop. p. 57; Fraas fl. class. p. 205; Clem. sert. p. 56; *(Atractylis)*; Less. syn. p. 12; Marg. et R. fl. Zante p. 59; Heldr. Nutzpfl. p. 26, Fl. cephalon. p. 46, chlor. Mykon p. 246; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 787; Bois. fl. or. III. p. 451; Bald. in nuov. giorn. bot. ital. 1894 p. 97, viagg. Creta p. 67. — *Acarna gummifera* Willd. sp. III. p. 1699; S. et S. pr. II. p. 159, Fl. gr. t. 838. — *C. acaulis* Dall. prosp. p. 110, non L. — Exsicc.: Orph. fl. gr. n. 208; Heldr. herb. norm. n. 424; Dörfl. fl. aeg. n. 430.

Subacaulis; foliis araneosis, fere omnibus basilaribus, in orbem expansis, oblongo-lanceolatis, in segmenta ovato-oblonga, pinnatifida, spinoso-dentata pinnatisectis; capitulis maximis, subsessilibus, solitariis vel 2—3 nis; involucri phyllis pluriserialibus, capitulum non superantibus, extimis coriaceis, lineari-lanceolatis, in spinam validam desinentibus, margine spinas digitatas gerentibus, mediis spinescentibus integris, intimis muticis, apice purpurascentibus; corollis purpureis; pappi pilis achenio 5 plo longioribus. — Rhizoma venenum homini mortiferum continet.

Ad vias, agrorum margines regionis inferioris. Epirus: ad Nicopolis pr. Prevesa (Bald.); Boeotia: pr. Lebadea (Guicc.); Attica: pr. Athenas (Orph.), Marussi, Kephissia, Tatoi (Heldr.); Corinthus (Chaub.); Argolis: pr. Nauplia (Sart.), Epidaurus Limera (Chaub.), Poros (Reinh.), Astros (Orph.); Laconia: pr. Monembasia (Chaub.), Skardamula (Heldr.); Cycladum insula: Andros (Fraas), Keos, Tenos, Syra, Mykonos (Heldr.), Rhenea (Tunt.), Naxos (Leon.); Creta: pr. Almyros (Tourn. voy. p. 13), Platania (Bald.), Canea, Retymo, insula Dia (Raul.); Zante: pr. Tsilivi (Marg.); Cephalonia: pr. Lasi, Gerolaki, Akoli, Kontogionata, Lixuri, Livatho (Heldr.). — Jul. Nov. ♃.

b. Involucri phylla intima radiantia; pappus uniserialis.

α. Caulis monocephalus; capitula maxima, 8—12 cm. diametro, solitaria, saepe in rosula subsessilia.

2. **C. utzka** Hacq. pl. alp. carn. p. 9, t. 1; Haussk. symb. p. 119; Form. in Ver. Brünn 1896 p. 54, 1897 p. 40. — *C. acanthifolia* All. fl. ped. I. p. 156, t. 51; Ch. et B. fl. pelop. p. 57; Bois. fl. or. III. p. 448; Form. in D. bot. Monat. 1890 p. 21, in Ver. Brünn 1895 p. 29; Hal. Beitr. fl. Epir. p. 28, Beitr. fl. thessal. p. 14. — Exsicc.: Heldr. it. gr. thessal. IV. a. 1885.

Acaulis; foliis rosulatis, coriaceis, subtus incano-lanatis, pinnatifidis, lobis spinoso-dentatis; involucri phyllis exterioribus lanceolatis, apice margineque breviter spinosis, mediis nigricantibus, linearibus, pectinatim-spinosis, intimis multo longioribus, linearibus, acuminatis, sulfureis; receptaculi palearum fibris acutis, longioribus sub apice paulo crassioribus; corollis luteis; pappi pilis achenio duplo longioribus.

In herbidis saxosis regionis montanae, subalpinae et alpinae. Epirus: mt. Micikeli, in colle Ardomista pr. Janina (Form.), mt. Peristeri, pr. Chaliki (Hal.); Thessalia: mt. Oxya, Zygos (Haussk.), pr. Lepenica, Vendista, Malakasi, Said Pascha, Gisel Tepe, Dokimi, Mandra Hodza Koturi, mt. Baba, Ochsa Despot (Form.), Kastania (Hal.) in Pindo, mt. Chassia, mt. Kerfi pr. Diodendron, mt. Godaman, pr. Baraktari, Lokatamburia et Tafilvris in Olympo (Form.); Messenia: pr. Arcadia (Chaub.). — Jul. Sept. ♃.

3. **C. simplex** W. et K. pl. rar. hung. II. p. 164, t. 152; Form in Ver. Brünn 1896 p. 55, *v. thessala* (sed quomodo haec a typo discedit ex descriptione vix patet, nisi foliorum lobis angustioribus et involucri phyllis intimis parte inferiore puberulis). —

Caulescens; caule erecto, foliato; foliis herbaceis, glabris vel subtus arachnoideis, pinnatifidis, lobis spinoso-dentatis; involucri phyllis nigricantibus, exterioribus lanceolatis, pinnatifidis, spinosis, mediis linearibus pectinatim-spinosis, intimis multo longioribus, linearibus, acuminatis, supra candidis, subtus parte inferiori fusco-purpurascentibus; receptaculi palearum fibris apice evidenter clavato-incrassatis; corollis albidis; pappi pilis achenio duplo longioribus. — Differt a praecedente foliis subtus non incanis, colore phyllorum involucri intimorum et palearum fibris clavatis.

In mt. Godaman in Olympo (Form.). — Jul. Sept. ♃. N. v.

β. Caulis pleiocephalus; capitula mediocria, 4—5 cm. diametro.

× Radius involucri pallidus.

○ Involucri phylla externa radio breviora vel aequilonga.

4. **C. vulgaris** L. sp. p. 828; Form. in D. bot. Monat. 1890 p. 21. — *C. rigida* Form. in Ver. Brünn 1896 p. 53, 1897 p. 38 cum var. *pallida*, *sordida*, *humilis* et *othryana*. — Icon: Fl. dan. 1174. — Exsicc.: Form. pl. thessal. a. 1896.

Caule erecto, araneoso, 1—2 cephalo vel corymboso; foliis subtus araneosis, dentato-spinosis, inferioribus oblongo-lanceolatis, in petiolum attenuatis, superioribus lanceolatis, basi semiamplexicauli sessilibus; involucri phyllis externis oblongo-lanceolatis, margine et apice spinosis, mediis spinis nigricantibus ramosis pectinatim marginatis, intimis longioribus, linearibus, acuminatis, scariosis, albidis; receptaculi palearum fibris omnibus subulatis; corollis lutescentibus. —

In rupestribus montanis. Thessalia: pr. Malakasi, mt. Ghavellu, Said Pascha, Zygos, Gisel Tepe in Pindo, mt. Phlamburo in mt. Oxya, mt. Mavrika et Katasara in mt. Othrys (Form.). — Jul. Aug. ⊙.

[illegible] Involucri phylla externa radio longiora, eum interdum duplo superantia.

5. **C. acanthophylla** Haussk. symb. p. 119; Form. in Ver. Brünn 1896 p. 53, 1897 p. 38. — *C. vulgaris* Bois. fl. or. suppl. p. 305, quoad locum, non L. — *C. semiamplexicaulis* Form. in D. bot. Monat. 1890 p. 21. in Ver. Brünn 1895 p. 29. 1896 p. 53. ex autore ipso, non *C. semiamplexicaulis* Form. in ö. b. Z. 1890 p. 86, quae ex autore speciem aliam sistere dicitur. — Huc quoque ex descriptione: *C. dolopica* Form. in D. bot. Monat. 1897 p. 74. in Ver. Brünn 1897 p. 39. — Exsicc.: Heldr. it. gr. septentr. a. 1879.

Caule erecto, araneoso, 3—4 cephalo, corymboso, dense folioso; foliis subtus araneosis, pinnatifidis, rhachide angusta, segmentis lanceolatis, spinoso-dentatis; involucri phyllis externis foliaceis, lanceolatis, subtus araneosis, pinnatim et tenuiter spinosis, radium superantibus, mediis lineari-lanceolatis, margine et apice spinosis, intimis radiantibus, lineari-lanceolatis, acuminatis, scariosis, albido-lutescentibus, in parte inferiore purpurascente tenuiter ciliatis; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis, corollis purpurascentibus. — Differt a praecedente foliis longioribus, pinnatisectis, rhachide angusta 3—4 mm. tantum lata, spinis longioribus validioribus, phyllis involucri externis elongatis, radio longioribus et palearum fibris longioribus superne tenuiter fusiformibus. — Specimina originalia quidem non vidi, tamen plantam a Heldreich in mt. Chelidoni lectam, cum diagnosi bene congruentem, huc spectare censeo.

In apricis saxosis regionis montanae et subalpinae. Epirus: mt. Prosgoli (Form.); Thessalia: mt. Baba inter Krania et Klinovo (Haussk.), pr. Velitsena, mt. Ghavellu, pr. Kastri, Tafilvris in mt. Olympo, pr. Selicany in mt. Ossa, pr. Kalyvia, Kapurna (Form.), mt. Pelion supra Volo (Haussk.); Eurytania: mt. Chelidoni supra Mikrochorio (Held.). — Jul. Aug. ⊖.

6. **C. frigida** Bois. et Heldr. diagn. VI. p. 109, Fl. or. III. p. 450; Heldr. chlor. Parn. p. 21. — Exsicc.: Heldr. herb. norm. n. 516; Pichl. pl. gr. a. 1876.

Caule erecto, glaberrimo, 2—6 cephalo, congeste corymboso, dense folioso; foliis glaberrimis, pinnatifidis, rhachide angusta, segmentis lineari-lanceolatis, in spinas longas attenuatis; involucri phyllis externis foliaceis, lanceolatis, glabris, pinnatim et tenuiter spinosis, radium duplo superantibus, mediis lineari-lanceolatis, inermibus, subtus araneosis, intimis radiantibus, linearibus, acuminatis, scariosis, flavescentibus, ad medium usque ciliatis; receptaculi palearum fibris omnibus subulatis; corollis flavis. — A praecedente glabritie, phyllorum forma et palearum fibris omnibus subulatis; a formis *C. corymbosae* glabritie, radio pallide stramineo et palearum fibris non incrassatis discedens.

In saxosis regionis mediae et superioris mt. Parnassus, Kyllene et Taygetos (Heldr.). — Jul. Aug. ⊖.

×× Radius involucri luteus.

○ Involucri phylla externa radio breviora vel aequilonga.

7. **C. elegans** Heldr. in herb. gr. norm. n. 1647. —

Caule erecto, glabriusculo, corymboso, folioso; foliis glaucis, subtus araneosis, tandem glabratis, pinnatifidis, divaricatim spinosis, rhachide latiuscula, inferioribus lanceolatis, subpetiolatis, ceteris oblongo-lanceolatis, auriculato-amplexicaulibus; capitulis parvis; involucri phyllis lanceolatis, externis foliaceis, spinoso-dentatis, mediis brevibus, integris, araneosis, intimis radiantibus, lineari-lanceolatis, acutis, scariosis, glabris; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis; corollis flavis. — Sequenti affinis, ab ea glaucedine, foliis et phyllis angustioribus, profundius divisis et capitulis parvis, eis nuce avellanae subminoribus specifice discedit.

In arenosis maritimis et in alveo fluminis Nedon pr. Kalamata Messeniae (Zahn). — Jul. Aug. ⊙

8. **C. corymbosa** L. sp. p. 828; Sieb. avis p. 5, rem. p. 5; Ch. et B. exp. p. 243, Fl. pelop. p. 57; Marg. et R. fl. Zante p. 59; Raul. cret. p. 787; Bois fl. or. p. 449; Heldr. fl. cephal. p. 46; Form. in D. bot. Monat. 1890 p. 11; in Ver. Brünn 1895 p. 29, 1896 p. 54; Hal. in ö. b. Z. 1892 p. 371. — Icon: Rchb. t. 13. — Exsicc.: Sint. et Bornm. it turc. n. 1312; Sint it. thessal. n. 1192.

Caule erecto, parce araneoso, corymboso, rarius simplici, folioso; foliis viridibus, subtus araneosis, tandem glabratis, pinnatifido-dentatis, divaricatim spinosis, rhachide lata, inferioribus lanceolatis, subpetiolatis, ceteris oblongo-lanceolatis, auriculato-amplexicaulibus; capitulis mediocribus; involucri phyllis externis foliaceis, oblongo-lanceolatis, spinoso-dentatis, mediis brevibus, integris, lanceolatis, araneoso-lanatis, spinula terminatis, intimis radiantibus, lineari-lanceolatis, acutis, scariosis, glabris vel ciliatis; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis; corollis flavis.

In campis, ruderatis regionis inferioris et montanae. Epirus: pr. Prevesa, Agiotoma, Nikopolis, Philippiades, Ephemiada, Han Mescho, Janina, Sadovica, mt. Micikeli, in valle Dipotami (Form.); Thessalia: pr. Sermeniko (Sint.), Han Kuraneos, Velestinos, Kapurna, in valle Tempe, pr. Rapsani, Ligaria, Miluna (Form.) et Litochori in Olympo (Sint.); Peloponnesus (Chaub.): Messenia (Bois.); Creta (Sieb.): pr. Drakona, mt. Ida (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli, mt. Phalaris (Heldr.); sed loca plura probabiliter ad speciem sequentem cum nostra iterum confusam spectant. — Jul. Aug. ⊙

○○ Involucri phylla externa radio longiora.

9. **C. graeca** Heldr. et Sart. in Heldr. herb. norm. n. 423, Fl. Aegina p. 305, in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 246; Haussk. symb. p. 119. — *C. corymbosa* S. et S. pr. II. p. 158, Fl. gr. t. 837; Link in Linnaea IX. p. 581; Friedr. Reise p. 284; Fraas fl, class. p. 206. — *C. corymbosa v. graeca* Bois. fl. or. III. p. 449; Form. in Ver. Brünn 1896 p. 54, 1897 p. 40. —

Caule erecto, glabro vel parce araneoso, simplici vel corymboso, folioso; foliis glabrescentibus, pinnatifido-dentatis, divaricatim-spinosis, oblongo-lanceolatis lanceolatisve, inferioribus subpetiolatis, caeteris auriculato-amplexicaulibus; involucri phyllis externis foliaceis, patentissimis, longe spinoso-dentatis, capitulo duplo longioribus, mediis brevibus, integris, lanceolatis, araneoso-lanatis, spinula terminatis, intimis radiantibus, lineari-lanceolatis, acutis, scariosis, glabris vel ciliatis; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis; corollis flavis. — A praecedente involucri phyllis externis elongatis, uti jam Link l. c. monuit, specifice differre videtur. — Variat:

α. **platyrhachis.** — Caulis elongatus, superne ramosus; foliorum et phyllorum extimorum rhachis latiuscula, circa 10—5 mm. lata, ideo folia eis *C. corymbosae* similia. — Exsicc.: Orph. fl. gr. n. 792.

β. **actinobola** Heldr. herb. gr. n. 1648 pro sp. — Caulis saepe a basi ramosus, depressus; foliorum et phyllorum rhachis circa 8—4 mm. lata; capitula majora, phyllis plerumqne explanatis.

γ. **stenorhachis.** — Huc forsan: *C. corymbosa r. gralilis* Form. in D. bot. Mon. 1890 p. 21, in Ver. Brünn 1896 p. 54. — Caulis erectus, superne ramosus; foliorum et phyllorum extimorum rhachis angusta, 5—3 mm. lata. — Exsicc.: Heldr. herb. norm. n. 423.

In campis, saxosis regionis inferioris et montanae. Thessalia: mt. Baba, pr. Chaliki, Klinovo (Haussk.), Malakasi in Pindo, in valle Tempe (Form.); Attica: frequens pr. Athenas, in mt. Parnes usque in regionem abietinam adscendens; insula Aegina, Mykonos, Delos, Naxos (Heldr.), Rhenia (Tunt.); Peloponnesus (Sibth.): pr. Patras (Marg.), Nauplia (Fried.), mt. Malevo (Orph.). — Jul. Aug. ⊙

10. **C. Rothii** Heldr. et Sart. pl. exsicc. a. 1855. — *C. corymbosa v. Rothii* Bois. fl. or. III. p. 449. — Exsicc.: Heldr. reliqu. Orph. a. 1886.

Involucri phyllis extimis spinis elongatis valde ramosis munitis, mediis toto margine vel saltem apicem versus infra spinam terminalem breviter spinuloso-scabris, ceteris ut in var β. praecedentis.

In mt. Parnassus pr. Rachova (Guicc.) et in mt. Chelmos (Orph.). — Jul. Aug. ⊙

11. **C. Curetum** Heldr. pl. in ins. Creta aestate 1870 coll. — *C. corymbosa v. involucrata* Bois. fl. or. III. p. 449; Bald. viagg. Creta p. 66; vix *C. involucrata* Poir. voy. II. p. 234, quae ex DC. pr. VI. p. 547 capitulis majoribus et foliis minus incisis, quam in *C. corymbosa*, cui simillima, gaudet. — Exsicc.: Heldr. l. c.; Bald. it. cret. n. 158.

Glabrescens; caule erecto, remotiuscule foliato, 1—2 cephalo; foliis elongatis, coriacissimis, pinnatifido-dentatis, divaricatim spinosis, spinis validis, rhachi angusto multo longioribus munitis; involucri phyllis externis foliis similibus, capitulo 2—2 ½ plo longioribus, mediis

brevibus integris, lanceolatis, araneosis, spinula terminatis, intimis pallide luteis, radiantibus, lineari-lanceolatis, acutis, scariosis, glabris; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis; corollis flavis. — A praecedentibus caule 1—2 cephalo, remote foliato, foliis coriaceissimis, longe et valide spinosis, capitulis minoribus et involucri phyllis externis longissimis egregie differt.

In regione silvatica mt. Ida Cretae (Heldr.). — Jul. Aug. ⊙.

××× Radius involucri purpureus.

12. **C. lanata** L. sp. p. 828; S. et S. pr. II. p. 158; Sieb. avis p. 5; Ch. et B. exp. p. 244, Fl. pelop. p. 57; Marg. et R. fl. Zante p. 58; Weiss in z. b. G. 1869 p. 42; Raul. cret. p. 787; Bois. fl. or. III. p. 451; Held. fl. cephal. p. 46, Fl. Aegina p. 305; Form. in D. bot. Monat 1890 p. 21, in Ver. Brünn 1895 p. 29, 1896 p. 54, 1897 p. 40; Haussk. symb. p. 119. — Icon: Fl. gr. t. 836. — Exsicc.: Dörfl. fl. aeg. n. 49.

Araneoso-lanata; caule erecto, simplici vel corymbose ramoso; foliis oblongis vel lanceolatis, sinuato-pinnatifidis, spinosis, inferioribus in petiolum attenuatis, ceteris semiamplexicaulibus; involucri phyllis externis foliaceis, oblongo-lanceolatis, spinoso-dentatis, tomentosis, mediis brevioribus, lineari-lanceolatis, spina elongata terminatis, intimis scariosis, radiantibus, linearibus, glabris; receptaculi palearum fibris longioribus superne anguste fusiformi-incrassatis. — Radio purpureo ab omnibus distinctissima.

In arenosis, ruderatis, rupestribus regionis inferioris et montanae. Epirus: pr. Prevesa, Luros (Form.); Thessalia: pr. Karditza (Haussk.), Kapurna, Larissa, Velestinos, Aivali, Persufli, Pharsalus, mt. Pelion pr. Portaria, Mavro Mandila, Lamia (Form.); Euboea: pr. Kurbatzi (Wild); Aetolia: pr. Mesolongion (Nied.), Klissura (Samar.); Attica: pr. Munychia, insula Aegina (Heldr.); Corinthus (Hal.); Argolis: pr. Nauplia (Sart.), Epidaurus Limera (Chaub.); Laconia: pr. Monembasia, Zarex, promontorium Malea (Chaub.); Cyclades (Sibth.): Syra (Weiss), Naxos (Leon.); Creta: ad promontorium Grabusa, pr. Kissamos, Selino (Raul.), Canea (Weiss); Zante (Marg.); Cephalonia: pr. Piscardo, Lixuri (Heldr.); Corcyra: pr. Kanali, Kontokali (Form.). — Jun. Aug. ⊙.

40. Atractylis L. gen. n. 930.

1. **A. cancellata** L. sp. p. 830; Ch. et. B. exp. p. 244, Fl. pelop. p. 57; Marg. et R. fl. Zante p. 59; Friedr. Reise p. 277 et 278; Clem. sert. p. 56; Weiss in z. b. G. 1869 p. 43; Raul. cret. p. 787; Bois. fl. or. III. p. 452; Heldr. fl. ceph. p. 46; Haussk. symb. p. 121; Hal. in ö. b. Z. 1895 p. 412; Form. in Ver. Brünn 1896 p. 55, 1897 p. 40, in D. bot. Monat. 1898 p. 78; Heldr. fl. Aegina p. 305. — *Acarna cancellata* All. fl. ped. I. p. 154; S. et S. pr. II. p. 159. Icon: Fl. gr. t. 839. — Exsicc.: Heldr. herb. norm. n. 616.

In collibus aridis. Thessalia: pr. Pharsalus, Konisko et Neraida Phthiotidis (Form.); Sporadum insula Jura (Reis.); Attica: pr. Athenas,

in colle Lycebettus, Philopappus (Heldr.), mt. Hymettus (Clem.); insula Aegina et Hydra (Heldr.); Argolis (Sibth.): pr. Poros (Friedr.), Nauplia (Friedr.); Achaia: pr. Patras (Hal.); Laconia: pr. Vitilos, Chimova, Scardamula, ad promontorium Malea (Chaub.); Cycladum insula: Cythnos (Tunt.), Syra (Weiss), Tenos, Naxos (Chaub.); Creta: ad Akroteri, Canea (Raul.); Zante (Marg.); Cephalonia (Mazz.). — Apr. Jun. ⊙.

41. Staehelina L. gen. n. 938.

a. Folia utrinque viridia, glandulosa; achenia sericea.

1. **S. fruticosa** L. sp. ed. 2 p. 1286 (*Centaurea*); L. syst. nat. ed. 12, II. p. 538; Sieb. avis p. 5, rem. p. 6; Raul. cret. p. 786; Bois. fl. or. III. p. 456. — *Serratula fruticosa* Lam. ill. t. 666, f. 3. — *Hirtellina lanceolata* Cass. dict. 50 p. 441. — Icon: Jaub. et Sp. ill. t. 397. — Exsicc.: Sieb. pl. cret.; Bald. it. cret. alt. n. 180.

Basi suffruticosa, glabriuscula; ramis flexuosis, crebre foliatis, composite corymbosis, foliisque glandulis sessilibus flavescentibus dense obsitis; foliis oblongo-spathulatis vel lanceolatis, acutis, mucronatis, integerrimis, infimis in petiolum attenuatis, ceteris sessilibus, summis valde diminutis; capitulis breviter pedunculatis, oblongo-cylindricis, paucifloris; involucri resinoso-pruinosi phyllis arcte imbricatis, acutis, mucronulatis, ciliatis, exterioribus ovatis vel oblongis, intimis elliptico-linearibus; flosculis roseis; pappo subscabrido, achenio sublongiore.

In rupestribus montanis Cretae: ad Theodori in mt. Sphacioticis (Sieb.), mt. Aphendi Kristo et Sarakeno (Bald.) in mt. Lassiti (Raul.). — Jun. Sept. ♃.

b. Folia subtus albo-tomentosa, eglandulosa; achenia glabra.

2. **S. arborescens** L. mant. p. 111; S. et S. pr. II. p. 162; Sieb. avis p. 5, rem. p. 6, in Flora I. p. 272; Raul. cret. p. 786; Bois. fl. or. III. p. 456; Spreitz. in z. b. G. 1890 p. 296; Bald. viagg. Creta p. 66. — *Barbellina sericea* Cass. dict. 50 p. 440. — Icon: Fl. gr. t. 845. — Exsicc.: Heldr. pl. cret. n. 1596.

Fruticosa, ramis erectis vel adscendentibus, tomentosis, foliatis, apice dense corymbosis; foliis ovatis vel ovato-oblongis, obtusis vel acutiusculis; integerrimis, supra viridibus, glabriusculis, subtus argenteo-sericeis, inferioribus longiuscule petiolatis, superioribus subsessilibus; capitulis subsessilibus, oblongo-cylindricis, paucifloris; involucri tomentosi phyllis arcte imbricatis, acutis, mucronulatis, ab infimis ovato-triangularibus ad intima elliptico-linearia auctis; flosculis roseis; pappi setis inferne barbellatis, achenio striato duplo longioribus. — Frutex foliis amplis discoloribus pulcherrimus.

In rupibus regionis inferioris et montanae Cretae: pr. Enneachoria, ad promontorium Grabusa (Raul.), in mt. Sphaciotis (Sibth.) pr. Theriso, Hagia Rumeli, Samaria, Aradhena, Lakus et Nipros, inter Selia et Hagios Joannes, Pharangi Khordaliotikon, mt. Kophino, insula Dia (Raul.). — Jul. Aug. ♃.

3. **S. unifloscculosa** S. et S. pr. II. p. 162; Bois. fl. or. III. p. 456; Heldr. chlor. Parn. p. 21; Hal. Beitr. fl. Epir. p. 28; Bald. riv. coll. bot. Alb. 1895 p. 49. — Icon: Fl. gr. t. 896. — Exsicc.: Heldr. herb. norm. n. 390 et 945; Orph. fl. gr. n. 100, in Magn. fl. sel. n. 2503, in Baen. herb. europ. n. 5526; Sint. et Bornm. it. turc. n. 1821; Bald. it alb. n. 144, it. alb. epir. III. n. 166; Dörfl. herb. norm. n. 3543.

Fruticosa; ramis erectis vel adscendentibus, foliatis, junioribus tomentosis, apice simpliciter vel composite corymbosis; foliis oblongis, acutis, breviter petiolatis, glanduloso-serratis, supra viridibus, subtus adpresse tomeutosis; capitulis subsessilibus, anguste cylindricis, 1—2 floris; involucri glabriusculi phyllis arcte imbricatis, acutis, mucronulatis, ciliatis, ab infimis oblongis ad intima lanceolata auctis; flosculis roseis; pappi setis inferne scabridis, achenio striato duplo longioribus. — Foliis multo minoribus, serrulatis, subtus non sericeo-nitentibus, capitulis multo minoribus, 1—2 floris, involucri phyllis glabriusculis, rubiginosis a praecedente diversissima. — Species elegans. —

In rupestribus regionis montanae et subalpinae. Epirus: in mt. Acroceraunicis, pr. Suli distr. Paramythia, pr. Krapsi distr. Janina, pr. Syraku (Bald.) et Kalarrytes ad radices mt. Peristeri, pr. Melisurgi ad radices mt. Tsumerka (Hal.); Thessalia: mt. Olympus pr. Hagios Dionysios (Heldr.); Euboea: mt. Dirphys (Leon.); Aetolia: mt. Tymphrestus (Nicolaides), mt. Korax (Oertzen); mt. Parnassus (Sibth.); Achaia: mt. Chelmos supra Klukines (Orph.). — Jul. Sept. ♃.

Obs. *S. dubia* L. sp. p. 840, in Ch. et B. fl. pelop. p. 56 in Peloponneso, inter Pyrgos et Androvista certe ex confusione indicatur.

10 Tribus. **CARDUINEAE** Cass. tabl. p. 9.

42. **Lappa** Tourn. inst. p. 450.

1. **L. major** Gaertn. fruct. II. p. 379, t. 162, f. 3; Raul. cret. p. 793; Bois. fl. or. III. p. 457; Heldr. fl. cephal. p. 46; Form. in D. bot. Monat. 1890 p. 18, in Ver. Brünn 1895 p. 26, 1896 p. 47, 1897 p. 35. — *Arctium lappa* L. sp. p. 816 p. p.; S. et S. pr. II. p. 147; Dallap. prosp. p. 109; Fraas fl. class. p. 203. — *A. majus* Bernh. verz. Erf. 1800 p. 154; Haussk. symb. p. 121. —

Caule erecto, ramoso, lanuginoso-puberulo; foliis petiolatis, ovatis vel cordato-ovatis, subtus lanuginoso-tomentellis; capitulis subcorymbosis, glabris; involucri phyllis subulatis, uncinatis, flosculos superantibus; flosculis purpureis; acheniis maculatis, superne praesertim transverse rugosis.

In ruderatis, ad vias, sepes regionis inferioris et montanae. Epirus: pr. Luros, Janina, Kacka, Han Balduma (Form.); Thessalia: mt. Zygos, ad monasterium Korona (Haussk.), pr. Han Kriavris in valle Penei, Trikala, Velestinos, Pharsalus, Koturi, mt. Phlamburo in mt. Oxya, mt. Cuka in mt. Chassia, pr. Tafilvris in Olympo, Lamia Phthiotidis (Form.);

mt. Korax Aetoliae, mt. Dirphys Euboeae (Fraas); Archipelagus (Sibth.); Creta: pr. Enneachoria (Raul.); Cephalonia (Dall.). — Jul. Aug. ⊙ h. v.

2. **L. minor** Bernh. verz. Erf. 1806 p. 154; Schk. Handb. III. p. 49, t. 227; Haussk. symb. p. 121 (*Arctium*); DC. fl. fr. IV. p. 77; Bois. fl. or. III. p. 457. — Exsicc.: Heldr. pl. fl. hellen. a. 1890 et 1899.

Differt a praecedente capitulis fere duplo minoribus, subracemosis, involucri phyllis flores vix aequantibus, acheniis non nisi basi leviter transverse rugulosis.

In locis ubi praecedens. Thessalia: ad monasterium Korona (Haussk.); Euboea: pr. Kastaniotissa (Heldr.), Aetolia: mt. Arapocephalo (Samar.); Achaia: mt. Kyllene (Heldr.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.) pr. Megali Anastasova (Zahn). — Jul. Aug. ⊙

43. Carduus L. gen. n. 925.

a. Capitula globosa vel ovoidea.

α. Capitula solitaria, magna, 4—8 cm. diametro, globoso-depressa; involucri phylla media supra basin ovatam vel oblongam, adpressam constricta, supra stricturam in appendicem lanceolatam vel subulatam, patulam vel refractam abeuntia.

1. **C. leiophyllus** Petr. add. fl. Nyss. p. 105. — Huc probabiliter pertinet ex descriptione: *C. pindicolus* Haussk. in Mitth. thür. bot. Ver. 1887 p. 86, symb. p. 123. — Exsicc.: Orph. herb. n. 3704.

Caule erecto, simplici vel ramoso, alis angustis lobatis spinosis marginato, inferne glabro, apice plus minus longe nudo, tomentoso; foliis oblongo-lanceolatis lanceolatisve, sinuato-pinnatifidis, glabris vel ad nervos parce puberulis, laciniis trifidis vel dentatis, dentibus spinosis; capitulis erectis vel cernuis, involucri glabri phyllis exterioribus in appendicem latissimam, basi ad 9 mm. latam, eis latiorem et sublongiorem, patulam, in spinam rigidulam abruptiuscule attenuatam, abeuntibus; flosculis purpureis. — Differt a *C. nutante*, cui valde affinis, glabritie, capitulis saepius majoribus et praesertim phyllorum involucri appendicibus abbreviatis, latissimis, spina breviori terminatis.

β. **vestitus.** — Foliis subtus araneoso-lanatis, involucri phyllorum appendicibus brevioribus, brevissime acuminatis, spinula tenui brevi terminatis. — Species forsan propria. — Exsicc.: Sint. it thessal. n. 476 sub *C. pindicolus*.

In regione montana et alpina. Thessalia: mt. Ghavellu in Pindo (Haussk.), mt. Olympus (Orph.); — β. ad Guvelsi pr. Kalabaka Thessaliae (Sint.). — Maio, Aug. ⊙

2. **C. nutans** L. sp. p. 821; S. et S. pr. II. p. 149; Ch. et B. exp. p. 240, Fl. pelop. p. 50; Bois. fl. or. III. p. 515 p. p.; Form. in Ver. Brünn 1895 p. 28. — Exsicc.: N. v.

Caule erecto, simplici vel ramoso, alis angustis lobatis spinosissimis marginato, araneoso, apice plus minus longe nudo, tomentoso; foliis oblongo-lanceolatis lanceolatisve, sinuato-pinnatifidis, araneosis, laciniis trifidis vel dentatis, dentibus spinosis; capitulis erectis vel cernuis; involucri plus minus araneosi phyllis exterioribus in appendicem lanceolatam, basi 3—5 mm. latam, eis latiorem et longiorem, patulo-reflexam, in spinam rigidam sensim attenuatam abeuntibus, flosculis purpureis.

β. **brachycentros** Haussk. symb. p. 123. — Caule apice longe nudiusculo; involucri glabri phyllis glabris vel intus subciliatulis, in appendicem reflexam, spina breviori terminatam abruptius contractis. — Exsicc.: Held. it thessal. IV. a. 1885.

γ. **Sporadum.** — *C. nutans* Hal. in ö. b. Z. 1895 p. 460, non L. — *C. macrocephalus* Hal. in ö. b. Z. 1897 p. 95, non Desf., qui phyllis involucri latissimis insignis. — Caule apice longe nudiusculo; foliis pinnatipartitis, laciniis longius spinosis; capitulis maximis, involucri glabri phyllis in appendicem stricte patentem, angustam, valde carinatam, eis multo longiorem, spina elongata valida revoluta terminatam, sensim abeuntibus. — Species forsan propria. — Exsicc.: Heldr. et Hal. fl. sporad. a. 1896. —

In ruderatis, sterilibus regionis inferioris et montanae. Epirus: in valle Dipotami (Form.); in Peloponnesi agris (Sibth.); — β. Thessalia: pr. Rivali, Pharsalus (Haussk.); — γ. Sporadum insula Jura (Leon). — Maio, Aug. ⊙

3. **C. taygeteus** Bois. et Heldr. diagn. ser. 2, III. p. 42; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Epir. p. 28, Beitr. fl. Achaia p. 25. — *C. nutans* Bois. fl. or. III. p. 516 p. p. — *C. nutans v. taygeteus* Heldr. chlor. Parn. p. 21. — Exsicc.: Heldr. herb. norm. n. 709.

Caule erecto, simplici vel ramoso, alis angustis lobatis spinosissimis marginato, araneoso, apice plus minus longe nudo, tomentoso; foliis oblongo-lanceolatis lanceolatisve, sinuato-pinnatifidis, parce araneosis, laciniis trifidis vel dentatis, dentibus spinosis, capitulis erectis vel cernuis; involucri glabriusculi vel araneosi phyllis exterioribus in appendicem lanceolato-subulatam, 2—4 mm. latam, eis vix latiorem, longiorem, refractam, in spinam validam revolutam sensim attenuatam, abeuntibus; flosculis purpureis. — Differt a praecedente, ad quam formis intermediis transire videtur, involucri phyllis in spinas refractas tenues, nec lanceolates patulo-reflexas abeuntibus.

β. **parnassicus.** — Involucri dense araneoso-lanati phyllis angustioribus, refractissimis, spina valde elongata rigida terminatis. — Exsicc.: Doerfl. fl. gr. n. 214.

In lapidosis regionis mediae. Epirus: mt. Tsumerka, Peristeri (Hal.); Doris: mt. Kiona (Hal.); mt. Parnassus (Heldr.); Peloponnesus: mt. Olenos (Hal.) Panachaicon, Taygetos (Heldr.). — Jul. Aug. ⊙

β. Capitula mediocria, 2—4 cm. diametro, subglobosa vel ovata; involucri phylla anguste lanceolato- vel lineari-subulata, non constricta.

× Perennes; folia praesertim superiora parenchymate fere destituta, longe et valide spinosa.

4. **C. armatus** Bois. et Heldr. diagn. VI. p. 104. — *C. cronius* Bois. fl. or. III. p. 515 p. p. — Exsicc.: Held. fl. gr. exs. a. 1855; Bald. it alb. alt. n. 142; Dörfl. fl. gr. n. 213.

Multiceps; caulibus erectis, simplicibus, glabris vel parce araneosis, ad apicem usque foliosis, alatis, alis in lobos lanceolatos, longe spinosos partitis; foliis glabriusculis glabrisve, in segmenta anguste et longe lanceolata, longe spinosa, pinnatipartitis; capitulis subsessilibus, foliis summis saepe superatis, approximatim corymbosis vel racemum thyrsoideum foliosum formantibus; involucri glabrescentis phyllis erectis, a basi anguste lanceolata longe subulato-spinosis, externis vix brevioribus, intimis subscariosis, longe acuminatis; flosculis roseis. — Foliis laete viridibus, fere substantia foliacea orbatis, spinis flavis, acerosis, tenuibus, ad 4 cm. longis dense armatis, insignis.

β. **cronius** Bois. et Heldr. diagn. VI. p. 105; Bois fl. or. III. p. 515 p. p., suppl. p. 307; Hal. in z. b. G. 1888 p. 759, in ö. b. Z. 1897 p. 285; Heldr. chlor. Parn. p. 21; Form. in Ver. Brünn 1896 p. 50; pro sp. — Caulis et folia araneoso-cana, subtus interdum niveolanata; phylla externa intimis saepius manifeste breviora. — Specimina extrema a formis fere omnino glabris typi valde recedunt. — Exsicc.: Heldr. herb. norm. n. 226; Orph. fl. gr. n. 466 et 796; Dörfl. fl. gr. n. 425.

In regione abietina et superiori, gregarie. Epirus: mt. Cika in monte Acroceraunicis (Bald.); Thessalia: mt. Dokimi (Form.), Olympus (Heldr.); Euboea: mt. Dirphys (Heldr.); Aetolia: mt. Chelidoni, Tymphrestus (Heldr.), Korax (Tunt.); mt. Kiona (Tunt.), Parnassus (Guicc.); Achaia: mt. Kyllene, Klokos (Orph.), Chelmos (Leon.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.). — Jul. Aug. ♃.

×× Biennes; folia parenchymatosa, minus valide spinosa.

○ Caulis ramosus, pleiocephalus; capitula solitaria vel 2—3 aggregata, pedunculis crispatis spinosis suffulta.

5. **C. acanthoides** L. sp. p. 821; S. et S. pr. II. p. 149; Raul. cret. p. 792. — Icon: Jacq. austr. t. 249. — Exsicc.: N. v.

Caule erecto, araneoso, ad apicem usque alis angustis, in lobulos crebre spinosos partitis marginato; foliis glabris vel subtus an nervos pilosulis, sinuato-pinnatifidis, segmentis inciso-dentatis, dentibus spinuloso-ciliatis, spinaque longiore terminatis; capitulis plerumque solitariis, erectis; involucri glabriusculi phyllis lineari-subulatis, erecto-patulis, rarius subsquarrosis, in spinam pungentem desinentibus; flosculis purpureis.

β. **thessalus** Bois. et Heldr. diagn. ser. 2. III. p. 46; Form. in D. bot. Monat. 1890 p. 20; pro sp.; Bois. fl. or. III. p. 518; Hal.

Beitr. fl. Thessal. p. 14; Haussk. symb. p. 124; Form. in Ver. Brünn 1895 p. 36, 1896 p. 50, 1897 p. 36. — Capitula brevius pedunculata, numerosiora, magis approximata, involucri phylla longius acuminata; planta interdum canescens. — Exsicc.: Orph. fl. gr. n. 795; Sint. it. thessal. n. 1193.

γ. **tymphaeus** Form. in Ver. Brünn 1896 p. 50, 1897 p. 36. — *C. armatus* Hal. Beitr. fl. Epir. p. 28; Sint. it. thessal. n. 1133 (f. *canescens*, foliis subtus araneosis). — Foliorum segmenta numerosiora, augustiora, elongata, crebre et longe spinosa, cetera ut in β. — Exsicc.: Hal. it. gr. II. a. 1893; Sint. it. thessal. n. 1133.

In campis, herbidis regionis inferioris in alpinam usque adscendens. Indicatur in Peloponneso et in Creta (Sibth.); — β. Epirus: pr. Janina (Form.); frequens in Thessalia a Pindo, per planitiem totam, usque ad Olympum; — γ. Epirus: mt. Tsumerka, Peristeri (Hal.); Thessalia: mt. Said Pascha, Zygos, Dokimi, Karava (Form.), Ghavellu (Sint.). — Jun. Sept. ⊙.

6. **C. crispus** L. sp. p. 821; S. et S. pr. II. p. 149; Ch. et B. exp. p. 241, Fl. pelop. p. 56. — Icon: Fl. dan. t. 621.

Caule erecto, superne lanato, ad apicem usque alis angustis, denticulatis, breviter spinulosis marginato; foliis supra sparsim pilosis, subtus lanato-tomentosis, oblongis, saepius parum profunde sinuato-lobatis, denticulatis, minute spinuloso-ciliatis, summis lineari-lanceolatis, saepius indivisis; capitulis solitariis vel saepius 2—3 aggregatis; involucri glabriusculi phyllis lineari-subulatis, breviter spinescentibus, exterioribus recurvo-patulis, intimis erectis; flosculis purpureis.

In Peloponneso (Sibth.). — Jul. Aug. ⊙. N. v.

○○ Caulis simplex, monocephalus vel in ramos virgatos, monocephalos, superne nudos partitus.

7. **C. hamulosus** Ehrh. Beitr. VII. p. 166; Haussk. symb. p. 123. — Icon: W. et K. pl. rar. hung. t. 233.

Caule erecto, araneoso, anguste alato, alis in lacinias minutas, ovatas, minute spinulosas, partitis; foliis glabriusculis vel subtus araneosis, in lacinias rotundatas vel oblongas, obtusas, minute spinuloso-dentatas pinnatifidis; capitulis majusculis, solitariis, subglobosis; involucri glabri phyllis anguste lineari-subulatis, tenuiter mucronatis, intermediis recurvatis; flosculis purpureis.

In fruticetis regionis superioris mt. Ghavellu in Pindo dolopico (Haussk.). — Jul. Aug. ⊙. N. v.

Obs. *C. intercedens* (*hamulosus* × *pindicolus*) Haussk. symb. p. 124 indicatur ab auctore in mt. Ghavellu. — Habitu *C. hamuloso* propius accedere dicitur, sed caules crassiores, latius alatae, alis longius sed parcius spinosae, folia et saepius capitula majora, involucri phylla latiora sunt. N. v.

8. **C. candicans** W. et K. pl. rar. hung. I. p. 85, t. 83; Bald. riv. coll. bot. Alb. 1896 p. 68. — *C. collinus* Form. in Ver. Brünn

1897 p. 36, non W. et K., quae pedunculis fere ad capitulum crispato-alatis, involucri glabriusculi phyllis exterioribus patulis parum discedit. — Exsicc.: Form. pl. thessal. a. 1896; Bald. it. alb. epir. IV. n. 177.

Caule erecto, araneoso, anguste alato, alis in lacinias minutas, ovatas, minute-spinulosas partitis; foliis supra araneosis, subtus cano-tomentosis, in lacinias ovatas, spinuloso-dentatas vel lobatas pinnatifidis; capitulis majusculis, solitariis, subglobosis; involucri araneosi phyllis lineari-subulatis, tenuiter mucronatis, adpressis patentibusque; flosculis purpureis. — Praecedenti habitu valde affinis, ab eo praesertim foliis bicoloribus, supra viridibus, subtus cano-tomentosis et pedunculis longius nudis discrepat.

In herbidis montanis, Epirus: mt. Kuruna supra Vonicko distr. Ljaskovik (Bald.); Thessalia: mt. Cuka in mt. Chassia (Form.). — Jul. Aug. ⊙.

b. Capitula oblonga, subcylindrica.

α. Acheniorum umbo crasse stipitatus, non angulatus.

9. **C. pycnocephalus** L. sp. ed. 2, p. 1151; Ch. et B. exp. p. 240, Fl. pelop. p. 56; Weiss in z. b. G. 1869 p. 44; Bois. fl. or. III. p. 520; Heldr. fl. cephal. p. 46, Fl. Aegina p. 305, Chlor. Mykon. p. 246; Spreitz. in z. b. G. 1887 p. 663; Hal. Beitr. fl. Epir. p. 28; Haussk. symb. p. 124; Form. in Ver. Brünn 1895 p. 28, 1897 p. 36. — *C. albidus* M. a. B. fl. taur. cauc. II. p. 269, qui sec. Bois. l. c. involucro magis tomentoso tantum discedit. — *C. tenuiflorus* S. et S. pr. II. 149; Ch. et B. exp. p. 241, Fl. pelop. p. 56; Marg. et R. fl. Zante p. 60; Fraas fl. class. p. 203; vix Curt. lond. fasc. t. 55, qui ramis latius et ad apicem usque alatis, capitulis numerosioribus angustioribus, involucri glabri phyllis angustioribus, exterioribus magis patulis, interioribus longius acuminatis, flosculos superantibus discedit et quem e Graecia nondum vidi. — *C. tenuiflorus v. albidus* Friedr. Reise p. 269. — *C. tenuiflorus v. pycnocephalus* Friedr. Reise p. 273 et 278. — *C. pycnocephalus v. albidus* Bois. fl. or. III. p. 521; Heldr. fl. Aegina p. 305. — Huc quoque probabiliter: *C. arabicus* Friedr. Reise p. 277; Weiss in z. b. G. 1869 p. 44; vix Jacq. coll. I. p. 56, qui involucri phyllis in spinulam subabrupte apiculatis differe videtur. — Icon: Jacq. hort. vind. I. t. 44. — Exsicc.: Heldr. pl. fl. hellen. a. 1886, 1892 et 1900; Baen. herb. europ. n. 9227; Sint. it. thessal. n. 412; Dörfl. pl. cret. n. 60.

Caule erecto, araneoso-lanuginoso, simplici vel ramoso, fere ad, apicem usque alato-spinoso; foliis subtus canescentibus vel tomentosis, sinuatis pinnatifidisque, laciniis ovatis, angulato-dentatis, spinosis; capitulis in caule ramisque terminalibus, oblongis, 2 pluribusve congestis, lateralibus sessilibus; involucri plus minus araneosi phyllis erectis, lanceolatis, in spinulam attenuatis, flosculos subaequantibus, flosculis purpureis, rarius albidis.

In ruderatis, herbidis regionis inferioris passim per totam Graeciam. Mart. Jul. ⊙

10. C. **acicularis** Bert. ann. stor. nat. Bologn. 1829 I. 1, p. 274. — *C. neglectus* Ten. fl. nap. IV. p. 118, t. 187; Marg. et R. fl. Zante p. 60. — Exsicc.: Heldr. pl. fl. hellen. a. 1877 et 1899.

Caule erecto, araneoso-lanuginoso, ramoso, alato-spinoso, ramis superne nudis; foliis subtus canescentibus vel tomentosis, sinuatis pinnatifidisque, laciniis ovatis, angulato-dentatis, spinosis; capitulis solitariis subsolitariisque, involucri plus minus araneosi phyllis erecto-patulis, lineari-subulatis, in spinulam attenuatis, flosculos aequantibus; flosculis purpureis. — Differt a praecedente capitulis solitariis, pedunculis longe nudis suffultis et involucri phyllis angustioribus, longe subulatis.

In herbidis, arenosis regionis inferioris. In regione litorali pr. Corinthum, ad fossas pr. Manolas Elidis (Heldr.); Zante: mt. Scopo (Marg.); et probabiliter alibi. — Maio, Jun. ⊙

β. Acheniorum umbo tenuiter stipitatus, radiatim sulcatus.

11. C. **marmoratus** Bois. et Heldr. diagn. ser. 2, III. p. 43, Fl. or. III. p. 521.

Caule erecto, araneoso, ramoso, fere ad apicem usque alato-spinoso; foliis supra albo-marmoratis, subtus araneoso-canescentibus, sinuatis pinnatifidisque, laciniis ovatis, angulato-dentatis, spinosis; capitulis in caule ramisque terminalibus, oblongis, 2 pluribusque congestis, lateralibus sessilibus; involucri plus minus araneosi phyllis erecto-patulis, lanceolatis, in spinulam attenuatis, flosculos aequantibus; flosculis pnrpureis. — Differt a *C. pgcnocephalo*, cui affinis, spinis validis, saepe pollicaribus alarum et phyllorum superiorum et praesertim acheniorum umbone sulcato. — An a *C. argyroa* Biv. specifice distinctus?

α. **typicus.** — Alis ramorum angustis foliisque longe et valide spinosis. — Exsicc.: Heldr. pl. fl. hellen. n. 190.

β. **intermedius.** — Alis ramorum angustis, foliisque brevius et minus valide spinosis. — Exsicc.: Heldr. herb. norm. n. 1446 sub *C. argyroa.*

γ. **Halácsyi** Heldr. herb. n. 1258 pro sp. — Alis ramorum latiusculis, foliisque breviter spinosis, spinis mollioribus. — Exsicc.: Heldr. l. c.

In herbidis, ad vias, olivetis regionis inferioris. Messenia: pr. Diabolitzi, Pylos (Heldr.); — β. pr. Kalamata (Zahn); — γ. Achaia: pr. Patras (Hal.), Psathopyrgos (Heldr.). — Maio, Jun. ⊙

12. C. **argentatus** L. mant. p. 288; Haussk. symb. p. 125. — Icon: Jacq. hort. vind. II. t. 192. — Exsicc.: Heldr. it thessal. IV. a. 1885.

Caule erecto, araneoso-lanuginoso, ramoso, anguste alato-spinoso, ramis superne nudis; foliis subtus canescentibus vel tomentosis sinuatis pinnatifidisque, laciniis ovatis, dentatis-spinulosis; capitulis solitariis subsolitariisque; involucri araneosi phyllis erecto-patulis, lineari-subulatis,

in spinulam attenuatis, flosculos aequantibus: flosculis purpureis. — *C. aciculari* simile a quo praesertim umbone tenuiter stipitato angulato-sulcato distinctus; a formis praecedentis ramis superne longe nudis et involucri phyllis angustis discedit.

In neglectis Thessaliae pr. Aivali, Orman Magula, Pharsalus, in consortio *C. pycnocephali* (Haussk.). — Jun. Jul. ⊙

44. Cirsium Tourn. inst. p. 447.

Dispositio specierum:

A. Corollae limbus ad medium 5 fidus; pappus corolla brevior.

1. Sectio. *Epitrachys* DC. in Duby bot. gall. I. p. 286. — Folia supra spinuloso-strigosa; achenia compressa, epicarpio membranaceo.

a. Folia basi amplexicauli sessilia, non decurrentia.

α. Involucri spinae sub apice spathulato-dilatatae, pectinato-spinulosae vel serrulatae.

× Capitula magna, 5—8 cm. lata.

1. C. spathulatum Mor. **2. C. ligulare** Bois. **3. C. armatum** Vel.

×× Capitula parva, nucis magnitudinem non excedentia.

4. C. Lobelii Ten. **5. C. morinaefolium** Bois. et Heldr.

β. Involucri spinae sub apice non dilatatae, minute serrulatae.

6. C. Heldreichii Hal. **7. C. leucocephalum** (Willd.).

b. Folia caulina decurrentia.

8. C. lanceolatum (L.). **9. C. italicum** (Savi).

2. Sectio. *Chamaeleon* DC. l. c. — Folia supra non strigosa; achenia compressa, epicarpio membranaceo.

a. Folia non decurrentia.

α. Flores purpurei.

× Folia subtus albo-tomentosa; involucri phylla in spinam validam, patulam, phyllo ipso multo longiorem, flosculos subsuperantem abeuntia.

10. C. cynaroides (Lam.).

×× Folia glabra vel subtus araneoso-canescentia; involucri phylla mucronata vel breviter spinulosa.

○ Folia pinnatipartita.

11. C. appendiculatum Griseb.

○○ Folia breviter lobulata.

12. C. tymphaeum Haussk. **13. C. pauciflorum** (W. et K.).

β. Flores ochroleuci vel albi.

14. C. oleraceum (L.). **15. C. candelabrum** Griseb.

b. Folia caulina decurrentia.

16. C. creticum (Lam.). **17. C. pindicolum** Haussk.

3. Sectio. *Lamyra* Cass. dict. 25 p. 218 pro gen.: Bois. fl. or. III. p. 524. — Folia supra non strigosa, achenia globosa, ovata vel pyriformia, non compressa, epicarpio duro.

a. Capitula magna; involucri phylla non gibbosa.

18. C. afrum (Jacq.).

b. Capitula parva; involucri phylla exteriora ad spinae basin internam gibbosa.

19. C. strictum (Ten.). **20. C. stellatum** (L.).

B. Corollae limbus ad basin 5partitus; pappus corolla demum longior.

4. Sectio. *Cephalonoplos* Neck. elem. p. 68. — Folia supra non strigosa; achenia lineari-oblonga, compressa, epicarpio membranaceo.

21. C. arvense (L.).

A. Corollae limbus ad medium 5 fidus; pappus corolla brevior.

1. Sectio. *Epitrachys* DC. in Duby bot. gall. I. p. 286. — Folia supra spinulose-strigulosa; achenia compressa, epicarpio membranaceo.

a. Folia caulina basi amplexiculi sessilia, non decurrentia.

α. Involucri spinae sub apice spathulato-dilatatae, pectinato-spinulosae vel serrulatae.

× Capitula magna, 5—8 cm. lata.

1. C. **spathulatum** Moretti pl. ital. dec. III. p. 6. *(Cnicus)*; Gaud. fl. helv. V. p. 202. — *C. decussatum* Janka in Linneaea XXX p. 582; Bois. fl. or. III. p. 529; Heldr. chlor. Parn, p. 21; Form. in D. bot. Monat. 1890 p. 21 cum?. — *Cnicus ferox* S. et. S. pr, II. p. 152; Fraas fl. class. p. 204; non L. sec. Bois. l. c. —? *Carduus ferox* Ch. et B. fl. pelop. p. 56. — Exsicc.: Sint. et Bornm. it. turc. n. 1939.

Caule erecto, ramoso, araneoso; foliis supra viridibus, subtus albo-lanuginosis, in lacinias lanceolatas, ad basin bipartitas, in spinas validas abeuntes, pinnatipartitis; capitulis globosis, foliis summis flores non superantibus bracteatis; involucri araneoso- lanati phyllis lineari-subulatis, recurvis vel patentibus, infra spinam dilatatis, lateraliter plus minusve spinulosis, in spinulam tenuem attenuatis; flosculis purpureis. — *C. eriophoro* affine, a quo involucri phyllis apice dilatatis et spinulosis differt.

β. **Parnassi.** — Differt a typo capitulis minoribus, valde araneoso-lanatis, foliis summis longioribus, capitulum saepius longe superantibus bracteatis, foliorum spinis longioribus validioribus. — An species propria?. — Exsicc.: Hal. it gr. a. 1888; Dörfl. fl. gr. n. 204.

In regione media. Thessalia: pr. Velitsena in Pindo (Form.), mt. Olympus (Heldr.); Aetolia: mt. Tymphrestus (Samar.); mt. Parnassus (Heldr.); Peloponnesus: inter Pyrgos et Olympia, si *Carduus ferox* Ch. et B. huc pertineat; — β. mt. Parnassus (Hal.). — Jul. Aug. ⊙

Obs. Ad hanc, vel ad sequentes spectare videtur *C. eriophorum* Form. in D. bot. Mon. 1890 p. 20 non (L.), quod in Thessalia pr. Kastania, Velitsena, Koturi, Chaliki, mt. Zygos, pr. Han Kriavris in valle Penei, pr. Spilia et Selicany in mt. Ossa indicatur.

2. C. **ligulare** Bois. fl. o.r III. p. 529. — *C. Pelii* Form. in Ver. Brünn 1896 p. 51, quod meo sensu a planta thracica nullo modo differt. — *C. odontolepis* Heldr. in Sitzungsb. Acad. Wiss. Berlin 1883 p. 4; Haussk. symb. p. 121; non Bois. voy. esp. p. 362 t. 110,

quod foliis supra grosse, sed minus crebre spinosis, etiam supra plus minus araneosis, foliis involucrantibus numerosis, magnis, capitulo adpressissimis, eo valde interdum fere duplo longioribus et involucri phyllis virescentibus, superne minus dilatatis eximie discedit. — Exsicc.: Heldr. it. thessal. n. 62, Sint. it. thessal. n. 1334.

Caule erecto, ramoso, araneoso, foliis supra viridibus, subtus albolanuginosis, in lacinias lanceolatas, ad basin bipartitas, in spinas validas abeuntes pinnatipartitis; capitulis globosis, foliis summis, involucrum saepe aequantibus bracteatis; involucri parce lanati phyllis laxe erectis, in ligulas lineares atro-purpurascentes, spinuloso-denticulatas, spina terminatas, ab inferioribus ad interiora elongatas, abeuntibus; flosculis purpureis. — Praecedenti affinis, a qua spinis foliorum validioribus et praesertim involucri phyllis erectis, in ligulam latam longam, spina validiori terminatam abeuntibus egregie differt. — Formas intermedias nunquam observavi.

In regione media et superiori mt. Pelion Thessaliae (Heldr.); *C. Pelii* indicatur a Form. in Ver. Brünn p. 37 praeterea pr. Konisko, mt. Cuka, pr. Phlambures et mt. Mitrica in mt. Chssaia, pr. Katasara in mt. Othrys, et (var. *angustilobum*) in mt. Phlamburo in mt. Oxya, an hae loca revera ad *C. ligulare* pertineant, dijudicare nequeo, cum plantam solam pr. Konisko lectam vidi, quae tamen ad speciem sequentem spectat. — Jul. Sept. ⊙

3. **C. armatum** Vel. in Sitzungsb. böhm. Ges. Wiss. 1888 p. 52. — *C. horridum* Form. in Ver. Brünn 1895 p. 28; non M. a. B. fl. taur. cauc. II. p. 278 (*Cnicus*), Bois. fl. or. III. p. 537. — *C. validum* Form in Ver. Brünn 1896 p. 51, cum var. *montanum et macrocephalum*, 1897 p. 37. — *C. Pelii* Form. in Ver. Brünn 1897 p. 37 saltem p. p. — Huc probabiliter: *C. odontolepis* Bois. fl. or. III. p. 529, non voy. esp. p. 362, quod foliis subtus virescentibus, supra spinulis longioribus strigulosis, foliis involucrantibus longioribus et capitulis densius araneosis a planta orientali etsi parum, tamen constanter differre videtur. — Exsicc.: Form. pl. Thessal. a. 1896; Sint. it. thessal. n. 1131.

Caule erecto, ramoso, araneoso; foliis supra viridibus, subtus cinereolanuginosis, in lacinias lanceolatas, ad basin bipartitas, in spinas validas abeuntes pinnatipartitis; capitulis globosis, foliis summis flores superantibus involucratis; involucri araneoso-lanati phyllis erectis, infra spinam dilatatis, lateraliter spinuloso-denticulatis, spina tenui abruptiuscule terminatis; flosculis purpureis. — Differt a praecedentibus praesertim foliis summis magnis, capitulum involucrantibus et eo longioribus, a *C. spathulato* insuper involucri phyllis latioribus erectis, sed non in ligulam longam dilatatis ut in *C. ligulare*. —

In regione montana et subalpina. Epirus: mt. Micikeli, Peristeri (Form.); Thessalia: in oropedio Neuropolis pr. Sermeniko (Sint.), mt. Said Pascha, Dokimi, Oxya, pr. Mandra Hodsa, Koniko in Pindo, pr. Patsios, Baraktari, Godaman, Diodendron, Nezeros et Rapsani in Olympo, mt. Alafovris in mt. Othrys (Form.); Peloponnesus: mt. Cotylus supra Andrizena (Heldr.). — Jul. Sept. ⊙.

×× Capitula parva, nucis magnitudinem non excedentia.

4. **C. Lobelii** Ten. fl. nap. V. p. 211, t. 189; Bois. fl. or. III. p. 530. — *C. hellenicum* Bois. et Orph. mss. ex Bois. fl. or. l. c. — Probabiliter huc pertinet: *Cnicus eriophorus* S. et S. pr. II. p. 152 ex Arcadia. — Exsicc.: Orph. fl. gr. n. 469; Sint. it. thessal. n. 1124.

Caule erecto, racemoso-corymboso, araneoso; foliis supra viridibus, subtus araneoso-canescentibus, in lacinias anguste lanceolatas, basi 2—3 partitas, spina validiuscula terminatas, abeuntes pinnatipartitis; capitulis ovatis, foliis summis, flores aequantibus vel superantibus bracteatis; involucri araneoso-lanati phyllis lineari-lanceolatis, erecto-patulis, infra spinam dilatatis, lateraliter spinuloso-denticulatis, spina tenui abruptiuscule terminatis; floribus sordide albis. — Differt a praecedentibus foliorum segmentis gracilioribus, spina minus valida terminatis et praesertim capitulis 2—3plo minoribus.

β. **hypopsilum** Bois. et Heldr. diagn. VI. p. 101 pro sp.; Bois. fl. or. III. p. 530. — Folia supra strigosa, subtus glaberrima, lucida. — A specie sequenti foliorum laciniis longioribus, floralibus magis parenchymatosis et phyllorum spinis longioribus (an satis?) differre dicitur. — Exsicc.: N. v.

In regione subalpina mt. Ghavellu Thessaliae (Sint.); mt. Clokos Achaiae et in mt. Malevo pr. Hagios Joannes Laconiae (Orph.); — β. ad limites regionis superioris regionis abietinae mt. Taygetos (Heldr.). — Jul. Aug. ⊙.

5. **C. morinaefolium** Bois. et Heldr. fl. or. III. p. 530; Bald. viagg. Creta p. 67. — *C. hypopsilum* Raul. cret. p. 792, non Bois. et Heldr. — Exsicc.: Heldr. pl. cret. a. 1870; Rev. pl. cret. n. 187; Bald. it. cret. alt. n. 230.

Caule erecto, subracemoso-corymboso, glabriusculo; foliis utrinque viridibus, glabris, in lacinias triangulares, basi 2—3 partitas, spina valida terminatas abeuntes pinnatipartitis; capitulis ovatis, foliis summis flores superantibus bracteatis; involucri araneosi phyllis lineari-lanceolatis, erectis, infra spinam subdilatatis, lateraliter minutissime serrulatis in spinam tenuem attenuatis; floribus albis. — A *C. Lobelii* foliis glaberrimis, nec subtus araneoso-canescentibus, laciniis triangularibus, involucri phyllis erectis, apice parum dilatatis, minute denticulatis discedit.

In regione media et superiori. Creta: mt. Drakona (Rev.), Theodori et Korfikastron (Raul.) in mt. Sphacioticis, mt. Ida (Heldr.). — Jul. Aug. ⊙.

β. Involucri spinae sub apice non dilatatae.

6. **C. Heldreichii** Hal. in ö. b. Z. 1890 p. 114. — *C. decussatum* Bois. fl. or. suppl. p. 308, non Janka. — *C. dolopicum* Form. in D. bot. Monat. 1897 p. 74, in Ver. Brünn 1897 p. 37. — Exsicc.: Heldr. it. gr. septentr. a. 1879; Form. pl. thessal. a. 1896.

Caule erecto, corymboso, araneoso; foliis supra viridibus, subtus albo-lanuginosis, in lacinias lineari-lanceolatas, basi 2—3 partitas, spina

valida terminatas abeuntes pinnatipartitis; capitulis ovatis, nucis magnitudinem non excedentibus, foliis summis, flores aequantibus vel superantibus bracteatis; involucri parce araneosi phyllis anguste lineari-subulatis, minutissime serrulatis, in spinam erectam vel erecto-patulam, tenuem, longam attenuatis; floribus sordide albidis. — A praecedentibus praeter alias notas, praesertim involucri spinis sub apice non dilatatis discedens; habitu *C. Lobelii* simile.

In regione media et superiori. Thessalia: mt. Ghavellu et Karava (Form.); Eurytania: mt. Tymphrestus (Heldr.). — Jul. Aug. ⊙.

7. **C. leucocephalum** Willd. sp. III. p. 1668 (*Cnicus*); Spreng. syst. III. p. 377; Raul. cret. p. 792. — *Carduus leucocephalus* Pers. syn. II. p. 387. — *Serratula leucocephala* Poir. dict. VI. p. 565. — *C. lappaceum v. leucocephalum* Bois. fl. or. III. p. 532. —

Caule erecto, subracemoso, araneoso; foliis supra viridibus, subtus tomentosis, in lacinias lanceolatas, basi bipartitas, in spinas tenues abeuntes pinnatipartitis; capitulis cylindraceis, parvis, solitariis vel binis, in racemum simplicem longum interruptum dispositis, foliis summis involucro brevioribus vel aequilongis bracteatis; involucri araneosi phyllis adpressis, ab infimis ad superiora valde elongatis, in spinam subrecurvo-patulam, tenuem, longam attenuatis; floribus purpurascentibus.

In Creta (Willdenow l. c.); a recentioribus non lectum. — Jul. Aug. ⊙. N. v.

b. Folia caulina decurrentia.

8. **C. lanceolatum** L. sp. p. 821 (*Carduus*); Scop. fl. carn. ed. 2, II. p. 130; Marg. et R. fl. Zante p. 60; Clem. sert. p. 55; Bois. fl. or. III. p. 538; Heldr. fl. cephalon. p. 47; Form. in D. bot. Monat. 1890 p. 20, 1897 p. 74 var. *brunneum*, in Ver. Brünn 1895 p. 28 cum *f. opacum*, 1896 p. 53, 1897 p. 38 cum var. *brunneum*; Hal. in ö. b. Z. 1897 p. 285. — *C. nemorale* Rchb. fl. germ. exc. p. 286, f. foliis hypoleucis. — Icon: Rchb. t. 95. — Exsicc.: Sint. et Bornm. it. turc. n. 1316.

Caule erecto, superne ramoso; foliis supra viridibus, subtus plus minus araneoso-canis, in lacinias inaequaliter et divaricatim lobatas, spinosas sinuatim pinnatifidis partitisve, caulinis in alas angustas lobatas decurrentibus; capitulis magnis, breviter et anguste bracteatis; involucri araneosi phyllis lanceolato-subulatis, apice non dilatatis, in spinam erecto-patulam sensim attenuatis; floribus purpureis.

In regione inferiori et montana. Frequens in Epiro et Thessalia a Pindo per planitiem usque ad Olympium et Pelion (Form.); Aetolia: pr. Prusso (Samar.); Locris: pr. Granitza (Tunt.); Boetia: ad Cephissum (Sart.); Attica: mt. Parnes, Pentelicon, pr. Kephissia (Heldr.), mt. Hymettus pr. Caesariani (Clem.); Zante (Marg.); Cephalonia: pr. Poros (Heldr.); Corcyra (Form.). — Jul. Aug. ⊙

9. **C. italicum** Savi bot. etrur. III. p. 140; Ch. et B. exp. p. 242, Fl. pelop. p. 56; (*Carduus*); DC. hort. monsp. p. 96; Link in Linnaea

IX. p. 580; Marg. et R. fl. Zante p. 60; Bois. fl. or. III. p. 538, Heldr. fl. cephal. p. 47, chlor. Parn. p. 21; Hal. Beitr. fl. Aetol. p. 7, in ö. b. Z. 1895 p. 412; Bald. in nuov. giorn. bot. 1894 p. 97, riv. coll. bot. Alb. 1895 p. 50; Haussk. symb. p. 121; Form in Ver. Brünn 1897 p. 38. — Icon: Rchb. t. 96. — Exsicc.: Orph. fl. gr. n. 349; Sint. it. thessal. n. 1231.

Caule erecto, superne ramoso; foliis utrinque plus minus araneoso-canis vel supra viridibus, in lacinias anguste lanceolatas, spina terminatas, pinnatipartitis, longe acuminatis, caulinis in alam brevem auriculaeformem integram decurrentibus; capitulis parvis, avellanae magnitudinem non excedentibus, foliis summis, flores superantibus bracteatis; involucri glabriusculi vel parum araneosi phyllis oblongo-lanceolatis, apice non dilatatis, in spinam erecto-patulam abruptiuscule abeuntibus; floribus purpureis.

In campis regionis inferioris et montanae, saepissime gregarie. Epirus: pr. Prevesa, Nicopolis, Philippiada (Bald.); Thessalia: ad monasterium Korona, pr. Kataphygi, Kalabaka, Ruso, Povelci (Form.), Trikala, Demirli, Larissa (Haussk.), Palaeokastro (Leon.), Mavro Mandila, Lamia, Limogardi (Form.), pr. Khani ad mt. Kukkos Phthiotidis (Heldr.); pr. Rachova in mt. Parnasso (Orph.); Acarnania: pr. Sorovigli (Hal.); Aetolia: pr. Mesolongion (Nied.); Euboea: pr. Kurbatzi (Wild); Argolis: pr. Nauplia (Link), Argos (Chaub.), Tyrins (Heider); Laconia: pr. Sparta (Heldr.), Xerocampos (Pichl.); Messenia (Bois.); Zante (Marg.); Cephalonia: in oropedio Omala pr. Francata, in valle Heracleon, in halipedo Livadi, pr. Lixuri (Heldr.). — Jun. Aug. ⊙

2. Sectio. *Chamaeleon* DC. l. c. — Folia supra non strigosa; achenia compressa, epicarpio membranaceo.

a. Folia non decurrentia.

α. Flores purpurei.

× Folia subtus albo-tomentosa; involucri phylla in spinam validam, patulam, phyllo ipso multo longiorem, flosculos subsuperantem abeuntia.

10. **C. cynaroides** Lam. dict. I. p. 705; Ch. et B. fl. pelop. p. 56; (*Carduus*); Spreng. syst. III. p. 372; Bois. fl. or. III. p. 546, suppl. p. 309; Heldr. fl. cephal. p. 47; Spreitz. in z. b. G. 1890 p. 296; Bald. viagg. Creta p. 68. — *Chamaepeuce cynaroides* DC. pr. VI. p. 659; Raul. cret. p. 793. — *Cnicus cynaroides* Willd. sp. III. p. 1670; Desf. cor. p. 46, t. 35. — *Cnicus afer* Sieb. avis p. 5, rem. p. 8, non Willd. — Exsicc.: Orph. fl. gr. n. 793; Heldr. herb. norm. n. 627; Rev. pl. cret. n. 87, in Baen. herb. europ. n. 4862.

Caule erecto, niveo-tomentoso, simplici vel ramoso, 1—3 cephalo; foliis lanceolatis, supra glabriusculis, viridibus, subtus adpresse niveo vel cano-tomentosis, in segmenta ovata, 3—5 lobata, spinosa, pinnatipartitis, caulinis diminutis amplexicaulibus; capitulis majusculis, terminalibus, solitariis; involucri parce araneosi phyllis in spinam lanceo-

latam, infimorum reflexam, intermediorum patulam abeuntibus, intimis purpureis scariosis, erectis. — Species pulchra, habitu ad sectionem sequentem accedens; sed ob acheniis compressis epicarpio membranaceo nec duro praeditis, huic sectioni adnumerandum.

In olivetis, pinetis, rupestribus regionis inferioris et montanae. Argolis: adversus insulam Poros, in insula Poros, Hydra (Heldr.); Laconia: mt. Malevo pr. Meligon (Orph.), mt. Taygetos supra Mistra (Bois.); Messenia: pr. Pylos (Chaub.); Cephalonia: pr. Vuttes, in eparchia Koroni (Heldr.); Zante: pr. Litakia (Letourn.); Cythaera: pr. Gatuni (Spreitz.); Creta: pr. Rumata (Bald.), Malaxa, Amalos (Rev.), Askyphos (Spreitz.), Candia (Raul.). — Apr. Jun. ♃.

×× Folia glabra vel subtus araneoso-canescentia, involucri phylla mucronata vel breviter spinulosa.

○ Folia pinnatipartita.

11. **C. appendiculatum** Griseb. spic. II. p. 250; Bois. fl. or. suppl. p. 309; Haussk. symb. p. 121; Bald. riv. coll. bot. Alb. 1895 p. 50; Form. in Ver. Brünn 1896 p. 51, v. *thessalum* (quod a typo nullomodo discedit). — Icon: Vis. fl. dalm. suppl. alt. t. 3, — Exsicc.: Bald. it. alb. epir. III. n. 164, Sint. it. thessal. n. 728.

Glabriusculum, virens; caule erecto, simplici vel parce ramoso, superne subnudo; foliis concoloribus, oblongis, amplexicaulibus, in lacinias oblongas, lobatas, breviter spinosas pinnatipartitis, acuminatis; capitulis mediocribus, terminalibus, saepius glomeratis; involucri glabriusculi phyllis ovato-lanceolatis, adpressis, dorso nigro-purpurascentibus, in spinulam erecto-patentem attenuatis, interioribus lanceolatis, inermibus. — *C. montano* (W. et K.) affine, ab eo spinis foliorum elongatis flavicantibus et involucri phyllis angustioribus longius spinosis diversum.

Ad rivulos regionis subalpinae. Epirus: in jugo Bresani mt. Tsumerka (Bald.), in valle Negerli supra Chaliki (Sint.), mt. Zygos (Haussk.); Phthiotis: mt. Oeta loco dicto Lupakia (Heldr.). — Jul. Aug. ♃

○○ Folia breviter lobulata.

12. **C. tymphaeum** Haussk. symb. p. 121. — Exsicc.: Heldr. it. IV. Thessal. a. 1885.

Caule erecto, leviter araneoso, superne ramoso, subnudo; foliis coriaceis, rigidis, glaucis, glaberrimis vel subaraneosis, oblongo-lanceolatis, late truncato-sessilibus, parum breviterque lobulatis, lobulis tenuiter spinulosis, in spinam longiusculam, stramineam, pungentem attenuatis; capitulis mediocribus, terminalibus, solitariis vel binis; involucri leviter araneosi phyllis exterioribus triangulari-lanceolatis, adpressis, apice nigricantibus, in spinulam recurvatam attenuatis, interioribus lanceolatis, inermibus, superne subconstrictis, in appendiculum purpureo-scariosum abeuntibus. — Caules pluripedales, crassitie pennae anserinae et ultra, pedunculi ad 30 cm. longi, folia circa 15 mm. lata.

In scaturiginosis supra Malakasi et in glareosis humidis fluminis Salamvrias pr. Uranaeos in Pindo tymphaeo Thessaliae (Haussk.). — Jul. Aug. ♃.

13. **C. pauciflorum** W. et K. pl. rar. hung. II. p. 175, t. 161 (*Cnicus*); Spreng. syst. III. p. 375; Rchb. icon. XXV. p. 63 et 80 n. 47, obs. t. 102; Bois. fl. or. III. p. 545; Heldr. fl. cephal. p. 47. —

Caule erecto, araneoso, simplici vel subramoso, superne subnudo; foliis membranaceis, supra pilosulis, subtus araneoso-canescentibus, amplissimis, ovatis, breviter lobatis dentatisque, spinuloso-ciliatis, inferioribus petiolatis, intermediis petiolo alato, basi auriculato amplexicaulibus, superioribus diminutis, basi cordata sessilibus; capitulis mediocribus, terminalibus, 2—4 conglobatis; involucri sub lente velutini phyllis nigricantibus, triangulari-lanceolatis, mucronatis, internis lanceolatis, scarioso-marginatis.

In silvaticis mt. Aenos Cephaloniae (Mazziari); postea a nemini lectum. — Jul. Aug. ♃. N. v.

β. Flores ochroleuci vel albi.

14. **C. oleraceum** L. sp. p. 826 (*Cnicus*); Scop. fl. carn. ed. 2, II. p. 124; Form. in D. bot. Monat. 1890 p. 21. —

Caule erecto, simplici vel superne parce ramoso, foliisque glabriusculo; foliis oblongis vel ellipticis, spinuloso-ciliatis, amplexicaulibus, inferioribus pinnatifidis, superioribus integris; capitulis mediocribus, terminalibus, congestis, foliis summis latis, pallidis, capitula superantibus involucratis; involucri glabri vel parce araneosi phyllis lanceolatis, in spinulam terminatis, apice patentibus; floribus ochroleucis.

In locis humidiusculis pr. Vendista in Pindo Thessaliae (Form.), si determinationi fides habenda. — Jul. Aug. ♃. N. v.

15. **C. candelabrum** Griseb. spic. II. p. 251; Bois. fl. or. III. p. 540; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 4; Form. in D. bot. Monat. 1890 p. 21, in Ver. Brünn 1895 p. 28, 1896 p. 53; Hal. Beitr. fl, Epir. p. 28, Beitr. fl. Thessal. p. 14; Haussk. symb. p. 121. — *C. chelmeum* Orph. in Bois. diagn. ser. 2, III. p. 41. — Exsicc.: Sint. it. thessal. n. 1201; Heldr. herb. norm. n. 1555.

Glabrum; caule erecto, folioso, longe racemoso-paniculato; foliis lanceolatis, sinuato-lobatis, creberrime et tenuiter spinoso-marginatis, infimis in petiolum attenuatis, ceteris sessilibus, basi auriculatis; capitulis parvis, ad racemi ramos subsessilibus, glomeratis, paniculam amplam, polycephalam formantibus, foliis floralibus, involucrum aequantibus bracteatis; involucri phyllis lanceolatis, in spinam brevem, erectam attenuatis, intimis scariosis, apice serrulatis; floribus albidis. — Species insignis, usque 2 m. alta, capitula eis *C. arvensis* haud minora, nutantia, saepe numerosissima.

In dumosis regionis montanae et subalpinae. Epirus: pr. Ephemiades, Safikbi, Han Balduma, in valle Dipotami, mt. Prosgoli (Form.), mt. Tsumerka, Peristeri, pr. Chaliki (Hal.); Thessalia: mt Baba, pr. Neuropolis (Haussk.), Sermeniko (Sint.), Kastania (Hal.), Krania, Klinovo, Vendista, Velitsena, Koturi, Malakasi, mt. Said Pascha, Borovikon, Han Kukleus et Kriavris (Form.), mt. Ossa, Pelion (Heldr.); Eurytania: mt. Chelidoni (Heldr.), Tymphrestus (Samar.), Korax pr. Musinitza (Heldr.); Peloponnesus: mt. Kyllene pr. Trikala (Heldr.), mt.

Chelmos pr. Zaruchla, mt. Malevo (Orph.), in distr. Alagonia pr. Megali-Anastasova et in faucibus Langada mt. Taygetos (Zahn). — Jul. Aug. ⊙.

b. Folia caulina decurrentia.

16. **C. creticum** Lam. enc. I. p. 701 (*Carduus*); Urv. enum. p. 107; Raul. cret. p. 792. — *Cnicus palustris* S. et S. pr. II. p. 151, non Willd. — *Cirsium siculum* DC. pr. VI. p. 642; Bois. fl. or. III. p. 548; Heldr. fl. cephal. p. 47, in Sitzungsb. acad. Wiss. Berl. 1883 p. 4; Hal. Beitr. fl. thessal. p. 14; Haussk. symb. p. 123; Bald. in nuov. giorn. bot. 1894 p. 97, viagg. Creta p. 67, riv. coll. bot. Alb. 1896 p. 68; Form. in Ver. Brünn 1895 p. 28, 1896 p. 50, 1897 p. 37; non Spreng. neue Entd. p. 36, quod = *Galactites tomentosa* Moench. (cf. Schultz in ö. b. Z. 1858 p. 246). — *Cnicus siculus* Guss. fl. sic. II. p. 442. — *Cirsium microcephalum* Schultz l. c. — *C. siculum v. longispinum* Bois. fl. or. III. p. 548; Heldr. chlor. Parn. p. 21 (forma spinis foliorum et praesertim alarum creberrimis elongatis). — *Cirsium palustre* Form. in D. bot. Monat. 1890 p. 21, non Scop. — Formae involucri phyllis spinula brevissima saepe ad mucronem reducta terminatis: *Cirsium polyanthos* Urv. enum. p. 107; Marg. et R. fl. Zante p. 60. — *Carduus polyanthos* Ch. et B. fl. pelop. p. 56. — *Carduus Hippolyti* Ch. et B. exp. p. 241, t. 29, Fl. pelop. p. 56, t. 31. — *Cirsium Hippolyti* Link in Linn. IX. p. 580. — Exsicc.: Orph. fl. gr. n. 790; Heldr. herb. norm. n. 437 et 946; Rev. pl. cret. n. 88 et 89, in Baen. herb. europ. n. 4861; Sint. et Bornm. it. turc. n. 1318; Sint. it. thessal. n. 1227.

Caule erecto, araneoso, superne stricte ramoso, polycephalo; foliis superne saepissime tuberculato-hirtis, subtus plus minus araneoso-canescentibus, in lacinias triangulares lobatas longe spinosas sinuato-pinnatifidis, caulis alis lobatis, spinosis; capitulis parvis, ovatis, approximatis, involucri phyllis margine araneosis, ovatis vel oblongis, spina eis aequilonga vel breviore terminatis, internis abrupte in appendicem scariosam triangularem purpurascentem abeuntibus; floribus purpureis. — *C. palustri* Scop. proximum, a quo foliorum spinis crassioribus longioribus praesertim discedit.

In humidis, secus rivulos regionis inferioris et montanae per totam Graeciam. — Jun. Sept. ⊙.

17. **C. pindicolum** Haussk. symb. p. 122. —

Caule erecto, leviter araneoso, superne breviter ramoso, mono-vel oligocephalo; foliis glaucis, supra glabriusculis, subtus araneoso-canescentibus, oblongo-lanceolatis, breviter triangulato-lobatis, spinosis, inferioribus ad medium internodium usque, intermediis brevius spinoso-decurrentibus; capitulis mediocribus, semiglobosis, subsolitariis vel ad apicem ramorum 2—3 congestis; involucri phyllis apice subaraneosis, oblongo-lanceolatis, in spinam brevem, intimis sensim in apicem setaceum purpurascentem angustatis; floribus purpureis. — Capitula magnitudine *C. cani*. — Species mihi ignota.

In scaturiginosis subalpinis mt. Baba inter Krania et Klinovo in Pindo (Haussk.). — Jul. Aug. ⊙ vel ♃? — N. v.

3. Sectio. *Lamyra* Cass. dict. 25 p. 218 pro gen.; Bois. fl. or. III. p. 524. — Folia supra non strigosa, achenia globosa, ovata vel pyriformia, non compressa, epicarpio duro.

a. Capitula magna; involucri phylla non gibbosa.

18. **C. afrum** Jacq. hort. schoenbr. II. p. 180, t. 145 (*Carduus*); DC. cat. hort. monsp. p. 96; Link in Linn. IX. p. 580; Bois. fl. or. III. p. 552; Heldr. chlor. Parn. p. 21. — *Cnicus afer* Willd. sp. III. p 1682; S. et S. pr. II. p. 154, Fl. gr. IX. t. 829. — *Chamaepeuce afra* DC. pr. VI. p. 659; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 4; Hal. in ö. b. Z. 1888 p. 759, in ö. b. Z. 1892 p. 371, Beitr. fl. Epir. p. 28, Beitr. fl. thessal. p. 14, Beitr. fl. Achaia p. 24; Form. in D. bot. Monat. 1890 p. 20, in Ver. Brünn 1895 p. 27, 1896 p. 50, 1897 p. 36; Haussk. symb. p. 121. — *Carduus Casabonae* Ch. et B. fl. pelop. p. 56, non L. -- Exsicc.: Sint. et Bornm. it. turc. n. 1322; Sint it. thessal. n. 1200.

Caule erecto, crasso, corymboso, polycephalo; foliis sessilibus, lineari-lanceolatis, supra parce arachneoideis, subtus niveo-tomentosis, in lacinias lanceolatas, ad spinas validas fere reductas pinnatipartitis; capitulis solitariis; involucri araneosi phyllis lanceolatis, acuminato-spinosis, intimis scariosis purpureis; floribus purpureis. — Species pulchra.

In saxosis regionis submontanae, in subalpinam adscendens. Frequens in Epiro et in Thessaliae montibus omnibus; dein in mt. Korax (Tunt.), Kiona (Hal.), Parnassus (Sibth.); Peloponnesus: mt. Olenos, Chelmos, Kyllene (Hal.), pr. Tripolis (Sart.), mt. Taygetos (Heldr.). — Jul. Aug. ⊙.

b. Capitula parva; involucri phylla exteriora ad spinae basin internam gibbosa.

19. **C. strictum** Ten. pr. p. 48 (*Cnicus*); Spreng. syst. III. p. 370; Bois. fl. or. III. p. 550. — *Chamaepeuce stricta* DC. pr. VI. p. 659; Hal. Beitr. fl. thessal. p. 14; Haussk. symb. p. 121; Form. in Ver. Brünn 1897 p. 37. — *Chamaepeuce stricta v. graeca* Bois. et Heldr. diagn. ser. 2, III. p. 47. — *Chamaepeuce atropurpurea* Bois. et Heldr. l. c.; Form. in D. bot. Monat. 1890 p. 20. — Exsicc.: Orph. fl. gr. n. 794; Heldr. herb. norm. n. 712; Sint. et Bornm. it. turc. n. 1315.

Caule erecto, simplici, anguste alato, superne breviter racemoso, oligocephalo; foliis supra glabris, subtus cano-tomentosis, decurrentibus, lanceolatis, sinuato-dentatis, dentibus triangularibus, in spinam tenuem attenuatis; capitulis solitariis vel 2—3 approximatis, involucri parce araneosi phyllis adpressis, in spinam erecto-patulam abeuntibus, intimis erectis, purpurascentibus; flosculis purpureis.

In nemorosis regionis montanae. Thessalia: pr. Tsungeri, Malakasi, mt. Oxya, pr. Chaliki, Klinovo (Haussk.), Kastania (Hal.), Han Kura-

neos in valle Penei superioris, pr. Neochorion in mt. Agrapha (Form.), pr. Mikrochorio Eurytaniae (Samar.), mt. Olympus ad Hagios Dionysios (Orph.). — Jul. Aug. ♃.

20. **C. stellatum** L. sp. p. 823; Ch. et B. exp. p. 242, Fl. pelop. p. 56; (*Carduus*); All. fl. ped. I. p. 153; Bois. fl. or. III. p. 550; Heldr. fl. cephal. p. 47. — *Chamaepeuce stellata* DC. pr. VI. p. 658; Clem. sert. p. 55; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Achaia p. 24; Form. in Ver. Brünn 1895 p. 27. — Icon: Fl. gr. t. 830. — Exsicc.: Orph. fl. gr. n. 263.

Caule erecto, exalato, simplici et superne breviter racemoso oligocephalo vel a medio ramoso pleiocephalo; foliis sessilibus, supra glabris, subtus cano-tomentosis, lanceolatis vel linearibus, integerrimis, utrinque ad axillam spinas 1—2 gerentibus; capitulis solitariis; involucri glabri phyllis adpressis, in spinam erecto-patentem abeuntibus, intimis erectis, viridibus; flosculis purpureis.

In nemorosis regionis montanae. Corcyra (Clem.): pr. Pyrgi (Bickn.), Scripero (Gelmi), Gasturi (Form.); Cephalonia: mt. Gerolaki, pr. Pylaros, Valeriano, Koroni, Skala, Elios (Heldr.); Achaia: pr. Megaspilaeon (Hal), Trikala (Orph.); Arcadia: pr. Tripolis (Bois.); Messenia: mt. Ithome (Sart.); Laconia: pr. Marathonisi, Vitilos, Chimova (Chaub.), Selitza (Zahn). — Maio, Jul. ⊙.

B. Corollae limbus ad basin 5partitus; pappus corolla demum longior.

4 Setio. *Cephalonoplos* Neck. elem. p. 68. — Folia supra non strigosa; achenia lineari-oblonga, compressa, epicarpio membranaceo.

21. **C. arvense** L. sp. p. 820 (*Serratula*); Scop. fl. carn. ed. 2, II. p. 126; Friedr. Reise p. 264; Bois. fl. or. III. p. 552; Hal. Beitr. fl. Epir. p. 28, in ö. b. Z. 1895 p. 412; Haussk. symb. p. 123; Form. in Ver. Brünn 1895 p. 28, 1896 p. 50, 1897 p. 37; Bald. riv. coll. bot. Alb. 1896 p. 69. — Icon: Rchb. XXV. t. 111. — Exsicc.: Orph. herb. n. 2031.

Caule erecto, superne ramoso; foliis subdecurrentibus, oblongo-lanceolatis, spinuloso-ciliatis, integris vel pinnatifido-sinuatis, apice lobisque spina validiore terminatis, glabriusculis vel utrinque subtus densius canis; capitulis parvis; involucri phyllis adpressis, margine araneosis, in spinam brevissimam acuminatis; flosculis purpureis. — Formam foliis hypoleucis validiuscule spinosis tantum e ditione vidi.

In arvis, cultis, lapidosis regionis inferioris, in alpinam usque adscendens. Epirus: pr. Kurenta (Bald.), Janina (Form.), Theodoriana, Mazuki (Hal.); passim per totam Thessaliam; Aetolia: mt. Korax, Arapokephalo (Heldr.); Achaia: pr. Patras (Friedr.), mt. Chelmos, Kyllene (Orph.) et probabiliter alibi. — Jul. Sept. ♃.

Obs. Erronee in Graecia indicantur: *C. rivulare* Jacq. fl. austr. I. t. 91; Ch. et B. exp. p. 241, Fl. pelop. p. 56 (*Carduus*); Link en. hort. berol. II. p. 301; = *Cnicus rivularis* Willd. sp. III. p. 1676

S. et S. pr. II. p. 153; in Peloponneso; dein *C. spinosissimum* L. sp. p. 826; Pieri corc. fl. p. 116 (*Cnicus*); Scop. fl. carn. ed. 2. II. p. 129; in Corcyra.

45. Picnomon Lob. ic. II. t. 14.

1. **P. acarna** L. sp. p. 820; Ch. et B. exp. p. 242, Fl. pelop. p. 56 (*Carduus*); Cass. dict. 40 p. 188; Clem. sert. p. 57; Raul. cret. p. 792; Weiss in z. b. G. 1869 p. 43; Spreitz. in z. b. G. 1887 p. 663, 1890 p. 296; Form. in D. bot. Mon. 1890 p. 20, in Ver. Brünn 1895 p. 27, 1896 p. 50, 1897 p. 37; Bald. viagg. Creta p. 68; Heldr. fl. Aegina p. 305, chlor. Thera p. 15. — *Cnicus acarna* L. sp. ed. 2 p. 1158; S. et S. pr. II. p. 151; Fraas fl. class. p. 205. — *Cirsium acarna* Moench. suppl. p. 226; Friedr. Reise p. 274; Bois. fl. or. III. 549; Heldr. fl. cephal. p. 47; Hal. in z. b. G. 1888 p. 759; Beitr. fl. Epir. p. 28, Beitr. fl. Aetol. p. 7, Beitr. fl. Thessal. p. 14; Haussk. symb. p. 123. — Icon: Fl. gr. t. 827. — Exsicc.: Rev. pl. cret. n. 190; Sint. et Bornm. it. turc. n. 1314.

Lanuginoso-tomentosum; caule erecto, ramoso, alato, alis spinuloso-ciliatis; foliis lanceolatis, sinuato et spinoso-dentatis, spinuloso-ciliatis; capitulis mediocribus, saepius 2—4 aggregratis, foliis summis ea superantibus bracteatis; involucri phyllis adpressis, praeter intima scarioso-cuspidata, spina pinnata terminatis; flosculis purpureis.

In ruderatis, sterilibus, agris, aridis regionis inferioris et montanae passim per totam Graeciam. — Jun. Aug. ⊙.

46. Notobasis Cass. dict. 25 p. 225 et 35 p. 170.

1. **N. syriaca** L. sp. p. 823; Ch. et B. exp. p. 242, Fl. pelop. p. 56 (*Carduus*); Cass. l. c.; Marg. et R. fl. Zante p. 60; Ung. Reise p. 124; Weiss. in z. b. G. 1869 p. 43; Raul. cret. p. 793; Bois. fl. or. III. p. 553; Heldr. fl. cephal. p. 47, Fl. Aegina p. 305; Hal. Beitr. fl. Thessal. p. 14, in ö. b. Z. 1897 p. 95; Haussk. symb. p. 121. Form. in Ver. Brünn 1896 p. 50, 1897 p. 37. — *Cirsium syriacum* Gaertn. fr. II. p. 383, t. 163; Urv. enum. p. 107. — *Cnicus syriacus* Willd. sp. III. p. 1683; S. et S. pr. II. p. 154, Fl. gr. t. 831. — *Cirsium bracteatum* Link in Linn. IX. p. 580. — Exsicc.: Rev. pl. cret. n. 91; Dörfl. fl. Aeg. n. 63.

Caule erecto, araneoso, simplici vel superne ramoso; foliis supra glabrescentibus, albo-venosis, subtus pubescentibus, oblongis, sinuato-lobatis, lobis spinoso-dentatis, inferioribus in petiolum attenuatis, ceteris amplexicauli-auriculatis, superioribus fere ad spinas validas reductis, floralibus capitula bracteantibus et superantibus; capitulis mediocribus, ad ramorum apicem glomeratis; involucri subaraneosi phyllis adpressis, spina brevi subpatula terminatis; flosculis dilute purpureis.

Ad vias, fossas, ad sepes, in ruderatis regionis inferioris, per Graeciam totam. — Maio, Jun. ⊙

47. Cynara L. gen. n. 928.

1. **C. cardunculus** L. sp. p. 827; Marg. et R. fl. Zante p. 60; Heldr. Nutzpfl. p. 27, Fl. cephal. p. 47; Raul. cret. p. 791; Bois. fl. or. III. p. 557; Hal. in z. b. G. 1888 p. 759 et 763, Beitr. fl. Aetol. p. 7, Beitr. fl. Achaia p. 24; Form. in Ver. Brünn 1895 p. 27. — *C. scolymus* S. et S pr. II. p. 157; Ch. et B. exp. p. 243; Fl. pelop. p. 57; non L. — *C. horrida* Ait. hort. Kew. III. p. 148; S. et S. pr. II. p. 157, Fl. gr. t. 834; Ch. et B. exp. p. 343; Fl. pelop. p. 57; Friedr. Reise p. 285. — *C. humilis* Heldr. Nutzpfl. p. 27, non L. — Exsicc.: Hal. it. gr. a. 1888.

Caule robusto, simplici vel ramoso, foliis supra glabrescentibus, subtus cano-tomentosis, oblongis, in lacinias lanceolatas, pinnatilobas, valide spinosas, secus rachidem decurrentes pinnatisectis; capitulis magnis, solitariis, involucri glabriusculi phyllis late oblongo-triangularibus, infimis spina brevi terminatis, intermediis in acumen longum, lanceolatum, supra canaliculatum, erecto-patulum abeuntibus, intimis subscariosis; flosculis coeruleis.

In arvis, pratis, collibus arenosis regionis inferioris. Epirus: pr. Nikopolis, Luros, Philippiades (Form.); Aetolia: pr. Mesolongion et copiosissime in planitie pr. Antirrhion, pr. Amphissa Doridis (Hal.); in planitie Thebana, pr. Lebadia Boeotiae; pr. Oropos Atticae (Heldr.); Corinthus (Hal.); Argolis (Chaub.); pr. Nauplia (Friedr.); Achaia: pr. Patras (Hal.); Elis, Messenia (Sibth.): pr. Navarin (Chaub.); Creta (Sibth.): in planitie Messara (Raul.); Zante: pr. Argassi, ad paludem Makri (Marg.); Cephalonia: pr. Lixuri, Lychata (Heldr.); Corcyra: pr. Alipu, Kanali (Form.); Cycladum insula Naxos (Sibth.). — Jul. Aug. ♃.

Obs. *C. scolymus* L. sp. p. 827; Dallap. prosp. p. 109, Marg. et Reut. fl. Zante p. 85; Fraas fl. class. p. 202; Raul. cret. p. 791; Heldr. chlor. Thera p. 29. — Forma culta praecedentis, a quo differt capitulis maximis, involucri phyllis inermibus, basi valde carnosis. — Colitur in hortis.

2. **C. Sibthorpiana** Bois. et Heldr. diagn. X. p. 94, Fl. or. III. p. 557; Raul. cret. p. 792, t. 15. — *C. humilis* S. et S. pr. II. p. 157, Fl. gr. IX. p. 25, t. 835; Ch. et B. exp. p. 243; Fl. pelop. p. 57; non L. —

Differt a praecedente foliorum breviter et tenuiter spinosorum segmentis minus dissectis, non decurrentibus, involucri phyllis in acumina contractis, flosculis albis.

In Peloponneso et Creta (Sibth.): pr. Piskokephali eparchiae Sitia (Heldr.). — Apr. Maio. ♃. N v.

48. Onopordon L. gen. n. 927.

a. Involucri phylla angusta, in spinas elongatas subulatas, basi adpresse manifeste longiores abruptiuscule angustata, exteriora interiorum fere longitudine.

α. Virescentia.

× Folia grosse et irregulariter lobata.

1. **O. acanthium** L. sp. p. 827; S. et S. pr. II. p. 157; Pieri corc. fl. p. 116; Hal. in ö. b. Z. 1895 p. 412; Haussk. symb. p. 121; Form. in Ver. Brünn 1895 p. 27, 1896 p. 50, 1897 p. 36. — Icon: Fl. dan. t. 909. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Adpresse araneosum; caule elato ramoso, late alato, alis sinuato-lobatis, spinosis; foliis ovato-oblongis, sinuato-lobatis, lobis spinoso-dentatis; capitulis globosis, basi truncato-concavis; involucri parce araneosi et glandulosi phyllis attenuato-subulatis, triquetris, margine scabris, exterioribus patenti-curvatis, interioribus erectis, flosculos non aequantibus; flosculis purpureis, glabris; acheniis griseis, nigro-maculatis, pappo dimidio brevioribus.

β. **parnassicum** Bois. et Heldr. diagn. ser. 2, VI p. 114, Fl. or. III. p. 559; Heldr. chlor. Parn. p. 21; pro sp.; Rouy rev. Onop. p. 10. — Indumento parco, tandem evanido, alis caulinis angustis, interruptis, involucro magis araneoso. — Exsicc.: Heldr. herb. norm. n. 666.

In incultis, ruderatis regionis inferioris et montanae rarius. Corcyra (Pieri); Epirus: pr. Janina, Katschka (Form.); Thessalia: ad monasterium Korona (Heldr.), pr. Sophates, Karditza, Pharsalus, Demirli (Haussk.), mt. Phlamburo, pr. Mavreli, Konisko, Palaeokastro, Trikala, Ligaria, Lamia (Form.); — β. mt. Parnassus pr. Gurna (Samar.). — Jun. Sept. ⊙.

2. **O. elatum** S. et S. pr. II. p. 156; Fl. gr. IX. p. 23, t. 833; Ten. annot. fl. gr. in rendic. real. acad. sc. n. 2 p. 17; Raul. cret. p. 791; Haussk. symb. p. 121. — *O. tauricum v. elatum* Bois. fl. or. III. p. 559; Spreitz. in z. b. G. 1890 p. 296. — Exsicc.: Heldr. pl. fl. hellen. a. 1885 et 1898.

Viride; pube brevi glandulosa obductum; caule elato, ramoso, anguste alato, alis lobatis, spinosis; foliis oblongo-lanceolatis, grosse et irregulariter lobatis, lobis spinoso-dentatis; capitulis globosis, basi truncato-concavis, involucri denudati, glanduloso-viscidi phyllis lanceolatis, triquetris, margine scabriusculis, exterioribus patentissimis, interioribus erectis, flosculos non aequantibus; flosculis purpureis, glabris; acheniis brunneis, pappo duplo brevioribus. — Ab *O. taurico* Willd. sp. III. p. 1687 vix distinctum. — Differt a praecedente indumento viscido et involucri phyllis latioribus.

β. **argolicum** Bois. diag. X. p. 91; Hal. Beitr. fl. Achaia p. 24; Heldr. chlor. Thera p. 15; pro sp.; Bois. fl or. III. p. 560. — Huc

forsan *O. arabicum* S. et S. pr. II. p. 156. — Folia et capitula plus minus araneosa, involucri phylla angustiora, longius subulata. — Formis intermediis ad typum transire videtur. — Exsicc.: Dörfl. pl. aeg. n. 81.

Ad vias, in incultis, ruderatis regionis inferioris. Boeotia (Sprun.); Achaia: pr. Trikala (Heldr.); Argolis: pr. Vromolimni in peninsula Methana (Haussk.), insula Hydra, Keos Cycladum (Heldr.); Creta: pr. Askyphos (Spreitz.), Kalyves, in planitie Messara (Raul.); — *β.* pr. Corinthum (Hal.); Argolis: pr. Nauplia (Bois.); Cycladum insula Naxos, Anaphi (Leon.), Thera, Therasia (Heldr.). — Maio, Jun. ⊙.

×× Folia pinnatisecta, rhachide angustissima.

3. **O. messeniacum.** — Exsicc.: Heldr. pl. fl. hellen. a. 1899 n. 874.

Parce et adpresse araneosum; caule humili, ramoso, alis angustissimis, interruptis, ad spinas fere reductis; foliis oblongo-lanceolatis, supra glabriuculis, subtus adpresse canis, in lacinias lanceolatas, breviter spinosas pinnatisectis; capitulis globosis, basi truncatis; involucri parce araneosi et glandulosi phyllis lanceolatis, triquetris, margine scabris, exterioribus reflexis, interioribus erectis, flosculos non aequantibus; flosculis purpureis, viscidulis; acheniis brunneis, pappo duplo brevioribus. — Species foliis profunde divisis, eis *Galactitis tomentosae* similibus, segmentis remotis angustis ab omnibus affinibus egregie diversa.

In aridis Messeniae pr. Kalamata (Zahn). — Maio, Jun. ⊙.

β. Albo-tomentosa.

4. **O. Sibthorpianum** Bois. et Heldr. fl. or. III. p. 561; Haussk. symb. p. 121; Heldr. fl. Aegina p. 305. — *O. macracanthum* S. et S. pr. II. p. 157, Fl. gr. IX. p. 23, t. 832; Raul. cret. p. 791; non Schousb. — *O. graecum* Ch. et B. exp. p. 243, Fl. pelop. p. 57, sec. Bois. l. c.; an quoque S. et S. pr. II. p. 156; Sieb. avis p. 5; Urv. enum. p. 105; non Gou. ill. t. 25. — ? *C. horridum* Haussk. symb. p. 121, vix Viv.; sec. Rouy. rev. Onop. p. 19. — Exsicc.: Heldr. herb. norm. n. 479; Orph. fl. gr. n. 768 (ambo sub *O. alexandrinum*); Dörfl. fl. aeg. n. 58.

Caule humili, simplici vel ramoso, albo-tomentoso, alis angustissimis, crebre spinoso-lobatis, ad spinas interdum reductis; foliis oblongo-lanceolatis, pinnatilobatis partitisve, supra arachnoideo-canis, subtus dense albo-tomentosis, lobis triangulari-lanceolatis, spinoso-dentatis, interdum ad spinas fere reductis; capitulis depresse globosis; involucri araneosi, vix glandulosi phyllis lanceolatis, margine scabriusculis, in spinam semiteretisubulatam, carinatam angustatis, exterioribus patentissimis vel reflexis, intimis erectis, flosculos aequantibus; flosculis purpureis, viscidulis; acheniis griseis, pappo duplo brevioribus.

In campis sterilibus. Thessalia: pr. Sophates, Pharsalus, si *O. horridum* Haussk. huc pertineat; Attica: pr. Athenas, insula Aegina (Heldr.); Laconia et Messenia (Sibth.): pr. Navarin, Methone (Chaub.); Cyclades

(Sibth.): insula Melos (Urv.), Anaphi (Leon.); Creta (Sibth.). — Maoi, Jun. ⊙.

5. **O. caulescens** Urv. enum. p. 105; Ch. et B. fl. pelop. p. 57; Heldr. in ö. b. Z. 1898 p. 184, chlor. Thera p. 15, chlor. Mykon. p. 246. — *O. Weissianum* Aschers. ap. Weiss. in z. b. G. 1869 p. 43; Rouy rev. Onop. p. 20, cum f. *brevicaule*, t. 24. — Exsicc.: Heldr. herb. norm. n. 1043; Dörfl. fl. aeg. n. 120.

Differt a praecedente foliis longioribus latioribusque, lobis majoribus, triangularibus, pinnatilobatis, spinis validioribus armatis; capitulis ovoideis; involucri phyllis parce glandulosis, basi latioribus, intimis flosculis brevioribus. — Certe species propria.

In locis aridis Cycladum, in insula Syra (Weiss), Cythnos, Mykonos, Naxos, Thera (Heldr.), Melos (Urv.). — Jun. Jul. ⊙.

b. Involucri phylla lata, in spinas breves attenuata, exteriora interioribus evidenter breviora.

α. Albo-tomentosa.

6. **O. illyricum** L. Sp. p. 827; S. et S. pr. II. p. 155; Ch. et B. exp. p. 243, Fl. pelop. p. 57; Friedr. Reise p. 278; Fraas fl. class. p. 205; Clem. sert. p. 56; Raul. cret. p. 791; Weiss in z. b. G. 1869 p. 43; Bois. fl. or. III. p. 560; Heldr. fl. cephal. p. 47; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Epir. p. 28, Beitr. fl. Aetol. p. 7, Beitr. fl. Thessal. p. 14, Beitr. fl. Achaia p. 24; Form. in D. bot. Monat. 1890 p. 20, in Ver. Brünn 1895 p. 27, 1897 p. 36; Haussk. symb. p. 121. — *O. elongatum* Lam. fl. fr. II. p. 6. — *O. graecum* Gou. ill. t. 25. — *O. orientale* Mill. dict. n. 4. — *O. illex* Janka in Termész. füz. II., 1 p. 2, quod ex descriptione vix differre videtur. — *O. myriacanthum* Hal. in z. b. G. 1888 p. 759, non Bois. — *O. euboeum* Heldr. pl. fl. hellen a. 1890. — Icon: Jacq. hort. vind. II. t. 148. — Exsicc.: Rev. pl. cret. n. 86; Sint. et Bornm. it. turc. n. 1308; Sint. it. thessal. n. 1023.

Caule elato, stricto, simplici vel ramoso, alis angustis dentato spinosissimis marginato; foliis oblongo-lanceolatis, utrinque albo-tomentosis, in lacinias lobatas spinosas pinnatifidis partitisve; capitulis depresse globosis; involucri glabriusculi phyllis ab infimis brevibus triangularibus ad interiora sensim auctis, apice purpurascentibus, margine scabris, praeter intima erecto-patula saepius recurvo-reflexis, parte reflexa inferiori adpressa aequilonga; flosculis purpureis, viscidulis; acheniis brunneis, pappo dimidio brevioribus. — Variat involucri phyllis in spinam abruptiuscule vel sensim attenuatis.

In incultis, ruderatis, campis regionis inferioris et montanae per totam Graeciam. — Jun. Aug. ⊙.

7. **O. myriacanthum** Bois. diagn. ser. 2 VI. p. 114, Fl. or. III. p. 561; Heldr. chlor. Parn. p. 21. — Exsicc.: Orph. fl. gr. n. 55; Bald. it. cret. alt. n. 32.

Differt a praecedente, cui maxime affine, foliis angustioribus, in lacinias angustiores profundius partitis et praesertim involucri phyllis longe attenuatis, parte reflexa inferiori adpressa duplo longiore; — an tamen hujus varietas subalpina? (Bois. l. c.).

In sterilibus, dumosis regionis montanae. Attica: ad radices mt. Parnes pr. Dekeleiam (Hal.); Euboea: mt. Dirphys pr. Steni; mt. Parnassus pr. Rachova (Heldr.); Laconia: mt. Malevo pr. Xerocampos (Orph.); Creta: mt. Aphendi-Kristo in mt. Lassiti (Bald.). — Jun. Aug. ⊙.

β. Virescens.

8. **O. laconicum** Heldr. et Sart. ap. Rouy rev. Onop. p. 9. — *O. tauricum v, elatum* Spreitz. in z. b. G. 1887 p. 662, non Bois. — Exsicc.: Heldr. herb. norm. n. 1554.

Parce araneosum et pube brevi glandulosa obductum; caule elato, ramoso, alis angustis dentato-spinosissimis marginato; foliis oblongo-lanceolatis lanceolatisve, subtus valde insculpto-nervosis, in lacinias lobatas valide spinosas pinnatifidis partitisve; capitulis subglobosis; involucri glabri phyllis lanceolatis, ab externis recurvo-reflexis ad interna erecto-patentia sensim auctis, apice purpurascentibus, margine scabris, parte reflexa inferiori adpressa breviore; flosculis purpureis, glabris; acheniis brunneis, pappo dimidio brevioribus. — Foliis vix araneosis, viridibus, subtus eximie reticulato-venosis ab affinibus distinctum.

In sterilibus, dumosis, ad vias regionis inferioris. Laconia: in distr. Alagonia l. d. Dyrrhemma (Zahn), inter Levetzova et Marathonisi (Sart.); Cythaera: pr. Katuni (Spreitz.). — Jun. Aug. ⊙.

Obs. Quid sit *O. acaulon* S. et S. pr. II. p. 157; Ch. et B. exp. p, 243, Fl. pelop. p. 57; Friedr. Reise p. 275; Fraas fl. class. p. 205; mihi ignotum est.

49. Silybum Vaill. act. acad. Par. 1718 p. 172.

1. **S. marianum** L. sp. p. 823; S. et S. pr. II. p. 150; Ch. et B. exp. p. 240, Fl. pelop. p. 56; (*Carduus*); Gaertn. fruct. II. p. 378. t. 168; Friedr. Reise p. 274; Fraas fl. class. p. 206; Ung. Reise p. 124; Raul. cret. p. 790; Bois. fl. or. III. p. 556; Heldr. fl. cephal. p. 47, prosth. chlor. Thera p. 3; Hal. in z. b. G. 1888 p. 759, Beitr. fl. thessal. p. 14; Haussk. symb. p. 121. — Exsicc.: Orph. fl. gr. n. 769 *f. pygmaea* (Chios); Sint. it. thessal. n. 471.

Glabrum; caule simplici vel ramoso; foliis sinuato-pinnatifidis, supra secus nervos albovittatis, lobis rotundatis, spinosis, inferioribus basi attenuatis, superioribus diminutis, auriculato-amplexicaulibus; capitulis globosis; involucri phyllis exterioribus ovato-oblongis, in appendicem spinuloso-pectinatam, spina flavescente terminatam abeuntibus, intimis integris, cuspidatis; flosculis purpureis; acheniis pappo triplo brevioribus.

In herbidis, cultis regionis inferioris, probabiliter per totam Graeciam in Thessalia: pr. Kalabaka (Sint.), Karditza, Pharsalus

(Haussk.), ad viam ferream (Hal.); Doris: pr. Amphissa (Hal.); Attica: in oliveto Athenarum (Heldr.), insulae Petali (Münter); Argolis: pr. Nauplia (Haussk.), Poros (Friedr.); Cephalonia (Ung.): pr. Argostoli (Heldr.); Cycladum insula Thera (Heldr.); Creta: in planitie Messara (Raul.). — Maio, Jul. ⊙.

50. Tyrimnus Cass. dict. 41 p. 335.

1. **T. leucographus** L. sp. p. 820; S. et S. pr. II. p. 148; Sieb. avis p. 5; Ch. et B. fl. pelop. p. 56; Friedr. Reise p. 283; (*Carduus*); Cass. l. c.; Weiss in z. b. G. 1869 p. 44; Raul. cret. p. 791; Bois. fl. or. III. p. 555; Heldr. fl. cephal. p. 47; Form. in D. bot. Mon. 1890 p. 18, in Ver. Brünn 1896 p. 47, 1897 p. 34; Hal. Beitr. fl. Epir. p. 28, Beitr. fl. Aetol. p. 7, in ö. b. Z. 1897 p. 96; Haussk. symb. p. 125. — *Cirsium maculatum* Lam. fl. fr. II. p. 22. — *Carduus seminudus* Ch. et. B. exp. p. 240, non M. a. B. — Icon: Jacq. vind. III. t. 23. – Exsicc.: Rev. pl. cret. n. 85; Sint. it. thessal. n. 411.

Araneosum; caule gracili, erecto, inferne foliato alato, superne in ramos virgatos nudos abeunte; foliis tenuibus, breviter lobatis, spinuloso-ciliatis, supra secus nervos albo-vittatis, inferioribus obovato-oblongis, in petiolum brevem attenuatis, superioribus lanceolatis, decurrentibus; capitulis parvis ovatis; involucri phyllis lanceolatis, in spinulam brevem attenuatis; flosculis dilute purpureis; acheniis fuscis, pappo triplo brevioribus. — Habitu *C. pycnocephalo* similis, a quo generice filamentis monadelphis differt.

In herbidis, arenosis regionis inferioris et montanae. Epirus: ad Kalentini pr. Arta (Hal.); Thessalia: pr. Kalabaka (Sint.), Trikala, Povelci, Kataphygi, Larissa, Velestinos (Form.), Pharsalus, Karditza (Haussk.), mt. Pelion pr. Milies (Heldr.); Sporadum insula Jura (Leon.); Euboea: pr. Oreus (Reinhold), Petalium insula Xeronisi (Holzm.); Acarnania: pr. Sorovigli (Hal.); Aetolia: pr. Mesolongion (Nied.), mt. Taphiassos (Hal.); mt. Parnassus pr. Pania (Heldr.); Attica: mt. Kerata (Heldr.), ad Phalerum (Haussk.); Achaia: pr. Patras, Psatopyrgos (Hal.), Xylocastron (Orph.); Arcadia (Sibth.); Argolis: pr. Nauplia (Haussk.), Argos (Chaub.); Laconia: mt. Malevo (Orph.); Messenia: mt. Ithome (Heldr.), pr. Methone, Kalamata (Chaub.); Cephalonia: pr. Argostoli (Heldr.); Creta: pr. Platania (Rev.), Canea (Weiss), Aradhena, Aludha, Kritza (Raul.). — Maio, Jun. ⊙.

51. Galactites Moench. meth. p. 558.

1. **G. tomentosa** Moench. l. c.; DC. pr. VI. p. 616, cum *β. alata* (f. foliis longe decurrentibus); Marg. et R. fl. Zante p. 59; Ung. Reise p. 124; Weiss in z. b. G. 1869 p. 43; Raul. cret. p. 791; Bois. fl. or. III. p. 556; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 47; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Epir. p. 28, Beitr. fl. Aetol. p. 7, Beitr. fl. Achaia p. 24, in ö. b. Z. 1899

p. 25. — *Centaurea galactites* L. sp. p. 919; S. et S. pr. II p. 206; Sieb. avis. rem. p. 6. — *Calcitrapa galactites* Lam. fl. fr. II. p. 30. — *Cnicus galactites* Lois. fl. gall. p. 538. — *Carduus galactites* Ch. et B. exp. p. 242, Fl. pelop. p. 56. — Icon: Rchb. t. 151. — Exsicc.: Baen. herb. europ. n. 9262.

Caule gracili, erecto, ramoso, albo-tomentoso; foliis supra viridibus, glabratis, subtus albo-tomentosis, in lacinias triangulares vel lanceolatas, breviter spinosas, pinnatilobis partitisve, caulinis plus minus decurrentibus; capitulis mediocribus, pedunculatis, corymbosis; involucri parce araneosi phyllis in spinas subulato-triquetras abeuntibus; flosculis roseis, rarissime albis, marginalibus valde radiantibus; acheniis fuscis, pappo 4 plo brevioribus.

β. **elegans** All. fl. ped. I. p. 163 (*Centaurea*), pro sp.; DC. pr. VI. p. 617. — Foliis subintegris. — Exsicc.: Reis. fl. gr. a. 1898.

In herbidis, collibus, ad vias, in cultis regionis inferioris et montanae. Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.); Aetolia: pr. Mesolongion (Heldr.), Antirrhion (Hal.); Achaia: pr. Patras (Hal.); Elis: pr. Bartholomeo, Olympia (Heldr.); Messenia: pr. Methone (Chaub.); Laconia: pr. Sparta (Sart.), ad Xerocampos (Pichl.), mt. Taygetos (Chaub.); Cyclades (Sibth.): insula Andros (Sart.); Creta: pr. Canea (Weiss), Askyphos, Anopolis, Francocastello (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli (Spreitz.), Pessada, Koroni, Poros (Heldr.); Corcyra (Gelmi): Fortezza Abramo (Baen.); — β. in Strophadum insula majore (Reiser). — Maio, Jul. ⊙.

52. Chamaepeuce Prosp. Alp. exot. 77.

1. **C. mutica** Cass. dict. 44 p. 59 (*Ptilostemon*); DC. pr. VI. p. 657; Marg. et R. fl. Zante p. 60; Clem. sert. p. 55; Raul. cret. p. 792; Haussk. symb. p. 121. — *Staehelina chamaepeuce* L. syst. nat. ed. 12 II. p. 538; S. et S. pr. II. p. 163; Fl. gr. t. 847; Sieb. avis p. 5, rem. p. 6, in Flora I. p. 272. — *Serratula chamaepeuce* L. sp. ed. 2. p. 1147; Ch. et B. exp. p. 239; Fraas fl. class. p. 210. — *Pteronia chamaepeuce* Spreng. syst. III. p. 440. — *C. Alpini* Jaub. et Sp. ill. or. V. p. 26, t. 425; Bois. fl. or. III. p. 554; Spreitz. in z. b. G. 1877 p. 663; Heldr. fl. cephal. p. 47; Bald. viagg. Creta p. 68. — *C. gnaphaodes* Gelmi in bull. soc. bot. 1889 p. 450, an DC. pr. VI. p. 658, quae axillis foliorum superiorum bispinulosis et involucri phyllis elongatis spinescentibus parum differre videtur. — Exsicc.: Orph. fl. gr. n. 261; Heldr. herb. norm. n. 433 et 1257; Rev. pl. cret. n. 80; Bald. it. cret. n. 23; Dörfl. fl. aeg. n. 61,

Fructicosa; ramis novellis parte inferiori foliosis, tomentosis, superiori sparsius et brevius foliatis saepe glabrescentibus; foliis lineari-filiformibus, margine revolutis, supra glabris, subtus incano-tomentosis; capitulis terminalibus, 1—9, ovatis, pedunculatis, corymbosis; involucri parce tomentelli vel glabrescentis phyllis imbricatis, ab inferioribus triangu-

laribus, muticis vel breviter spinescentibus, ab intima lanceolata elongatis; flosculis purpureis.

α. **typica.** — *C. Alpini α. mutica* Bois. fl. or. III. p. 554. — Involucri phylla adpressa.

β. **polycephala** DC. pr. VI. p. 658 pro sp. — *Ch. Alpini β. camptolepis* Bois. fl or. III. p. 554, suppl. p. 309; Heldr. in ö. b. Z. 1898 p. 184. — Involucri phylla apice breviter patulo-curvata, saepe manifestius spinescentia. — Intermediis ad typum transit.

In rupium fissuis regionis inferioris et montanae. Aetolia: pr. Mesolongion (Nied.); mt. Parnassus (Fraas); Attica: mt. Parnes, Hymettus (Heldr.); Euboea: mt. Dirphys (Pichl.); Argolis (Sibth.); Messenia: pr. Pylos (Daenzer); Laconia: mt. Taygetos (Psarid.); Cyclades (Chaub.): insula Amorgos (Heldr.); Creta: pr. Malaxa (Rev.), secus rivulum Tavroniti infra Vukolies (Bald.), ad Palaeokastro pr. Kissamos, Aradhena, Pharangi Khordaliotikon, Spili et Smili ad radices mt. Kedros (Raul.) et ad Omalos (Rev.) in mt. Sphacioticis (Sieb.); Zante: inter urbem et Krionero (Marg.); Cephalonia: pr. Pharsa, Steno, Poros, Kipuria (Heldr.); Leucas: ad promontorium Zuana (Spreitz.); Corcyra: pr. Garuna (Letourn.), Dukades (Grimb.), Palaeokastrizza (Gelmi). — Maio, Jul. ♃.

2. **C. fruticosa** Desf. cat. hort. Par. p. 91 (*Carduus*); DC. pr. VI. p. 658; Bois. fl. or. III. p. 554; Hal. Beitr. fl. Achaia p. 5; Haussk. symb. p. 121. — *Ptilostemon appendiculatum* Cass. dict. 44 p. 59. — Exsicc.: Orph. fl. gr. n. 262.

Differt a praecedente statura robustiore, foliis longioribus, capitulis majoribus, phyllis in cuspidem apice spinosam multo longioribus attenuatis. — Flosculi in speciminibus a me in Aetolia lectis albi erant, an quoque in iis Argolidis e planta siccata dijudicare nequeo. — Certe species propria.

In rupibus mt. Chalkis ad sinum Patranum Aetolia (Hal.) et mt. Palamidi pr. Nauplia (Zuccar.). — Maio, Jun. ♄.

53. Jurinea Cass. in bull. philom. 1821 p. 140.

1. **J. mollis** L. amoen. IV. p. 328; S. et S. pr. II. p. 150: Ch. et B. exp. p. 240; Fl. pelop. p. 56 (*Carduus*); Rchb. fl. germ. exp. p. 290; Ung. Reise p. 124; Bois. fl. or. III. p. 575; Hal. in ö. b. Z. 1892 p. 371; Haussk. symb. p. 125. — Icon: Jacq. fl. austr. I. t. 18. — Exsicc.: Orph. fl. gr. n. 1118; Sint. et Bornm. it. turc. n. 1307.

Caule erecto, simplici, monocephalo, araneoso-floccoso, inferne tantum folioso; foliis supra araneosis, subtus albo-tomentosis, in lacinias oblongas vel lineari-lanceolatas, margine subrevolutas pinnatipartitis, rarius indivisis, radicalibus in petiolum attenuatis, caulinis sessilibus vel subdecurrentibus; capitulo majusculo, hemisphaerico; involucri plus minus arachnoideo-lanati phyllis herbaceis, lanceolatis, parte superiori

patentibus vel refractis, intimis chartaceis, acuminatis, rectis; flosculis purpureis; acheniis transverse lamellatis; pappi barbellati paleis intimis paulo longioribus et sublatioribus.

In saxosis montanis. Thessalia: pr. Litochoron ad radices Olympi (Sint.); Euboea: mt. Dirphys (Sprun.), pr. Kurbatzi (Wild.); Attica: mt. Parnes (Bois.), Pentelicon (Ung.), Hymettus, pr. Pikermi, Eleusis, mt. Pateras, pr. Megara (Heldr.); Peloponnesus (Sibth.): pr. Corinthum Chaub.). — Maio, Jun. ♃.

2. **J. glycacantha** S. et S. pr. II. p. 150; Fl. gr. IX. t. 826; Ch. et B. exp. p. 240; Fl. pelop. p. 56 (*Carduus*); DC. pr. VI. p. 677; Bois. fl. or. III. p. 576; Heldr. chlor. Parn. p. 21; Hal. Beitr. fl. Achaia p. 25, in ö. b. Z. 1897 p. 285; Haussk. symb. p. 125; Bald. riv. coll. bot. Alb. 1896 p. 69. — Exsicc.: Heldr. reliqu. Orph. a. 1887; Bald. it. alb. epir. IV. n. 178.

Praecedenti nimis forte affinis, ab ea differt foliis minus divisis, saepe integris, capitulo majore et praesertim involucri dense araneoso-lanati phyllis longe subulato-acuminatis.

In collibus herbidis, rupestribus montanis. Epirus: mt. Profetae Elias, ad Vromonero et mt. Kuruna distr. Ljaskovik (Bald.); Thessalia: pr. Pharsalus, Aivali, Orman Magula (Haussk.); Aetolia: mt. Korax (Tunt.); mt. Parnassus (Sibth.); Achaia: pr. Megaspilaeon (Hal.), mt. Chelmos (Orph.); Laconia (Sibth.): mt. Taygetos, pr. Megali Anastasova (Zahn). — Maio, Jul. ♃.

3. **J. anatolica** Bois. diagn. IV. p. 21, Fl. or. III. p. 574; Heldr. in ö. b. Z. 1898 p. 184. — *J. chia* Bois. et Orph. in Orph. fl. gr. exs. n. 771. — Exsicc.: Heldr. reliqu. Orph. a. 1887; Dörfl. fl. aeg. n. 133 a.

Caule erecto, simplici vel parce ramoso, 1—5 cephalo, araneoso-floccoso, inferne praesertim folioso; foliis supra araneosis, subtus albo-tomentosis, in lacinias oblongas vel lineari-lanceolatas, margine subrevolutas pinnatipartitis, infimis interdum indivisis, radicalibus in petiolum attenuatis, caulinis sessilibus vel subdecurrentibus; capitulis mediocribus, subglobosis; involucri plus minus arachnoideo-lanati phyllis herbaceis, lanceolatis, apice patentibus vel reflexis, intimis chartaceis, subulato-acuminatis, subpungentibus, rectis; flosculis purpureis; acheniis scrobiculatis; pappi barbellati setis 4—5 intimis elongatis, latioribus et longioribus. — Differt ab antecedentibus caule saepius ramoso, capitulis minoribus et pappi serie intima longiori latiorique.

In rupestribus montanis pr. Apiranthos insulae Naxos (Orph.) et probabiliter in aliis Cycladum insulis. — Maio, Jul. ♃.

11. Tribus. **CENTAUREAE** Less. Syn. p. 53.

54. Phaeopappus Bois. diagn. VI. p. 122.

1. **P. saxatilis** C. Koch in Linn. XXIV. p. 419 (*Centaurea*); Bois. fl. or. III. p. 602. —

Glabrescens; caulibus a collo fibroso simplicibus, monocephalis, folia radicalia non superantibus; foliis viridibus, margine papillari-ciliatis, radicalibus crebris, oblongis, acutis, obsolete denticulatis, in petiolum longe attenuatis, caulinis paucis, brevius petiolatis, capitulum superantibus; capitulo mediocri, ovato-haemisphaerico; involucri glabri phyllis paucis, viridi-lutescentibus, adpressis, infimis breviter lanceolatis, integris, breviter spinosis, ceteris oblongo-lanceolatis, sub spina brevi margine angusto, decurrenti, brevissime ciliolato obsitis; flosculis . . . . ; ovario dense villoso.

Creta (Gundelsh.); a recentioribus non lectum. — ♃. N. v.

55. Centaurea L. gen. n. 984.

Dispositio specierum:

1. Sectio. *Aplolepideae* DC. pr. VI. p. 565 p. p. — Appendix phyllorum, involucri scariosa, hyalina, obtusa vel molliter cuspidata, integra vel denticulata, saepius lacera, raro decurrens, nunquam ciliata; achenia saepissime papposa, hilo nudo. — Perennes vel biennes, capitulis mediocribus (1—2 cm. diam.), raro minoribus (*C. subciliaris*), flosculis purpureis vel flavis.

1. Subsectio. *Centaurium* DC. l. c. p. 566. — Involucri phylla adpressa, appendice obtusiuscula, anguste decurrente; achenia (in nostra) transverse rugulosa, papposa; pappi series exteriores setosae, intima brevior, late paleacea.

1. C. amplifolia Bois. et Heldr.

2. Subsectio. *Phalolepis* DC. l. c. p. 568. — Involucri phylla saepius subpatula, appendice obtusiuscula vel molliter cuspidata, non decurrente; achenia laevia, saepissime papposa; pappus setosus, serie intima brevissima.

a. Plantae caespitosae, saepius albo-tomentosae.

α. Flosculi flavi.

2. C. musarum Bois. et Orph.

β. Flosculi purpurei.

× Folia saltem inferiora bipinnatisecta.

3. C. Heldreichii Hal. **4. C. ustulata** Hal.

×× Folia lyrata vel pinnatisecta.

5. C. cadmea Bois. **6. C. epirotica** Bald.

b. Plantae non caespitosae, plus minus pilosae.

α. Biennes.

× Phyllorum appendices centro pallide fuscae.

7. C. princeps Bois. et Heldr. **8. C. brunnea** Hal.

×× Phyllorum appendices centro nigrae.

9. C. deusta Ten.

××× Phyllorum appendices immaculatae.

10. C. alba L.

β. Perennes.

11. C. subciliaris Bois. et Heldr. **12. C. amara** L.

2. Sectio. *Jaceineae* DC. l. c. p. 570. — Appendix phyllorum involucri scariosa, fuscidula, exteriorum saltem pectinatim-ciliata, phylla tegens, non decurrens; achenia papposa vel epapposa, hilo nudo. — Perennes, capitulis mediocribus, flosculis purpureis.

1. Subsectio. *Jacea* Cass. dict. 24 p. 88. — Appendix phyllorum subrotunda, obtusa, saltem exteriorum pectinatim-ciliata.

13. C. nigrescens Willd.

2. Subsectio. *Lepteranthus* DC. diss. comp. I. p. 24. — Appendix phyllorum lanceolato-subulata, crebre pectinatim-ciliata.

14. C. plumosa Lam.

3. Sectio. *Cyaneae* DC. pr. VI. p. 577. — Appendix phyllorum involucri scariosa, plerumque nigricans, pectinatim-ciliata, saepius mucrone vel spina terminata, phylla raro tegens, utrinque decurrens; achenia papposa vel epapposa, hilo barbato vel nudo. — Annuae, biennes vel perennes, capitulis parvis vel magnis, flosculis varie coloratis.

1. Subsectio. *Cyanus* DC. l. c. p. 578. — Appendix plerumque usque ad basin phyllorum decurrens, obtusa vel acutiuscula; achenia papposa, hilo barbato. — Annuae vel perennes, capitulis mediocribus, flosculis plerumque azureis, rarius flavis, albis vel purpureis.

a. Annuae.

15. C. cyanus L. **16. C. depressa** M. a. B.

b. Perennes.

α. Flosculi azurei vel purpurei.

17. C. variegata Lam. **18. C. epirota** Hal.

β. Flosculi ochroleuci.

19. C. pindicola Griseb. **20. C. Baldaccii** Deg.

2. Subsectio. *Acrolophus* DC. l. c. p. 581. — Appendix phyllorum plerumque breviter vel vix decurrens, saepius mucronata; achenia saepissime papposa, hilo nudo. — Biennes vel perennes, capitulis parvis vel mediocribus, flosculis purpureis, raro flavis vel albis.

§. Biennes vel perennes.

a. Capitula solitaria, pedunculata.

α. Capitula ovata vel ovato-globosa. — Caulis plerumque parce ramosus: phyllorum mucro plerumque brevis (ciliis saepe non longior), raro validior.

× Pappus achenio sesquilongior; planta albo-tomentosa.

21. C. Niederi Heldr.

×× Pappus achenio plerumque brevior, raro sublongior.

○ Flosculi lutei; planta albo-tomentosa.

22. C. argentea L.

○○ Flosculi purpurei, rarissime (*C. laureotica*) ochroleuci.

. Folia caulina vel earum laciniae latiusculae, ovatae vel ellipticae; appendix phyllorum pallida, mucrone debili, ciliis breviori terminata.

23. C. sublanata (DC.). **24. C. pallida** Friv.

.. Folia caulina vel earum laciniae angustae, oblongae vel lineares.

; Capitula ovata, circa 8 mm. diametro; appendix phyllorum pallida vel fuscidula, triangulari-lanceolata, mucronata, ad basin hyalino-marginata, ciliis mucrone brevioribus, saltem apice hyalinis.

, Flosculi purpurei.

.— Folia viridia vel leviter canescentia, plus minus asperula.

— Folia radicalia lyrata, lobo terminali oblongo, 5—8 mm. lato.

25. C. kalambakensis Fr. et Sint. **26. C. transiens** Hal.

= Folia radicalia pinnatisecta, laciniis anguste linearibus.

27. C. pentelica Haussk. **28. C. asperula** Hal.

:= Folia albo-tomentosa, vix asperula.

29. C. attica Nym.

,, Flosculi ochroleuci.

30. C. laureotica Heldr.

;; Capitula ovata vel ovato-globosa, 5—12 mm. diametro; appendix phyllorum pallida, fusca vel nigricans, triangulari-rotundata et breviter mucronata vel semilunata, ad basin hyalino-marginata, ciliis mucrone aequilongis vel longioribus, raro brevioribus (et tunc capitula ovato-globosa 12 mm. diametro).

, Caulis et folia dense albo-lanata, non aspera; appendix phyllorum nigra, mucrone recurvato.

31. C. ossaea Hal.

,, Caulis et folia puberula vel leviter araneoso-lanata, plus minus aspera; appendix pallida vel nigra, mucrone erecto.

32. C. lacerata Haussk. **33. C. affinis** Friv. **34. C. macedonica** Griseb.

β. Capitula ovato-oblonga, parva. — Caulis valde ramosus; phyllorum mucro validior, ciliis plerumque longior et patens vel recurvatus.

× Flosculi purpurei.

○ Appendix phyllorum in spinulam erectam attenuata; capitula 10 mm. longa.

35. C. tymphaea Haussk. **36. C. brevispina** Haussk.

⊙⊙ Appendix phyllorum in spinulam patulam attenuata; capitula 10 mm. longa.

37. C. Orphanidea Heldr. et Sart.

⊙○○ Appendix phyllorum in spinulam recurvatam attenuata, capitula 10—15 mm. longa.

38. C. drakiensis Fr. et Sint.

×× Flosculi lutei.

39. C. pelia DC.

××× Flosculi albi.

40. C. lactiflora Hal. **41. C. diffusa** Lam.

b. Capitula sessilia, in glomerulos terminales aggregata.

42. C. Zuccariniana DC.

§§ Suffruticosa, erinaceo-dumosa, caulibus spinosis.

43. C. spinosa L.

3. Subsectio. *Acrocentron* DC. l. c. p. 586. — Appendix phyllorum breviter vel vix decurrens, plerumque spina valida terminata, raro inermis; achenia papposa, hilo barbato. — Perennes capitulis magnis vel mediocribus, flosculis purpureis, raro luteis.

a. Flosculi purpurei.

α. Caulescentes, saepius elatae.

× Appendix phyllorum mutica vel breviter mucronata.

44. C. Oliveriana DC. **45. C. ebenoides** Heldr.

×× Appendix phyllorum spina valida terminata (cf. *C. Guicciardii var. β.*).

○ Appendix phyllorum magna, ovata, phylla tegens.

46. C. Sibthorpii Hal. **47. C. achaia** Bois. et Heldr.

CO Appendix phyllorum semilunata, saepe angustissima, phylla non tegens.

. Appendix phyllorum nigra.

48. C. redempta Heldr.

.. Appendix phyllorum pallida vel fusca.

; Folia inferiora bipinnatisecta.

49. C. subarachnoidea (Bois. et Heldr.). **50. C. psilacantha** Bois. et Heldr.

;; Folia inferiora pinnatisecta vel indivisa.

, Folia laevia, arachnoideo-cana, segmentis petiolatis.

51. C. graeca Griseb.

,, Folia scabrida puberula, saepius asperula, segmentis sessilibus.

.— Folia inferiora lyrato-pinnatisecta vel pinnatisecta.

52. C. Spruneriana Schultz. **53. C. Guicciardii** Bois.

54. C. Halácsyi Dörfl.

:= Folia inferiora obovata integra vel alia lyrata.

55. C. minor Heldr.

β. Caulis nullus vel brevis, capitula radicalia.

56. C. mixta DC. **57. C. myconia** Bois. et Sart. **58. C. raphanina** S. et S.

b. Flosculi flavi.

α. Caulis 20—60 cm. altus.

59. C. Tuntasia Heldr. **60. C. salonitana** Vis.

β. Caulis 1—3 cm. altus.

61. C. parnonia Hal.

4. Sectio. *Calcitrapeae* DC. l. c. p. 592. — Appendix phyllorum involucri cornea, pallida vel fuscidula, non decurrens, spina saepe utrinque spinulosa terminata, phylla non tegens; achenia papposa vel epapposa, hilo nudo. — Annuae vel biennes, capitulis mediocribus, flosculis flavis vel purpureis.

a. Folia caulina decurrentia; flosculi flavi.

α. Flosculi eglandulosi.

62. C. solstitialis L. **63. C. idaea** Bois. et Heldr.

β. Flosculi glandulosi.

64. C. melitensis L.

b. Folia non decurrentia.

α. Flosculi purpurei.

65. C. calcitrapa L. **66. C. iberica** Trev.

β. Flosculi sulfurei.

67. C. hyalolepis Bois.

5. Sectio. *Seridieae* DC. l. c. p. 598. — Appendix phyllorum involucri cornea, pallida vel nigrescens, non decurrens, palmato-spinosa, phylla non tegens: achenia omnia aut saltem disci papposa, hilo nudo. — Annuae vel perennes, capitulis mediocribus, flosculis purpureis vel flavis.

a. Flosculi purpurei.

α. Folia caulina basi auriculato-amplexicaulia, non decurrentia.

68. C. sphaerocephala L.

β. Folia caulina decurrentia.

69. C. sonchifolia L. **70. C. napifolia** L.

b. Flosculi lutei.

71. C. lancifolia L.

1. Sectio. *Aplolepideae* DC. pr. VI. p. 565 p. p. —

1. Subsectio. *Centaurium* DC. l. c. p. 566. —

1. **C. amplifolia** Bois. et Heldr. diagn. pl. or. ser. 2, III. p. 68, Fl. or. III. p. 623; Hal. in bull. herb. Bois. VI. p. 571. Exsicc.: Heldr. herb. n. 2067; Orph. fl. gr. n. 125.

Caule elato, glabro vel inferne hirsuto, in ramos 2—4, nudos abeunti; foliis cartilagineis, inferioribus amplis, petiolatis, lobato-pinnatisectis, glabris vel ad petiolum et subtus ad nervum hirsutis, segmentis 3—5, ovato-rotundatis vel oblongis, duplicato-serratis, secus rhachidem decurrentibus, terminali plerumque multo majore, foliis superioribus 2—3 diminutis, sessilibus, pinnatipartitis vel lyratis; capitulis mediocribus, ovatis, nudis; involucri glabri phyllis striatis, apicem versus nigro-marginatis, exterioribus orbiculatis, scarioso-marginatis, mediis ovatis, cum intimis lineari-ellipticis, appendice suborbiculata, hyalina, saepissime lacera, concava, mutica terminatis; flosculis pallide purpureis, valde radiantibus; acheniis transverse rugulosis, glabris; pappi serie penultima achenio vix breviore, interiore tertia parte breviore, late paleacea. — Planta spectabilis, 50—150 cm. alta.

In faucibus umbrosis regionis abietinae rarissime, hucusque tantum in mt. Kyllene pr. Flamburitza (Heldr.). — Jun. Jul. ♃.

2. Subsectio. *Phalolepis* DC. l. c. p. 568.

a. Plantae caespitosae, saepius albotomentosae.

α. Flosculi flavi.

2. **C. musarum** Bois. et Orph. diagn. ser. 2, V. p. 112, Fl. or. III. p. 623; Heldr. chlor. Parn. p. 21; Hal. in bull. herb. Bois. VI. p. 572. — Exsicc.: Orph. fl. gr. n. 308; Heldr. herb. norm. n. 321.

Adpressissime albo-tomentosa; caulibus pumilis, simplicibus vel parce ramosis, 1—4 cephalis, tota longitudine foliatis; foliis oblongis vel obovato-oblongis, lyratis vel pinnatisectis, inferioribus in petiolum attenuatis, superioribus sessilibus, summis diminutis, plerumque indivisis, segmentis ovatis vel oblongis, obtusiusculis; capitulis mediocribus, subglobosis, nudis; involucri glabri phyllis fuscescentibus, striatis, appendicibus orbiculatis, hyalinis, cucullatis, obtusis, erosulo-denticulatis obtectis; flosculis vix radiantibus; acheniis glabris, pappo dimidio brevioribus. — Species pulcherrima, caulibus 10—20 cm. altis, inter affines floribus luteis insignis.

In rupestribus regionis abietinae mt. Parnassi loco Klistozasteno dicto, rarissime (Orph.). — Jul. Aug. ♃.

β. Flosculi purpurei.

× Folia saltem inferiora bipinnatisecta.

3. **C. Heldreichii** Hal. Beitr. fl. Aetol. p. 7, t. 2, in bull. herb. Bois. VI. p. 573. — Exsicc.: Hal. it. gr. II. a. 1893; Heldr. herb. norm. n. 1259.

Adpresse albo-tomentosa; caulibus erectis vel adscendentibus, composite et corymbose ramosissimis, polycephalis, foliatis, foliis rosularum sterilium et caulinis inferioribus longe petiolatis, bipinnatisectis, laciniis oblongo-linearibus, mucronulatis, superioribus sursum diminutis, pinnatisectis; capitulis mediocribus, subglobosis, nudis; involucri glabri phyllis striatis in appendices scariosas, late-orbiculatas, centro pallide fuscas, margine hyalinas, denticulatas, laceras, breviter molliterque aristatas abeuntibus; flosculis radiantibus; acheniis puberulis, pappo aequilongis. — Species elegans, 20—30 cm. alta, habitu *C. cinereae* Lam. non absimilis.

In rupibus calcareis mare imminentibus mt. Chalkis — Varassova hodie — pr. Krioneri ad sinum Patranum Aetoliae (Hal.). — Apr. Maio. ♃.

4. **C. ustulata.** — *C. deusta* Bald. riv. coll. bot. Alb. 1896 p. 69, non Ten. — Exsicc.: Bald. it. alb. epir. IV. n. 179.

Canescens, caulibus erectis vel adscendentibus, parce ramosis, 2—4 cephalis, foliatis, petiolisque papillis crispulis patulis crebre obsitis, viscidulis; foliis hirtis, rosularum sterilium et caulinis inferioribus petiolatis, bipinnatisectis, laciniis anguste linearibus, mucronulatis, superioribus sursum diminutis, pinnatisectis; capitulis mediocribus, subglobosis, nudis; involucri glabri phyllis vix striatis, in appendices scariosas, late orbiculatas, centro pallide fuscas, margine hyalinas, denticulatas, laceras, breviter molliterque aristatas abeuntibus; flosculis parum radiantibus; acheniis puberulis, pappo subbrevioribus. — Praecedenti affinis, ab ea indumento viscidulo, non albo-tomentoso, caulibus gracilioribus, parce ramosis, foliorum laciniis multo angustioribus, involucri phyllis vix striatis, margine latius hyalinis specifice differt. — A. *C. deusta* Ten. toto coelo diversa.

In praeruptis mt. Konitza supra Gorica secus viam ad Papingon in Epiro boreali (Bald.). — Jul. ♃.

×× Folia pinnatisecta vel lyrata.

5. **C. cadmea** Bois. diagn. VI. p. 16; Hal. in bull. herb. Bois. VI. p. 572. — Exsicc.: Guicc. pl. graec. a. 1856.

Adpresse albo-tomentosa; rhizomate indurato; caulibus extrarosularibus, adscendentibus vel decumbentibus, 15—30 cm. altis, dichotome et subdivaricatim 5—7 cephalis; foliis radicalibus petiolatis, lyratis, segmentis subintegris, mucronulatis, lateralibus utrinque 3—5, oblongis, terminali ovato, 2—3plo majore, foliis caulinis sessilibus, pinnatipartitis, summis oblongo-linearibus, minimis; capitulis mediocribus, ovatis; involucri phyllis glabris, appendicibus orbiculatis, centro fuscis, margine late albo-hyalinis, denticulatis vel laceris, setaceo-cuspidatis obtectis; flosculis purpureis; acheniis glabriusculis, pappo subbrevioribus.

In regione abietina mt. Cithaeron Atticae, ubi Guicciardi a. 1856 specimen unicum legit. — Jun. Jul. ♃.

6. **C. epirotica** Bald. riv. coll. bot. Alb. 1896 p. 69 pro var. *C. deustae.* — Exsicc.: Bald. it. alb. epir. IV. n. 243.

Floccoso-lanata, canescens; rhizomate indurato; caulibus extrarosularibus, 7—12 cm. altis, monocephalis; foliis pectinato-pinnatifidis, segmentis utrinque 4—8, brevibus, integris vel parce denticulatis, mucronulatis, inferioribus breviter petiolatis, superioribus sursum diminutis, sessilibus; capitulis parvis, ovatis; involucri phyllis glabris, appendicibus orbiculatis, centro nigris, margine late albo-hyalinis, laceris, setaceo-cuspidatis, obtectis; flosculis? purpureis; acheniis . . . — Species pumila, foliis pectinatis, capitulis parvis, involucri phyllis nigris egregia; a. *C. deusta* jam rhizomate caespitoso diversissima.

In lapidosis mobilibus summis mt. Smolika supra Kerasovo distr. Konitza in Epiro (Bald.). — Jul. Aug. ♃.

b. Plantae non caespitosae, plus minus pilosae.

α. Biennes.

× Phyllorum appendices centro pallide fuscae.

7. **C. princeps** Bois. et Heldr. diagn. ser. 2, VI. p. 111; Hal. in bull. herb. Bois. VI. p. 573. — *C. alba γ. princeps* Bois. fl. or. III. p. 621. — Exsicc.: Heldr. pl. gr. n. 3298.

Caule erecto, crassiusculo, scabrido et papillis crispulis patulis viscidulo, paulo supra basin composite et corymbose ramosissimo, polycephalo; foliis papilloso-scabridis, punctatis, radicalibus sub anthesi jam emarcidis, petiolatis, bipinnatisectis, laciniis oblongo-linearibus, mucronulatis, caulinis sessilibus, late lanceolatis, pinnatilobatis, summis oblongo-lanceolatis, integris, capitulo approximatis; capitulis ovatis; involucri phyllis omnino scarioso-membranaceis, glabris, pallidis, squarroso-patentibus, appendicibus orbiculatis vel ovatis, cucullatis, centro pallide fuscescentibus, margine hyalinis, denticulatis, saepe fissis, longiuscule setaceo-cuspidatis obtectis; flosculis albidis, parum radiantibus; acheniis glabriusculis, pappo subtriplo longioribus. — Indumento viscido, caulibus et ramis crassiusculis strictis, foliis caulinis vix dissectis, capitulis fere duplo majoribus et appendicibus centro fuscis ab affini *C. alba* specifice differt.

In summo cacumine Petra mt. Veluchi Eurytaniae (Samarit.). — Jul. Aug. ⊙.

8. **C. brunnea** Hal. in bull. herb. Bois. VI. p. 575 pro var. *C. deustae.* — *C. alba v. princeps* Bald. riv. coll. bot. Alb. 1895 p. 50, non Bois. — Exsicc.: Bald. it. alb. epir. III. n. 72.

Caule erecto, crassiusculo, scabro et parce subfloccoso, non viscidulo, a medio composite et corymbose ramosissimo, polycephalo; foliis coriaceis, valde scabris, punctatis, inferioribus petiolatis, pinnatipartitis, segmentis elongatis, oblongo-linearibus, mucronulatis, summis lanceolatis, integris; capitulis ovatis; involucri phyllis omnino scarioso-membranaceis, glabris, pallidis, subpatentibus, appendicibus ovatis, convexis, centro fuscescentibus, margine hyalinis, integris, saepe fissis, breviter setaceo-cuspidatis, obtectis; flosculis roseis, vix radiantibus; acheniis glabriusculis, pappo subtriplo longioribus. — A praecedente indumento non papilloso-

viscidulo, caulibus elatioribus, foliis coriaceis, eis *C. scabiosae* similibus, capitulis minoribus, phyllis angustioribus, appendicibus integris, breviter cuspidatis; a sequente caule corymboso, ramis brevioribus crebre foliatis, foliis amplioribus, segmentis elongatis, capitulis subminoribus ovatis, phyllis pallide maculatis distinguitur.

In rupestribus mt. Zalongos distr. Prevesa in Epiro (Bald.). — Jun. Jul. ⊙.

×× Phyllorum appendices centro nigrae.

9. **C. deusta** Ten. fl. nap. pr. p. 57, Fl. neap. t. 84; Bald. in bull. herb. Bois. IV. p. 205; Hal. in bull. herb. Bois. VI. p. 574. — *C. alba β. deusta* DC. pr. VI. p. 569; Bois. fl. or. III. p. 621, suppl. p. 312. — Exsicc.: Orph. fl. gr. n. 1086 (Macedonia); Heldr. pl. fl. hellen. a. 1887.

Caule erecto, scabrido, inferne saepe papilloso-subviscidulo, patule ramoso, ramis elongatis, parce foliatis; foliis scabridis, inferioribus petiolatis, pinnatipartitis, segmentis oblongo-lanceolatis, mucronulatis, ceteris diminutis, sessilibus, pinnatisectis, summis lanceolatis, integris; capitulis subglobosis; involucri phyllis scarioso-membranaceis, glabris, fuscis, subpatentibus, appendicibus ovatis, centro nigris, margine hyalinis, integris vel denticulatis, saepe fissis, breviter setaceo-cuspidatis obtectis; flosculis roseis, vix radiantibus; acheniis glabriusculis, pappo subtriplo longioribus. —

In dumosis regionis montanae. Thessalia: mt. Olympus (Auch.), mt. Pelion pr. Zagora (Aphentulis); Leucas: pr. Euklivi (Bald.); Cephalonia: pr. Livadi (Letourn.). — Jun. Aug. ⊙.

××× Phyllorum appendices immaculatae.

10. **C. alba** L. sp. p. 914; Clem sert. p. 58; Hal. in bull. herb. Bois. VI. p. 575. — ? *C. princeps* Haussk. symb. p. 125; Form. in Ver. Brünn 1897 p. 35; vix Bois. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Sint. it. thessal. n. 145.

Differt a praecedente caulibus corymbose ramosis, ramis brevioribus, crebrius foliatis, capitulis minoribus, involucri phyllis concoloribus, albidis.

In dumosis regionis montanae. Thessalia: mt. Baba pr. Klinovo in Pindo (Heldr.), ad monasterium Hagios Stephanos pr. Kalabaka (Heldr.), mt. Pirgo et Jurti in mt. Othrys (Form.); Corcyra (Clem.). — Jun. Aug. ⊙.

β. Perennes.

11. **C. subciliaris** Bois. et Heldr. fl. or. III. p. 627; Heldr. fl. cephal. p. 47; Hal. in bull. herb. Bois. VI. p. 576. — Exsicc.: Heldr. pl. cephal. a. 1872.

Adpresse araneoso-canescens; caulibus erectis, simplicibus vel divaricatim ramosis; foliis inferioribus oblongo-spathulatis, in petiolum attenuatis, integris vel sublyratis, lobis utrinque 1—4, minutis, mucronatis, foliis intermediis in lacinias lineares breves pinnatipartitis, summis squa-

maeformibus; capitulis oblongis, basi nudis; involucri phyllis scariosis, glabris, adpressis, appendicibus ovatis, centro fuscidulis, margine late hyalinis, denticulatis, cuspide brevi subspinosa terminatis; flosculis roseis, breviter radiantibus; acheniis glabris, pappo 4plo longioribus. — Caulis 2—25 cm. altus, capitula 5—10 mm. diametro. — Habitu *C. amarae*, ab ea foliis, capitulis minoribus, appendicibus cuspidatis et acheniis papposis distincta.

In regione media mt. Aeni pr. Balsamata oropedii Omala, nec non in regione abietina mt. Rhudi Cephaloniae (Heldr.). — Aug. ♃.

12. **C. amara** L. sp. ed. 2, p. 1292; Haussk. symb. p. 125; Form. in Ver. Brünn 1896 p. 48; Hal. in bull. herb. Bois. VI. p. 576. — *C. jacea* Form. in D. bot. Mon. 1890 p. 19, in Ver. Brünn 1895 p. 27. — *C. jacea v. montana* Form. in Ver. Brünn 1896 p. 48, 1897 p. 35. — Exsicc.: Sint. it. thessal. n. 1074.

Adpresse araneoso-canescens; caulibus erectis vel adscendentibus, simplicibus vel divaricatim ramosis; foliis oblongo-vel lineari-lanceolatis, integris, mucronatis, inferioribus in petiolum attenuatis, interdum pinnatifidis, superioribus sessilibus; capitulis ovato-globosis, foliis summis suffultis; involucri phyllis scariosis, glabriusculis, appendicibus orbiculatis vel ovatis, concavis, fuscidulis, margine anguste hyalinis, denticulatis, muticis; flosculis roseis, breviter radiantibus; acheniis puberulis, calvis. — Subspecies australis *C. jaceae* L., a qua indumento araneoso, foliis angustioribus et ramis virgatis discrepat et cum praecedente ad sectionem sequentem transitum praebet.

In herbidis montanis. Epirus: pr. Prevesa, Han Balduma, in valle Dipotami (Form.); Thessalia: in oropedio Neuropolis pr. Sermenikon (Haussk.), pr. Korona, Koturi, Velitsena, mt. Dokimi in Pindo, mt. Godaman in Olympo (Form.). — Jul. Sept. ♃.

2 Sectio. *Jaceineae* DC. pr. VI. p. 570.

1. Subsectio *Jacea* Cass. dict. 24 p. 88.

13. **C. nigrescens** Willd. sp. III. p. 2288; Bois. fl. or. suppl. p. 313; Form. in Ver. Brünn 1896 p. 48; Hal. in bull. herb. Bois. VI. p. 577. — Icon: Rchb. XXV. t. 25.

Papilloso-asperula, viridis; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis oblongis oblongo-lanceolatisve, integris, dentatis vel lyrato-sinuatis, inferioribus petiolatis, superioribus sessilibus, capitulis ovato-globosis, foliis summis suffultis; involucri nigricantis phyllis appendice scariosa orbiculari-ovata, profunde pectinatim ciliata superatis; flosculis roseis, radiantibus; acheniis puberulis, calvis. — Habitu praecedentis, ab ea virescentia, ramis brevioribus, foliis latioribus et praecipue appendicibus crebre et profunde pectinatim-ciliatis distincta.

In herbidis subalpinis. Thessalia: mt. Dokimi in Pindo (Form.), mt. Oeta Phthiotidis (Heldr.). — Jul. Sept. ♃. N. v.

2. Subsectio. *Lepteranthus* DC. diss. comp. I. p. 24.

14. **C. plumosa** Lam. fl. fr. II. p. 51 (*Jacea*): Kern in ö. b. Z. 1872 p. 44; Hal. in bull. herb. Bois. VI. p. 578. — *C. nervosa* Willd. en. hort. berol. II. p. 925; Haussk. symb. p. 125. — Icon: Rchb. IV. t. 554.

Papilloso-asperula, viridis; caulibus erectis, simplicibus, monocephalis; foliis lanceolatis, denticulatis, inferioribus in petiolum attenuatis, superioribus sessilibus; capitulis subrotundis, foliis summis suffultis; involucri fuscescentis phyllis exterionibus appendice e basi lanceolata, longe subulata, recurvata, pinnato-fimbriata, phyllis intimis appendice subrotunda, lacero-dentata superatis; flosculis roseis, radiantibus; acheniis glabris, pappo triplo longioribus.

In nemorosis supra monasterium Korona in Pindo (Haussk.). — Jul. Aug. ♃. N. v.

3 Sectio. *Cyaneae* DC. pr. VI. p. 577.

1. Subsectio. *Cyanus* DC. l. c. p. 588.

a. Annuae.

15. **C. cyanus** L. sp. p. 911; S. et S. pr. II. p. 198; Ch. et B. exp. p. 255. Fl. pelop. p. 59; Fraas fl. class. p. 204; Ung. Reise p. 124; Bois. fl. or. III. p. 634; Spreitz. in z. b. G. 1877 p. 713, 1887 p. 663; Heldr. chlor. Parn. p. 22; Hal. Beitr. fl. Aetol. p. 8, Beitr. fl. Achaia p. 25, in ö. b. Z. 1897 p. 285, in bull. herb. Bois. VI. p. 578; Form. in Ver. Brünn. 1897 p. 35; Bald. riv. coll. Alb. 1896 p. 70. — Icon: Fl. dan. t. 993. — Exsicc.: Heldr. herb. norm. n. 623.

Arachnoideo-tomentella; caulibus erectis vel adscendentibus, saepe a basi ramosis; foliis radicalibus obovato-lanceolatis vel lyratis, in petiolum attenuatis, caulinis lineari-lanceolatis, sessilibus; capitulis ovatis; involucri phyllis laxe arachnoideis, pallidis, appendice scariosa, fuscescenti, in dentes acute triangulares, apice argenteos, phylli latitudine duplo breviores partita marginatis; flosculis azureis, valde radiantibus; acheniis puberulis, pappo sublongioribus.

In rupestribus regionis inferioris et montanae. Epirus: pr. Janina (Bald.); Thessalia: mt. Cuka et Hagios Elias mt. Chassia (Form.), mt. Agrapha in Pindo (Heldr.); pr. Pagna ad radices Parnassi (Samar.); Aetolia: pr. Krioneri ad radices mt. Chalkis (Hal.), pr. Agrinion, Mesolongion (Nicl.); Arcadia: pr. Zatuna (Orph.); Laconia: mt. Malevo (Orph.); Taygetos (Psarid.); Messenia: pr. Kalamata (Zahn); Achaia: mt. Omplo pr. Patras (Heldr.); Ithaca (Ung.); pr. Vathy (Spreitz.); Leucas: ad monasterium Hagios Georgios (Spreitz.); Corcyra: mt. Pantocrator et Hagios Deca (Baen.). — Apr. Jul. ⊙.

16. **C. depressa** M. a Bieb. fl. taur. cauc. II. p. 346; Haussk. symb. p. 125; Hal. in bull. herb. Bois. VI. p. 579. — Exsicc.: Heldr. herb. norm. n. 947.

Adpresse-canescens; caulibus erectis vel adscendentibus, parce ramosis; foliis inferioribus petiolatis, oblongis, integris vel lyrato-pinnatifidis, superioribus lineari-lanceolatis; capitulis ovatis; involucri phyllis glabris,

pallidis, appendice scariosa, alba vel fuscescenti, in dentes lanceolatos, albos, phylli latitudinem subaequantes partita marginatis; flosculis azureis, valde radiantibus; acheniis puberulis, pappo aequilongis vel subbrevioribus. — Foliis latioribus, caulinis etiam inferioribus pinnatifidis, capitulis majoribus, appendicibus latioribus, longius dentatis et pappo longiore a praecedente diversa.

Inter segetes in arvis. Thessalia: pr. Pharsalus (Haussk.), Volo (Heldr.); Attica: pr. Paliokundura, Amarysia, Liosia (Heldr.); Arcadia: pr. Tripolis, Zatuna (Orph.). — Mart. Apr. ⊙.

Obs. *C. cyanoides* Bergg. et Wahlenb. in Isis 1828 p. 21; Heldr. herb. norm. n. 1261. — A praecedente acheniis praesertim calvis distincta, a Charrel pr. Thessalonicam lecta, in Thessalia inquirenda.

b. Perennes.

α. Flosculi azurei vel purpurei.

17. **C. variegata** Lam. dict. I. p. 668; Hal. in bull. herb. Bois. VI. p. 580. — *C. seusana* Chaix in Vill. fl. Dauph. III. 52. — *C. cana* S. et. S. pr. II. p. 198; Hal. Beitr. fl. Achaia p. 25; Bald. riv. coll. bot. alb. 1895 p. 50, 1896 p. 70. — *C. lingulata* Lag. elench. p. 32. — *C. axillaris* Ch. et B. fl. pelop. p. 59; non Willd. — C. *axillaris v. cana* Bois. fl. or. III. p. 636; Haussk. symb. p. 125; Form. in Ver. Brünn 1897 p. 35. — Huc. probabiliter quoque: *C. montana* S. et S. pr. II. p. 198; Ch. et B. exp. p. 254, Fl. pelop. p. 60. — *C. uniflora* S. et S pr. II. p. 197; Ch. et B. fl. pelop. p. 59. — *C. axillaris v. angustifolia* Form. in D. bot. Monat. 1890 p. 19. — Exsicc.: Heldr. herb. norm. n. 318 et 1260; Orph. fl. gr. n. 121.

Adpresse-lanuginosa; rhizomate multicipite, nodoso; caulibus adscendentibus vel erectis, mono-vel oligocephalis; foliis oblongis lanceolatisve, integris vel repando-dentatis, anguste et breviter decurrentibus; capitulis ovatis; involucri phyllis pallidis, glabris vel puberulis, appendice scariosa, nigrescenti, in dentes lanceolatos, argenteos, phylli latitudine subaequantes partita, marginatis; flosculis azureis, valde radiantibus; acheniis puberulis, pappo triplo longioribus.

In rupestribus regionis subalpinae et alpinae. Epirus: mt. Micikeli, Olycika (Bald.); Thessalia: ad monasterium Korona, pr. Sermenikon (Haussk.), mt. Mitrica in mt. Chassia, mt. Ossa (Form.); mt. Oeta Phthiotidis, mt. Tymphrestus Eurytaniae (Heldr.); mt. Parnassus (Guicc.); Helicon (Orph.), Pateras et Parnes Atticae (Heldr.); Euboea: mt. Kandyli (Orph.); Peloponnesus: mt. Panachaicon, Chelmos (Hal.), Kyllene (Heldr.), Malevo (Orph.), Taygetos, pr. Tripolis, Zatuna (Nied.), Navarin (Chaub). — Apr. Juli. ♃.

18. **C. epirota** Hal. in bull. herb. Bois. VI. p. 581. — *C. cana v. pindicola* Bald. riv. coll. bot. Alb. 1895 p. 50, 1896 p. 70, non Griseb. — Exsicc.: Bald. it. alb. epir. III. n. 73, IV. n. 82.

Differt a praecedente rhizomate repente tenui, foliis lyrato-pinnatifidis, lobis utrinque 2—3 majusculis oblongis, terminali ovato vel orbiculato, multo majore et flosculis purpureis.

In declivibus mt. Micekili et in jugo Kakardista mt. Tsumerka in Epiro, nec non extra ditionem nostram in mt. Kudesi distr. Valona Albaniae (Bald.). — Maio, Jun. ♃.

β. Flosculi ochroleuci.

19. **C. pindicola** Griseb. it rum. II. p. 164, Bois. fl. or. III. p. 637; Hal. in bull. herb. Bois. VI. p. 581. — *C. cana v. pindicola* Griseb. spic. II. p. 236. — Exsicc.: Orph. pl. gr. a. 1857; Heldr. pl. gr. a. 1884.

Adpresse araneoso-cana; rhizomate brevi, truncato; caule erecto, pumilo, monocephalo; foliis lyrato-pinnatisectis, inferioribus in petiolum attenuatis, superioribus breviter decurrentibus, lobis utrinque 2—3, majusculis, oblongis, terminali ovato, multo majore; capitulis ovatis, involucri phyllis pallidis, glabris vel puberulis, appendice scariosa, nigrescenti, in dentes lanceolatos, apice pallidos, phylli latitudine breviores partita, marginatis; flosculis ochroleucis, valde radiantibus; acheniis puberulis, pappo triplo longioribus. — Rhizomate truncato et flosculis ochroleucis a praecedente discedit.

In regione media mt. Olympus supra Hagios Dionysios (Orph.) et in regione superiori mt. Ossa (Heldr.). — Jul. Aug. ♃.

20. **C. Baldaccii** Deg. ap. Bald. viagg. Creta p. 69; Hal. in bull. herb. Bois. VI. p. 582. —. *C cana v. albiflora* Raul. cret. p. 788, non *C. albiflora* C. Koch. — Exsicc.: Bald. it. cret. n. 76.

Differt a praecedente caule fere nullo, capitulis duplo minoribus subsessilibus, foliis lineari-lanceolatis, capitulum superantibus et flosculis albidis.

In summis mt. Volakia in mt. Sphacioticis Cretae (Heldr.). — Jul. ♃.

Obs. *C. orbelica* Vel. in Sitzungsber. böhm. Ges. Wiss. 1890 p. 51. — *C. variegata v. albida* Ces. in Griseb. spic. II. p. 235, = *C. albida* Heldr herb. norm. n. 1154 non *C. montana v. albida* DC. pr. VI. p 579. — A Charrel in mt. Korthiati pr. Thessalonicam lecta, in montibus Thessaliae borealis inquirenda. A praecedentibus fibris radicis napuliformi-incrassatis et foliis lineari-elongatis, inferioribus longe et tenuiter petiolatis discedit.

2. Subsectio. *Acrolophus* DC. XI. p. 581. —

§. Biennes vel perennes.

a. Capitula solitaria, pedunculata.

α. Capitula ovata vel ovato-globosa. — Caulis plerumque parce ramosus; phyllorum mucro plerumque brevis (ciliis saepe non longior), raro validior.

× Pappus achenio sesquilongior; planta albo-tomentosa.

21. **C. Niederi** Heldr. in Bois. fl. or. III. p. 641, suppl. p. 313; Hal. in bull. herb. Bois. VI. p. 583. — Exsicc.: Heldr. herb. fl. hell. a. 1878.

Adpresse albo-tomentosa; caulibus erectis vel adscendentibus, parce ramosis, ramis monocephalis; foliis radicalibus petiolatis, bipinnatipartitis, ceteris sessilibus, pinnatipartitis, laciniis oblongis vel oblongo-linearibus, basi attenuata decurrentibus, summis saepe indivisis vel trilobis; capitulis ovato-globosis, foliis summis bracteatis; involucri phyllis pallidis, in appendicem scariosam pallide fuscam, erecto-patentem, longe triangularem, pectinato-ciliatam, aristatam, abeuntibus, ciliis argenteis, diametro appendicis longioribus, arista terminali ciliis sublongiore, phyllorum intimorum appendice ovata, denticulata; flosculis laete purpureis, radiantibus; acheniis puberulis, pappo dimidio brevioribus. — Caules 20—40 cm. alti, fragiles, capitula 2 cm. diametro.

In faucibus Klissura dictis mt. Arakynthos Aetoliae (Nied.). — Apr. Maio. ♃.

Obs. Ex affinitate praecedentis esse videtur *C. incompleta* Hal. in bull. herb. Bois. VI. p. 583. — Adpresse albo-tomentosa; caulibus adscendentibus, superne parce ramosis, ramis monocephalis; foliis pinnatipartitis, laciniis anguste lineari-lanceolatis; capitulis 1 cm. diametro; involucri straminei phyllis in appendicem patule-recurvam, aristatam, longe ciliatam abeuntibus; flosculis et acheniis ignotis. — In rupibus pr. Bitos vallis Megarema in Olympo (Sint. et Bornm. it. turc. n. 1332). —

×× Pappus achenio plerumque brevior, raro sublongior.

○ Flosculi lutei; planta albo-tomentosa.

22. **C. argentea** L. sp. p. 912; Sieb. avis p. 5, rem. p. 6; Raul. cret. p. 788; Bois. fl. or. III. p. 642; Bald. viagg. Creta p. 70; Hal. in bull. herb. Bois. VI. p. 589. — *C. ragusina* S. et S. pr. II. p. 199, Fl. gr. X. p. 2, t. 903; Sieb. in Flora I. p. 272. — *C. cineraria* Sieb. in Flora I. p. 272; Raul. cret. p. 788. — Exsicc.: Heldr. pl. cret. a. 1846; Bald. it. cret. n. 25, it. cret. alt. n. 128.

Adpresse albo-pannosa; caulibus erectis vel adscendentibus, parce ramosis, ramis brevibus, monocephalis; foliis inferioribus petiolatis, lyrato-pinnatisectis, segmentis utrinque 3—4, oblongis, integris vel subdentatis, terminali majori, ovato, foliis superioribus sessilibus, paucilobis vel indivisis; capitulis parvis, 5—7 mm. diametro, minute subbracteatis; involucri phyllis pallidis, puberulis, appendice scariosa, pallide fusca, adpressa, breviter pectinato-ciliata terminatis, phyllorum intimorum appendice denticulata; flosculis luteis, non radiantibus; acheniis puberulis, pappo aequilongis.

In rupibus regionis inferioris et montanae Cretae: ad promontorium Grabusa (Sieb.), mt. Hagios Elias distr. Kissamos, mt. Kophino distr. Monofatri (Bald.), in fauce Hagia Rumeli (Heldr.), pr. Muliana (Raul.). — Jun. Jul. ♃.

⊙⊙ Flosculi purpurei; rarissime (*C. laureotica*) ochroleuci.

. Folia caulina vel earum laciniae latiusculae, ovatae vel ellipticae; appendix phyllorum pallida, mucrone debili ciliis breviore terminata.

23. **C. sublanata** DC. pr. VI p. 584 pro var. *C. paniculatae*; Bois. fl. or. III. p. 645; Hal. in bull. herb. Bois. VI. p. 585. — Exsicc.: Orph. fl. gr. n. 745; Sint. it. or. a. 1889 n. 1828 et 1533.

Adpresse araneoso-cana; caulibus fere a basi divaricato-ramosis; foliis inferioribus petiolatis, lyratis, segmentis ovatis vel oblongis, integris vel parce dentatis, superioribus sessilibus, ovatis vel ellipticis, integris vel basi unijugis; capitulis solitariis, ovatis, nudis vel subbracteatis; involucri phyllis adpressis, oblongis, pallidis, araneoso-puberulis, in appendicem pallide fuscam, 5—7 pectinatim ciliatam, mucrone ciliis non longiore terminatam abeuntibus; flosculis purpureis, breviter radiantibus; acheniis puberulis, pappo duplo longioribus. — Foliis caulinis latiusculis, canis 10—15 mm. longis, 5—7 mm. latis, capitulis araneoso-puberulis insignis.

In arenosis maritimis Thessaliae pr. Katerina (Orph.) et Litochori (Sint.). — Jul. Sept. ♃.

24. **C. pallida** Friv. in Flora 1835 p. 333; Haussk. symb. p. 126; Form. in Ver. Brünn 1896 p. 48; Hal. in bull. herb. Bois. VI. p. 585. — *C. ciliata* Friv. l. c. p. 334, f. capitulis minoribus. — *C. macedonica* Hal. Beitr. fl. Epir. p. 28, Beitr. fl. thessal. p. 15, non alior. — Exsicc.: Sint. it. thessal. n. 997.

Glabriuscula, viridis, asperula; caulibus saepe a basi divaricato-ramosis; foliis inferioribus petiolatis, lyratis, segmentis ovatis vel oblongis, integris vel parce dentatis, superioribus sessilibus, ovatis vel ellipticis, integris vel basi unijugis; capitulis solitariis, ovatis; involucri phyllis adpressis, ovatis, pallidis, glabriusculis, in appendicem pallide fuscam, 5—7 pectinatim ciliatam, mucrone ciliis non longiore terminatam abeuntibus; flosculis pallide purpureis, breviter radiantibus; acheniis glabris, pappo triplo longioribus. — Praecedenti valde affinis et ab ea praesertim indumento subnullo discedit.

In vinetis, fruticetis, saxosis. Epirus: in regione inferiori mt. Peristeri (Hal.); Thessalia: pr. Velitsena, Koturi (Form.), Kastania (Hal.), Klinovo, Malakasi, Tsungeri, Meteora, Kalabaka (Haussk.). — Jul. Aug. ♃.

24 × 35. **C. pallida × tymphaea** Haussk. symb. p. 127. — Habitu *C. pallidae*, sed capitula minora, phyllorum appendices angustiores et firmiores; a *C. tymphaea* capitulis majoribus, ovatis, ciliis magis decurrentibus differre dicitur. — Thessalia: pr. Klinovo (Haussk.). N. v.

.. Folia caulina vel earum laciniae angustae, oblongae vel lineares.

: Capitula ovata, circa 8 mm diametro; appendix phyllorum pallida vel fuscidula, triangulari-lanceolata, mucronata, ad basin hyalino-marginata, ciliis mucrone brevioribus, saltem apice hyalinis.

, Flosculi purpurei.

.— Folia viridia vel leviter canescentia, plus minus asperula.

— Folia radicalia lyrata, lobo terminali oblongo, 5—8 mm. lato.

25. **C. kalambakensis** Freyn. et Sint. in bull. herb. Bois. V. p. 784; Hal. in bull. herb. Bois. VI. p. 586. — Exsicc.: Sint. it. thessal. n. 409.

Glabriuscula, viridis, subasperula; caulibus erectis, foliosis, superne ramosis; foliis radicalibus petiolatis, lyratis, ceteris sessilibus, pinnatisectis, laciniis lineari-lanceolatis, summis integris vel basi unijugis; capitulis solitariis, ovatis; involucri phyllis pallidis, glabris, in appendicem stramineam, pectinatim-ciliatam, mucrone subsetacea, ciliis validiore et 2—3 plo longiore terminatam abeuntibus; flosculis roseis, vix radiantibus; acheniis glabris, pappo aequilongis vel subbrevioribus. — Caulis 25—50 cm. altus, capitula 1 cm. diametro. — Colore laete viridi, caulibus foliosis, flosculis vix radiantibus et pappo longo insignis.

In rupibus ad Guwelzi pr. Kalabaka Thessaliae (Sint.). — Maio, Jun. ♃.

26. **C. transiens** Hall. in bull. herb. Bois. VI. p. 587. — Exsicc.: Sint. it. or. p. 1889 n. 1829.

Parce araneosa, canescens, asperula; caulibus erectis, remote foliatis, a medio ramosis; foliis radicalibus petiolatis, lyratis vel subindivisis, ceteris sessilibus, pinnatisectis, laciniis oblongo-lanceolatis, summis integris vel basi unijugis; capitulis solitariis, ovatis; involucri phyllis pallidis, glabris, in appendicem pallide fuscam, pectinato-ciliatam, mucrone ciliis validiore et 2—3 plo longiore terminatam abeuntibus; flosculis purpureis, radiantibus; acheniis glabris, pappo 4 plo longioribus. — Caulis 15—30 cm. altus, capitula 6—8 mm. diametro. — A praecedente habitu, indumento, caule non folioso, flosculis radiantibus et pappo brevi discedens et notis nonnullis *C. macedonicae* accedens, a qua autem jam phyllorum mucrone elongato statim diagnoscitur.

In declivibus mt. Olympus pr. Hagios Dionysios Thessaliae (Sint.). — Aug. Sept. ♃.

= Folia radicalia pinnatisecta, laciniis anguste linearibus.

27. **C. pentelica** Haussk. symb. p. 128; Hal. in bull. herb. Bois. VI. p. 588. — Exsicc.: Heldr. herb. norm. n. 1351.

Parce araneosa, canescens, asperula; caulibus erectis, superne remote foliatis, a medio vel a basi ramosis; foliis pinnatisectis, inferioribus petiolatis, superioribus sessilibus, laciniis anguste linearibus, summis indivisis; capitulis solitariis, ovatis; involucri phyllis pallidis, glabriusculis, in appendicem pallide fuscam, pectinato-ciliatam, mucrone ciliis validiore et subduplo longiore terminatam abeuntibus; flosculis roseis, vix radiantibus; acheniis glabris, pappo duplo longioribus. — Ex affinitate *C. kalambakensis* et *C. atticae*, a priori foliis radicalibus non lyratis,

superioribus remotis, phyllorum mucrone validiori et pappo breviori; ab altero, quacum formis intermediis conjuncta esse videtur, habitu robustiore, indumento non albo-tomentoso, involucri phyllis brevius mucronatis discedit.

In saxosis mt. Pentelicon (Haussk.), Parnes et Hymettus (Heldr.) Atticae. — Maio, Jun. ♃.

28. **C. asperula** Hal. in bull. herb. Bois. VI. p. 588. — Exsicc.: Heldr. pl. fl. hellen. a. 1886 sub *C. Parlatoris v. Boissieri.*

Differt a praecedente caulibus crassioribus, rigidis, foliis valde asperis, radicalibus exceptis, oblongis, integris, rarius nonnullis basi 1—2 jugis, et pappo acheniis aequilongo; ab affini *C. attica* insuper tomenti defectu. Ceterum ulterius observandum est in loco natali anne omnes tres formas varias, intermediis forsan conjunctas unius speciei sistent.

In regione Laurii pr. Kamariza Atticae (Heldr.). — Jun. ♃.

: = Folia albo-tomentosa, vix asperula.

29. **C. attica** Nym. syll. fl. europ. p. 33; Hal. in bull. herb. Bois. VI. 589 — *C. graeca* Bois. et Spr. diagn. XI. p. 128, Fl. or. III. p. 664; Haussk. symb. p. 128; non Griseb. — *C. Parlatoris v. Boissieri* Heldr. et Sart. in Bois. diagn. ser. 2, III. p. 74. — *C. Boissieri* Walp. rep. VI. p. 298, non DC. — ? *C. paniculata* S. et S. pr. II. p. 199. — Exsicc.: Orph. fl. gr. n. 122.

Albo-tomentosa, sublaevis; caulibus erectis vel adscendentibus, saepe caespitosis, a basi vel paulo supra ramosis; foliis pinnatisectis, radicalibus petiolatis, ceteris sessilibus, laciniis anguste linearibus, summis indivisis; capitulis solitariis, ovatis; involucri phyllis pallidis, puberulis, in appendicem pallide fuscam, pectinato-ciliatam, mucrone ciliis validiore et duplo longiore terminatam abeuntibus; flosculis roseis, vix radiantibus, glabriusculis, pappo duplo longioribus.

β. **pateraea** Hal. in bull. herb. Bois. VI. p. 590. — Phyllorum mucro ciliis 3—4 plo longior, phyllo ipso sublongior. — Exsicc.: Pichl. pl. gr. a. 1876; Heldr. pl. fl. hellen. a. 1876.

In saxosis submontanis montanisque. Attica: pr. Eleusis (Sprun.); mt. Kerata (Haussk.), Hymettus (Orph.), Pentelicon, Parnes (Heldr.); Euboea (Auch.); — β. mt. Pateras Atticae (Heldr.). — Maio, Jul.

,, Flosculi ochroleuci.

30. **C. laureotica** Heldr. ap. Hal. in bull. herb. Bois. VI. p. 590. — Huc forsan pertinet: *C. graeca v. ochroleuca* Heldr. et Sart. in Bois. diagn. ser. 2, III. p. 75, e mt. Anhydros (Hymetti pars meridionalis). — Exsicc.: Heldr. herb. norm. n. 948.

Parce araneoso-canescens, aspera; caulibus erectis, a medio divaricatim ramosis; foliis inferioribus petiolatis, superioribus sessilibus, pinnatisectis, laciniis oblongis, breviusculis, summis subintegris; capitulis solitariis, ovatis; involucri phyllis pallidis, glabriusculis, in appendicem pallide fuscam pectinato-ciliatam, mucrone ciliis validiore et 2—3 plo

longiore terminatam abeuntibus; flosculis ochroleucis, vix radiantibus; acheniis glabriusculis, pappo aequilongis. — Caulis 20—35 cm. altus, foliorum laciniae eis praecedentium latiores brevioresque; hac nota et flosculis ochroleucis facile diagnoscenda.

In pinetis apricis montium Laurii circa Kamariza Atticae (Heldr.). — Maio, Jun. ♃.

;; Capitula ovata vel ovato- globosa, 5—12 mm. diametro; appendix phyllorum pallida, fusca vel nigricans, triangulari-rotundata et breviter mucronata vel semilunata, ad basin hyalino-marginata, ciliis mucrone aequilongis vel longioribus, raro brevioribus (et tunc capitula ovato-globosa 12 mm. diametro).

, Caulis et folia albo-lanata, non aspera; appendix phyllorum nigra, mucrone recurvato.

31. **C. ossaea** Hal. in bull. herb. Bois. VI. p. 591. — *C. dissecta v. alpina* Heldr. pl. exs. a. 1884, non Heldr. in Bois. fl. or. III. p. 645.

Caulibus pumilis, adscendentibus, parce ramosis; foliis inferioribus petiolatis, bipinnatisectis, superioribus sessilibus, pinnatisectis, laciniis anguste linearibus vel lanceolatis; capitulis solitariis, ovatis; involucri phyllis pallidis, puberulis, in appendicem breviter triangularem, nigram, pectinato-ciliatam, mucrone ciliis hyalinis subvalidiore, aequilongo, recurvato terminatam abeuntibus; flosculis purpureis; acheniis puberulis, pappo triplo longioribus. — Caulis 8—15 cm. altus, ramis 2—6 brevibus, erecto-patulis. — Species pulchra, *C. dissectae* affinis, a qna differt indumento dense lanato, foliorum laciniis angustioribus, phyllorum mucrone recurvato et ciliis setaceo-elongatis.

In regione superiori mt. Ossa Thessaliae (Heldr.) — Jul. ♃.

,, Caulis et folia puberula vel leviter araneoso-lanata, plus minus aspera; appendix pallida vel nigra, mucrone erecto.

. — Capitula plerumque ovato-globosa, 10—12 mm. diametro; appendix phyllorom plerumque nigra, raro fusca; folia asperula.

32. **C. lacerata** Haussk. symb. p. 126 pro var. *C. affinis*; Hal. in bull. herb. Bois. VI. p. 591. — *C. affinis* Haussk. l. c.; Form. in Ver. Brünn 1896 p. 48, non Friv. — Exsicc.: Haussk. it gr. a. 1885; Form. pl. thessal. a. 1895.

Asperula, pubescens; caulibus adscendentibus vel erectis, a basi vel a medio ramosis; foliis pinnatisectis, inferioribus petiolatis, superioribus sessilibus, laciniis anguste linearibus, summis unijugis vel integris, capitulis solitariis, ovatis; involucri phyllis pallidis, glabris, in appendicem nigram, breviter setaceo-mucronatam, membrana plus minus lata, hyalina, lacerata, denticulata vel pectinato-ciliata marginatam abeuntibus; flosculis

purpureis, parum radiantibus; acheniis glabris, pappo 4 plo longioribus — Caulis 10—30 cm. altus, folia 1—3 cm. longa, capitulis vix 1 cm. diametro. — Phyllorum appendices in speciminibus nonnullis regulariter pectinato-ciliatae, ad basin tantum hyalino-marginatae, ciliis hyalinis, mucrone longioribus, in aliis autem circumcirca late hyalino-marginatae et irregulariter laceratae, fere ut in speciebus sectionis *Phalolepis*, sunt. Has formas ultimas distinxit Haussknecht sensu stricto sub nomine *C. affinis v. lacerata*; cum autem hae notae valde variant, nomen pro specie adoptavi. — A *C. affini* Friv. praeter characteres phyllorum, indumento tenuiore aspero, foliis minoribus, minus divisis, laciniis brevioribus et angustioribus et capitulis minoribus, specifice differre videtur.

In schistosis pr. Klinovo, Malakasi (Haussk.), mt. Dokimi (Form.) in Pindo tymphaeo. — Jul. Aug. ♃.

33. **C. affinis** Friv. in Flora XIX. p. 435; Hal. in bull. herb. Bois. VI. p. 592. — *C. Parlatoris* Heldr in ann. acad. aspir. nat. I. p. 286, in Flora XXVII. p. 68 (ex specim. Tod. fl. sic. n. 915). — *C. dissecta* Bois. fl. or III. p. 644; Heldr. chlor. Parn. p. 22; Hal. Beitr. fl. Epir. p. 29; non Ten. fl. nap. I. p. 51, quae ex specim. a Groves in Aprutio lectis capitulis globosis, duplo majoribus, appendicibus intense nigris longe argenteo-ciliatis differt. — Huc forsan: *C. paniculata* Ch. et B. exp. p. 254, Fl. pelop. p. 59. — Exsicc.: Heldr. herb. norm. n. 27 et 1556; Dörfl. fl. gr. n. 195.

Asperula, plus minus araneoso-canescens; caulibus erectis, a basi vel a medio ramosis; foliis inferioribus petiolatis, pinnatisectis, laciniis oblongo-lanceolatis, superioribus sessilibus, minus divisis vel integris; capitulis solitariis, ovatis; involucri phyllis pallidis, glabris, in appendicem nigram, breviter mucronatam, pectinato-ciliatam, basi membrana hyalina angusta marginatam, abeuntibus; flosculis purpureis, parum radiantibus, acheniis glabris, pappo 4 plo longioribus. — Caulis 20—40 cm. altus, plerumque ramosus, raro simplex et humilis vix 10 cm. altus (*C. dissecta v. alpina* Heldr. in Bois. fl. or. III. p. 645). — Species polymorpha, variat praesertim indumento, foliorum divisione, capitulorum magnitudine et colore appendicorum. — Formae sequentes intermediis transientes seorsim notandae.

β. **denudata** Hal. in bull. herb. Bois. VI. p. 593. — Caulis superne tantum et brevius ramosus; folia viridia, aspera; capitula angustiora. — Exsicc.: Sint. it thessal. n. 1126.

γ. **peloponnesiaca** Hal. l. c. — Virescens; caulis a basi divaricatim ramosus; folia caulina indivisa vel unijuga; appendix fusca, mucrone validiore. — Exsicc.: Orph. pl. exs. a. 1850 et 1852.

δ. **pallidior** Bois. fl. or. III. p. 645 pro forma *C. dissectae*. — *C. pallidior* Hal. l. c. p. 594. — Virescens, caulis a basi divaricatim ramosus; appendix phyllorum pallescens. — Variat insuper capitulis minoribus (*C. affinis v. microcephala* Hal. l. c., = *C. Parlatoris* Haussk. symb [illegible] ex exsicc., = *C. paucijuga v. melanolepis*

Hal. l. c. p. 596); capitulis itidem minoribus, et foliis inferioribus paucijugis, caeteris fere omnibus indivisis (*C. paucijuga* Hal. l. c.). — Exsicc.: Heldr. herb. norm. n. 710, it gr. septentr. a. 1879.

In siccis regionis montanae et abietinae. Epirus: mt. Tsumerka (Hal.); mt. Parnassus (Heldr.); in faucibus Langada mt. Taygetos (Zahn); — *β*. pr. Sermenikon in Pindo (Sint.); — *γ*. mt. Malevo, Chelmos (Orph.), Kyllene (Heldr.); — *δ*. pr. Karpenisi Eurytaniae (Samar.), mt. Korax Aetoliae (Heldr.), *f. microcephala* in collibus inter Kalabaka et Malakasi in Pindo (Haussk.), pr. Segditza in mt. Kiona, cum *f. paucijuga* (Hal.). — Jul. Aug. ♃.

34. **C. macedonica** Griseb. spic. II. p. 240 pro var. *C. paniculatae*; Form. in Ver. Brünn 1895 p. 27; Hal. in bull. herb. Bois. VI. p. 595 (non *C. macedonica* Bois. diagn. VI. p. 130). — *C. graeca v. macedonica* Bois. fl. or. III. p. 644. — *C. Grisebachii* Nym. consp. p. 427; Form. in Ver. Brünn 1896 p. 48, 1897 p. 35; Haussk. symb. p. 126. — Forsan huc etiam: *C. affinis* × *Grisebachii* Haussk. l. c. p. 127. — *C. graeca v. Grisebachii* Bald. riv. coll. bot. Alb. 1895 p. 50. — ? *C. Biebersteinii* Form. in D. bot. Monat. 1896 p. 19. — *C. pallidior v. pallidissima* Hal. in bull. herb. Bois. VI. p. 594. — *C. confusa* Hal. l. c. p. 596. — Exsicc.: Sint. et Bornm. it. turc. a. 1891 n. 1331; Bald. it. alb. epir. III. n. 165; Orph. fl. gr. n. 746 e mt. Korfiati; Heldr. herb. norm. n. 1155, Kavala Macedoniae.

Adpresse pubescens, pallide virens, asperula; caulibus erectis, a medio vel superne ramosis; foliis inferioribus petiolatis, pinnatisectis, laciniis oblongo-lanceolatis linearibusve, superioribus sessilibus, minus divisis vel integris; capitulis solitariis, ovatis, apice plus minus constrictis; involucri phyllis pallidis, glabris vel puberulis, in appendicem pallide fuscam, breviter mucronatam, pectinato-ciliatam, basi membrana hyalina angusta marginatam abeuntibus; flosculis purpureis, parum radiantibus; acheniis puberulis, pappo 3—4 plo longioribus. — Species polymorpha, ulterius observanda; a praecedentibus capitulis praesertim minoribus, appendicibus pallidis, longius ciliatis discedit, sed transitus ad var. *δ*. speciei praecedentis adesse videntur.

In dumosis montanis. Epirus: in valle Dipotamo, mt. Micikeli, pr. Govrica, mt. Peristeri (Form.), pr. Janina (Bald.); Thessalia: pr. Koturi, Malakasi, mt. Zygos, Dokimi, Ghavellu (Form.), pr. Klinovo, mt. Baba (Haussk.), pr. Phlamburo in mt. Oxya, pr. Vlachava, Asproklisia, Hagios Paraskevi, mt. Hagios Elias et Mitrica in mt. Chassia (Form.), pr. Paligriatzano (Heldr.), pr. Metochi (Heldr.), Hagios Dionysios (Sint.) et Rapsani in Olympo, pr. Longici, Stylida (Form.); Doris: pr. Segditza ad radices mt. Kiona (Hal.). — ? ⊙ et ♃.

β. Capitula ovato-oblonga, parva. — Caulis valde ramosus; phyllorum mucro validior, ciliis plerumque longior et patens vel recurvatus.

× Flosculi purpurei.

○ Appendix phyllorum in spinulam erectam attenuata; capitula 10 mm. longa.

35. **C. tymphaea** Haussk. symb. p. 127; Form. in Ver. Brünn 1896 p. 48; Hal. in bull. herb. Bois. VI. p. 597. — *C. brevispina* Form. l. c. p. 49 p. p. — Exsicc.: Haussk. it. gr. a. 1885.

Asperula, glabrescens; caulibus erectis, a basi vel a medio ramosissimis; foliis in lacinias oblongo-lanceolatas vel lineares pinnatipartitis, inferioribus petiolatis, ceteris sessilibus minus divisis, summis integris; capitulis solitariis, ovato-oblongis, 5—8 mm. longis; involucri phyllis pallidis, glabris, in appendicem pallide fuscam, pectinato-ciliatam, mucrone subspinescente, ciliis aequilonga terminatam, abeuntibus; flosculis purpureis, parum radiantibus; acheniis glabris, pappo triplo longioribus.

In herbidis regionis inferioris et montanae. Thessalia: pr. Malakasi (Heldr.), Prevenda, Klinovo (Haussk.) in Pindo, pr. Kucuro et Vanaluka in mt. Chassia, pr. Tafilvris et Lokatamburia in mt. Olympo (Form.). — Jul. Aug. ⊙.

36. **C. brevispina** Haussk. symb. p. 129; Form. in Ver. Brünn 1896 p. 49, 1897 p. 35; Hal. in bull. herb. Bois. VI. p. 598. — *C. macedonica* Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 4, non (Griseb). — *C. Orphanidea* Hal. in bull. herb. Bois. VI. p. 599 p. p. — Exsicc.: Haussk. it. gr. a. 1885; Heldr. it. Thessal. n. 21; Sint. et Bornm. it. turc. n. 1330. —

Asperula, araneoso-subcanescens; caulibus erectis, a medio vel superne stricte ramosis, foliis in lacinias lineari-ellipticas pinnatisectis, inferioribus petiolatis, ceteris sessilibus, minus divisis vel unijugis, summis integris; capitulis solitariis, 8—10 mm. longis; involucri phyllis pallidis, glabris, in appendicem pallide fuscam, pectinato-ciliatam, spinula pungente, ciliis validiore terminatam, abeuntibus; flosculis purpureis, parum radiantibus; acheniis glabris, pappo triplo longioribus. — Praecedenti similis, ab ea caulibus elatioribus, 30—60 cm. altis, indumento canescente, ramis superne tantum ramulosis, minus divaricatis, foliorum laciniis brevioribus latioribusque et capitulis majoribus discedit. An a sequente caulibus minus divaricatim ramosis, phyllorum spinulis vix patulis et pappi setis brevioribus specifice diversa, ulterius observandum.

β. **fusconigra** Haussk. l. c.; Hal. l. c. — Appendicibus intensius fuscis vel fusconigris. — Exsicc.: Heldr. it. thessal. n. 22.

In nemorosis Thessaliae: pr Litochoron (Sint.), mt. Pelion supra Volo cum var. β. (Haussk.), pr. Koryza, Kastri, pr. Neraida et Lamia Phthiotidis (Form.). — Jul. Aug. ⊙.

36 × 39. *C. brevispina* × *pelia* Haussk. symb. p. 130. — Habitus *C. peliae*, sed flores flavorubentes. — Thessalia: mt. Pelion (Haussk.). N. v.

○○ Appendix phyllorum in spinulam patulam attenuata; capitula 10 mm. longa.

37. **C. Orphanidea** Heldr. et Sart. in Bois. diagn. ser. 2, III. p. 73, fl. or. III. p. 650; Hal. in bull. herb. Bois. VI. p. 599 p. p. —

? *C. squarrosa* S. et S. pr. II. p. 203. — Exsicc.: Orph. fl. gr. n. 124; Heldr. herb. norm. n. 227 et 949.

Asperula, araneoso-canescens; caulibus erectis, a basi divaricatim ramosissimis; foliis in lacinias oblongas, parce lobatas vel integras pinnatipartitis, inferioribus petiolatis, caeteris sessilibus, minus divisis, summis integris; capitulis solitariis, ovato-oblongis, 8—10 mm. longis; involucri phyllis pallidis, glabris vel arachnoideo-puberulis, in appendicem fuscam, pectinato-ciliatam, spinula pungente, ciliis validiore terminatam, abeuntibus; flosculis purpureis, parum radiantibus; acheniis puberulis, pappo sublongioribus.

β. **thessala** Haussk. in Mittheil. thür. bot. Ver. V. p. 86, symb. p. 130; Form. in D. bot. Monat. 1890 p. 49; pro sp.; Hal. in bull. herb. Bois. VI. p. 599. — Indumento tenuiore, caulibus altioribus, ramis minus divaricatis, appendicibus plerumque pallidioribus. — Intermediis ad typum transit. — Exsicc.: Haussk. it. gr. a. 1885.

In arvis, derelictis, ad vias regionis inferioris. Attica: pr. Athenas, Eleusis, Hagia Glycaria, Heracleon, Dekeleia, Kephissia, Patissia, Kato Liosia, Chelidonu (Heldr.); — *β.* Thessalia: pr. Tyrnavos (Haussk.), Patsios, Drepani, Kapurna (Form.). — Maio, Aug. ⊙.

⊙⊙⊙ Appendix phyllorum in spinulam recurvatam attenuata; capitula 10—15 mm. longa.

38. **C. drakiensis** Freyn et Sint. in bull. herb. Bois. V. p. 783; Hal. ibid. VI. p. 600. — Exsicc.: Sint. it. thessal. n. 1316.

Asperula, virens vel araneoso-canescens; caulibus erectis, a medio breviter ramosis; foliis in lacinias oblongo-lanceolatas pinnatipartitis, inferioribus petiolatis, caeteris sessilibus, sensim diminutis, summis integris; capitulis solitariis, ovato-oblongis; involucri phyllis pallidis, glabris, in appendicem pallide fuscam, spinuloso-ciliatam, spinula pungente, ciliis validiore, recurvata terminatam, abeuntibus; flosculis purpureis, parum radiantibus; acheniis glabris, pappo triplo longioribus. — Ab affinibus caulibus plerumque racemoso-ramosis, ramis ab infimis brevissimis, capitulo brevioribus vel paulo longioribus, ad summa capitula saepe 5—6plo longiora sensim elongatis et praecipue capitulis majusculis et appendicibus recurvatis insignis.

In aridis mt. Pelion pr. Drakia (Sint.), pr. Koryza ad Volo, pr. Patsios in Olympo (Form.). — Jul. Sept. ⊙.

×× Flosculi lutei.

39. **C. pelia** DC. pr. VI. p. 586; Bois. fl. or. III. p. 649, suppl. p. 313; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 4; chlor. Parn. p. 22; Hal. in z. b. Ges. Wien 1888 p. 759, Beitr. fl. thessal. p. 15; Form. in D. bot. Monat. 1890 p. 19, in Ver. Brünn 1896 p. 48, 1897 p. 35; Haussk. symb. p. 29. — *C. xanthina* Bois. et Heldr. in Heldr. pl. exs. a. 1852, non Spreng. — Exsicc.: Orph. fl. gr. n. 307 et 736; Heldr. herb. norm. n. 323; Sint. it. thessal. n. 1068.

Asperula, parce arachnoidea; caulibus a basi divaricatim ramosissimis; foliis in lacinias lineari-oblongas, parce lobatas vel integras pinnatipartitis, inferioribus petiolatis, caeteris sessilibus, sensim diminutis, summis integris; capitulis solitariis, ovato-oblongis; involucri phyllis pallidis, flavidis, in appendicem pallidam, spinuloso-ciliatam, spinula pungente, ciliis validiore, recta vel patula terminatam abeuntibus; flosculis luteis, non radiantibus; acheniis glabris, pappo sublongioribus. — Flosculis luteis insignis.

β. **refracta** Haussk. symb. p. 129. — Appendicibus hamato-refractis. — Exsicc.: N. v.

In aridis saxosis regionis inferioris et submontanae. Thessalia: ad radices mt. Pelion pr. Volo (Aucher), Portaria, Koryza, Bubulifra, in planitie pr. Ambelakia, Velestinos, Larissa (Form.), Tyrnavos, Aivali, Pharsalus, pr. Meteora, Tsungeri, Malakasi in Pindo (Haussk.), pr. Hypata ad radices mt. Oeta (Heldr.); pr. Rachova et Livadi in Parnasso (Heldr.), pr. Segditza ad radices mt. Kiona (Hal.); pr. Lutraki in isthmo Corinthiaco (Samar.); — β. pr Meteora (Haussk.). — Jun. Aug. ⊖.

××× Flosculi albi.

40. **C. lactiflora** Hal. in bull. herb. Bois. VI. p. 601. — *C. ovina* Form. in Ver. Brünn 1897 p. 35, non Pall. — Exsicc.: Form. pl. thessal. a. 1896.

Asperula, parce arachnoidea, virens; caulibus a medio divaricatim ramosis; foliis in lacinias lineares pinnatipartitis, inferioribus petiolatis, caeteris sessilibus, minus divisis, summis integris; capitulis solitariis, ovato-oblongis; involucri phyllis pallidis, glabris, in appendicem pallidam, spinuloso-ciliatam, spinula subpungente, ciliis aequilonga, subpatula terminatam abeuntibus; flosculis albis, parum radiantibus; acheniis glabris, pappo duplo longioribus. — Caulis 20—40 cm. altus, foliorum laciniae angustae, capitula 1 cm. longa, 5—6 mm. lata. Flosculis albis a praecedentibus distincta; *C. ovinae* Pall. similis, a qua indumento tenuiori, capitulis minoribus et pappo longiore discedit.

Thessalia: pr. Konisko (Form.). — Jul. Aug. ⊖.

41. **C. diffusa** Lam. dict. I. p. 675; Bois. fl. or. III. p. 650; Hal. in ö. b. Z. 1890 p. 39; Form. in D. bot. Monat. 1890 p. 19; in Ver. Brünn 1896 p. 48. — *C. parviflora* S. et S pr. II. p. 203; Fl. gr. X. p. 9. t. 912. — Exsicc.: Orph fl. gr. n. 743 (Thessalonica); Sint. it. or. a. 1889 n 1837; Sint. et Bornm. it turc. n. 1324 et 1325.

Asperula, parce arachnoidea, virens; caulibus a basi divaricatim et intricatim ramosissimis; foliis subpetiolatis vel sessilibus, in lacinias oblongas pinnatipartitis, caulinis abbreviatis, paucilaciniatis, summis integris; capitulis solitariis, ovato-oblongis; involucri phyllis coriaceis, flavidis, pruinosis, in appendicem pallidam, spinuloso-ciliatam, spinula pungente, ciliis validiore et multo longiore, patulo-recurva terminatam, abeuntibus; flosculis albis, vix radiantibus; acheniis glabris, calvis. — Praeter alias notas, pappi defectu ab omnibus affinibus distinctissima.

In sterilibus regionis inferioris. Thessalia: pr. Katerina et Litochori (Sint.), Makrichori, Larissa (Form.) — Jul. Sept. ⊙.

b. Capitula sessilia, in glomerulos terminales aggregata.

42. **C. Zuccariniana** DC. pr. VI p. 574; Bois. fl. or. III. p. 653; Form. in D. bot. Monat. 1890 p. 19, in Ver. Brünn 1896 p. 48, 1897 p. 35; Hal. Beitr. fl. Thessal. p. 15, Beitr. fl. Achaia p. 25; Bald. riv. coll. bot. Alb. 1895 p. 51, 1896 p. 70. — Exsicc.: Orph. fl. gr. n. 126; Heldr. herb. norm. n. 625 et 950; Bald. it. alb. epir. III. n. 74, IV. n. 35; Sint. it. thessal. n. 561; Dörfl. fl. gr. n. 397. —

Papilloso-hirtella, asperula; caulibus erectis, superne breviter ramosis; foliis in lacinias oblongo-lineares pinnatipartitis, inferioribus petiolatis, caeteris sessilibus, summis indivisis, glomerulos bracteantibus; capitulis cylindricis, paucifloris; involucri phyllis flavidis, glabris, in cilia longissima, setacea, sursum directa pinnatipartitis, plumosis, patenti-subrecurvis; flosculis roseis, non radiantibus; acheniis glabris, pappo triplo longioribus. — Species elegantissima, nulla cum alia comparanda.

In montosis. Epirus: mt. Murga et Kurenta pr. Janina (Bald.); Thessalia: pr. Meteora, Trikala (Haussk.), Kalabaka, Kataphygi, Kastania, Povelci, Kastri, Verenci, Vlachava, Aspro-Klisia, Neochorion (Form.); Eurytania: pr. Karpenisi et Hagios Sostis (Samar.), Achaia: pr. Megaspilaeon, in valle Voreikos pr. Kalavryta (Orph.), pr. Sudena (Hal.). — Jul. Aug. ⊙.

§§. Suffruticosa, erinaceo-dumosa, caulibus spinosis.

43. **C. spinosa** L. sp. p. 912; S. et. S. pr. II. p. 199, Fl. gr. X. p. 1. t. 902; Urv. enum. p. 115; Sieb. avis p. 5, rem. p. 6; Ch. et B. exp. p. 255, Fl. pelop. p. 60; Raul. cret. p. 788; Weiss in z. b. G. 1869 p. 44; Hal. in ö. b. Z. 1895 p. 460; Haussk. symb. p. 131; Bald. viagg. Creta p. 70; Heldr. fl. Aegina p. 306. — Exsicc.: Heldr. herb. norm. p. 428 et 1044; Orph. fl. gr. n. 304; Rev. pl. cret. n. 79, in Baen. herb. europ. p. 4858; Bald. it. cret. n. 24, it. cret. alb. n. 129.

Adpresse albo-tomentosa vel glabrescens; caulibus pumilis, fere a basi intricatim ramosissimis, apice demum spinosis; foliis radicalibus oblongo-lanceolatis, in petiolum attenuatis, integris vel lyratis, caulinis sessilibus, in lacinias lineares pinnatipartitis, summis squamaeformibus; capitulis sub spina terminali solitariis, sessilibus vel breviter pedunculatis, oblongo-cylindricis, paucifloris; involucri glabri phyllis pallidis, in appendicem breviter mucronatam, vix spinescentem, fimbriato-ciliatam abeuntibus; flosculis pallide roseis, vix radiantibus; acheniis glabris, calvis. — Caules 10—25 cm. alti, caespites pulviniformes formantes, capitula parva, cum flosculis 1 cm. longa, facile secedentia.

α. **tomentosa** Hal. in bull. herb. Bois. VI. p. 634. — Caulis et folia dense adpresse albo-tomentosa. —

β. **glabrata** Heldr. herb. norm. n. 1045. — Caulis et folia fere glabra. Intermediis ad typum transit.

In araneosis maritimis, nec non in collibus aridis saxosis. Attica: ad Phalerum, promontorium Colias (Heldr.), Sunium (Haussk.), insula Aegina (Sprun.); Euboea: pr. Kurbatzi (Wild.), Petalium insula Pontikonisi (Holzm.); Cycladum insula Cythnos (Krinos), Syra (Weiss), Andros, Tenos, Naxos, usque ad alt. 1000 m. adscendens, Siphnos, Seriphos (Heldr.), Melos (Urv.), Antimilos (Reis.); Laconia: pr. Gythion (Psarid.); Creta: pr. Kissamos (Rev.), Gonia, Gurnes (Bald.), Platania, Canea (Heldr.). — Jun. Jul. ♃.

3. Subsectio. *Acrocentron* DC. l. c. p. 586.

a. Flosculi purpurei.

α. Caulescentes, saepius elatae.

× Appendix phyllorum mutica vel breviter mucronata.

44. **C. Oliveriana** DC. pr. VI. p. 590; Bois. fl. or. III. p. 657; Hal. in bull. herb. Bois. VI. p. 634. — *C. Oliveriana β. amorgina* Bois. et Orph. in Bois. fl. or. suppl. p. 314, f. phyllorum appendice minima, ciliis nigris brevioribus obsita, subinermi. — *C. atropurpurea* Oliv. voy. I. p. 313, non Willd. — *Hyalea Oliveri* Jaub. et Sp. ill. t. 292. — Exsicc.: Orph. fl. gr. n. 1149; Dörfl. fl. aeg. n. 51.

Caulibus erectis, simplicibus vel 2—3 ramosis, inferne adpresse albo-tomentosis, crebre foliatis, superne glabrescentibus, nudiusculis; foliis ovatis vel ellipticis, indivisis vel lyrato-pinnatisectis, demum glabrescentibus, inferioribus petiolatis, caeteris sessilibus, segmentis utrinque 3—7 parvis, terminali magno, ovato, obsolete repando-dentato, foliis summis 1—3 parvis, denticulatis; capitulis solitariis, ovato-globosis; involucri phyllis ovatis, coriaceis, pallidis, parce araneosis, appendice minuta, fusca, breviter pectinatim ciliata, submutica vel brevissime mucronata terminatis; flosculis atropurpureis, non radiantibus; acheniis pubescentibus, pappo aequilongis. — Species pulchra, caulibus 30—50 cm. altis, capitulis 2—3 cm. diametro.

In rupibus mt. Dzia in insula Naxos (Oliv.), nec non in rupestribus maritimis insulae Mykonos et Amorgos (Orph.). — Jun. Jul. ♃.

45. **C. ebenoides** Heldr. in journ. of bot. 1878 p. 188; Bois. fl. or. suppl. p. 314; Hal. in bull. herb. Bois. VI. p. 635. — Exsicc.: Mill in Heldr. pl. fl. hellen. a. 1862.

Araneoso-cana, demum glabrescens; caulibus gracilibus, erectis, simplicibus vel subsimplicibus, remote foliatis; foliis petiolatis, pinnatisectis vel lyratis, segmentis lateralibus lanceolatis vel deltoideis, integris vel obscure paucilobis, terminali ovato; capitulis solitariis, ovatis; involucri phyllis ovatis, coriaceis, pallidis, subaraneosis, in appendicem minutam, fuscam, breviter pectinatim ciliatam, mucrone ciliis triplo longiore terminatam abeuntibus; flosculis roseis, breviter radiantibus; acheniis pubescentibus, pappo triplo brevioribus. — Caules 15—30 cm. alti,

capitula 15 mm. diametro, pappus conspicue longus. — Habitu *C. scabiosae* similis.

In silvis Euboeae septentrionalis (J. Stuart-Mill), loco speciali ignoto. — Maio. ♃.

×× Appendix phyllorum spina valida terminata (cf. *Guicciardii* var. *β.*).

○ Appendix phyllorum spina magna, ovata, phylla tegens.

46. **C. Sibthorpii** Hal. in bull. herb. Bois. VI. p. 635. — *C. eryngioides* S. et S. pr. II. p. 204, Fl. gr. X. p. 9, t. 913, an etiam Urv. enum. p. 116 ex insula Melos; non Lam. — *C. achaia γ. ferox* Bois. et Heldr. diagn. ser. 2, III. p. 79, non *C. ferox* Desf. — *C. achaia* Bois. fl. or. III. p. 661 p. p. — Exsicc.: Heldr. herb. norm. n. 426.

Caulibus erectis, adpresse araneosis, asperulis, stricte ramosis; foliis coriaceis, valde asperis, in lacinias oblongas vel lanceolatas, integras, dentatas vel pinnatipartitas pinnatisectis; capitulis ovato-globosis; involucri phyllis ovatis, coriaceis, glabris, pallidis, in appendicem ovatam, duram, flavidam, pectinato-ciliatam, intermediis spina latissima canaliculata, erecto-patula, inferne interdum parce spinulosa, flosculos aequante vel superante terminatam abeuntibus, ciliis albidis latitudine appendicis multo brevioribus; flosculis roseis, non radiantibus; acheniis adpresse hirsutis, pappo duplo brevioribus. — Species pulchra capitulis 3 cm. diametro, spinis phyllorum intermediorum 15—20 mm. longis; a sequente ramis brevioribus, foliis magis asperis, laciniis latioribus, phyllis pallidis breviter ciliatis diversa.

In campis et vinetis pr. Mesoitis Atticae (Heldr.). — Jun. Jul. ♃.

47. **C. achaia** Bois. et Heldr. diagn. ser. 2, III. p. 79, Fl. or. III. p. 661 p. p.; Hal. in bull. herb. Bois. VI. p. 636. — Exsicc.: Orph. fl. gr. n. 305.

Caulibus erectis, adpresse-araneosis, asperulis, stricte ramosis; foliis coriaceis, asperulis, in lacinias lanceolatas, integras vel dentatas pinnatisectis; capitulis ovato-globosis; involucri phyllis ovatis, coriaceis, glabris, pallidis, in appendicem ovatam, duram, pallidam, pectinato-ciliatam, intermediis spina lata, canaliculata, erecto-patula, inferne interdum parce spinulosa, flosculis breviore terminatam abeuntibus, ciliis flavidis, latitudine appendicis aequilongis; flosculis roseis, non radiantibus; acheniis adpresse hirsutis, pappo duplo brevioribus.

β. **submutica** Bois. fl. or. III. p. 661. — Phyllorum spina brevis, ciliis parum validior. — Exsicc.: N. v.

γ. **corinthiaca** Bois. et Heldr. diagn. ser. 2, VI. p. 113 pro sp.; Bois. fl. or. III. p. 661. — Minus elata, foliorum segmenta minuta, oblongo-linearia approximata, flosculi basi aurantiaci. — Exsicc.: N. v.

In regione subalpina mt. Chelmos pr. Klukines, Peristera (Orph.), in vinetis pr. Tripolis (Sart.); — *β.* pr. Sudena ad radices mt. Chelmos

(Heldr.); — γ. inter Kalamaki et Lutraki in isthmo Corinthiaco (Samar.). — Jun. Jul. ♃.

○○ Appendix phyllorum semilunata, saepe angustissima, phylla non tegens.

. Appendix phyllorum nigra.

48. C. **redempta** Heldr. in bull. soc. bot. Fr. XXXVII. p. 242. — *C. scopulorum et eryngioides* Raul. cret. p. 788 et 789. — Exsicc.: Reverch. pl. cret. n. 82.

Caulibus erectis, simplicibus vel parce ramosis, ad collum dense albo-lanatis, superne arachnoideis; foliis detersiliter arachnoideis, demum glabrescentibus, inferioribus petiolatis, in lacinias oblongo-lanceolatas, dentatas vel dentato-lobatas pinnatisectis, superioribus sessilibus diminutis, minus divisis, summis integris; capitulis ovato-globosis; involucri phyllis orbiculatis, glabris, pallide virentibus, in appendicem semilunatam, nigram, breviter pectinato-ciliatam, spina canaliculata, erecto-patula, fusca, inferne interdum parce spinulosa, flosculis breviore terminatam abeuntibus; flosculis atropurpureis, non radiantibus; acheniis adpresse hirsutis, pappo duplo brevioribus. — Caulis 20—40 cm. altus, folia inferiora 20—25 cm. longa, 5 – 9 cm. lata, capitula 3—4 cm. diametro. — A sequente collo valde bombycino, foliis pinnatisectis, laciniis amplioribus, capitulis majoribus, appendicibus nigris et flosculis atropurpureis specifice discedit.

In rupestribus Cretae: pr. Kissamos (Heldr.), Malaxa et in oropedio Homalos in mt. Sphacioticis (Reverch.). — Jun. Jul. ♃.

.. Appendix phyllorum pallida vel fusca.

; Folia inferiora bipinnatisecta.

49. C. **subarachnoidea** Bois. et Heldr. diagn. ser. 2, III. p. 80 pro var. *C. eryngioidis*; Hal. in bull. herb. Bois. VI. p. 638. — *C. laconica* Bois. fl. or. III. p. 660. — ? *C. eryngioides* Ch. et B. exp. p. 256, Fl. pelop. p. 60. — Exsicc.: Orph. pl. exs. a. 1857; Heldr. herb. norm. n. 1557.

Caulibus erectis, simplicibus vel parce ramosis, arachnoideis, inferne foliosis, superne nudiusculis; foliis parce arachnoideis, inferioribus petiolatis, sublyrato-subbipinnatisectis, laciniis oblongis dentatis, superioribus sessilibus, pinnatisectis; capitulis ovato-globosis, 2—3 cm. diametro; involucri phyllis ovatis, puberulis, pallidis, in appendicem semilunatam, fuscam, breviter pectinato-ciliatam, spina canaliculata, patula, inferne parce spinulosa, flosculos superante, 20—25 mm. longa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis adpresse hirsutis, pappo duplo brevioribus.

In saxosis regionis inferioris mt. Malevo pr. Kastanitza (Orph.), in districtu Alagonia l. d. Dyrrhemma (Zahn) et in mt. Taygetos (Heldr.). — Maio, Jul. ♃.

50. C. **psilacantha** Bois. et Heldr. diagn. ser. 2, III. p. 82, Fl. or. III. p. 663; Heldr. chlor. Parn. p. 22; Hal. in z. b. G. Wien

1888 p. 759, in bull. herb. Bois. VI. p. 639. — Exsicc.: Orph. fl. gr. n. 309.

Caulibus erectis, stricte ramosis, arachnoideis, inferne foliatis, superne nudiusculis; foliis parce arachnoideis, inferioribus petiolatis, in lacinias parvas ovatas vel oblongas bipinnatisectis, superioribus sessilibus, pinnatisectis; capitulis ovatis, 10—15 mm. diametro; involucri phyllis ovatis, puberulis, pallide-virentibus, in appendicem semilunatam, pallidam, breviter pectinato-ciliatam, spina aciculari, patula, inferne parce spinulosa, flosculis aequilonga vel longiore, 10—30 mm. longa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis parce hirsutis, pappo subduplo brevioribus. — Praecedenti multo gracilior, capitula duplo minora, phyllorum spina acicularis.

In dumosis regionis submontanae et montanae, rarissime. Ad radices mt. Parnassus pr. Rachova (Guicc.) et in montosis supra Amphissa (Hal.). — Jul. Aug. ♃.

;; Folia inferiora pinnatisecta vel indivisa.

, Folia laeviuscula, arachnoideo-cana, segmentis petiolulatis.

51. **C. graeca** Griseb. spic. II. p. 242; Hal. in bull. herb. Bois. VI. p. 639. — *C. Ceccariniana β. brachycentra* Bois. fl. or. III. p. 663. — *C. Cordonis* Bois. et Orph. l. c. . — *C. Ceccariniana* Haussk. symb. p. 131. — *C. Guicciardii* Form. in Ver. Brünn 1896 p. 49, 1897 p. 35, non Bois. — *C. Guicciardii v. minutispina* Hal. in bull. herb. Bois. VI. p. 642. — Exsicc.: Sint. it. thessal. n. 1125.

Caulibus erectis, stricte ramosis, arachnoideis; foliis arachnoideo-canis, tandem glabrescentibus, inferioribus petiolatis, oblongis, indivisis vel in segmenta ovata vel ovato-oblonga, integra vel angulato-dentata, basi attenuata, petiolulata pinnatisectis, superioribus valde diminutis, sessilibus, dentatis integrisve; capitulis ovato-globosis, 2—3 cm. diametro; involucri phyllis suborbicularibus, glabris, pallidis, in appendicem semilunatam, albidam, breviter pectinato-ciliatam, spina aciculari vel subulata, ciliis vix validiori vel usque 15 mm. longa, erecto-patula, inferne interdum parce spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis adpresse hirsutis, pappo aequilongis. —

β. **Ceccariniana** Bois. et Heldr. diagn. ser. 2, III. p. 81, Fl. or. III. p. 662; Heldr. chlor. Parn. p. 22; pro sp.; Hal. in bull. herb. Bois. VI. p. 640. — Phyllorum spinae validiores, usque 20 mm. longae. — Exsicc.: Heldr. herb. n. 2725.

In submontosis, in regionem abietinam usque adscendens. Thessalia: pr. Aivali (Haussk.), ad Acropolin pr. Pharsalum (Heldr.), ad Gionscala pr. Sermeniko (Sint.), mt. Ghavellu, ad monasterium Korona, pr. Kastri, Asproklisia, Patsios in Olympo; — *β.* pr. Livadi supra Delphi in Parnasso (Heldr.). Jun. Aug. ♃.

,, Folia puberula, saepius asperula, segmentis sessilibus.

.— Folia inferiora lyrato-pinnatisecta vel pinnatisecta.

52. **C. Spruneriana** C. H. Schultz in Flora XXV., Beibl. p. 160 pro var. *C. saxicolae*; Hal. in bull. herb. Bois. VI. p. 641. — *C. Spruneri* Bois. et Heldr. diagn. VI. p. 132 p. p.; Bois. Fl. or. III. p. 663; Hal. in z. b. Ges. Wien 1888 p. 763; Haussk. symb. p. 131. — *C. Guicciardii* Form. in D. bot. Monat. 1898 p. 78, non Bois. — Exsicc.: Heldr. herb. norm. n. 322.

Caule erecto, crasso, arachnoideo, breviter et stricte ramoso; foliis parce arachnoideis, asperulis, inferioribus sub anthesi emarcidis, petiolatis lyrato-pinnatisectis, segmentis integris vel repandis, terminali ovato vel oblongo, lateralibus utrinque 3—6 triangularibus vel subrotundis, lobulis intermixtis, foliis superioribus diminutis pinnatisectis vel subintegris; capitulis maximis, 30—35 mm. diametro, ovato-globosis; involucri phyllis ovatis, glabris, pallide virentibus, in appendicem semilunatam, fuscidulam, breviter pectinato-ciliatam, spina validissima, 25—35 mm. longa, erecto-patula, canaliculata, basi spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis adpresse hirsutis, pappo triplo brevioribus. — Omnium validissima, characteribus sequenti affinis, sed omnibus partibus multo major.

In campis et vinetis Atticae: inter Athenas et Phalerum (Heldr.), pr. Chalandra (Heid.), Laurion (Haussk.). — Jun. Jul. ♃.

53. **C. Guicciardii** Bois. fl. or. III. p. 661, suppl. p. 314; Hal. in z. b. G. Wien 1888 p. 759, Beitr. fl. Aetol. p. 8, in bull. herb. Bois. VI. p. 641; Form. in Ver. Brünn 1895 p. 27; Bald. riv. coll. bot. Alb. 1895 p. 51, in bull. herb. Bois. IV. p. 205. — Exsicc.: Orph. fl. gr. n. 467; Heldr. it. gr. septentr. a. 1879; Hal. it. gr. II. a. 1893.

Caule erecto, crassiusculo, arachnoideo, breviter et stricte ramoso; foliis parce arachnoideis, asperis, lyrato-pinnatifidis, segmentis ovatis vel oblongis, repandis vel obtuse-dentatis, terminali paulo majore, foliis inferioribus petiolatis, superioribus sursum diminutis, sessilibus; capitulis ovato-globosis, 15—20 mm. diametro; involucri phyllis ovatis, puberulis, pallidis, in appendicem semilunatam, fuscidulam, breviter pectinato-ciliatam, spina valida, 2—3 cm. longa, canaliculata, basi spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis adpresse hirsutis, pappo triplo brevioribus. — *C. graecae* similis, quae foliis laeviusculis, arachnoideo-canis, phyllis spina tenuiori et breviori armatis et pappo breviori discedit.

β. **lineariloba** Hal. et Dörfl. in Jahrescat. 1900 der Wien. bot. Tauschanst. p. 130. — Foliorum segmenta multo angustiora, plerumque anguste-linearia. — An species propria. — Exsicc.: Dörfl. fl. aeg. n. 62.

In collibus apricis, lapidosis, ad agrorum margines. Epirus: pr. Janina, ad monasterium Hagios Paraskevi (Form.); Acarnania: pr. Agri-

nion (Hal.); Aetolia: mt. Arapocephala pr. Prustoro et Mikrochorio, pr. Karpenisi (Heldr.); Doris: pr. Segditza (Hal.); Boeotia: pr. Livadia (Guicc.); Achaia: pr. Zugra in mt. Kyllene (Orph.), pr. Megaspilaeon (Heldr.); Cephalonia: pr. Dilinata (Heldr.); Leucas: pr. Karya (Bald.); — β. Cycladum insula Cythnos (Tuntas), Paros (Leon.). — Maio, Aug. ♃.

54. **C. Halácsyi** Dörfl. in ö. b. Z. 1901 p. 204. — Exsicc.: Dörfl. fl. aeg. n. 42.

Caule erecto, crassiusculo, parce arachnoideo, breviter ramoso; foliis parce arachnoideis, laeviusculis, inferioribus petiolatis, lyratis, segmentis lateralibus oblongis, integris vel remote dentatis, terminali multo majore, ovato, acute-dentato, foliis superioribus sursum diminutis, in petiolum attenuatis, lobato-dentatis; capitulis ovato-globosis, 2 cm. diametro; involucri phyllis ovatis, glabris, pallidis, in appendicem semilunatam, fuscidulam, breviter pectinato-ciliatam, spina aciculari, ciliis vix validiori, erecto-patula terminatam, abeuntibus; flosculis purpureis, basi aurantiacis, non radiantibus; acheniis adpresse hirsutis, pappo subbrevioribus. — Praecedenti affinis, ab ea foliis laeviusculis, segmento terminali multo majore et spina aciculari brevi phyllorum distincta; nota ultima accedit ad *C. graecam*, quae autem foliorum forma et indumento facile distinguitur.

In Cycladum insula Amorgos (Leon.). — Maio, Jun. ♃.

:= Folia inferiora obovata integra vel alia lyrata.

55. **C. Minoa** Heldr. in Bois. fl. or. III. p. 662; Hal. in bull. herb. Bois. VI. p. 642. — Exsicc.: Heldr. pl. cret. a. 1870, herb. gr. norm. n. 1046.

Caule erecto, crassiusculo, parce arachnoideo, breviter et stricte ramoso; foliis parce arachnoideis, asperulis, inferioribus petiolatis, aliis ovatis oblongisve indivisis dentatis, aliis lyratis, segmentis laterialibus rotundatis, terminali multo majore, foliis superioribus sessilibus, pinnatisectis, summis subintegris; capitulis ovato-globosis, 20—25 mm. diametro; involucri phyllis ovatis, glabriusculis, pallide virentibus, in appendicem semilunatam, pallidam, breviter pectinato-ciliatam, spina valida, 15—25 mm. longa, canaliculata, basi spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis adpresse hirsutis, pappo triplo brevioribus. — *C. Sprunerianae* maxime affinis et ab ea statura humiliori, foliis infimis indivisis, foliorum pinnatisectorum lobis obtusioribus et capitulis minoribus (an specifice) differt.

In campis, agris derelictis. Attica: pr. Eleusis (Heider); Creta: pr. Candia (Heldr.). — Jun. Jul. ♃.

β. Caulis nullus vel brevissimus, capitula radicalia.

56. **C. mixta** DC. pr. VI. p. 594 excl. loc. asiat.; Hal. in bull. herb. Bois. VI. p. 665. — *C. raphanina* Ch. et B. exp. p. 256, Fl. pelop. p. 60; Link in Linn. IX. p. 581. — ? *C. saxicola* β. *Fraa-*

siana C. H. Schultz in Flora XXV Beibl. p. 160, e descriptione „caule humillimo folia vix superante, capituli longitudine, 1 cephalo". — *C. hellenica* Bois. et Spr. diagn. VI. p. 131, Fl. or. III. p. 675; Clem. sert. p. 59; Ung. Reise p. 124; Raul. cret. p. 789; Weiss in z. b. G. Wien 1869 p. 44; Hal. Beitr. fl. Achaia p. 25; Boissieu in bull. soc. bot. Fr. 1896 p. 286; Heldr. fl. Aegina p. 306, chlor. Thera p. 15. — Exsicc.: Heldr. herb. norm. n. 320, 1047, 1047 b (f. intermedia) et 1448; Orph. fl. gr. n. 123.

Radice crassa, verticali; caule subnullo; foliis radicalibus, rosulatis, parce papilloso-puberulis, sublyrato-pinnatisectis, segmentis parvis, lanceolatis, acutis, denticulatis, basi attenuatis, lobulis intermixtis; capitulis ad collum subsessilibus, 1—4 aggregatis, ovatis; involucri phyllis ovatis, puberulis, pallide-virentibus, in appendicem semilunatam, fuscidulam, pectinato-ciliatam, spina 5—15 mm. longa, erecto-patula, basi spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis glabris vel sericeis, pappo subbrevioribus. — Radix circa 2 cm. crassa, folia 3—15 cm. longa, capitula 15—20 mm. diametro.

β. **caulescens** Hal. in bull. herb. Bois. VI. p. 644. — Caulis 2—7 cm. longus. —

In rupestribus regionis montanae et subalpinae. Boeotia: pr. Thebas (Zuccar.); Attica: pr. Athenas, mt. Pentelicon, Hymettus (Orph.), Laurion (Heldr.), in Petalium insulis (Holzm.), insula Aegina, pr. Kumi Euboeae (Heldr.); Peloponnesus: mt. Kyllene, Malevo, Taygetos (Heldr.), Chelmos, Panachaicon, Olenos (Hal.), pr. Patras (Link), Messene (Chaub.), insula Poros (Wied.), Hydra (Heldr.); Cycladum insula: Keos, Andros (Heldr.), Syra, Tenos (Weiss), Thera (Letourn.); Creta: pr. Askyphos (Raul.); Cephalonia (Mazz.). — Apr. Juli. ♃.

57. **C. myconia** Bois. et Sart. diagn. ser. 2, VI. p. 113; Hal. in bull. herb. Bois. VI. p. 643; Heldr. chlor. Mykon. p. 246. — *C. Urvillei* Bois. fl. or. III. p. 665 p. p., non DC. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889.

Differt (an specifice) a praecedente caule longiore usque 8 cm. alto, foliorum segmentis majoribus, petiolulatis, capitulis submajoribus et phyllorum spinis longioribus.

In Cycladum insulis Tenos (Leon.) et Mykonos (Sart.). — Apr. Maio. ♃.

58. **C. raphanina** S. et S. pr. II. p. 205, Fl. gr. X. p. 12, t. 917; Sieb. avis p. 5, rem. p. 6; Raul. cret. p. 789; Weiss. in z. b. G. Wien 1869 p. 741; Bois. fl. or. III. p. 675; Bald. viagg. Creta p. 71; Heldr. chlor. Mykon. p. 246, Hal. in bull. herb. Bois. VI. p. 645; — *C. nana* Sieb. in Flora V. 2 p. 639, sec. Raul. cret. p. 789. — *C. pumila* Urv. enum. p. 118 non L. et *C. carduncella* DC. pr. VI. p. 591, sec. Bois. l. c. — Exsicc.: Rev. pl. cret. n. 255, in Baen. herb. europ. n. 5222.

Duabus praecedentibus nimis affinis, ab eis involucri phyllis brevissime spinosis, breviter et parce ciliatis, interdum ciliis fere omnino destitutis specifice differt.

In saxosis regionis inferioris et montanae. Creta: in mt. Sphacioticis (Sibth.), mt. Volokia (Rev.) et Hagion Pneuma (Bald.), in mt. Ida (Sieb.), Lassiti (Heldr.), dein pr. Candia, Strombolo, Karadagh, Kaenurio-Khorio, Spinalonga, Hagios Joannes in distr. Sitia (Raul.); ? Cycladum insula Melos (Urv.) et Mykonos (Heldr.). — Jun. Jul. ♃.

Obs. *C. exscapa* Urv. enum. p. 117, species subacaulis flaviflora, indicatur a Bois. fl. or. III. p. 678 in insula Naxos, ubi ab Urville lecta esse dicitur, sed probabiliter erronee, nam autor speciem suam in insulis asiaticis Cos et Leros tantum indicat. — *C. crocodylium* L. sp. p. 919 indicatur in Lam. dist. I. p. 677 in Creta, sed a recentioribus ibi non lecta fuit.

b. Flosculi flavi.

α. Caulis 20—60 cm. altus.

59. **C. Tuntasia** Heldr. ap. Hal. in bull. herb. Bois. VI. p. 646. — Exsicc.: Heldr. herb. norm. n. 1447.

Papilloso-scabrida; caulibus erectis, stricte ramosis, rarius simplicibus; foliis coriaceis, canescentibus, in segmenta oblonga, integra vel parce denticulata sublyrato-pinnatisectis, inferioribus petiolatis, superioribus sessilibus, summis subintegris; capitulis ovatis, 15—20 mm. diametro; involucri phyllis ovatis, glabris, in appendicem late semilunatam, pectinato-ciliatam, phylla subtegentem, phyllorum intermediorum adpressam, spina valida, canaliculata, inferne spinulosa terminatam, superiorum convexam, muticam, abeuntibus; flosculis ochroleucis, non radiantibus; acheniis sericeis, pappo 4 plo brevioribus. — Sequenti affinis, ab ea canescentia, phyllorum appendicibus majoribus, phylla subtegentibus, superioribus convexis et pappo longiore distinctissima.

In locis cultis, ad vineta pr. Liosia Atticae (Tuntas). — Jun. Jul. ♃.

60. **C. salonitana** Vis. in Flora XII. Erg.Bl. I. p. 23, Fl. dalm. II. p. 35, t. 13; Bois. fl. or. III. p. 666; Heldr. in Sitzungsb. acad. Berlin 1883 p. 4 et 8, chlor. Parn. p. 22; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Thessal. p. 15, in ö. b. Z. 1897 p. 285, in bull. herb. Bois. VI. p. 645; Form. in D. bot. Monat. 1890 p. 19, in Ver. Brünn 1895 p. 27, 1896 p. 49, 1897 p. 35; Haussk. symb. p. 131; Bald. riv. coll. bot. Alb. 1895 p. 51, 1896 p. 70. — *C. latisquama* DC. pr. VI. p. 589. — *C. collina* S. et S. pr. II. p. 204, Fl. gr. X. p. 10 t. 914; Sieb. in Flora I. p. 275; Ch. et B. fl. pelop. p. 60; Raul. cret. p. 789; non L. — *C. centauroides* S. et S. pr. II. p. 204; Ch. et B. exp. p. 255, Fl. pelop. p. 60; non L. — Exsicc.: Orph. fl. gr. n. 306; Heldr. herb. norm. n. 624; Sint. it. thessal. 1228.

Papilloso-scabrida; caulibus erectis, stricte ramosis; foliis coriaceis, viridibus, in segmenta oblonga vel oblongo-lanceolata, integra vel rarius parce denticulata pinnatisectis, inferioribus petiolatis, superioribus sessilibus, summis indivisis; capitulis ovatis, 15—20 mm. diametro; involucri phyllis ovatis, glabris, in appendicem adpressam, anguste semilunatam, pectinato-ciliatam, phylla non tegentem, muticam vel spina usque 4 cm. longa, canaliculata, inferne spinulosa terminatam, abeuntibus; flosculis suffureis, non radiantibus; acheniis sericeis, pappo subbrevioribus.

α. **subinermis** Bois. fl. or. III. p. 666. — Involucri phylla mutica vel spina ciliis non vel parum longiore terminata. — Intermediis ad

β. **macracantha** Bois. diagn. ser. 2, III. p. 78, spinis validioribus usque 4 cm. longis, transit.

In campis, arvis, vinetis regionis inferioris et montanae. Epirus: in valle Dipotamo, pr. Han Kanberga (Form.), mt. Smolika, Prosgoli (Bald.); frequens in Thessalia a Pindo, mt. Oxya et Chassia, per planitiem usque ad mt. Olympus, Ossa et Pelion, Oeta et in Phthiotide; Aetolia: mt. Korax (Heldr.); mt. Kiona pr. Segditza, mt. Parnassus pr. Livadi (Hal.) et Rachova (Orph.); Attica: pr. Mesoitis, Amarysia, Kephissia, Dekeleia (Heldr.), mt. Pentelicon (Leutwein); Arcadia: pr. Tripolizza (Sart.); Cycladum insula Melos (Armenis); Creta: pr. Candia, Pedhiada, in oropedio Nida in mt. Ida (Raul.). — Jul. Aug. ♃.

Obs. *C. macedonica* Bois. diagn. VI. p. 130, Orph. fl. gr. exs. n. 747; Heldr. herb. gr. norm. n. 1156 (nomen specificum ceteroqui mutandum est ob homonymum Grisebachianum in *C. thessalonica*), praecedenti similis, sed ab ea foliorum segmentis longis angustis, phyllis araneosis parce et brevissime pectinato-ciliatis diversissima, ad confines ditionis nostrae pr. Thessalonicam crescens, in Thessalia septentrionali probabiliter occurrit. — Ibidem quoque forsitan inquirenda species altera, flaviflora, *C. chrysolepidae* Vis. affinis, splendidissima, a Charrel pr. Vodena Macedoniae detecta et a nobis sub nomine *C. Charrelii* Hal. et Dörfl. in Dörfl. Jahrescat. 1894 Wien. bot. Tauschver. p. 6 descripta (Caule elato, 50—70 cm. alto, subarachnoideo, breviter ramoso, folioso; foliis oblongo-lanceolatis, integris, asperulis, secus caulem longe lateque decurrentibus, summis capitula subbracteantibus; capitulis ovato-globosis, 4 - 5 ½ cm. diametro; involucri pallidi phyllis ovatis, in appendicem magnam, stramineam, brevissime puberulam, ovato-lanceolatam, pectinato-ciliatam, phylla tegentem, spina ciliis latitudine appendicis aequilongis subvalidiore et subduplo longiore terminatam, abeuntibus; flosculis flavis, non radiantibus; acheniis glabris, pappo subduplo brevioribus).

β. Caulis 1—3 cm. altus.

61. **C. parnonia** Hal. in bull. herb. Bois. VI. p. 648. — *C. rupestris* v. *minor* Bois. fl. or. III. p. 668, non *C. minor* Willd. — Exsicc.: Orph. pl. gr. a. 1858.

Arachnoidea, cana; caulibus erectis, nanis, simplicibus, monocephalis; foliis in lacinias oblongas, angustas, breves pinnatisectis, radicalibus petiolatis, caulinis sessilibus; capitulis ovatis, 10—15 mm. diametro; involucri phyllis ovatis, parce araneosis, pallidis, in appendicem ovatam, centro fuscidulam, margine hyalinam, pectinato-ciliatam, spina brevi, ciliis validiore terminatam, abeuntibus; flosculis luteis, non radiantibus; acheniis subsericeis, pappo aequilongis. — Affinis *C. rupestris* L. differt ab ea caulibus saepissime elatis et ramosis, glabrescentia, foliis viridibus, laciniis anguste linearibus, longis et pappo breviore.

In cacumine mt. Malevo — Parnon veterum — Laconiae (Orph.), rarissime. — Jul. Aug. ♃.

4. Sectio. *Calcitrapeae* DC. l. c. p. 592. —

a. Folia caulina decurrentia; flosculi flavi.

α. Flosculi eglandulosi.

62. **C. solstitialis** L. sp. p. 917; S. et S. pr. II. p. 202, Fl. gr. X. p. 6 t. 908; Ch. et B. exp. p. 255, Fl. pelop. p. 60; Marg. et R. fl. Zante p. 59; Friedr. Reise p. 273 et 284; Fraas fl. class. p. 204; Raul. cret. p. 789; Weiss in z. b. G. Wien 1869 p. 44; Bois. fl. or. III. p. 685; Heldr. fl. cephal. p. 48, in ö. b. Z. 1898 p. 184, var. *insularis* (f. ramulis elongatis); Spreitz. in z. b. G. Wien 1887 p. 663, 1890 p. 296; Hal. in z. b. G. Wien 1888 p. 759, Beitr. fl. Epir. p. 29, Beitr. fl. thessal. p. 15, in bull. herb. Bois. VI. p. 649; Form. in D. bot. Monat. 1890 p. 19, in Ver. Brünn 1895 p. 27, 1896 p. 49, 1897 p. 35; Haussk. symb. p. 131; Bald. viagg. Creta p. 71. — ? *C. sicula* S. et S. pr. II. p. 202. — Exsicc.: Heldr. herb. norm. n. 1449 var. *pycnoclada* (f. capitulis numerosioribus in ramulis abbreviatis; Dörfl. fl. gr. n. 210 et 394).

Araneoso-canescens; caule erecto, paulo supra basin divaricatim ramosissimo; foliis radicalibus petiolatis, lyratis, laciniis oblongis, integris vel dentatis, caulinis sessilibus, lineari-lanceolatis, integris, decurrentibus; capitulis solitariis, ovatis, 6—10 mm. diametro; involucri phyllis virescentibus, parce araneosis, in appendicem parvam, stramineam, spina patula, acerosa, 10—20 mm. longa, basi utrinque 1—2 spinulosa terminatam, abeuntibus; flosculis luteis, non radiantibus; acheniis glabris, pappo aequilongis vel subbrevioribus.

β. **Adami** Willd. sp. III. p. 2310; Friedr. Reise p. 273; Hal. in ö. b. Z. 1890 p. 39; pro sp.; Bois. fl. or. III. p. 685; Haussk. symb. p. 131; Hal. in bull. herb. Bois. VI. p. 650. — Phyllorum spina tenuior et brevior, spinulas laterales saepe parum superans. — Exsicc.: Sint. it. or. 1889 n. 1835; Heldr. it. IV. Thessal. a. 1885.

In aridis, ruderatis, arvis regionis inferioris, per totam Graeciam vulgaris; — β. Thessalia: pr. Trikala (Heldr.), Megala Kalyvia (Haussk.), Katerina (Sint.); Argolis: pr. Poros (Friedr.). — Jun. Jul. ⊙.

63. **C. idaea** Bois. et Heldr. diagn. X. p. 119, Fl. or. III. p. 686 suppl. p. 316; Raul. cret. p. 789; Hal. in bull. herb. Bois. VI.

p. 650. — *C. solstitialis v. idaea* Bald. viagg. Creta p. 71. — Exsicc.: Heldr. pl. cret. a. 1870; Rev. pl. cret. n. 81; Bald. it. cret. alt. n. 33.

Araneoso-cana; caulibus a collo pluribus, pumilis, centrali brevissimo, monocephalo, lateralibus divaricatim ramosis, pleiocephalis; foliis radicalibus rosulatis, petiolatis, in segmenta ovato-triangularia pinnatipartitis, caulinis inferioribus minoribus, decurrentibus, summis integris diminutis, oblongo-linearibus; capitulis solitariis, ovatis, 6—10 mm. diametro; involucri phyllis virentibus, araneosis, in appendicem parvam stramineam vel fuscidulam, spina patula, acerosa, 10—20 mm. longa, basi utrinque 1—2 spinulosa terminatam abeuntibus; flosculis luteis, non radiantibus; acheniis glabris, pappo aequilongis. — A praecedente radice crassiuscula bienni, caulibus a collo numerosis, centrali brevissimo simplici, foliis radicalibus rosulatis, segmentis parvis numerosis specifice diversissima.

In aridis regionis inferioris et montanae. Cretae: pr. Kissamos, Malaxa (Reverch.), Stylo, Kephala, Anopolis, Aradhena in mt. Sphacioticis, mt. Ida (Raul.), pr. Spilia ultra Temenos (Bald.). — Jun. Aug. ⊙.

β. Flosculi glandulosi.

64. **C. melitensis** L. sp. p. 917; S. et S. pr. II. p. 202, Fl. gr. X. p. 6 t. 909; Ch. et B. exp. p. 256, Fl. pelop. p. 60; Heldr. fl. Aegina p. 306; Hal. in bull. herb. Bois. VI. p. 651. — *C. apula* Lam. dict. I. p. 674; ? Raul. cret. p. 789. — ? *C. lyrata* Sieb. avis. p. 5. — Exsicc.: Heldr. pl. fl. hellen. n. 2069.

Asperulo-pubescens, tandem glabrata; caule erecto, ramoso; foliis in petiolum attenuatis, inferioribus lyrato-pinnatifidis, laciniis paucis brevibus obtusis, ceteris oblongo-linearibus linearibusve, integris, decurrentibus; capitulis ovatis, solitariis vel aggregatis; involucri phyllis virescentibus, parce araneosis, in appendicem parvam, fuscidulam, spina patulo-recurva, acerosa, utrinque 2—3 spinulosa terminatam, abeuntibus; flosculis pallide luteis, glandulosis, non radiantibus; acheniis glabris, pappo sublongioribus.

In collibus aridis, ruderatis, rarissime. Prope Perivolia et Hagia Marina in insula Aegina (Heldr.); dein in Laconia (Sibth.) et si planta Sieberi huc spectet, in Creta pr. Canea. — Jun. Aug. ⊙.

b. Folia non decurrentia.

α. Flosculi purpurei.

65. **C. calcitrapa** L. sp. p. 917; S. et S. pr. II. p. 201; Sieb. avis p. 5, rem. p. 6; Ch. et B. exp. p. 255, Fl. pelop. p. 60; Fraas fl. class. p. 204; Raul. cret. p. 789; Bois. fl. or. III. 689; Heldr. fl. cephal. p. 48; Hal. in z. b. G. Wien 1888 p. 763, Beitr. fl. Epir. p. 29, Beitr. fl. Achaia p. 25, in bull. herb. Bois. VI. p. 651; Spreitz. in z. b. G. Wien 1890 p. 296; Form. in D. bot. Monat. 1890 p. 19, in Ver. Brünn 1895 p. 27, 1896 p. 49, 1897 p. 35; Haussk. symb.

p. 131. — *C. calcitrapa v. longispina* Aschers. in z. b. G. 1889 p. 44. — Icon: Rchb. t. 67. — Exsicc.: Dörfl. fl. gr. n. 420. —

Caule erecto, a basi divaricatim ramosissimo; foliis viridibus, asperulo-puberulis, radicalibus rosulatis, in lacinias lineares dentatas vel pinnatilobatas pinnatisectis, caulinis sessilibus, laciniis paucis, elongatis, summis indivisis; capitulis solitariis, ovatis, terminalibus et lateraliter ad ramos vel in dichotomiis subsessilibus; involucri phyllis ovatis, glabris, pallidis, anguste hyalino-marginatis, in appendicem parvam, spina patula, basi utrinque 2—3 spinulosa terminatam, abeuntibus; flosculis purpureis, non radiantibus; acheniis glabris, calvis.

In ruderatis, ad vias, sepes regionis inferioris et montanae per omnem fortasse Graeciam continentalem frequens; dein in Corcyra, Cephalonia et Creta, probabiliter etiam in insulis ceteris. — Jun. Sept. ⊙.

66. **C. iberica** Trev. in Spreng. syst. III. p. 406; Bois. fl. or. III. p. 690; Haussk. symb. p. 131; Form. in Ver. Brünn 1896 p. 49, 1897 p. 35. — *C. aegyptiaca* S. et S. pr. II. p. 201, Fl. gr. X. p. 5 t. 907; Ch. et B. fl. pelop. p. 60; non L. — *C. calcitrapoides* DC. pr. VI. p. 597, an L. — Exsicc.: Heldr. it. IV. Thessal. a. 1885.

Praecedenti simillima, ab ea praesertim acheniis pappo eis dimidio breviore superatis distincta.

β. **Holzmanniana** Heldr. herb. fl. hellen. n. 111 pro sp.; Bois. fl. or. suppl. p. 316; Hal. in bull. herb. Bois. VI. p. 653. — Involucri spinae validiores, 3 cm. longae, basi 5 mm. latae, pappus achenio parum brevior.

In ruderatis, ad vias. Thessalia: pr. Trikala (Heldr.), ad monasterium Korona, pr. Tyrnavos, Larissa, Karditza, Volo (Haussk.), Ilias Parapansa, Palaeokastro, Sermeniko, Mavro Mandila, Limogardi, Velestino, Ligaria, Godaman, Nezeros (Form.); Attica: in oliveto Athenarum (Heldr.); Achaia (Sibth.); — *β.* pr. Hagios Mercurios in mt. Parnes Atticae (Heldr.). — Jul. Sept. ⊙.

β. Flosculi sulfurei.

67. **C. hyalolepis** Bois. diagn. VI. p. 133; Haussk. symb. p. 131; Hal. in bull. herb. Bois. VI. p. 653. — *C. pallescens v. hyalolepis* Bois. fl. or. III. p. 691, suppl. p. 316. — Exsicc.: Reverch. pl. cret. n. 80; Heldr. herb. norm. n. 852.

Caule erecto, a basi divaricatim ramosissimo; foliis viridibus, puberulis, radicalibus rosulatis, in segmenta oblonga vel lanceolata, dentata vel lobata pinnatisectis vel lyratis, superioribus sessilibus, linearibus, dentatis; capitulis solitariis, ovatis, terminalibus et lateraliter ad ramos vel in dichotomiis subsessilibus; involucri phyllis ovatis, glabris, pallidis, in appendicem semilunatam, hyalinam, spina patula, interdum utrinque 1 spinulosa terminatam, abeuntibus; flosculis sulfureis, non radiantibus; acheniis glabris, pappo aequilongis. — A *C. iberica* praesertim florum colore discedit. —

In valle Cephissi pr. Myli oliveti Athenarum (Heldr.) et pr. Kissamos et Canea Cretae (Reverch.). — Maio, Jun. ⊙.

5. Sectio. *Seridieae* DC. l. c. p. 598.

a. Flosculi purpurei.

α. Folia caulina basi auriculato-amplexicaulia, non decurrentia.

68. **C. sphaerocephala** L. sp. p. 916; S. et S. pr. II. p. 201, Fl. gr. X. p. 3, t. 904; Raul. cret. p. 789; Bois. fl. or. III. p. 691; Hal. in bull. herb. Bois. VI. p. 654. — Icon: Rchb. t. 69.

Subaraneosa, asperula; caulibus erectis vel procumbentibus, ramosis; foliis lyratis vel sinuato-lobatis, lobis obtusis dentatis, infimis petiolatis, ceteris basi rotundato-auriculatis, summis saepe integris; capitulis bracteatis, solitariis, ovato-globosis; involucri phyllis ovatis, glabris vel puberulis, in appendicem parvam, pallidam, patulo-reflexam, in 5—7 spinulas tenues, phyllo subaequilongas palmatipartitam abeuntibus; flosculis purpureis, radiantibus; acheniis glabris, exterioribus pappo nullo, centralibus pappo brevissimo coronatis.

In arenosis Archipelagi (Sibth.), in Creta (Raul.). — Apr. Jun. ♃. N. v.

β. Folia caulina decurrentia.

69. **C. sonchifolia** L. sp. p. 915; Marg. et R. fl. Zante p. 59; Bois. fl. or. III. p. 692; Hal. in bull. herb. Bois. VI. p. 655. — Huc probabiliter *C. romana* Ch. et B. exp. p. 256, Fl. pelop. p. 60, e Methone et Kardamyle. — Exsicc.: Heldr. herb. norm. n. 1558.

Subaraneosa, asperula; caulibus erectis vel adscendentibus, ramosis; foliis lyratis vel obtuse lobatis, spinuloso-denticulatis, infimis petiolatis, ceteris lanceolatis, oblongisve, remote-dentatis, decurrentibus; capitulis bracteatis, solitariis, ovatis; involucri phyllis ovatis, glabris, in appendicem parvam, pallidam, reflexam, in 5—7 spinulas tenues, phyllo subaequilongas vel breviores palmatipartitam abeuntibus; flosculis purpureis, subradiantibus; acheniis glabris, pappo 2—3 plo longioribus. — Habitus praecedentis, sed folia decurrentia.

In arenosis maritimis, rarissime. Corcyra: pr. Melikia (Bicknell); Zante: pr. Tsilivi (Marg.); Elis: pr. Lintzi (Heldr.); Messenia: pr. Navarin (Heldr.). — Maio, Jun. ♃.

70. **C. napifolia** L. sp. p. 916; S. et S. pr. II. p. 201, Fl. gr. X. p. 3 t. 905; Sieb. avis rem. p. 6; Friedr. Reise p. 203; Raul. cret. p. 789; Bois. fl. or. III. p. 694; Hal. in bull. herb. Bois. VI. p. 655. —

Lanuginosa; caulibus erectis, ramosis; foliis inferioribus petiolatis, lyratis, segmentis obtuse dentatis, lateralibus utrinque 1—3, terminali ovato-rotundo, multo majore, mediis et superioribus sublyratis, decurrentibus, summis lanceolatis; capitulis vix bracteatis, solitariis, ovatis; involucri phyllis ovatis, glabris, in appendicem pallidam, semiorbicularem, erecto-patulam, pectinatim breviter 5—9 spinulosam abeuntibus; flosculis

purpureis, valde radiantibus; acheniis glabris, pappo duplo longioribus. — Differt a praecedente radice annua, foliorum segmento terminali multo majore, appendicibus phyllorum transverse latioribus et flosculis valde radiantibus.

Teste Linné, Sibthorp et Sieber in Creta et secundum Friedrichsthal in valle fluminis Eurotas inter Leondari et Misitra, sed a recentioribus non lecta. — Maio, Jun. ⊙. N. v.

b. Flosculi lutei.

71. **C. lancifolia** Sieb. in Spreng. syst. III. p. 405, in avis p. 5, rem. p. 6; Bois. fl. or. III. p. 694; Hal. in bull. herb. Bois. III. p. 656. — *Amberboa lancifolia* DC. pr. VI. p. 561; Raul. cret. p. 788. — *Chartolepis lancifolia* Fenzl. in Tchih. As. min. III. p. 327. — Exsicc.: Sieb. pl. cret.; Reverch. pl. cret. n. 90.

Minute asperula; caulibus erectis, simplicibus vel superne parce ramosis; foliis rigidulis, lanceolatis, elevatim nervosis, infimis in petiolum attenuatis, ceteris subauriculatis, brevissime decurrentibus; capitulis solitariis, ovatis, foliis summis bracteatis; involucri phyllis ovatis, glabriusculis, in appendicem fusco-nigram, erectam, palmato-subpectinatim 9—13 spinulosam abeuntibus; flosculis luteis, non radiantibus; acheniis glabris, pappo aequilongis. — Caulis 10—20 cm. altus, folia 5—6 cm. longa, 4—6 mm. lata, eis *Inulae salicinae* similia; capitula 15 mm. diametro.

In montibus pr. Sphakia (Sieb.) et Drakona Cretae (Reverch.). — Jul. ♃.

Species dubiae:

C. armoracifolia S. et S. pr. II. p. 205, e mt. Peloponnesi. Cf. Spreng. syst. III. p. 396; Ch. et B. exp. p. 256, Fl. pelop. p. 60; DC. pr. VI. p. 603; Bois. fl. or. III. p. 656. —

C. coronopifolia S. et S. pr. II. p. 204, e Graecia, sine loci specialis indicatione.

C. dalmatica Fraas fl. class. p. 204, in Euboea pr. Mantudi et Limni (fibris radicis napuliformi-incrassatis).

C. eumorpha Sieb. avis rem. p. 6 (solum nomen), e Creta.

56. Microlonchus Cass. bull. phil. 1818 p. 142.

1. **M. salmanticus** L. sp. p. 918 *(Centaurea)*; DC. pr. VI. p. 565; Form. in Ver. Brünn 1896 p. 49; Heldr. fl. Aegina p. 306. Icon: Jacq. vindob. t. 64. — Exsicc.: Heldr. pl. fl. hellen. a. 1848 et 1877.

Caule erecto, virgatim ramoso, inferne parce puberulo; foliis inferioribus petiolatis, runcinato-vel lyrato-pinnatipartitis, mucronato-denticulatis, superioribus sessilibus, linearibus; capitulis solitariis, ovato-conicis; involucri glabri phyllis adpressis, ovatis, apice nigro-maculatis, mucrone brevissimo caduco terminatis; flosculis purpureis, non radiantibus;

acheniis inter costas transverse rugulosis; pappo sordido, achenio subbreviore.

In sterilibus, incultis, vineis, ad vias, rarissime. Thessalia: pr. Velestinos (Form.); Attica: ad ostia Cephissi pr. Phalerum (Kyriakos); pr. Corinthum et in insula Aegina (Heldr.); Argolis: pr. Nauplia (Sart.). — Maio, Jul. ♃.

57. Crupina Cass. dict. 12 p. 67.

1. **C. vulgaris** Cass. dict. 44 p. 39; Marg. et R. fl. Zante p. 59; Ung. Reise p. 124; Form. in D. bot. Monat. 1890 p. 20; in Ver. Brünn 1896 p. 49 et 1897 p. 36 *v. serrata* (f. foliorum segmentis denticulatis); Haussk. symb. p. 131; Bald. riv. coll. bot. Alb. 1896 p. 70. — *Centaurea crupina* L. sp. p. 909; Clem. sert. p. 57. — *Serratula crupina* Vill. dauph. III. p. 38. — Icon: Rchb. XXV. t. 18. — Exsicc.: Sint. it. thessal. n. 910.

Asperula; caule erecto, stricte ramoso, ramis tenuibus, nudis; foliis primordialibus indivisis, oblongis, basi attenuatis, ceteris in segmenta tenuiter linearia, integra vel denticulata pinnatisectis; capitulis anguste oblongis, 3—5 floris, basi attenuatis; involucri pruinosi phyllis pallide virentibus, membranaceo marginatis, apice saepe purpurascentibus, ab infimis ovatis ad superiora lanceolata acuminata auctis; flosculis purpureis, involucrum parum superantibus; acheniis subcylindricis, basi non compressis, hilo basilari orbiculari; pappi interni paleis triangularibus, acutis.

In collibus siccis regionis inferioris et submontanae. Epirus: pr. Vromonero distr. Ljaskovik (Bald.); Thessalia: pr. Kastania, Velitsena, Koturi, Malakasi, Chilniades in valle Penei, Kucuro in mt. Chassia, pr. Trikala, Velestinos, Kapurna, Kalyvia, Portaria, Volo, mt. Pelion (Form.), Pharsalus, Aivali, ad monasterium Korona (Haussk.); Euboea: pr. Kurbatzi (Wild); Attica: pr. Athenas (Ung.); Zante: mt. Scopo (Marg.); Corcyra: pr. Spartilla (Clem.); — sed loca nonnulla probabiliter ad sequentem spectant. — Maio, Jul. ⊙.

2. **C. crupinastrum** Mor. en. sem. hort. taur. 1842 p. 12 (*Centaurea*); Vis. fl. dalm. II. p. 42, t. 51; Weiss in z. b. G. 1869 p. 45; Bois. fl. or. III. p. 699; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 48, Fl. Aegina p. 306, chlor. Thera p. 15; Hal. Beitr. fl. Epir. p. 29, Beitr. fl. Achaia p. 25, in ö. b. Z. 1895 p. 460, 1897 p. 96; Haussk. symb. p. 131; Form. in Ver. Brünn 1895 p. 27, 1896 p. 49, 1897 p. 36. — *Centaurea crupina* S. et S. pr. II. p. 197, Fl. gr. t. 900; Urv. enum. p. 115; Ch. et B. exp. p. 254, Fl. pelop. p. 59; Friedr. Reise p. 283. — *Crupina Morisii* Bor. fl. centr. II. p. 292; Raul. cret. p. 788. — Huc probabiliter *Serratula attica* Fraas fl. class. p. 210. — Exsicc.: Heldr. herb. norm. n. 319 et 1352; Orph. fl. gr. n. 468; Reverch. pl. cret. n. 196.

Differt a praecedente, cui habitu simillima, capitulis basi rotundatis, saepe plurifloris (9—15), flosculis involucro fere sesquilongioribus,

acheniis basi acute compressis, hilo obliquo lineari, pappi interni paleis oblongis denticulatis.

In iisdem locis, praecedenti multo frequentior, passim per Graeciam totam. — Maio, Jul. ⊙.

58. Aegialophila Bois. et Heldr. diag. X. p. 105.

1. **A. cretica** Bois. et Heldr. diagn. X. p. 106, Fl. or. III. p. 704; Raul. cret. p. 790. — *Centaurea pumila* S. et S. pr. II. p. 206, Fl. gr. X. p. 13 t. 918; Sieb. avis. p. 5, rem. p. 6; Raul. cret. p. 789; Ch. et B. fl. pelop. p. 60; non L. — Exsicc.: Heldr. pl. cret. n. 1451; Sint. et Rigo it. cypr. n. 296 (Cyprus).

Araneoso-canescens; radice crassa; caulibus brevissimis, parce ramosis; foliis petiolatis, capitula superantibus, aliis indivisis, ovato-oblongis, aliis lyratis, segmento terminali multo majore; capitulis ovatis; involucri glabriusculi phyllis ovatis, membranaceo-marginatis, in spinulam phyllo 4 plo breviorem attenuatis; flosculis purpureis, subradiantibus; pappi setis externis achenio sericeo subtriplo longioribus, seriei intimae paleolis pappi externi tertiam partem aequantibus. — *A. pumilae* (L.) Bois. foliis pinnatisectis, phyllis brevius spinosis et seriei intimae paleolis pappi externi sextam partem aequantibus, diversae, valde affinis.

In arenosis maritimis ad portum Sitia Cretae orientalis (Heldr.). — Apr. Maio. ♃.

59. Carthamus L. gen. n. 411.

a. Flosculi flavi.

1. **C. lanatus** L. sp. p. 830; S. et S. pr. II. p. 160, Fl. gr. t. 841; Sieb. avis. p. 5, rem. p. 5; Ch. et B. exp. p. 244, Fl. pelop. p. 57; Friedr. Reise p. 284; Fraas fl. class. p. 206; Bois. fl. or. III. p. 706; Heldr. fl. cephal. p. 48, Fl. Aegina p. 305, in ö. b. Z. 1898 p. 184; Hal. in z. b. G. 1888 p. 759, 1899 p. 186, Beitr. fl. Epir. p. 28, Beitr. fl. thessal. p. 14; Haussk. symb. p. 125; Form. in Ver. Brünn 1895 p. 27, 1896 p. 49, 1897 p. 36. — *Kentrophyllum lanatum* Duby bot. gall. I. p. 293; Marg. et R. fl. Zante p. 59; Weiss in z. b. G. 1896 p. 45; Raul. cret. p. 790; Spreitz. in z. b. G. 1887 p. 663; Form. in D. bot. Monat. 1890 p. 20. — Exsicc.: Sint. it. thessal. n. 908; Dörfl. fl. gr. n. 395.

Caule erecto, corymboso, plus minus arachnoideo; foliis coriaceis, glanduloso-viscidulis, elevatim nervosis, inferioribus pinnatipartitis, superioribus pinnatifidis vel sinuatis, omnibus inaequaliter spinoso-dentatis; capitulis ovatis; involucri phyllis externis foliaceis, erecto-patentibus, pinnatim dentato-spinosis, flores aequantibus vel parum superantibus, intimis scariosis, lineari-lanceolatis, integris; flosculis luteis; acheniis turbinato-tetragonis, superne excavato-punctatis; pappi paleis ab externis brevissimis retusis, ad intermedias acutas, achenio triplo longiores auctis, abbreviatis.

β. **creticus** L. sp. ed. 2 p. 1163, syst. nat. ed. 12, II. p. 533 Sieb. avis p. 5, rem. p. 5; Ch. et B. fl. pelop. p. 57; Celak. in Sitzungsb. böhm. Ges. Wiss. 1885 p. 8; Hal. in z. b. G. 1899 p. 185; pro sp. — *C. lanatus v. graecus* Heldr. herb. norm. n. 1152. — *C. cycladum* Heldr. pl. fl. hellen. a. 1897. — Vix arachnoidea, foliis nitidis, angustioribus longioribusque, capitulis submajoribus, involucri phyllis externis patentissimis, flores pallide-luteos manifeste superantibus. — Formae ad typum transientes adesse videntur. — Exsicc.: Heldr. l. c.; Orph. fl. gr. n. 789; Dörfl. fl. aeg. n. 48.

In ruderatis, cultis, campis regionis inferioris per totam Graeciam, varietas typo ut videtur multo frequentior. — Maio, Aug. ⊙.

b. Flosculi purpurei.

α. Pappi series intima serie intermedia 3—4 plo brevior.

× Involucri phylla intermedia apice non constricta, sensim in spinam integram vel utrinque spinula instructam abeuntia.

2. **C. Boissieri** Hal. in z. b. G. 1899 p. 186. — *Kentrophyllum creticum* Bois. diagn. X. p. 99; Weiss in z. b. G. 1869 p. 741; Raul. cret. p. 790. — *C. glaucus v. syriacus* Bois. fl. or. III. p. 707, saltem quoad pl. cret. — *C. glaucus* Bois. fl. or. suppl. p. 317, non M. B. — *K. tauricum* Raul. cret. p. 790, non (M. B.). — *C. ruber* Bald. viagg. cret. p. 68, non Link. — Exsicc.: Reverch. pl. cret. n. 83, in Baen. herb. europ. n. 4883; Bald. it. cret. n. 126.

Caule erecto, corymboso, arachnoideo; foliis oblongo-lanceolatis, coriaceis, arachnoideis, glanduloso-viscidulis; elevatim nervosis, spinoso-dentatis; capitulis oblongis; involucri phyllis externis foliaceis, patentibus, spinoso-dentatis, capitulum superantibus; acheniis ovato-tetragonis, superne parce excavato-punctatis; pappi paleis ab extimis brevissimis retusis, ad intermedias acutas sensim auctis, intimis abbreviatis.

In locis aridis, collibus, cultis. Creta: pr. Kissamos, Canea, Khalepa (Reverch.), Armeni (Bald.), Melidoni (Raul.); Cycladum insula Amorgos (Heldr.). — Jun. Aug. ⊙.

3. **C. leucocaulos** S. et S. pr. II. p. 160, Fl. gr. t. 842; Sieb. avis p. 5, rem. p. 5; Urv. enum. p. 104; Bois. fl. or. III. p. 708; Bald. viagg. cret. p. 68; Heldr. fl. Aegina p. 305, in ö. b. Z. 1898 p. 184, chor. Thera p. 15. — *Kentrophyllum leucocaulon* DC. pr. VI. p. 610; Weiss in z. b. G. 1869 p. 45; Raul. cret. p. 790. — Exsicc.: Bald. it. cret. n. 159; Dörfl. fl. aeg. n. 265.

Differt a praecedente caule nitido, glaberrimo, candido, foliis angustis, nitidis glabris, longius spinoso-dentatis, involucri phyllis externis capitulo plus duplo longioribus.

In locis aridis regionis inferioris. Argolis: pr. Nauplia (Sart.); insula Aegina, Poros, Naxos, Siphnos, Amorgos, Thera (Heldr.), Thermia (Auch.), Syra (Urv.); Creta: pr. Asomatos distr. Amari (Bald.), insula Dia (Sieb.). — Jul. Aug. ⊙.

XX Involucri phylla intermedia apice constricta, in appendicem oblongam, pectinato-spinulosam, apice spinosam dilatata.

4. **C. ambiguus** Heldr. ap. Hal. in z. b. G. 1899 p. 186. — Huc probabiliter *C. dentatus* Urv. enum. p. 104, Heldr. in ö. b. Z. 1898 p. 184 et *Kentrophyllum dentatum* Weiss in z. b. G. 1869 p. 45. — Exsicc.: Heldr. pl. fl. hellen. a. 1897; Dörfl. fl. aeg. n. 266.

Caulo erecto, arachnoideo, mox supra basin patule ramosissimo; foliis coriaceis, pubescenti-viscidulis, elevatim nervosis, semiamplexicaulibus, oblongo-lanceolatis, spinoso-dentatis; capitulis oblongis; involucri phyllis externis foliis similibus, patentibus, capitulo duplo longioribus, interioribus scariosis, in appendicem oblongam, pectinato-spinulosam, apice tenuiter spinosam dilatatis; acheniis turbinato-tetragonis; pappi paleis ab extimis brevissimis retusis, ad intermedias acuminatas, serrulato-dentatas, achenio triplo longiores elongatis, seriei intimae eis 2—3 plo brevioribus. — Sequenti valde affinis et ab eo forsan vix specifice distinctus.

In aridis regionis inferioris insulae Aegina (Tunt.), Paros, Naxos (Heldr.), ? Syra (Urv.). — Jun. Oct. ⊙.

5. **C. dentatus** Vahl symb. I. p. 69, t. 17; Bois. fl. or. III. p. 708 p. p.; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 8, in ö. b. Z. 1898 p. 184; Celak. in böhm. Ges. Wiss. 1885 p. 2; Haussk. symb. p. 125; Form. in Ver. Brünn 1896 p. 49, 1897 p. 36; Hal. in z. b. G. 1899 p. 188. — *Kentrophyllum dentatum* DC. pr. VI. p. 611; Form. in D. bot. Monat. 1890 p. 19. — Exsicc.: Heldr. it. thessal. n. 95; Sint. et Bornm. it. turc. n. 1313; Sint. it. thessal. n. 1018; Dörfl. fl. gr. n. 190.

Differt a praecedente caule elatiore, minus arachnoideo, superne vel a medio ramoso, interdum simplici, monocephalo et involucri phyllis externis brevioribus, capitulum aequantibus vel parum superantibus.

In locis aridis, collibus, ad vias. Frequens in Thessalia a Malakasi in Pindo, per planitiem usque ad radices mt. Olympus, Ossa et Pelion; pr. Agoriani in mt. Parnasso (Leon.); ? pr. Athenas (Haussk.). — Jun Aug. ⊙.

β. Pappi series intima serie intermedia longior vel aequilonga (involucri phylla intermedia uti duorum antecedentium).

6. **C. ruber** Link in Linnaea 1834 p. 580; Ch. et B. fl. pelop. p. 57; Celak. in böhm. Ges. Wiss. 1885 p. 2; Hal. in z. b. G. 1899 p. 188. — *Kentrophyllum rubrum* Link. l. c. — *C. dentatus* Bois. fl. or. III. p. 708 p. p.; Hal. in z. b. G. 1888 p. 759; non Vahl. — *C. Sartorii* Heldr. fl. Aegina p. 305. — Exsicc.: Heldr. herb. norm. n. 430 et 1153.

Caule erecto, parce arachnoideo, patule corymboso; foliis coriaceis, parce pubescenti-viscidulis, elevatim nervosis, inferioribus in lacinias lineari-lanceolatas pinnatipartitis, superioribus semiamplexicaulibus,

spinoso-dentatis; capitulis oblongis; involucri phyllis externis foliis similibus, capitulo duplo longioribus, interioribus scariosis, in appendicem oblongam, pectinato-spinulosam, apice tenuiter spinosam dilatatis; acheniis turbinato-tetragonis; pappi paleis ab extimis brevissimis, ad intermedias acuminatas, serrulato-dentatas, achenio triplo longiores elongatis, seriei intimae eis aequilongis vel longioribus. — Praecedenti affinis, ab eo colore virenti, pubescentia parciori, foliis longioribus, angustioribus, inferioribus profunde pinnatipartitis, capitulis subminoribus, flosculis paucioribus, pallidioribus et pappi series intima elongata differt.

In locis aridis, arvis, campis regionis inferioris. Frequens in Attica pr. Athenas; Euboea: pr. Steni (Orph.); insula Aegina (Heldr.); Argolis: pr. Nauplia (Sart.); Laconia: pr. Sparta (Heldr.), ad radices mt. Taygetos (Psarid.); Creta: pr. Skylus distr. Temenos (Heldr.); et probabiliter alibi. — Jun. Aug. ⊙.

Obs. *C. tinctorius* L. sp. p. 830; Dallap. prosp. p. 111; Fraas fl. class. p. 206; Bois. fl. or. III. p. 709; Heldr. fl. cephal. p. 48. Hinc inde colitur.

60. Carduncellus Ad. fam. II. p. 116.

1. **C. coeruleus** L. sp. p. 830; S. et S. pr. II. p. 161, Fl. gr. t. 843; Sieb, avis p. 5, rem. p. 5; Ch. et B. exp. p. 244, Fl. pelop. p. 57; Bois. fl. or. III. p. 710; Heldr. fl. cephal. p. 48; (*Carthamus*); DC. pr. VI. p. 615; Weiss in z. b. G. 1869 p. 43; Raul. cret. p. 790; Hal. Beitr. fl. Aetol. p. 7; Bald. viagg. Creta p. 69. — Exsicc.: Reverch. pl. cret. n. 15.

Araneoso-puberulus; caule erecto, plerumque simplici; foliis subcoriaceis, elevatim nervosis, inferioribus lanceolatis, in petiolum attenuatis, superioribus sessilibus, e basi cordata oblongo-lanceolatis; capitulis ovatis; involucri phyllis externis foliaceis, breviter spinoso-dentatis, internis sublongioribus, scariosis, in appendicem pectinato-ciliatam, dilatatis; flosculis coeruleis; acheniis obscure tetragonis, pappo dimidio brevioribus.

α. **dentatus** DC. l. c.; Weiss l. c. — Foliis indivisis, spinoso-dentatis. — Exsicc.: Reverch. pl. cret. n. 15.

β. **tingitanus** L. sp. ed. 2, p. 1163 pro sp. *(Carthamus)*. — *Carduncellus coeruleus* v. *incisus* DC. l. c.; Marg. et R. l. c.; Weiss l. c. — Foliis pinnatifidis vel pinnatipartitis, spinoso-dentatis. — Exsicc.: Reverch. pl. cret. n. 84.

In herbidis, cultis regionis inferioris, rare. Corcyra (Bickn.); Cephalonia: pr. Lixuri (Heldr.); Zante (Marg.); Aetolia: pr. Antirrhion (Hal.); Messenia: pr. Methone (Chaub.); Creta: pr. Kissamos (Reverch.), Canea (Weiss), Akroteri (Bald.), Khalepa, Candia, Aludha (Raul.). — Maio, Jul. ♃.

61. Cnicus Vaill. act. acad. 1718 p. 163.

1. **C. benedictus** L. sp. p. 826; Heldr. Nutzpfl. p. 27, Fl. Aegina p. 306; Raul. cret. p. 790; Bois. fl. or. III. p. 705; Hal. Beitr. fl. Thessal. p. 14; Haussk. symb. p. 125. — *Centaurea benedicta* L. sp. ed. 2 p. 1296; S. et S. pr. II. p. 201, Fl. gr. t. 906; Ch. et B. exp. p. 255, Fl. pelop. p. 60; Friedr. Reise p. 274. — *Carduus benedictus* Camer. epit. 566 ic.; Fraas fl. class. p. 203. — Exsicc.: Sint. it. thessal. n. 401.

Villoso-lanuginosus; caule erecto, interdum brevissimo, divaricatim ramoso; foliis mollibus, oblongo-lanceolatis, sinuato-pinnatifidis partitisve, segmentis spinuloso dentatis, radicalibus in petiolum attenuatis, caulinis breviter decurrentibus, summis capitulum involucrantibus; involucri phyllis in spinam pinnatam abeuntibus; flosculis flavidis; acheniis cylindricis, pappum subaequantibus.

In arvis, cultis, herbidis regionis inferioris. Thessalia: pr. Kastania (Hal.), Kalabaka (Sint.); Attica: pr. Eleusis (Haussk.), Athenas, insula Salamis, Aegina (Heldr.); in Petalium insula Megalonisi (Holzm.); pr. Kalamaki in isthmo Corinthiaco (Haussk.); Peloponnesus: Argolis (Fraas), Poros (Friedr.), pr. Kalamata Messeniae (Zahn); Creta (Sibth.): pr. Akroteri (Raul.); et certe alibi. — Maio, Jul. ⊙.

II. Divisio. LIGULIFLORAE DC. pr. VII. p. 74.

12. Tribus SCOLYMEAE Less. syn. p. 126.

62. Scolymus L. gen. n. 922.

1. **S. hispanicus** L. sp. p. 813; S. et S. pr. II. p. 147, Fl. gr. t. 825; Ch. et B. exp. p. 239, Fl. pelop. p. 56; Marg. et R. fl. Zante p. 60; Friedr. Reise p. 263; Clem. sert. p. 57; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 793; Bois. fl. or. III. p. 713; Heldr. fl. cephal. p. 48, Fl. Aegina p. 307, chlor. Thera p. 16; Spreitz. in z. b. G. 1887 p. 663, 1890 p. 296; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Epir. p. 30, Beitr. fl. Thessal. p. 15, in ö. b. Z. 1899 p. 25; Haussk. symb. p. 142; Form. in Ver. Brünn 1896 p. 49. — *S. maculatus* Fraas fl. class. p. 201, non L. — Exsicc.: Orph. fl. gr. n. 763.

Crispule-pubescens; caule erecto, ramoso, ramis subracemosis; foliis oblongo-lanceolatis, sinuato-pinnatipartitis, spinosis, ala deorsum attenuata decurrentibus; capitulis axillaribus, solitariis vel 2—4 congestis, foliis arcte involucratis; involucri phyllis lineari-lanceolatis; receptaculi paleis apice coarctatis; ligulis flavis, glabris; antheris flavis; acheniis clavatis; pappi corona intus aristis binis aucta.

In incultis, ruderatis, derelictis per totam Graeciam. — Jun. Aug. ⊙.

2. **S. maculatus** L. sp. p. 813; S. et S. pr. II. p. 147, Fl. gr. t. 824; Sieb. avis rem. p. 5; Marg. et R. fl. Zante p. 60; Raul. cret. p. 794; Bois. fl. or. III. p. 713; Bald. viagg. Creta p. 69. —

Differt a praecedente praecipue foliis ala aequali lobata decurrentibus, floralibus pectinatis, capitulis terminalibus, solitariis vel 2—3 aggregatis, receptaculi paleis apice dilatatis, ligulis extus nigro-pilosis, antheris fuscis, acheniis obovatis, corona brevissima, denticulata, inermi superatis.

In ruderatis, incultis, rare. Creta (Sieb.): ad Almyros (Raul.), pr Kalyves et Kalamia (Bald.); in insula Zante et in Archipelago (Sibth.). — Jun. Aug. ⊙ N. v.

13. Tribus **CATANANCHEAE** Schultz in Bonpl. 1860 p. 367.

63. Catananche Vaill. act. Par. 1721 p. 215.

1. **C. lutea** L. sp. p. 812; Ch. et B. exp. p. 238, Fl. pelop. p. 55; Raul. cret. p. 795; Bois. fl. or. III. p. 714; Heldr. fl. Aegina p. 308; Hal. in ö. b. Z. 1899 p. 25. — Icon: Fl. gr. t. 821. — Exsicc.: Sart. pl. gr. a. 1835; Reis. fl. gr. a. 1898.

Villosula; caulibus erectis, parce ramosis; foliis lanceolatis, integris vel parce dentatis, radicalibus in petiolum attenuatis, caulinis paucis, sessilibus; capitulis solitariis, longe pedunculatis; involucri phyllis albo-hyalinis, ab infimis ovatis ad intima lanceolata elongatis; flosculis luteis, involucro brevioribus; pappi paleis 5—6, achenio 5 plo longioribus.

In collibus regionis inferioris rarissime Peloponnesus (Chaub.); insula Aegina (Sart.); in Strophadum insula majore (Reiser); Creta (Raul.). — Apr. Maio. ⊙.

64. Hymenonema Cass. bull. philom. 1817, dict. 22 p. 316.

1. **H. graecum** L. sp. p. 813; *(Catananche)*; DC. pr. VII. p. 116; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 797; Bois. fl. or. III. p. 714; Hal. in ö. b. Z. 1895 p. 460; Heldr. in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 247. — *H. Tournefortii* Cass. dict. 22 p. 316. — *Scorzonera elongata* Willd. sp. III. p. 1508; S. et S. pr. II. p. 124; Urv. enum. p. 99. — Icon: Tourn. voy. p. 86; Fl. gr. t. 789. — Exsicc.: Heldr. herb. norm. n. 1055 et 1660; Dörfl. fl. aeg. n. 56 et 122.

Asperum, inferne plus minus villosum; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis pinnatisectis, sublyratis, segmentis oblongis, denticulatis, rhachide lobulata, radicalibus in petiolum attenuatis, caulinis paucis, diminutis, sessilibus; capitulis ovatis, longe pedunculatis; involucri phyllis coriaceis, albo-marginatis, ciliatulis, ab externis ovatis obtusis, ad interna oblonga acutiuscula elongatis; ligulis luteis, concoloribus; receptaculi paleis fructiferis auctis, deciduis, seriei exterioris ovato-lanceolatis, facie interna, basi plicata achenia radii in-

cludentibus, seriei interioris spongioso-incrassatis, rugosis; acheniis radii cylindricis, glabris, palea inclusis, pappi paleolis 5—7, trifidis, laciniis lateralibus brevibus, intermedia in setam longam abeunti, superatis; acheniis disci turbinatis, villosis, liberis, pappi paleolis 10—12, indivisis, aequilongis, achenio multo longioribus, setaceo-attenuatis, fulvis, scabris, superatis.

In montosis, collibus, ad vias. Cyclades: insula Tenos, Keos (Heldr.), Kythnos, Syra, Mykonos, Siphnos (Sart), Naxos (Tourn.), Melos, Amorgos (Sibth), Anaphi (Leon.); Creta: pr. Akrotori, Malaxa (Raul.). — Maio, Jul. ♃.

2. **H. laconicum** Bois. et Heldr. fl. or. III. p. 715; Rouy ill. p. 29 t. 90 *v. minus* (f. caulibus monocephalis, foliis radicalibus brevioribus, saepe ovatis, basi sublobatis, lobo terminali semper lateralibus multo majore). — *Catananche graeca* Ch. et B. exp. p. 238, Fl. pelop. p. 55, non L. — Exsicc.: Orph. fl. gr. n. 1152; Pichl. pl. gr. a. 1876; Heldr. herb. norm. n. 1355.

Differt a praecedente, cui habitu maxime affinis, indumento copiosiore, ligulis basi purpureis et pappi paleolis acheniorum disci 15—20, inaequalibus, externis 4—5 brevioribus setiformibus, ceteris setaceo-attenuatis, scabris, superne subplumosis, purpureis.

In collibus apricis, herbidis regionis inferioris. Arcadia: mt. Maenalus (Sart.); Laconia: mt. Malevo pr. Platanos (Orph.), mt. Taygetos pr. Androvista, Anavryti, Mistra, Palaeochori (Heldr.), Golass (Pichl.), mt. Selitza (Zahn), pr. Kalamata (Chaub.). — Maio, Jul. ♃.

14. Tribus. **HYPOCHAERIDEAE** Less. syn. p. 130.

65. **Hypochaeris** L. gen. n. 918.

a. Pappus conformis, biserialis, radiis externis brevioribus scabris, internis plumosis.

1. **H. glabra** L. sp. p. 811; Weiss in z. b. G. 1869 p. 46; Bois. fl. or. III. p. 783. — Icon: Fl. dan. t. 424. — *H. dimorpha* Brot. fl. lusit. I. p. 332. — Exsicc.: Heldr. pl. fl. hellen. a. 1887 et 1900.

Radice annua, tenui; caulibus erectis vel adscendentibus, glabris, simplicibus vel furcato-ramosis, aphyllis, squamatis; foliis radicalibus, rosulatis, glabris vel setulosis, oblongo-spathulatis, obtuse sinuato-dentatis; pedunculis apice subincrassatis; involucri phyllis glabris; flosculis luteis, marginalibus involucrum aequantibus; acheniis muricatis, radii erostribus, disci rostratis.

β. **minima** Cyr. pl. rar. neap. fasc. I. p. 10; S. et S. or. II. p. 143, Fl. gr. t. 816; pro sp.; DC. pr. VII. p. 90; Heldr. fl. Aegina p. 307. — Caulibus pumilis, simplicibus; acheniis saepe omnibus erostribus. — Exsicc.: Heldr. pl. fl. hell. a. 1879.

In arvis, collibus aridis, campis regionis inferioris. Attica: in regione Laurii pr. Spilazeza (Heldr.); Cycladum insula Cythnos (Tunt.), Tenos (Sart.), Syra (Weiss); — *β.* in Peloponneso (Sibth.); insula Aegina, mt. Pentelicon Atticae (Heldr.). — Apr. Jun. ⊙.

2. **H. radicata** L. sp. p. 811; S. et S. pr. II. p. 143; Ch. et B. exp. p. 237, Fl. pelop p. 55; Marg. et R. fl. Zante p. 61; Weiss in z. b. G. 1869 p. 742; Heldr. fl. cephal. p. 49; Form. in D. bot. Monat. 1890 p. 15; Bald. riv. coll. bot. Alb. 1896 p. 73. — Icon: Fl. dan. t. 150. — Exsicc.: N. v.

Rhizomate perenni, ramoso, crasso; caulibus erectis, glabris vel inferne parce setulosis, furcato-ramosis, raro simplicibus, aphyllis squamatis; foliis radicalibus, rosulatis, setulosis, oblongo-lanceolatis, dentatis vel runcinato-pinnatifidis; pedunculis apice subincrassatis; involucri phyllis glabris vel parce setulosis; flosculis luteis, marginalibus involucro longioribus; acheniis striatis, rugulosis, rostratis.

β. **neapolitana** DC. pr. VII. p. 91; Raul. cret. p. 796; Hal Beitr. fl. Epir. p. 30; Heldr. fl. Aegina p. 307; pro sp. — *H. dimorpha* Ten. syn. p. 407; Sieb. avis p. 5; non Brot. — *H. radicata v. heterocarpa* Mor. fl. sard. II. p. 487; Bois. fl. or. III. p. 784; Haussk. symb. p. 142; Form. in Ver. Brünn 1896 p. 42, 1897 p. 31. — Achenia radii erostria vel brevissime rostrata, disci longe rostrata. — Exsicc.: Sint. et Bornm. it. turc. n. 1340; Dörfl. fl. aeg. n. 80; Heldr. herb. norm. n. 1659.

In herbidis regionis inferioris et montanae. Epirus: pr. Janina (Bald.); Thessalia: pr. Malakasi, Velestinos, mt. Pelion (Form.); Peloponnesus (Sibth.); Cycladum insula Tenos (Weiss); Cephalonia: inter Mavrata et Foramine (Heldr.); Zante (Marg.); — *β.* Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.); Thessalia: pr. Konisko, mt. Cuka et Mitrica in mt. Chassia (Form.), pr. Litochori in Olympo (Sint.), mt. Pelion (Haussk.); Euboea: mt. Telethrion pr. Kastaniotissa (Heldr.); Attica: mt. Parnes pr. Dekeleia (Hal.), mt. Pentelicon, pr. Geraka (Haussk.), ad Phalerum (Sprun.), insula Aegina (Heldr.); Laconia: mt. Malevo supra Castanitza (Orph.); Cycladum insula Keos (Heldr.), Naxos (Leon.); Creta: pr. Nerokuru (Raul.). — Maio, Sept. ♃.

b. Pappus uniserialis, biformis, acheniorum radii coroniformis, disci plumosus.

3. **H. cretensis** L. sp. ed. 2 p. 1139; Sieb. avis rem. p. 5; (*Seriola*); Ch. et B. exp. p. 237, Fl. pelop. p. 55; Bois. fl. or. III. p. 784; Hal. Beitr. fl. Aetol. p. 8, Beitr. fl. Achaia p. 26, in ö. b. Z. 1895 p. 460; Bald. riv. coll. bot. Alb. 1896 p. 73; Heldr. fl. Aegina p. 307. — *Achyrophorus serioloides* DC. pr. VII. p. 93. — *Metabasis hymettia* DC. l. c. p. 97. — *M. cretensis* DC. l. c. p. 307; Marg. et R. fl. Zante p. 61; Raul. cret. p. 796; Gelmi giorn. bot. ital. 1889 p. 450; Bald. giorn bot. ital. 1894 p. 98; Haussk. symb. p. 142. — *Achyrophorus cretensis* Bois. ap. Weiss in z. b. G. 1869

p. 742. — *Seriola urens* S. et S. pr. II. p. 143; Ch. et B. fl. pelop. p. 55; Friedr. Reise p. 269, 270 et 276; vix L. sp. ed. 2 p. 1139, cujus species e Sicilia dubia videtur. — *Fabera hispida* Schultz in nov. act. Leop. Carol. 1845 p. 131. — *F. cretensis* Schultz l. c. p. 137. — *H. pinnatifida* Hal. Beitr. fl. Epir. p. 30, non Cyr. — Exsicc.: Heldr. herb. norm. n. 855; Sint. it. thessal. n. 248.

Perennis; caulibus erectis, glabris vel inferne parce setulosis, furcato-ramosis, foliatis; foliis parce setulosis, radicalibus rosulatis obovato-oblongis, dentatis vel saepius in lacinias triangulari-lanceolatas pinnati-partito-lyratis, caulinis inferioribus sursum minoribus, superioribus lanceolato-linearibus, integris; pedunculis apice subincrassatis; involucri phyllis tomentellis et saepe uniserialiter setosis; flosculis luteis, marginalibus involucro longioribus, fructiferis dorso induratis; acheniis rugulosis, radii apice attenuatis, disci longe rostratis. — Species forma foliorum et indumento summopere variabilis.

In collibus, herbidis, vineis, ad vias regionis inferioris et montanae. Epirus: ad Skafidaki pr. Prevesa, pr. Muspina distr. Janina (Bald.), pr. Vulgarelion ad radices mt. Tsumerka (Hal.); Thessalia: pr. Kalabaka (Sint.), Malakasi, ad monasterium Korona (Haussk.); Aetolia: pr. Karpenisi (Fraas), Aetolikon (Reis.), Antirrhion (Hal.); Boeotia: pr. Lebadea (Fraas); Attica: in valle Cephissi pr. Athenas (Orph.), Kukuvanes, pr. Marathon, mt. Parnes, Pentelicon, pr. Liopesi ad radices Hymetti, pr. Spata, Phaleron (Heldr.), Turcovuni, Laurion, Corinthus (Haussk.); Achaia (Sibth.); Elis: mt. Olenos (Hal.), pr. Kunupeli, Olympia (Heldr.); Messenia: a Kyparissia usque Gargaliano, pr. Halmyron, Kytriais (Chaub.); Argolis (Sprun.): pr. Nauplia (Borga), insula Poros; Aegina (Friedr.); Cycladum insula Keos, Naxos (Heldr.), Tenos (Weiss), Syra (Sart.); Creta: pr. Nerokuru, Khalepa, Aludha, Kritza, insula Dia (Raul.); Zante (Marg.); Corcyra (Gelmi). — Apr. Jul. ♃.

c. Pappus conformis, uniserialis, plumosus.

4. **H. tenuiflora** Bois. diagn. XI. p. 35; Raul. cret. p. 796 (*Robertia*); Bois. fl. or. III. p. 785. — Huc forsitan: *H. acaulis* Sieb. avis rem. p. 5. —

Perennis, nana; caulibus tenuissimis, monocephalis, squamatis, folia vix duplo superantibus; foliis radicalibus, rosulatis, parce setulosis, runcinato-pinnatifidis, lobo terminali longiore; capitulis parvis; involucri phyllis paucis, lineari-lanceolatis, albo-puberulis, basi parce crinitis; ligulis concoloribus; acheniis tenuiter rugulosis, exterioribus apice subattenuatis, internis breviter attenuato-rostratis. — Planta bipollicaris, ex unico specimine descripta, habitu *Robertiae taraxacoidis*, ab ea foliis minus profunde divisis, capitulis minoribus et insuper involucri phyllis imbricatis nec uniserialibus differre dicitur.

In regione subalpina superiore mt. Hellinoseli distr. Sphakia Cretae (Raul.). — Jun. ♃. N. v.

66. Seriola L. gen. n. 927.

1. **S. aethnensis** L. sp. ed. 2 p. 1139; S. et S. pr. II. p. 143, Fl. gr. t. 815; Sieb. avis p. 5; Ch. et B. fl. pelop. p. 55; Link in Linn. IX. p. 582; Friedr. Reise p. 277; Weiss in z. b. G. 1869 p. 46; Bois. fl. or. III. p. 785; Heldr. fl. cephal. p: 49, Fl. Aegina p. 307; Haussk. symb. p. 142; Hal. in ö. b. Z. 1897 p. 96; Form. in D. bot. Monat. 1898 p. 78. — *Metabasis aethnensis* DC. pr. VII. p. 307; Raul. cret. p. 796. — Exsicc.: Sint. it. thessal. n. 1471.

Caulibus erectis vel adscendentibus, furcato-ramosis, rarius simplicibus, glabris vel saepius saltem superne setoso-hispidis; foliis fere omnibus radicalibus, rosulatis, oblongis vel obovatis, in petiolum brevem attenuatis, repandis vel dentatis, hirtis; involucri phyllis setoso-hispidis; ligulis flavis; acheniis fusiformibus, rugosis, longe rostratis.

In herbidis, arenosis, glareosis regionis inferioris per totam ut videtur Graeciam. Indicatur: Thessalia: pr. Kalabaka (Sint.), Pharsalus (Haussk.); Aetolia: pr. Aetolikon (Reis.), Antirrhion (Hal.); Attica: mt. Lycabettus, Hymettus, pr. Eleusis, ad Phalerum, Rhaphti, insula Salamis, Aegina (Heldr.); Argolis: pr. Nauplia (Haussk.), Poros (Friedr.), insula Hydra (Heldr.); Achaia: pr. Patras (Link); pr. Messene (Chaub.); Sporadum insula Skiathos (Leon.); Cyclades: Cythnos (Tunt.), Syra, Tenos (Weiss), Melos, Kimolos (Leon.); Creta: pr. Canea, Akroteri, Malaxa, Askyphos, Anopolis, Istovai in promontorio Sidero (Raul.); Cephalonia: pr. Argostoli, Lixuri (Heldr.); Corcyra (Haussk.). — Mart. Jun. ⊙.

15. Tribus. RODIGIEAE DC. pr. VII. p. 98.

67. Rodigia Spreng. neue Entd. I. p. 273.

1. **R. commutata** Spreng. l. c.; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 796; Bois. fl. or. III. p. 880; Haussk. symb. p. 142. — *Seriola commutata* Less. syn. p. 131. — Exsicc.: Rev. pl. cret. n. 98; Heldr. herb. norm. n. 1054; Dörfl. pl. cret. n. 11.

Parce setulosa; caulibus adscendentibus, furcato-ramosis; foliis inferioribus oblongis vel obovatis, in petiolum attenuatis, dentatis vel lyrato-pinnatipartitis; involucri phyllis setoso-hispidis, externis dimidio brevioribus, demum patulo-incurvis, internis demum induratis, subcurvatis; ligulis flavis; acheniis fusiformibus, subrugosis, interioribus in rostrum eis 4 plo longius attenuatis, externis brevius rostratis.

In herbidis, olivetis, ad vias, in cultis regionis inferioris. Thessalia: ad monasterium Korona (Haussk.); Attica: pr. Palaeokundura, Eleusis, Geraka ad radices mt. Pentelicon (Heldr.), pr. Phalerum, Laurion (Haussk.); Euboea (Pichl.), Petalium insula Xeronisi (Holzm.); Argolis: in peninsula Methana (Heldr.), pr. Nauplia (Sart.), Poros (Reinh.); Cycladum insula Keos, Kythnos (Tunt.), Syra, Tenos (Weiss), Amorgos (Heldr.); Creta: pr. Kissamos (Rev.), Canea, Nerokuru (Raul.), pr.

Males distr. Hierapetra (Leon.); Zante: in colle Hagios Joannes (Mazz.). Apr. Jun. ⊙.

68. Lagoseris M. B. fl. taur. cauc. III. p. 538.

1. **L. bifida** Vis. stirp. dalm. p. 19 t. 7 (*Trichocrepis*); Koch. syn. p. 435; Bois. fl. or. III. p. 881; Gelmi in bull. soc. bot. ital. 1869 p. 450; Hal. Beitr. fl. Achaia p. 25; Haussk. symb. p. 138. — *Crepis nemausensis* Sieb. avis p. 5; Ch. et B. exp. p. 234, Fl. pelop. p. 54, non Gou. — *Pterotheca bifida* Fisch. et Mey. ind. hort. petrop. 1837 n. 1736. — *P. nemausensis* Rchb. fl. exc. p. 269, Raul. cret. p. 800; non Cass. — Exsicc.: Heldr. herb. norm. n. 1263.

Hispidula; caulibus erectis vel adscendentibus, nudis, superne ramosis; foliis radicalibus, rosulatis, oblongis, in petiolum attenuatis, dentatis vel sublyrato-pinnatifidis; involucri phyllis externis calyculiformibus, internis 3—4 plo longioribus, glanduloso-pilosis vel glabratis; ligulis flavis; acheniis apice attenuatis, marginalibus laevibus, oblongis, corticosis, dorso convexis, ventre obtuse carinatis, bisulcatis, interioribus cylindricis, striatis, aliis magis externis scabridis, ceteris laevibus. — *L. nemausensi* Koch affinis, quae differt acheniis marginalibus ventre 3—5 alatis.

In collibus, saxosis, incultis regionis inferioris et montanae. Thessalia: mt. Ghavellu, ad monasterium Korona (Haussk.); Boeotia: pr. Thebas (Heldr.); Attica: pr. Athenas, mt. Lycabettus, Turcovuni, Parnes, pr. Palaeokundura ad mt. Pateras (Heldr.); Achaia: mt. Panachaicon (Hal.); Arcadia: pr. Tripolis, Alonistena, Nemnitza (Heldr.), Phigalea (Chaub.); Laconia: ad radices mt. Taygetos (Zahn); Creta (Raul.). — Mart. Jun. ⊙.

16. Tribus. HYOSERIDEAE Less. syn. p. 127.

69. Cichorium L.

1. Sectio. *Eucichorium* DC. pr. VII. p. 84. — Capitula multiflora; rami inermes.

1. **C. intybus** L. sp. p. 813; S. et S. pr. II. p. 146; Pieri corc. fl. p. 115; Dallap. prosp. p. 108; Ch. et B. exp. p. 239, Fl. pelop. p. 56; Fraas fl. class. p. 197; Raul. cret. p. 795; Bois. fl. or. III. p. 715; Heldr. fl. cephal. p. 48; Hal. in z. b. G. 1888 p. 759, Beitr. Fl. Epir. p. 30, Beitr. fl. thessal. p. 15; Form. in D. bot. Mon. 1890 p. 14, in Ver. Brünn 1895 p. 25, 1896 p. 39, 1897 p. 30; Haussk. symb. p. 142. — Icon: Fl. dan. t. 907. — Exsicc.: Sint. et Bornm. it. turc. n. 1341.

Perenne, laete virens; caule erecto, aspero, divaricatim ramoso; foliis radicalibus rosulatis, runcinatis, caulinis paucis, lanceolatis, semiamplexicaulibus; capitulis lateralibus sessilibus, geminis ternisve, termi-

nalibus solitariis, in apice ramulorum fistulosorum subincrassatorum involucri phyllis externis ovato-lanceolatis, acutis, internis linearibus, duplo longioribus, omnibus glanduloso-ciliatis; acheniis foveolatis, transverse striolatis, pappo paliaceo 8—10 plo longioribus.

β. **glabratum** Presl fl. sic. I. p. 32; Haussk. symb. p. 142; pro sp.; Gr. et Godr. fl. fr. II. p. 286. — Subglabrum; capitulis geminis, alio sessili, alio longe pedunculato; involucri phyllis glabris vel glanduloso-ciliatis; flosculis dimidio minoribus. — Exsicc.: Heldr. it IV. Thessal. a. 1885.

In arvis, ruderatis, ad vias regionis inferioris et montanae ut videtur per totam Graeciam; — β. Thessalia: pr. Pharsalum (Heldr.), Volo (Haussk.); Cycladum insula Rhenea (Tunt.) et certe alibi. — Maio, Sept. ♃.

2. **C. pumilum** Jacq. obs. bot. IV. p. 3 t. 80; Ung. Reise p. 125; Haussk. symb. p. 142. — *C. divaricatum* Schousb. Beob. Gewächsr. Marokko p. 184; Ch. et B. fl. pelop. p. 56; Bois. fl. or. III. p. 716; Heldr. Nutzpfl. p. 28, Fl. cephal. p. 48, Fl. Aegina p. 308, chlor. Mykon. p. 247. — *C. minimum* Port. en. pl. Dalm. p. 16, t. 12. — *C. endivia v. pumilum* Vis. fl. dalm. II. p. 97; Weiss in z. b. G. 1869 p. 46. — *C. intybus v. divaricatum* DC. pr. VII. p. 84; Marg. et R. fl. Zante p. 61; Raul. cret. p. 795. —

Differt a praecedente radice annua; foliis glauco-virentibus, caulinis ovato-lanceolatis ovatisve; capitulis terminalibus in apice ramulorum fistulosorum valde incrassatorum, subtubaeformium insidentibus; involucri phyllis externis ovatis, obtusis, longe ciliatis, eglandulosis, interiora subaequantibus; acheniis laevibus, pappo 3 plo longioribus.

Ad vias, in herbosis, arvis regionis inferioris. Indicatur: Attica: pr. Dramesi (Ung.), Kamariza Laurii (Haussk.), insula Aegina (Heldr.); Cycladum insula Syra (Weiss), Rhenea (Heldr.); Messenia: pr. Kalamata (Gittard.); Creta: pr. Canea (Weiss), Guves (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli, Lixuri (Heldr.); sed loca forsan plurima ad var. β. speciei praecedentis spectant. — Maio, Sept. ⊙. N. v.

Obs. *C. endivia* L. sp. p. 813; S. et S. pr. II. p. 146; Dall. prosp. p. 109; Ch. et B. exp. p. 239, Fl. pelop. p. 56; Marg. et R. fl. Zante p. 85; Fraas fl. class. p. 197; Heldr. Nutzpfl. p. 28. — Differt a praecedente foliis caulinis basi hastatis, involucri phyllis externis angustioribus glanduloso-ciliatis et pappo achenio quater breviori. — Colitur in hortis ad usum culinarem.

2. Sectio. *Acanthophyton* Less. syn. p. 128 pro gen.; DC. pr. VII. p. 84. — Capitula 5—6 flora; rami spinescentes.

3. **C. spinosum** L. sp. p. 813; S. et S. pr. II. p. 146, Fl. gr. t. 823; Sieb. avis p. 5, rem. p. 5; Ch. et B. exp. p. 239, Fl. pelop. p. 56; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 795; Bois. fl. or. III. p. 717; Heldr. fl. cephal. p. 48, Fl. Aegina p. 308, in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 247; Haussk. symb. p. 142; Bald.

viagg. Creta p. 71. — Exsicc.: Heldr. pl. gr. n. 1474, herb. norm. n. 1661; Rev. pl. cret. n. 93, in Baen. herb. europ. n. 4860; Dörfl. fl. aeg. n. 50.

Perenne, glaucovirens; caulibus nanis, a basi intricatim ramosissimis, ramis demum apice spinescentibus; foliis radicalibus rosulatis, sinuato-dentatis vel pinnatifido-lyratis, caulinis linearibus, saepius integris; capitulis in dichotomiis sessilibus, solitariis, rarius 2—3 aggregatis, terminalibus paucis, in apice ramulorum non incrassatorum insidentibus; involucri glabriusculi phyllis externis ovatis, obtusis, internis linearibus, triplo longioribus; pappo setoso-subpaleaceo, achenio 8—10 plo breviore.

In saxosis et arenosis maritimis. Attica: pr. Eleusis, Megara (Sprun.), ad Phalerum, pr. Laurion (Haussk.); in isthmo Corinthiaco pr. Kalamaki, Corinthus; insula Metopi pr. Aegina (Heldr.); Cycladum insula Syra (Weiss), Naxos (Leon.), Tenos, Mykonos (Heldr.); Laconia: a Marathonisi usque promontorium Tenare (Chaub.); Creta: pr. Canea (Tourn. voy. p. 10), Kissamos (Rev.), Khalepa, Suda, Askyphos, insula Gaidaronisi (Raul.); Cythaera: pr. Kapsali, San Nicolo (Spreitz.); Cephalonia: pr. Drapano (Heldr.). — Jun. Aug. ♃.

2 × 3. **C. pumilum × spinosum.** — *C. hybridum* Heldr. herb. gr. norm. n. 1662. — Habitu inter parentes intermedia, a *C. pumilo* caulibus intricatim ramosissimis, ramulis demum non incrassatis, capitulis paucifloris; a *C. spinoso* radice annua, caulibus elatioribus, ramulis vix spinescentibus differt. Achenia abortiva. — In oropedio Anoraera insulae Mykonos Cycladum, inter parentes (Heldr.). —

70. Hyoseris L. gen. n. 916.

1. **H. radiata** L. sp. p. 808; Weiss in z. b. G. 1869 p 46. — Huc probabiliter: *H. lucida* Sieb. avis p. 5; Ch. et B. exp. p. 237, Fl. pelop. p. 55; vix L. mant. p. 103, quae calyculo biseriali, involucro fructifero erecto, nec stellatim expanso et foliorum segmentis imbricatis integris obtusioribus a *H. radiata* distinguitur. — Icon: Rchb. t. 9, f. 1. — Exsicc.: Dörfl. fl. aeg. n. 19.

Perennis; radice crassa verticali; foliis radicalibus, rosulatis, glabris vel ciliatis, saepe pulverulentis, in segmenta rhomboidea, dentata vel incisa runcinatim pinnatipartitis; scapis monocephalis, folia superantibus, fructiferis vix incrassatis; calyculi squamis involucro triplo brevioribus; involucri phyllis oblongo-linearibus, fructiferis stellatim expansis; acheniis marginalibus compressis, exalatis, oblongo-linearibus, striatis, pappo paleolato brevissimo, ceteris oblongo-linearibus, late alatis, costatis, margine scabris, pappo radiis numerosis, aliis setiformibus, aliis basi paleaceo-dilatatis constante superatis, intimis anguste linearibus, exalatis. — Planta graeca, saltem illa e Naxos, foliis crassiusculis, glaucovirentibus, segmentis parvis, dentatis, involucri phyllis latioribus et pappi setis basi parum paleaceo-dilatatis, magis piliformibus, ab occidentali discedit et subspeciem propriam — *graecam* — sistere videtur.

In arenosis maritimis, cultis regionis inferioris. Cycladum insula Naxos: pr. Haliki (Leon.); Creta: pr. Canea (Weiss); ? Peloponnesus: pr. Pylos, Methone, Kalamata, Mavrovuni in Maina (Chaub.), si *H. lucida* Ch. et B. huc spectet. — Mart. Maio. ♃.

2. **H. scabra** L. sp. p. 809; Sieb. avis rem. p. 5; Weiss in z. b. G. 1869 p. 47; Bois. fl. or. III. p. 718, suppl. p. 318; Spreitz. in z. b. G. 1877 p. 714; Heldr. fl. cephal. p. 48, Fl. Aegina p. 308; Haussk. symb. p. 143; Hal. in ö. b. Z. 1895 p. 461. — *H. microcephala* Cass. dict. 22 p. 338; Marg. et R. fl. Zante p. 61; Clem. sert. p. 51; Raul. cret. p. 794. — Icon: Rchb. t. 9, f. 9. — Exsicc.: Dörfl. fl. gr. n. 457.

Annua; radice tenui, filiformi; foliis radicalibus, rosulatis, scabriusculis, in segmenta ovata, dentata runcinatim pinnatipartitis; scapis monocephalis, foliis aequilongis vel vix longioribus, clavato-incrassatis; calyculi squamis involucro 5 plo brevioribus; involucri phyllis lanceolatis, fructiferis erecto-conniventibus; acheniis marginalibus compressis, linearibus, exalatis, pappo paleolato brevissimo, ceteris oblongo-lanceolatis, late alatis, costatis, margine scabris, pappo radiis aliis setiformibus, aliis 5 paleaceo-dilatatis constante superatis, intimis linearibus, exalatis.

In cultis, arenosis maritimis, ad vias regionis inferioris. Aetolia: pr. Krioneri ad sinum Patranum (Reiser); Attica: pr. Piraeum (Clem.), Eleusis, Athenas, mt. Lycabettus, pr. Patissia, Kephissia, Phaleron, promontorium Colias, insula Salamis, Aegina (Heldr.); Corinthus: pr. Asso (Heldr.), Acrocorinthus (Haussk.); Peloponnesus (Bois.); Cycladum insula Keos, Kythnos (Tunt.), Syra, Tenos (Weiss), Melos (Leon.); Creta: pr. Sitia (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli (Spreitz.). — Febr. Maio. ⊙.

71. Hedypnois Tourn. inst. p. 478.

1. **H. rhagadioloides** L. sp. p. 809; Ch. et B. exp. p. 257, Fl. pelop. p. 55; (*Hyoseris*); Willd. sp. III. p. 1617; S. et S. pr. II. p. 142, Fl. gr. t. 812. — *H. tubaeformis* Ten. fl. nap. II. p. 173, t. 73; Clem. sert. p. 51, Raul. cret. p. 794; Heldr. fl. Aegina p. 308, chlor. Thera p. 16, chlor. Mykon. p. 247. — *H. cretica* Bois. fl. or. III. p. 719 p. p.; ? Haussk. symb. p. 143. — Exsicc.: Heldr. herb. fl. hellen. 1878.

Setuloso-scabra; caulibus erectis vel diffusis, parce ramosis vel simplicibus; foliis radicalibus rosulatis, obovato-oblongis, sinuato-dentatis vel pinnatifidis, caulinis minoribus, summis lanceolatis, integris; pedunculis fistulosis, fructiferis in tubam insignem ampliatis (diam. 4—6 mm.); capitulis majusculis, globosis; involucri phyllis setoso-hispidis, arcte inter se adpressis; acheniis scabridis, interioribus cum pappo subaequilongo involucrum parum superantibus.

In herbidis, sterilibus aridis, regionis inferioris. Indicatur in Aetolia: pr. Mesolongion (Nied.); Attica: insula Salamis, Pharmacusarum insula Leros, pr. Marathon, Tatoi, Piraeus, Phaleron, Eleusis, mt. Lycabettus,

pr. Athenas, insula Aegina (Heldr.); Cycladum insula Keos, Kythnos, Rhenea (Tunt.), Thera (Heldr.); Messenia: pr. Methone (Chaub.); Creta: ad Kalyves (Raul); et certe alibi. — Mart. Jul. ⊙.

2. **H. cretica** L. sp. p. 810; Urv. enum. p. 100 (*Hyoseris*); Willd. sp. III. p. 1617; S. et S. pr. II. p. 142, Fl. gr. t. 813; Marg. et R. fl. Zante p. 61; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 794; Spreitz in z. b. G. 1877 p. 713, 1890 p. 296; Heldr. fl. cephal. p. 48, chlor. Mykon. p. 247; Gelmi in bull. soc. bot. ital. 1889 p. 450. — *H. pendula* Willd. sp. III. p. 1618; Marg. et R. l. c.; Weiss l. c.; Raul. l. c. — *H. polymorpha* DC. pr. VII. p. 81; Marg. et R. l. c.; Raul. l. c. — *H. cretica v. gracilior* Bois. fl. or. III. p. 719; Haussk. symb. p. 143. — Exsicc.: Heldr. herb. fl. hellen. a. 1878, 1889, 1897.

Differt a praecedente pedunculis fructiferis siquidem fistulosis ac incrassatis, sed minime clavaeformibus, capitulis minoribus, involucri phyllis fructiferis inter se remotiusculis, demum subpatentibus.

β. **monspeliensis** Willd. sp. III. p. 1616; S. et S. pr. II. p. 142; Bald. cret. p. 794; pro sp. — *Hyoseris hedypnois* L. sp. p. 809. — Pedunculis et involucri phyllis glabris. — Intermediis ad typum transit. — Exsicc.: Heldr. herb. fl. hellen. 1878; Orph. herb. n. 618.

In arenosis, herbidis, incultis regionis inferioris per totam Graeciam. — Mart. Jul. ⊙.

17. Tribus. **LAPSANEAE** Less. syn. p. 126.

72. **Tolpis** Gaertn. fr. II. p. 371.

1. **T. umbellata** Bert. rar. lig. pl. dec. I. p. 13; Weiss in z. b. G. 1869 p. 46; Bois. fl. or. III. p. 725; Hal. Beitr. fl. Aetol. p. 8; Haussk. symb. p. 142; Boissieu in bull. soc. bot. Fr. 1896 p. 286; Heldr. fl. Aegina p. 308, chlor. Mykon. p. 247. — *T. quadriaristata* Biv. monogr. p. 9, t. 1. — *T. barbata* S. et S. pr. II. p. 140; Friedr. Reise p. 285; Raul. cret. p. 735; non L. sp. p. 805 *(Crepis)*, Gaertn. fr. II. p. 372, quae differt capitulis duplo majoribus, ligulis involucro duplo longioribus, centralibus purpurascentibus, calyculi phyllis eximie squarroso-patentibus et pappo subbiseto. — Icon: Rchb. t. 8. — Exsicc.: Heldr. herb. norm. n. 191 et 953; Dörfl. fl. aeg. n. 115.

Scabrido-puberula; caule erecto, patule ramoso, rarius simplici; foliis radicalibus rosulatis, oblongo-spathulatis, integris, dentatis vel pinnatifidis, caulinis paucis, linearibus; capitulis solitariis, in dichotomiis pedunculo brevi, tandem fistuloso, incrassato suffultis, lateralibus longe tenuiterque pedunculatis; calyculi phyllis setaceis, subarcuatis, involucri phylla lanceolata superantibus; ligulis flavis, involucrum vix superantibus; acheniis striatis, externis corona setulis brevissimis constante, ceteris 4—5 setis eis triplo longioribus superatis.

In collibus siccis, incultis, arenosis regionis inferioris et submontanae. Thessalia: ad monasterium Korona, Meteora, pr. Kalabaka

(Haussk.), Palaeokastron pr. Karditza, mt. Pelion pr. Milies (Heldr.); Euboea: mt. Dirphys (Heldr.), Petalium insula Megalonisi (Holzm.); Aetolia: ad radices mt. Taphiassos (Hal.); Attica: pr. Marathon, mt. Pentelicon, pr. Raphina, Laurion, insula Aegina (Heldr.); Argolis: pr. Nauplia (Friedr.), Vromolimni peninsulae Methana (Heldr.), Poros (Reinh.); mt. Chelmos Achaiae, mt. Malevo (Orph.), mt. Taygetos pr. Golas (Sart.); Cycladum insula: Andros (Sart.), Tenos (Weiss), Keos (Boissieu), Kythnos, Mykonos, Seriphos (Tunt.), Naxos, Kimolos (Leon.); Creta (Sibth.): pr. Canea (Weiss). — Mart. Jul. ⊙.

2. **T. virgata** Desf. fl. atl. II. p. 230 (*Crepis*); Bert. rar. lig. pl. dec. I. p. 15; Weiss in z. b. G. 1869 p. 46; Haussk. symb. p. 143; Hal. in ö. b. Z. 1895 p. 460. — *Crepis virgata* Desf. fl. atl. II. p. 230; Ch. et B. exp. p. 236, Fl. pelop. p. 55. — *Crepis altissima* Balb. cat. pl. hort. Taur. 1804 p. 15. — *T. altissima* Pers. ench. II. p. 377; Marg. et R. fl. Zante p. 61; Raul. cret. p. 795; Bois. fl. or. III. p. 725. — *T. quadriaristata* S. et S. fl. gr. IX. p. 6 t. 810; non Biv. — Huc forsan: *T. cretica* Sieb. avis p. 5, rem. p. 5. — Exsicc.: Orph. fl. gr. n. 758; Heldr. it. thessal. n. 42; Reverch. pl. cret. n. 94.

Differt a praecedente radice bienni; caulibus elatis, strictiuscule ramosis; pedunculis dichotomiarum vix incrassatis; calyculi phyllis linearibus, erectiusculis, involucro multo brevioribus; ligulis involucro quarta parte longioribus; acheniis omnibus 6—12 setis, rarius 4—6 setis terminatis.

In iisdem locis, ubi praecedens. Thessalia: ad monasterium Korona (Haussk.), Palaeokastron pr. Karditza, mt. Pelion pr. Portaria; pr. Kastaniotissa Euboeae (Heldr.); Achaia: pr. Patras (Reinh.); ad lacum Pheneon Arcadiae (Heldr.); Messenia: pr. Kube, Phigalea (Chaub.); Laconia: mt. Malevo pr. Kastanitza (Orph.); Cycladum insula Andros, Tenos, Mykonos (Sart.), Keos, Kythnos (Tunt.), Naxos (Reiser); Creta: pr. Canea (Weiss), inter Chersonesos et Malia, pr. Suda, Akroteri, Aradhena, Lutro, Embaro in mt. Lassiti (Raul.); Zante (Marg.). — Maio, Aug. ⊙.

73. **Lapsana** L. gen. n. 919.

1. **L. communis** L. sp. p. 811; S. et S. pr. II. p. 144; Ch. et B. exp. p. 237, Fl. pelop. p. 55; Bois. fl. or. III. p. 720; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1897 p. 30; Hal. Beitr. fl. Thessal. p. 15, in ö. b. Z. 1897 p. 286; Haussk. symb. p. 143, *v. glandulosa* Freyn. Fl. Südistr. p. 125 (pedicellis et saepe involucris parce pilis glanduliferis obsitis, forma ut videtur in ditione frequentior). — Icon: Fl. dan. t. 500. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Glabrescens vel inferne pubescens, superne saepe parce glandulosa; caule erecto, paniculato-ramoso; foliis inferioribus petiolatis, lyratis, intermediis subsessilibus, ovatis, grosse dentatis, superioribus sessilibus, ob-

longis vel lanceolatis, saepius integris; ligulis flavis, involucro glabro vel parce glandulifero sesquilongioribus; acheniis nitidis.

In silvaticis montanis, rare. Thessalia; in fagetis mt. Oxya supra Chaliki (Hal.), ad monasterium Korona, pr. Sermenikon (Haussk.), pr. Velitsena, pr. Hepdominta Aderfia, pr. Vlachava, mt. Hagios Paraskevi, mt. Cuka, ad Phlambures et Mitrica in mt. Chassia (Form.); Peloponnesus (Sibth.): mt. Malevo pr. Castanitza (Orph.), mt. Taygetos pr. Anavryti (Heldr.); Cyclades (Chaub.). — Jun. Aug. ⊙.

Obs. *L. tubiflora* Friedr. Reise p. 269 ex insula Aegina, mihi ignota et a Heldreichio ipso in sua Flore d'Égine non commemorata.

74. Rhagadiolus Tourn. inst. p. 479.

1. **R. stellatus** L. sp. p. 811; S. et S. pr. II. p. 144, Fl. gr. t. 817; Pieri corc. fl. p. 115; Ch. et B. exp. p. 237, Fl. pelop. p. 55; (*Lapsana*); Willd. sp. III. p. 1625; Ung. Reise p. 125; Weiss in z. b. G. 1869 p. 47; Raul. cret. p. 794; Spreitz. in z. b. G. 1877 p. 714; Heldr. fl. cephal. p. 48, Fl. Aegina p. 308; Haussk. symb. p. 143. —

Glaber vel puberulus; caule erecto, furcato et patule ramoso; foliis subintegris, dentatis vel lyratis; capitulis laxissime cymoso-paniculatis; involucri phyllis externis minimis, calyculiformibus, internis 5—8, fructiferis induratis, cylindricis, stellato-expansis, achenia marginalia omnino involventibus et cum eis persistentibus; ligulis flavis; acheniis centralibus 1—3, liberis, incurvatis. —

α. **typicus**. — *R. stellatus v. leiocarpus* DC. pr. VII. p. 77. — *R. stellatus v. leiolaenus* Bois. fl. or. III. p. 722. — *R. intermedius* Ten. fl. med. II. p. 25 (f. foliis inferioribus sinuato-lyratis) = *R. stellatus v. intermedius* DC. l. c.; Weiss l. c. — Achenia interna glaberrima; folia dentata vel sinuato-lyrata. — Exsicc.: N. v.

β. **edulis** Gaertn. fr. II. p. 354; Friedr. Reise p. 272; Hal. in ö. b. Z. 1897 p. 94; pro sp.; DC. pr. VII. p. 77; Marg. et R. fl. Zante p. 61; Raul. cret. p. 794; Weiss in z. b. G. 1869 p. 47; Heldr. fl. Aegina p. 308. — *R. stellatus v. hebaelenus* DC. l. c.; Weiss l. c.; Bois. fl. or. III. p. 722; Heldr. fl. Aegina p. 308; (f. foliis sinuato-dentatis). — *Lapsana rhagadiolus* L. sp. p. 812. — Achenia interna pubescentia, folia lyrata vel sinuato-dentata. — Exsicc.: Heldr. pl. fl. hellen. a. 1878.

In cultis, arvis, ad vias regionis inferioris per totam Graeciam; varietas ut videtur multo frequentior. — Mart. Jun. ⊙.

Obs. *Koelpinia linearis* Pall. it. III. app. p. 755 a Raulia cret. p. 794 ex Tourn. cor. 36 in Creta probabiliter erronee indicata.

18. Tribus. **LEONTODONTEAE** Schultz in Koch syn. p. 417.

75. **Thrincia** Roth cat. bot. I. p. 97.

1. **T. hirta** L. sp. ed. 2 p. 1123; Ch. et B. exp. p. 232, Fl. pelop. p. 54; (*Leontodon*); Roth. l. c. p. 98. — *Apargia hirta* Sm. engl. fl. III. p. 552; S. et S. pr. II. p. 131. — Icon: Rchb. t. 24.

Rhizomate truncato, fibris filiformibus; foliis oblongis, sinuatis, dentatis vel runcinato-pinnatifidis, plus minus setulosis; scapis nudis, monocephalis; involucri phyllis glabris vel hispidulis; ligulis flavis; acheniis marginalibus vix rugosis, subattenuatis, cupula brevi coronatis, disci rugosis, in rostrum eis 4 plo brevius attenuatis, pappo plumoso superatis.

In Arcadia (Sibth.); Messenia: pr. Navarin, Methone (Chaub.); a recentioribus non lecta. — Jun. Aug. ♃. N. v.

2. **T. tuberosa** L. sp. p. 799; Ch. et B. exp. p. 233, Fl. pelop. p. 54; (*Leontodon*); DC. fl. fr. IV. p. 52; Marg. et R. fl. Zante p. 62; Friedr. Reise p. 286; Ung. Reise p. 124; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 796; Bois. fl. or. III. p. 726; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. in ö. b. Z. 1897 p. 96; Heldr. fl. Aegina p. 307, chlor. Thera p. 16, chlor. Mykon. p. 247. — *Apargia tuberosa* Willd. sp. III. p. 1549; S. et S. pr. II. p. 130, Fl. gr. t. 797; Sieb. avis p. 5.

Rhizomate truncato, fibris napiformibus; foliis oblongis, sinuatis, dentatis, runcinatis vel lyratis, plus minus setulosis; scapis nudis, monocephalis; involucri phyllis glabris vel hispidis; ligulis flavis; acheniis marginalibus vix rugosis, apice attenuatis, muticis vel cupula fimbriata coronatis, disci muricato-rugosis, in rostrum eis dimidia vel tertia parte brevius attenuatis, pappo plumoso superatis. — Fibris napiformibus a praecedente statim diagnoscenda.

α. **typica.** — Glabrescens; involucri phylla glabra vel parce hispida; achenia disci crassiora, breviora, 4 mm. longa, in rostrum $1^1/_2$ —3 mm. longum abruptiuscule attenuata. — Exsicc.: Sint. it. thessal. n. 247; Heldr. herb. norm. n. 1560.

β. **Oliverii** DC. pr. VII. p. 100. — *T. Olivieri* Haussk. symb. p. 141. — Folia magis utrinque setulosa; capitula saepius eximie majora; involucri phylla hirsutissima; achenia disci 5 mm. longa, in rostrum 3—6 mm. longum sensim attenuata. — Etsi in formis extremis a typo valde discedit, tamen intermediis ad eum transit, quamobrem dubito, quin revera specifice differat. Exsicc.: Heldr. herb. norm. n. 1053; Dörfl. pl. cret. n. 86.

In herbidis, dumosis, cultis regionis inferioris et montanae, varietas ut videtur typo frequentior. Thessalia: pr. Vitomo in Pindo tymphaeo (Sint.); Attica: pr. Athenas (Friedr.), insula Salamis, mt. Hymettus, Parnes, pr. Tatoi, Kephissia (Heldr.), pr. Eleusis, Laurion (Haussk.),

insula Aegina (Heldr.); Argolis: pr. Argos (Heldr.), peninsula Methana (Haussk.); Laconia: pr. Gythion (Psarides); Messenia: pr. Kalamata (Zahn), Pylos, Methone, insula Sapienza (Chaub.); Sporadum insula Skopelos (Leon.); Cycladum insula: Tenos, Syra (Weiss), Kythnos, Mykonos, Rhenia (Tunt.), Melos (Leon.), Thera (Heldr.); Creta: pr. Canea, Askyphos (Raul.), Anatoli distr. Hierapetra (Leon.); Zante (Sibth.); Corcyra (Ung.). — Nov. Maio. ♃.

76. Leontodon L. gen. n. 912.

1. Sectio. *Perennes* Bois. fl. or. III. p. 727. — Pappi radii 15—40.

a. Rhizoma truncatum, fibris napiformibus.

1. **L. fasçiculatus** Biv. manip. II. p. 7 t. 2 (*Apargia*); Nym. syll. p. 53; Haussk. symb. p. 140. — *Apargia cichoracea* Ten. fl. nap. pr. p. 46. — *L. cichoraceum* Sanguin. cent. p. 111; Bois. fl. or. III. p. 729; Hal. Beitr. fl. Epir. p. 30, Beitr. fl. Achaia p. 26; non Schleich. — *Millina leontodontoides* Cass. dict. 31 p. 90; Raul. cret. p. 796. — Exsicc.: Sint. it. thessal. n. 1464.

Glabriuscula vel parce pilosa; foliis oblongo-vel lanceolato-spathulatis, dentatis vel runcinatis; scapis squamulosis, monocephalis, apice incrassatis; involucri phyllis lanceolatis, puberulis et pilis longioribus hirsutis, externis abbreviatis; ligulis flavis; acheniis rugulosis, in rostrum eis longius attenuatis; pappi sordidi, achenio brevioris radiis plumosis, basi paululum dilatatis, externis nonnullis subinde brevioribus. — Habitu *Thrinciae tuberosae* persimilis, ab ea praeter pappum omnium acheniorum conformem, scapis squamulosis, infra capitulum eximie incrassatis et acheniis longius rostratis facile diagnoscitur.

In herbidis regionis mediae et superioris. Epirus: mt. Tsumerka (Hal.); Thessalia: mt. Karava, Ghavellu in Pindo (Haussk.), ad Metochi pr. Kalabaka (Sint.), mt. Pelion pr. Zagora (Heldr.); Attica: mt. Parnes (Sprun.); Achaia: mt. Panachaicon, Olenos (Hal.); mt. Taygetos pr. Androvista (Heldr.); Creta: pr. Askyphos (Raul.). — Maio, Jul. ♃.

b. Rhizoma truncatnm vel fusiforme, fibrosum.

α. Rhizoma truncatum.

2. **L. hispidus** L. sp. p. 799; Ch. et B. exp. p. 233, Fl. pelop. p. 54. — *Apargia hispida* Host syn. p. 424; S. et S. pr. II. p. 130. — *L. hastilis v. hispidus* Bois. fl. or. III. p. 730. — Huc etiam, sed vix varietas dicenda: *L. hastilis v. banaticus* Heuf. en. pl. Ban. p. 109; Haussk. symb. p. 141. — Icon: Fl. dan. t. 862. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Totus vel ad folia tantum pilis 2—3 furcatis hirsutus; foliis oblongo-lanceolatis, in petiolum attenuatis, dentatis pinnatifidisve; scapis monocephalis, nudis vel apice 1—2 squamulosis, subincrassatis; involucri phyllis lanceolatis, hirtis; ligulis flavis; acheniis superne tenuiter rugu-

loso-muricatis, apice subattenuatis; pappi sordidi, achenium aequantis vel longioris, radiis plumosis, basi dilatatis, extimis brevibus, scabris.

β. **hastilis** L. sp. ed. 2 p. 1123; Heldr. chlor. Parn. p. 22; Hal. Beitr. fl. Thessal. p. 15; pro sp. — *Apargia hastilis* Host. syn. p. 423; Friedr. Reise p. 272. — *L. hastilis v. glabratus* Koch syn. p. 419; Bois. fl. or. III. p. 730. — Foliis, scapis involucrisque glabris vel pilis paucis adspersis. — Exsicc.: Heldr. fl. gr. a. 1855.

In pascuis regionis montanae in alpinam usque adscendens, rarior. Thessalia: mt. Oxya supra Chaliki (Hal.), pr. Malakasi, mt. Zygos, Karava, Ghavellu (Haussk.) in Pindo, mt. Olympus, Pelion ad Megarheuma, mt. Oeta Phthiotidis (Heldr.); Aetolia: mt. Tymphrestus (Samarit.), Korax (Tunt.); mt. Parnassus (Guicc.); Peloponnesus (Sibth.): pr. Troezene Argolidis (Friedr.). —

β. Rhizoma fusiforme, elongatum.

× Capitula magna; achenia 20—22 mm. longa; pappi setae omnes conformes elongatae.

3. **L. asperrimus** Willd. sp. III. p. 1507 (*Scorzonera*); Bois. fl. or. III. p. 732; Bald. riv. coll. bot. Alb. 1896 p. 71. — *Apargia strigosa* M. B. fl. taur. cauc. II. p. 249. — *Asterothrix asperrima* Cass. dict. 48 p. 434. — *Hypochaeris setosa* Form. in D. bot. Monat. 1897 p. 197, in Ver. Brünn 1897 p. 31. — Exsicc.: Bald. it. alb. epir. IV. n. 244; Form. pl. thessal. a. 1896.

Pilis rigidis, confertis, stellato-furcatis scaberrimus; foliis anguste oblongo-lanceolatis lanceolatisve, remote dentatis; scapis monocephalis vel inferne ramulo auctis, validis, squamulis linearibus vel setaceis obsitis, superne incrassatis; involucri cylindracei phyllis imbricatis, ab externis oblongis ad interna lanceolata elongatis, praeter indumentum brevissimum margine et ad nervum pilis albis, validis, apice stellatis pectinato-ciliatis; ligulis flavis; acheniis scabridis, a medio sursum attenuatis; pappi sordidi, achenio subbrevioris setis numerosis, 30—35, omnibus plumosis, basi non dilatatis. — Species capitulis maximis insignis.

In herbidis saxosis regionis mediae mt. Smolika distr. Konitza in Epiro (Bald.) et in mt. Cuka in mt. Chassia Thessaliae (Form.). — Jun. Jul. ♃.

×× Capitula mediocria; achenia 10—12 mm. longa; pappi setae externae nonnullae multo breviores.

4. **L. asper** W. et K. pl. rar. hung. II. p. 114 t. 110 (*Apargia*); Poir. enc. suppl. III. p. 453; Rchb. fl. germ. exc. p. 252; Form. in D. bot. Monat. 1890 p. 15; Hal. Beitr. fl. Epir. p. 30; Haussk. symb. p. 140; Bald. riv. coll. bot. Alb. 1896 p. 71. —

Pilis 3—4 furcatis, crebris asper; foliis oblongo-lanceolatis, sinuato-dentatis pinnatifidisve; scapis monocephalis, parce squamulosis vel sub nudis, rarius furcato 2—3 cephalis et tunc ad bifurcationes folio in structis, apice incrassatis; involucri phyllis imbricatis, ab externis triang

laribus ad interiora lanceolata elongatis, pilis brevibus albis, 4—5 furcatis eximie fimbriato-ciliatulis, dorso glabris vel ad nervum setulosis; ligulis flavis; acheniis scabridis, a parte interiori sursum attenuatis; pappi sordidi, achenio subbrevioris setis 15—20, omnibus plumosis, basi non dilatatis.

α. **typicus.** — Indumentum foliorum e pilis longioribus et brevioribus mixtis, apice 4—3 furcatis, radiis erecto-patentibus constans; involucri phylla omnia vel saltem exteriora eleganter ciliata, dorso glabra. — Exsicc.: Bald. it. alb. epir. IV. n. 181 et 245.

β. **setulosus** Hal. in ö. b. Z. 1892 p. 372. — *L. asper v. Huetii* Form. in Ver. Brünn 1896 p. 42, non Bois. — Indumentum foliorum ut in *α.*, sed copiosius; involucri phylla praeter marginem ciliatam, dorso pilis albis longis, flexuosis setulosa. — Transitum praebere videtur ad var. *β.* speciei sequentis. — Exsicc.: Sint. et Bornm. it. turc. n. 870 e mt. Athos; Heldr. it. thessal. IV. a. 1885.

γ. **Haussknechtii** Uechtr. ap. Haussk. in Mitth. thür. bot. Ver. V. p. 87, symb. p. 140. — Indumentum foliorum e pilis brevibus conformibus, basi dilatatis, apice brevissime uncinato-trifurcatis constans; indumentum phyllorum ut in *α.*, sed brevissimus. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

δ. **saxatilis** Ten. syll. pl. p. 393, Fl. nap. II. p. 168 t. 72 (*Apargia*); Rchb. fl. germ. exc. p. 252; Bald. riv. coll. bot. Alb. 1895 p. 52; pro sp. — Huc forsan *L. squamosum* Pieri corc. fl. p. 114, non Lam. — Indumentum foliorum ut in *α.*; involucri phylla pilis albis, flexuosis setulosa, margine (interdum phylla extima excepta) non ciliata. — Exsicc.: Baen. herb. europ. n. 9281.

In saxosis, praeruptis regionis montanae et sabalpinae. Epirus: mt. Kuruna distr. Ljaskovik, mt. Smolika distr. Konitza, mt. Vratedon distr. Zagorion (Bald.), mt. Tsumerka supra Vulgarelion (Hal.); Thessalia: mt. Baba supra Klinovo, mt. Karava, Gionscala supra Sermenikon (Haussk.), pr. Kalabaka (Form.); — *β.* mt. Karava in Pindo dolopico (Heldr.), mt. Zygos (Form.); — *γ.* pr. Malakasi, mt. Zygos in Pindo (Haussk.); — *δ.* mt. Peristeri in Pindo (Bald.); Corcyra: mt. Deka supra Garuna (Baen.). — Maio, Jul. ♃.

5. **L. graecus** Bois. et Heldr. diagn. XI. p. 39, Fl. or. III. p. 731; Heldr. fl. cephal. p. 48, Fl. Aegina p. 307; Hal. Beitr. fl. Achaia p. 26, in ö. b. Z. 1897 p. 96; Haussk. symb. p. 140; Bald. riv. coll. bot. Alb. 1895 p. 52, 1896 p. 71. — *L. incanus* Ch. et B. exp. p. 233, Fl. pelop. p. 54, non (L.) Schrank. — *L. crispus* Ung. Reise p. 124, non Vill. — *L. abietum* Heldr. herb. norm. n. 1561. — Exsicc.: Heldr. herb. norm. n. 145; 1052 et 1561; Orph. fl. gr. n. 1085; Sint. et Bornm. it. turc. n. 1335; Bald. it. alb. epir. III. n. 168, f. phyllorum indumento ad *L. asperum* vergens.

Pilis brevissimis, crebris, stellatis, et saepe longioribus 3—4 furcatis plus minus asper; foliis oblongo-lanceolatis, sinuato-dentatis vel in

lacinias triangulares vel lanceolatas pinnatipartitis, interdum sublyratis; scapis monocephalis, rarius ramo auctis, nudis vel parce squamulosis, apice incrassatis; involucri phyllis imbricatis, ab externis triangularibus ad interiora lanceolata elongatis, brevissime pulverulento-canis, praetereaque secus nervum plus minus et breviter setulosis, margine non fimbriato-ciliatulis; ligulis flavis; acheniis scabridis, superne subattenuatis; pappi sordidi, achenio subbrevioris setis 15—25, omnibus plumosis, basi subdilatatis. — Praecedenti maxime affinis et ab eo praesertim indumento discedit, attamen formis intermediis ad eum transire videtur.

β. **Heldreichianus** Bois. fl. or. III. p. 731; Heldr. chlor. Parn. p. 22; Hal. in ö. b. Z. 1897 p. 286. — *L. Heldreichianus* Hal. in z. b. G. 1888 p. 759. — *L. graecus v. alpinus* Bois. diagn. XI. p. 39, non *L. alpinus* Jacq. — Indumentum densius canum; involucri phylla praeter indumentum breve pilis longis flexuosis obsita. — Intermediis ad typum transit. — Exsicc.: Orph. fl. gr. n. 331.

In saxosis regionis montanae et alpinae; Epirus: mt. Kuruna, Tsumerka (Bald.); Thessalia: mt. Olympus pr. Litochori (Sint.); Sporadum insula Jura (Leon.); Euboea: mt. Dirphys (Heldr.); Boeotia: pr. Thebas (Heldr.); Attica: mt. Lycabettus (Orph.), Kerata (Haussk.), Pateras, Parnes, Pentelicon, Hymettus, pr. Laurion, Kamariza, insula Salamis, Aegina (Heldr.); Elis: mt. Olenos (Heldr.); Achaia: pr. Lopesi (Hal.), mt. Chelmos, Kyllene (Heldr.); ad Acrocorinthum, in peninsula Methana Argolidis (Haussk.); Arcadia: pr. Carithena (Heldr.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.); — *β.* Aetolia: mt. Korax (Heldr.); mt. Kiona (Hal.), Parnassus (Orph.); mt. Taygetos (Zahn). — Apr. Jul. ♃.

2. Sectio. *Annui* Bois. l. c. — Pappi radii 10.

6. **L. taraxacifolius** Cass. dict. 31 p. 89 (*Deloderium*). — *Apargia hyoseroides* Sieb. avis p. 5. in Spreng. syst. III. p. 664, non Vest. — *Millina hyoseroides* DC. pr. VII. p. 110; Raul. cret. p. 796. — *L. creticum* Bois. fl. or. III. p. 720.

Crispule et minute setulosum; scapis numerosis, tenuibus, 1—2 cephalis, folia parum superantibus, foliis lineari-oblongis, sinuato-pinnatifidis; capitulis parvis; involucri phyllis imbricatis, lanceolatis, praeter setulas secus nervum paucas glabris, ab externis apice sphacelatis ad intima elongatis; acheniis rugulosis, in rostrum eis subduplo brevius attenuatis; pappi albidi, achenio brevioris, setis 10, plumosis, basi paleaceo-dilatatis.

Creta (Sieb.). — N. v.

Obs. Species non satis nota: *L. daedaleum* Sieb. in Flora V. I. p. 243. „Squamis calycinis omnibus patentibus, lineari-lanceolatis; scapo abbreviato; foliis glabris, runcinato-pinnatis, pinnis runcinatis, contortuplicatis, inaequalibus. — In pascuis montosis in omni Cretae campo Cydoniensi. — Mart. Apr.“ N. v.

77. Picris L. gen. n. 907.

a. Biennis; caulis superne stricto breviterque ramosus, racemoso-paniculatus; capitula fructifera sub apice subconstricta.

1. **P. spinulosa** Bert. in Guss. syn. II. p. 400; Form. in Ver. Brünn 1895 p. 25. — *P. stricta* Jord. cat. Dijon 1848 p. 19; Bois. fl. or. III. p. 735; Haussk. symb. p. 140; Form. in Ver. Brünn 1896 p. 42, 1897 p. 31. — *P. hieracioides* S. et S. pr. II. p. 132; Ch. et B. fl. pelop. p. 55; Friedr. Reise p. 263; Form. in D. bot. Monat. 1890 p. 15; non L., quae caule patule ramoso, corymboso, interdum subumbellato, capitulis majoribus et acheniis subtiliter rugosis discedit. — Exsicc.: Orph. fl. gr. n. 1084 (Macedonia); Sint. et Bornm. it. turc. n. 1334.

Pilis simplicibus et glochidiatis aspera; caule erecto; foliis oblongo-lanceolatis, dentatis vel sinuatis, radicalibus basi attenuatis, caulinis semiamplexicaulibus; pedunculis brevibus, apice non incrassatis; involucri phyllis linearibus, pulverulento-canis et saepe ad carinam setis glochidiatis hispidis, exterioribus brevibus, patulis; ligulis flavis; acheniis subincurvis, transverse rugosis, apice in umbonem contractis.

In ruderatis, vineis, dumetis, saxosis regionis inferioris et montanae. Epirus: in valle Dipotami (Form.); Thessalia: pr. Chaliki (Haussk.), Malakasi, Koturi in Pindo, pr. Trikala, Volo (Form.), pr. Litochori (Sint.) et Hagios Dionysios in Olympo (Heldr.); Attica: pr. Athenas (Sibth.); Argolis: pr. Nauplia (Haussk.); Zante (Sibth.); Corcyra (Friedr.). — Jun. Sept. ⊙

b. Annuae; caulis divaricato-ramosus, laxe corymbosus; capitula fructifera sub apice strangulata.

2. **P. Sprengeriana** L. Sp. p. 804 (*Hieracium*); Lam. dict. V. p. 310; Marg. et R. fl. Zante p. 62, Raul cret. p. 797, Bois. fl. or. III. p. 738; Heldr. fl. Cephal. p. 49, Flor Aegina p. 307, chlor. Thera p. 6; Spreitz. in z. b. G. 1887 p. 663; Haussk. symb. p. 140; Form in Ver. Brünn 1895 p. 25. — *Crepis rhagadioloides* L. mant. p. 108. — *C. Sprengeriana* Willd. sp. III p. 1598. — *Hieracium ciliatum* Willd. sp. III p. 1585; Raul. cret. p. 801. — *P. rhagadioloides* Pers. ench. II p. 370. — Icon: Jacq. hort. Schönbr. t. 144. — Exsicc.: Reverch. pl. cret. n. 97; Dörfl. fl. aeg. n. 18, pl. cret. n. 104.

Pilis simplicibus et glochidiatis aspera; caule erecto; foliis oblongo-lanceolotis, sinuato-dentatis vel integris, radicalibus basi attenuatis, caulinis aurito-amplexicaulibus; pedunculis elongatis, apice non incrassatis; involucri phyllis exterioribus lineari-setaceis, brevibus, interioribus demum cymbaeformibus, acuminatis, dorso medio calloso-carinatis, glochidiato-setosis; ligulis flavis; acheniis curvatis, transverse rugosis, apice breviter attenuatis, non rostratis.

In herbidis, collibus siccis, ad vias regionis inferioris. Acarnania: pr. Bonitza (Lakon); Thessalia: pr. Aivali, Orman Magula, Pharsalus (Haussk.); Euboea: pr. Kastaniotissa (Held.); Aetolia: pr. Guritza (Reis.);

Corinthus: ad Lechaeon; insula Aegina (Heldr.); Elis: pr. Bartholomeo (Heldr.); Achaia: pr. Psatopyrgos (Hal.), Trikala (Heldr.); Argolis: pr. Nauplia (Heldr.); Messenia: pr. Kalamata (Zahn); Cycladum insula: Kythnos (Tunt.), Naxos (Leon.), Amorgos, Thera (Heldr.); Creta: pr. Kissamos (Rev.), Malaxa, Canea, Akroteri (Raul.), pr. Males distr. Hierapetra (Leon.); Cythaera: pr. Katuni, Kapsali (Spreitz.); Zante (Marg.); Cephalonia: pr. Pessada, Piscardo (Heldr.); Corcyra: pr. Kephalomanduchion (Form.). — Maio, Jul. ⊙

3. **P. pauciflora** Willd. sp. III p. 1557; S. et S. pr. II p. 132; Weiss in z. b. G. 1869 p. 46; Bois. fl. or. III p. 737; Hal. Beitr. fl. Achaia p. 26; Haussk. symb. p. 140; Boissieu in bull. soc. bot. Fr. 1896 p. 286; Bald. riv. coll. bot. Alb. 1896 p. 71; Heldr. fl. Aegina p. 307. — Icon: Rchb. t. 22. — Exsicc.: Sint. it. thessal. n. 399.

Differt a praecedente, cui habitu similis, pedunculis fructiferis valde inflatis, sub apice strangulatis, praeter glochides dorso albo-tomentellis et acheniis in rostrum crassum eis 4 plo brevius attenuatis.

In iisdem locis, ubi praecedens. Epirus: mt. Micikeli (Bald.); Thessalia: pr. Kalabaka (Sint.); Euboea: pr. Steni (Pichl.); Attica: pr. Sunium, ad Ilyssum, mt. Lycabettus, Hymettus, Corydalus, Penteli-con, insula Salamis, Lero Pharmacusarum, Aegina (Heldr.); Boeotia: pr. Larmes (Boissieu); pr. Kalamaki in isthmo Corinthiaco, in peninsula Methana Argolidis (Haussk.); insula Poros, Keos (Heldr.) et Syra (Weiss) Cycladum; Achaia: pr. Patras (Hal.). — Apr. Jun. ⊙

78. **Helminthia** Juss. gen. p. 170.

1. **H. echioides** L. sp. p. 792; S. et S. pr. II p. 131; Ch. et B. fl. pelop. p. 55; (*Picris*); Gärtn. fr. II p. 368; Sieb. avis rem. p. 5; Friedr. Reise p. 264; Fraas fl. class. p. 198; Clem. sert. p. 50; Raul. cret. p. 798; Bois. fl. or. III p. 742; Haussk. symb. p. 140; Form. in Ver. Brünn 1895 p. 25, 1897 p. 31; Heldr. fl. Aegina p. 307. — Icon: Rchb. t. 27. — Exsicc.: Heldr. pl. fl. hellen. a. 1882.

Setis simplicibus et glochidiatis hispida; caule erecto vel decumbente, a basi divaricatim ramoso; foliis oblongo-lanceolatis, sinuato-dentatis vel integris, radicalibus in petiolum attenuatis, caulinis cordato-amplexicaulibus; capitulis breve-pedunculatis, approximatis; involucri phyllis externis cordato-ovatis, interiora lanceolata subaequantibus; ligulis flavis; acheniis rugulosis, in rostrum tenue eis sublongius abrupte attenuatis.

In herbidis, arvis, cultis regionis inferioris et submontanae. Thessalia: pr. Pharsalus (Haussk.); Phthiotis: pr. Stylida (Form.): Aetolia: pr. Klissura (Samar.); Phocis: pr. Amplissa (Fraas); Attica: pr. Athenas (Sibth.), Cephissia (Heldr.), Tatoi (Hal.), Myli (Haussk.); insula Aegina (Heldr.); Argolis: pr. Poros (Cadet); Achaia: pr. Patras (Friedr.); Messenia: pr. Kalamata (Zahn): Creta: pr. Canea, Nerokuru, Enneachoria (Raul.); Corcyra: pr. Cephalomanduchion (Form.). — Maio, Jul. ⊙.

Obs *H. asplenioides* L. sp. p. 793; S. et S. pr. II. p. 132; (*Picris*); DC. pr. VII. p. 132; Marg. et R. fl. Zante p. 62; Bois. fl. or. III. p. 742. — *Leontodon muricatum* L'Her. stirp. p. 173 t. 82; Ch. et B. exp. p. 233, Fl. pelop. p. 54. — Indicatur probabiliter ex confusione a Sibthorp in insula Zante et a Chaubard et Bory pr. Messene, sed ibi a recentioribus non reperta fuit.

79. Urospermum Juss. gen. p. 170.

1. **U. picroides** L. sp. p. 790; Ch. et B. fl. pelop. p. 53; Fraas fl. class. p. 97; (*Tragopogon*); Desf. cat. hort. par. p. 90; Marg. et R. fl. Zante p. 62; Friedr. Reise p. 262; Clem. sert. p. 51; Ung. Reise p. 124; Weiss in z. b. G. 1869 p. 46; Raul. cret. p. 797; Bois. fl. or. III. p. 743; Heldr. fl. cephal. p. 49, Fl. Aegina p. 307, chlor. Mykon. p. 247; Haussk. symb. p. 140. — *Tragopogon asperum* L. sp. p. 790. — *U. asperum* DC. fl. fr. n. 2987. — *U. picroides β. asperum* DC. et Duby bot. gall. I. p. 295; Weiss, in z. b. G. 1869 p. 46. — *Arnopogon picroides* Willd. sp. III. p. 1496; S. et S. pr. II. p. 121, t. 781; Sieb. avis p. 4, rem. p. 5 (loco ultimo err. typogr. *Andropogon*). — ? *U.echioides* Fraas fl. class. p. 198. — Exsicc.: Baen. herb. europ. n. 9365.

Parce setuloso-hispidum; caule erecto, parce ramoso, rarius simplici; foliis teneris, inferioribus oblongo-spathulatis, in petiolum attenuatis, superioribus auriculato-amplexicaulibus, omnibus dentatis, pinnatifidis vel lyratis, sumnis lanceolato-linearibus; involucri phyllis ovato-lanceolatis, basi connatis; ligulis luteis; acheniis transverse lamellatis, rostro basi inflato superatis. — *U. asperum* (L.) formam caule submonocephalo, foliis superioribus subintegris sistit.

In herbidis, dumetis, cultis regionis inferioris. Acarnania: pr. Astakos (Nied.); Attica: pr. Athenas, mt. Pentelicon (Clem.), Hymettus (Hal.), ad Phalerum, Sunium, insula Lero Pharmacusarum (Heldr.), Aegina (Friedr.); Corinthus (Fraas); Argolis; peninsula Methana, pr. Nauplia (Haussk.); Achaia: pr. Patras (Hal.); Elis: pr. Lintzi (Heldr.); Messenia: pr. Methone (Chaub.); Cycladum insula: Keos, Kythnos (Heldr.), Tenos, Syra (Weiss), Mykonos (Sart.); Creta: pr. Nerokuru, Malaxa, Candia (Raul.); Zante (Marg.); Cephalonia (Ung.): pr. Argostoli, Pessada (Heldr.); Corcyra: pr. Kastrades (Baen.). — Apr. Maio. ⊙.

Obs. *U. Dalechampii* L. sp. p. 790 (*Tragopogon*); Desf. l. c.; Bois. fl. or. III. p. 742. — *Arnopogon Dalechampii* Willd. sp. III. p. 1496; S. et S. pr. II. p. 121. — Indicatur a Sibthorp cum ? in Archipelogi insulis, sed recenter non lectum est.

19. Tribus. SCORZONEREAE Schultz in Koch. syn. p. 422.

80. Geropogon L. gen. n. 904.

1. **G. glaber** L. sp. ed. 2. p. 1109; S. et S. pr. II. p. 119; Pieri corc. fl. p. 113; Ch. et B. exp. p. 230, Fl. pelop. p. 53; Raul.

cret. p. 797; Bois. fl. or. III. p. 744; Heldr. fl. Cephal. p. 49, Fl. Aegina p. 307; Haussk. symb. p. 138. — Icon: Jacq. hort. vind. t. 33. — Exsicc.: Heldr. pl. fl. hellen. a. 1881; Bald. it. erct. alt. n. 35.

Glaber; caule erecto, simplici vel stricte ramoso; foliis elongato-linearibus, semiamplexicaulibus; pedunculis fistulosis, demum incrassatis; involucri phyllis lineari-lanceolatis, ligulas violaceo-roseas superantibus; acheniis scabridis. — Habitu omnino *Tragopogonis*.

In collibus herbidis, rupestribus regionis inferioris. Boeotia (Bois.); Attica: mt. Lycabettus, ad Piraeum, insula Salamis, Aegina (Heldr.); Acrocorinthus (Haussk.); Argolis: pr. Nauplia (Sart.), insula Hydra (Heldr.); Elis, Laconia (Sibth.); Messenia: pr. Arcadia, Messene (Chaub.); Creta: pr. Kalyves, Francocastello (Raul.), Candia (Bald.); Cephalonia: pr. Argostoli, Aphrata (Heldr.); Corcyra (Pieri). — Apr. Maio. ⊙.

81. Tragopogon L. gen. n. 905.

a. Ligulae purpureae vel lilacinae.

1. **T. porrifolius** L. sp. p. 789; Pieri corc. fl. p. 113; Ch. et B. exp. p. 230, Fl. pelop. p. 53; Friedr. Reise p. 276; Fraas fl. class. p. 196; Ung. Reise p. 124; Raul. cret. p. 797; Bois. fl. or. III. p. 745; Heldr. fl. cephal. p. 49, chlor. Mykon. p. 247; Spreitz. in z. b. G. 1887 p. 663; Haussk. symb. p. 138; Bald. riv. coll. bot. Alb. 1896 p. 71; Hal. in ö. b. Z. 1897 p. 96; non Jord. cat. Dijon 1848 p. 32, nec Gr. et G. fl. Fr. II. p. 312, nec Willk. et L. pr. fl. hisp. II, p. 226, quorum stirps a Linnaeana ligulis involucrum superantibus. acheniis fulvis subito in rostrum tenue eis brevius contractis distiguitur. — *T. australis* Jord. cat. Dijon 1848 p. 32; Weiss in z. b. G. 1869 p. 45; Hal. Beitr. fl. Achaia p. 25; Heldr. fl. Aegina p. 307. — *Geropogon hirsutus* Urv. enum. p. 98, non L., sec. Bois. fl. or. III. p. 745. — Huc forsan: *T. Cupani* Guss. in DC. pr. VII. p. 113, saltem quoad plantam e Zacyntho; Marg. et R. fl. Zante p. 62. — Fxsicc.: Heldr. herb. norm. n. 395; Rev. pl. cret. n. 96; Sint. it. thessal. n. 402.

Glaber vel subfloccosus; caule erecto, crassiusculo, simplici vel saepius ramoso; foliis a basi plus minus dilatata, semiamplexicauli, lanceolato-linearibus; pedunculis valde incrassato-clavatis; capitulis magnis; involucri phyllis suboctonis, flores superantibus; acheniis testaceis, exterioribus muricato-squamulosis, interioribus sublaevibus, omnibus in rostrum crassiusculum, laeve, eis sublongius vel aequilongum, apice non incrassatum, sub pappo constrictum et hic interdum lanatum, sensim abeuntibus.

β. **brachyphyllum** Bois. fl. or. III. p. 745. — Involucri phylla flores non vel vix superantia. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889.

In herbidis, collibus, incultis regionis inferioris. Epirus: pr. Kusmera distr. Janina (Bald.); Thessalia: ad monasterium Korona (Haussk.),

pr. Kalabaka (Sint.), pr. Milies, Volo (Heldr.); Sporadum insula Skopelos (Leon.); Euboea: pr. Kurbatzi, Kumi (Wild), in Petaliarum insulis (Münter); Aetolia: pr. Mesolonghion (Nieder); Attica: pr. Athenas, in Acropoli, in colle Lycabetto, ad Piraeum, Phalerum, in colle Ardetto, pr. Heracleon, Tatoi, Menidi, mt. Pentelicon, Hymettus, Eleusis, Rhaphti, Kyprianos, Laurion, insula Salamis, Lero, Aegina (Heldr.); Achaia: pr. Patras (Hal.); Acrocorinthus (Haussk.); Argolis: pr. Poros (Friedr.); in Tzakonia (Fraas); Messenia: pr. Kalamata (Zahn), Petalidi, Andruso, Arcadia (Chaub.); Cycladum insula Keos, Kythnos (Heldr.), Tenos, Syra (Weiss), Mykonos (Sart.), Melos (Urv.); Creta: pr. Canea, Akroteri, Malaxa, Rhodakino, Selia, in oropedio Omalos (Raul.); Zante (Marg.); Cephalonia: pr. Drapano (Heldr.); Corcyra: ad Cap Bianco (Spreitz.), mt. Hagios Deka (Baen.). — Apr. Maio. ⊙.

2. **T. balcanicus** Vel. in böhm. Ges. Wiss. VII. p. 28, Fl. bulg. p. 355; Hal. Beitr. fl. Epir. p. 30. — Exsicc.: Hal. it. gr. II. a. 1893.

Inferne floccosus; caule erecto vel adscendente, gracili, saepius a collo in ramos elongatos diviso, raro subsimplici; foliis angustissime linearibus, gramineis, basi parum dilatatis, vix semiamplexicaulibus; pedunculis parum incrassatis; capitulis sub anthesi parvis, tenuibus, paucifloris; involucri phyllis 4—7 nis, flores aequantibus; acheniis fuscescentibus, a basi fere ad rostri apicem muricato-squamulosis, in rostrum eis brevius vel aequilongum, apice clavato-incrassatum, sub pappo constrictum sensim abeuntibus. — Subspecies orientalis *T. crocifolii* L., qui foliis latioribus, involucri phyllis longioribus et acheniorum rostro laevi a nostro parum differt. Maxime affinis quoque *T. nebrodensi* Guss. syn. II. p. 384, qui a *T. crocifolio* foliis angustis, involucri phyllis flores aequantibus et a nostro acheniorum rostro laevi tantum discedit.

In saxosis regionis mediae mt. Tsumerka supra Vulgarelion et Theodoriana in Epiro (Hal.). — Jul. ⊙.

Obs. Quid sint *T. crocifolius* Ch. et B. fl. pelop. p. 53 e Methone Messeniae et planta ex Attica et Tzakonia sub eodem nomine a Fraas fl. class. p. 197 enumerata, nescio.

b. Ligulae flavae.

α. Rostrum achenio longius vel subaequilongum.

× Folia angustissime linearia, graminea, basi vix dilatata.

3. **T. Samaritanii** Heldr. et Sart. in Bois. diagn. ser. 2, V. p. 116, Fl. or. III. p. 748; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Epir. p. 30, Beitr. fl. Achaia p. 25, in ö. b. Z. 1897 p. 285; Heldr. chlor. Parn. p. 22; Haussk. symb. p. 138. — Exsicc.: Heldr. herb. norm. n. 690 et 1559.

Inferne floccosus; caule erecto vel adscendente, saepius a collo in ramos elongatos diviso, rarius simplici; foliis angustissime linearibus, gramineis, basi parum dilatatis, vix semiamplexicaulibus; pedunculis vix incrassatis; capitulis sub anthesi parvis, tenuibus, paucifloris; involucri

phyllis 5—7 nis, flores aequantibus; acheniis fuscescentibus, muricato-squamulois, in rostrum subaequilongum, apice clavato-incrassatum sensim abeuntibus. — Praecedenti valde affinis et ab eo flosculis flavis (sed marginalibus ad apicem interdum violaceis) et rostro laevi vix specifice distinctus.

In herbidis, saxosis regionis mediae et superioris. Epirus: mt. Tsumerka supra Theodoriana (Hal.); Thessalia: in oropedio Neuropolis ad monasterium Korona, pr. Sermenikon (Haussk.) in Pindo, mt. Olympus (Orph.); Aetolia: mt. Tymphrestus (Samarit.), Korax (Tuntas); mt. Kiona (Hal.), Parnassus pr. Gurna (Guicc.) et Livadi (Hal.); Attica: mt. Parnes (Heldr); Achaia: mt. Panachaicon (Hal.), Kyllene (Heldr.); mt. Taygetos (Reis.). — Maio, Aug. ⊙

×× Folia e basi dilatata semiamplexicauli lanceolata.

○ Pedunculi non aut leviter incrassati.

4. **T. pratensis** L. sp. p. 789; Dallap. prosp. p. 107, sed secundum Heldr. fl. cephal. p. 49 determinatio incerta et planta cephalonica forte ad *Scorzoneram crocifoliam* pertinet; Haussk. symb. p. 138. — Icon: Fl. dan. t. 424. — Exsicc.: Heldr. it. thessal. IV a. 1895; Sint. it. thessal. n. 474.

Glaber, junior interdum subfloccosus; caule erecto, stricte ramoso; foliis a basi dilatata semiamplexicauli lanceolatis, strictis, vel apice reflexis contortisve; pedunculis sub capitulo parum incrassatis; involucro octophyllo, flores aequante vel superante; acheniis marginalibus squamuloso-muricatis, in rostrum filiforme aequilongum vel paulo longius, apice vix incrassatum, attenuatis.

β. **orientalis** L. sp. p. 789; Friedr. Reise p. 278 et 285; Haussk. symb. p. 138, cum β. *tortilis* (f. foliis undulatis, apice contortis); Form. in Ver. Brünn 1897 p. 31; pro sp.; Garcke Fl. Deutschl. ed. 13. p. 234. — Capitula majora; involucrum floribus tertia parte brevius; flores saturatiores, achenia rostro apice subclavato-incrassato, $^1/_3$ longiora. Sed omnes hae notae valde variare videntur; quamobrem iis libenter assentior, qui in *T. orientali* varietatem tantum agnoscunt. — Exsicc.: N. v.

In pratis regionis inferioris et montanae. Thessalia: in oropedio Neuropolis (Haussk.), ad Guvelzi pr. Kalabaka (Sint.), ad ripas Enipei pr. Ormanmagula (Heldr.); Cephalonia (Dallap.); — β. Thessalia: pr. Trikala (Form.), Ormanmagula (Haussk.); Argolis: pr. Nauplia (Friedr.). — Maio, Jul. ⊙.

5. **T. Tommasinii** Schultz in Bisch. Beitr. Fl. Deutschl. I. p. 97; Rchb. ic. fl. germ. XIX. p. 19 t. 40; Bois. fl. or. III. p. 748. —

Floccoso-tomentosus, floccis ad foliorum basin persistentibus; caule erecto, stricte ramoso; foliis a basi dilatata semiamplexicauli lanceolatis, flexuosis contortisve; pedunculis leviter subclavato-incrassatis; involucro octophyllo, flores aequante; acheniis marginalibus squamuloso-muricatis, in rostrum filiforme, eis sesquilongius, apice clavato-incrassatum, attenu-

atis. — Differt a praecedente, cui habitu persimilis, indumento persistente, acheniis tenuius costatis et longius rostratis.

In pratis siccis Atticae: pr. Marathon (Vrioni). — Maio, Jun. ⊙ N. v.

○○ Pedunculi valde incrassato-clavati.

6. **T. majus** Jacq. fl. austr. I. p. 19 t. 29; S. et S. p. II. p. 120; Ch. et B. exp. p. 230, Fl. pelop. p. 53; Marg. et R. fl. Zante p. 62; Haussk. symb. p. 138; Form. in Ver. Brünn 1897 p. 31. — Exsicc.: Sint. it. thessal. n. 468.

Glaber; caule erecto, parce ramoso; foliis a basi dilatata semiamplexicauli linearibus; involucri phyllis 8—12, flores superantibus; acheniis squamuloso-muricatis, in rostrum aequilongum vel sublongius, apice clavato-incrassatum attenuatis.

In collibus et herbidis siccis. Thessalia: pr. Kalabaka (Sint.), Karditza, Sophates, Pharsalus, Aivali (Haussk.), ad Hagios Paraskevi et Phlambures in mt. Chassia (Form.); Laconia et Arcadia (Chaub.); Zante (Sibth.). — Jun. Jul. ⊙.

β. Rostrum achenio 3—4 plo brevius.

7. **T. floccosus** W. et K. pl. rar. hung. II. p. 116 t. 112; Bald. riv. coll. bot. Alb. 1896 p. 71. — Icon: Rchb. t. 42.

Floccoso-tomentosus, floccis subpersistentibus; caule erecto vel adscendente, saepe a basi ramoso; foliis anguste linearibus, canaliculatis, flexuosis, inferioribus valde elongatis, superioribus basi dilatata semiamplexicaulibus, abbreviatis; pedunculis vix incrassatis; involucro octophyllo, flores subaequante; acheniis marginalibus muricatis, in rostrum brevissimum attenuatis, centralibus submuricatis, longius rostratis.

In humidis ad margines lacus Janina (Panajotis). — Jun. Aug. ⊙ N. v.

8. **T. longifolius** Heldr. et Sart. in Bois. diagn. ser. 2, III. p. 90, Fl. or. III. p. 749; Haussk. symb. p. 138; Form. in D. bot. Monat. 1898 p. 78. — Exsicc.: Heldr. herb. norm. n. 396 et 1157.

Glaber; caule erecto vel adscendente, simplici vel ramoso; foliis anguste linearibus, planis, flexuosis, inferioribus saepe elongatis, superioribus abbreviatis, basi vix dilatatis; pedunculis vix incrassatis; involucro octophyllo, flores subaequante; acheniis muriculatis vel laeviusculis in rostrum eis 3—4 plo brevius attenuatis.

In pascuis subhumidis, juncetis regionis inferioris et montanae. Attica: ad Phalerum (Heldr.); Argolis: pr. Nauplia (Haussk.), Tirynthum (Orph.); Laconia: mt. Malevo (Orph.); Achaia: mt. Panachaicon (Hal.). — Apr. Jun. ⊙.

82. Scorzonera L. gen. n. 986.

a. Achenia glabra.

α. Rhizoma cylindricum.

× Ligulae flavae.

1. **S. crocifolia** S. et S. pr. II. p. 123, Fl. gr. VIII. p. 63 t. 786; Marg. et R. fl. Zante p. 62; Ung. Reise p. 124; Bois. fl. or. III. p. 767; Heldr. fl. cephal. p. 49, Fl. Aegina p. 307; Hal. Beitr. fl. Achaia p. 25, in ö. b. Z. 1895 p. 460, 1897 p. 96; Haussk. symb. p. 138. — *S. graminifolia* Ch. et B. exp. p. 230, Fl. pelop. p. 53; probabiliter etiam S. et S. pr. II. p. 122 quoad pl. Zacynthiam nec non Marg. et R. fl. Zante p. 62; non L. — *S. angustifolia* Friedr. Reise p. 275 et 284, non L. — Exsicc.: Heldr. herb. norm. n. 180 et 1051.

Rhizomate crasso, elongato, verticali, ad collum squamato-lanato; caulibus erectis vel adscendentibus, glabris, simplicibus vel una basi furcatis, superne subnudis; foliis radicalibus numerosis, caespitosis, anguste linearibus, crasse uninerviis, basi dilatatis, duris, late vaginaeformibus, hirtulis, caulinis paucis, abbreviatis; involucri cylindrici phyllis ciliolatis, infimis ovato-lanceolatis, brevibus, intimis lanceolatis, acuminatis; ligulis involucro sesquilongioribus; acheniis eximie muriculatis; pappi sordidi, achenio aequilongi radiis basi dilatatis, liberis, longioribus apice scabris.

In collinis, rupestribus regionis inferioris et montanae. Thessalia: pr. Kalabaka (Haussk.); Sporadum insula Scopelos (Leon.); Attica: mt. Pentelicon, Hymettus (Sprun.), pr. Kephissia (Pichl.), mt. Kerata (Haussk.), mt. Parnes, pr. Bari, Piraeus, Phanari, Laurion, insula Aegina, Lero (Heldr.), Salamis (Reis.), pr. Megara (Orph.); Argolis: pr. Nauplia, Poros (Friedr.), insula Hydra (Heldr.); Achaia: mt. Omplo pr. Patras, pr. Psathopyrgos (Heldr.), Megaspilaeon (Hal.); Arcadia: pr. Zulatika (Nied.); Laconia: mt. Malevo pr. Kastanitza (Orph.); Messenia: Kalamata (Zahn), Messene, Methone, Arcadia (Chaub.); Zante (Sibth.): mt. Scopo (Marg.); Cephalonia: pr. Argostoli (Heldr.); Cycladum insula: Keos (Heldr.), Poros, Naxos (Wiedem.), Thera, Therasia (Letourn.). — Apr. Jun. ♃.

2. **S. parviflora** Jacq. fl. austr. IV. p. 3 t. 305; Haussk. symb. p. 138. —

Rhizomate crassiusculo, praemorso, obliquo, fibras cylindricas crassas edente, ad collum nudo vel vix squamato; caulibus erectis vel adscendentibus, glabris, simplicibus vel parce ramosis, nudiusculis; foliis radicalibus lanceolatis vel lineari-lanceolatis, plurinerviis, basi vix dilatata vaginaeformibus, glabris, caulinis paucis, abbreviatis; involucri breviter cylindrici phyllis glabris, ab infimis triangulari-lanceolatis ad intima lanceolata elongatis; ligulis involucro vix longioribus; acheniis laevibus; pappi albidi, achenio duplo longioris radiis basi in annulum cito secedentem coalitis, longioribus apice scabris.

In nemorosis oropedii Neuropolis in Pindo dolopico (Haussk.). — Maio, Jul. ♃ N. v.

×× Ligulae purpureae.

3. **S. rosea** W. et K. pl. rar. hung. II. p. 127 t. 121. — Exsicc.: Sint. it. thessal. n. 720.

Rhizomate crassiusculo, cylindrico, ad collum fibroso-comato; caule elato, 20—40 cm. alto, foliato, glabro, simplici vel basi ramo uno alterove instructo; foliis lineari-lanceolatis, planis, basi villosulis, ceterum glabris; involucri breviter cylindrici, araneosi, glabrescentis, phyllis ab externis ovato-lanceolatis ad interna lanceolata elongatis; ligulis involucro sesquilongioribus; acheniis superne ad costas scabriusculis, ceterum laevibus, pappo aequilongis; pappi setis longioribus apice scabris. — *S. purpureae* L., in Graecia vix obviae, foliis anguste linearibus et acheniis omnino laevibus tantum diversae, affinis, sed ab ea certe specifice distincta.

In pascuis montanis, hucusque solum ad Said Pascha in Pindo tymphaeo (Sint.). — Jun. Jul. ♃.

4. **S. rhodantha** Haussk. in Mitth. thür. bot. Ver. V. p. 86, symb. p. 139; Bald. riv. coll. bot. Alb. 1895 p. 52, 1896 p. 72, Form. in Ver. Brünn 1897 p. 31. — *S. purpurea* Bois. fl. or. suppl. p. 321; forsan quoque S. et S. pr. II. p. 123 e Creta. — Exsicc.: Heldr. it. gr. septentr. a. 1879; Bald. it. alb. epir. III. n. 167, IV. n. 180.

Rhizomate crasso, tortuoso, ad collum plus minus fibroso-comato; caule pumili, 3—10 cm. alto, nudo vel paucifoliato, glabro vel araneoso, simplici; foliis anguste linearibus, plicatis, saepe recurvis, glabris vel basi villosulis; involucri breviter cylindrici, araneosi vel glabri, phyllis ab externis ovato-lanceolatis ad interna lanceolata elongatis; ligulis involucro longioribus; acheniis superne ad costas vix scabridulis, pappo aequilongis; pappi setis longioribus apice scabris. — Vix specifice a *S. purpurea* distincta.

In aridis alpinis. Epirus: mt. Gamila distr. Zagorion, mt. Peristeri (Bald.); Thessalia: mt. Karava, Ghavellu (Haussk.); Aetolia: mt. Korax (Heldr.). — Jul. Aug. ♃.

β. Rhizoma ovato-tuberosum.

5. **S. mollis** M. a. B. fl. taur. cauc. III p. 522; Bois. fl. or. III p. 761; suppl. p. 320, *v. euboea* (f. caulibus abbreviatis capitulis paucifloris); Bald. riv. coll. bot. Alb. 1896 p. 72 v. *glabrata*; Heldr. fl. Aegina p. 307. — *S. tomentosa* Pall. it. III p. 513 non L. — *Podospermum molle* Fisch. et Mey. ind. petrop. VI p. 21. — *P. villosum* Stev. in DC. pr. VII p. 111; Raul. cret. p. 796. — *S. araneosa* Urv. enum. p. 98, non S. et S. — *S. undulata* Vahl symb. II p. 86, quoad. pl. graec. — *S. ambigua* DC. pr. VII p. 126. — *S. villosa* Friedr. Reise p. 269, non Scop. — Exsicc.: Orph. fl. gr. n. 399; Heldr. herb. norm. n. 371.

Adpresse araneoso-canescens; caulibus brevibus, simplicibus vel una basi ramosis, inferne foliosis, scapiformibus, folia parum superantibus; foliis linearibus, flexuosis, basi dilatatis, vaginantibus; involucri phyllis infimis ovatis, acutis, brevibus, intimis lanceolatis; ligulis luteis, extus rubellis, involucro sesquilongioribus; acheniis sulcatis, externis muricato-squamulosis, internis laevibus; pappi sordidi eis subbrevioris setis longi-

oribus apice scabris. — Radice tuberosa, caulibus scapiformibus, indumento cano et toto habitu a praecedentibus distinctissima.

In aridis, saxosis regionis inferioris et montanae. Epirus: mt. Smolika (Bald.); Attica: in colle Lycabetto (Orph.), Turcovuni, Corydalus, ad Ilyssum, pr. Phaleron, mt. Hymettus, insula Aegina (Heldr.); Euboea: mt. Dirphys (Pichl); Argolis (Sprun.); Cycladum insula Melos (Urv.); Creta: ad Katharos in mt. Lassiti (Raul.); Corcyra: in cacumine mt. Pantocrator (Bickn.). — Mart. Apr. ♃

b. Achenia villosissima.

α. Rhizoma in tuberem ovato-globosum incrassatum.

6. **S. lanata** L. amoen. IV p. 287 (*Leontodon*); M. a. B. fl. taur. cauc. II p. 237; Link in Linn. IX p. 582; Ch. et B. fl. pelop. p. 53; Bois fl. or. III p. 776; Haussk. symb. p. 138; Heldr. fl. Aegina p. 307. — *Tragopogon lanatus* Willd. sp. III p. 1495. — Exsicc.: Orph. fl. gr. n. 260; Heldr. herb. norm. n. 278, in Magn. fl. sel. n. 3313.

Sericeo-lanuginosum; caulibus brevibus, simplicibus vel una basi ramosis, inferne foliosis, scapiformibus, folia parum superantibus; foliis linearibus, flexuosis, basi subdilatatis, vaginantibus; involucri phyllis lanceolatis, externis brevibus vel saepius interna subaequantibus; ligulis luteis, extus rubellis, involucro vix longioribus; acheniis obconicis, basi acutis, pappo rufo quadruplo longiore, plumoso superatis.

In collibus saxosis, locis aridis regionis inferioris et montanae. Attica: in colle Lycabetto (Orph.), mt. Corydalus, Pentelicon, Hymettus, pr. Phanari, in Pharmacusarum insula Lero, insula Aegina (Heldr.); Euboea: mt. Dirphys (Pichl.); Argolis: pr. Nauplia (Link), Astros (Bois.). — Mart. Maio. ♃

β. Rhizoma crassum, cylindricum.

7. **S. cretica** Willd. sp. III p. 1504; S. et S. pr. II p. 123; Sieb. avis. p. 5, rem. p. 5; Raul. cret. p. 797; Bois. fl. or. III p. 779. — Exsicc.: Sieb. pl. cret.; Heldr. pl. cret. a. 1846; Dörfl. pl. cret. n. 67.

Parce lanata vel glabrescens; caulibus crassiusculis, adscendentibus, ramosis, inferne foliosis, superne nudis; foliis linearibus, 5—7 nerviis, saepe elongatis, capitula subaequantibus; capitulis majusculis, sub anthesi 20—25 mm. longis; involucri plus minus lanuginosi phyllis lanceolatis, margine membranaceis, externis brevibus, acuminatis, subpatulis; ligulis flavis, extus saepe rubellis, involucro sesquilongioribus; acheniis oblongis, basi attenuatis, pappo duplo longiore, rufescente, plumoso, apice scabro superatis. — Sequenti valde affinis et ab ea caulibus crassioribus, foliis elongatis latioribus, capitulis majoribus, involucro saepius dense lanuginoso, vix specifice distincta; specimina quidem sicula *S. hirsutae* a Huet du Pavillon pr. Mazzara lecta, cum planta cretica fere omnino congruunt.

In saxosis collinis Cretae: in mt. Sphacioticis (Sibth.) pr. Selia et Strombolo (Heldr.), pr. Parsas distr. Viano (Leon.) — Apr. Maio. ♃

8. **S. hirsuta** L. mant. p. 278; Freyn in bull. herb. Bois. V p. 785. — Icon: Jacq. hort. vind. t. 106. — Exsicc.: Sint. it. thessal. n. 404.

Parce lanata vel glabrescens; caulibus tenuibus, adscendentibus, simplicibus vel parce ramosis, inferne foliosis, superne longe nudis; foliis anguste linearibus, 3—5 nerviis, carinatis, curvatis, sursum valde decrescentibus; capitulis mediocribus, sub anthesi circa 15 mm. longis; involucri glabri phyllis lanceolatis, margine membranaceis, externis brevibus, acutis, subpatulis; ligulis flavis, involucro dimidio longioribus; acheniis oblongis, pappo triplo longiore rufescente, plumoso, apice scabro, superatis.

In glareosis pr. Kalabaka et in vineis pr. Kastreiki Thessaliae (Sint.). — Maio, Jul. ♃

9. **S. Doriae** Deg. et Bald. in ö. b. Z. 1896 p. 417. — *S. hirsuta v. Doriae* Deg. et Bald. riv. coll. bot. Alb. 1896 p. 72. — Exsicc.: Bald. it. alb. epir. IV n. 128.

Differt a praecedente capitulis parvis, sub anthesi 8—10 mm. longis; acheniis obconicis, brevioribus, 5 mm. longis, pappo violaceo sesquilongiore, 7—8 mm. longo superatis. Pappi radii insuper magis distant, ita ut capitulum fructiferum hemisphaericum, *Pterocephalo plumoso* simile (nec oblongum, ut in *S. hirsuta*), pappo violaceo ab acheniis albo-villosis valde discrepante insigne, evadit.

In saxosis alvei Sarandaporos ad Vromonero dist. Ljaskovik et in summo mt. Smolika distr. Konitza in Epiro (Bald.). — Jun. Jul. ♃.

Obs. *S. austriaca* Willd. sp. III p. 1498 in Graeciae montibus et *S. tomentosa* L. sp. ed. 2 p. 1112 in S. et S. pr. II p. 122 in scopulo Caloyero, porro *S. humilis* L. sp. p. 790 in Dallap. prosp. p. 107 in Cephalonia, probabiliter erronee in ditione nostra indicantur.

83. Podospermum DC. fl. fr. IV p. 61.

1. **P. canum** C. A. Mey. Verz. Pflanz. Cauc. und Kasp. Meer. 1831 p. 62; Weiss in z. b. G. 1869 p. 742. — *Scorzonera messeniaca* Ch. et B. exp. p. 231 t. 28, Fl. pelop. p. 53 t. 30. — *S. laciniata* Jacq. en. p. 40, Fl. austr. IV t. 356; S. et S. pr. II p. 124 t. 788; non L. — *P. Jacquinianum* Koch. syn. p. 425; Haussk. symb. p. 139, cum var. *subintegra* Bois. fl. or. III p. 758 (f. humilis, foliis linearibus integris vel laciniis paucis saepe brevibus auctis, manifeste ad typum transiens). — *P. octangulare* DC. pr. VII p. 110. — *S. Jacquiniana* Bois. fl. or. III p. 757. — *S. Jacquiniana v. messeniaca* Hal. Beitr. fl. Achaia p. 25. — *P. pindicolum* Haussk. symb. p. 139 (f. nana, magis caespitosa, caulibus simplicibus). — Exsicc.: Sint. it. thessal. n. 592 et 969; Bald. it. alb. epir. IV n. 129.

Perenne, glabrescens vel plus minus canescens; collo caules floriferos et fasciculos foliorum steriles edente; caulibus erectis vel adscendentibus, ramosis, rarius simplicibus, superne sulcatis; foliis, exceptis

interdum inferioribus anguste linearibus indivisis, pinnatipartitis, laciniis linearibus vel lineari-lanceolatis, terminali elongata; involucro ligulis flavis duplo breviore; acheniis glabris, pappo sublongiore superatis.

β. **alpinum** Bois. fl. or. III. p. 758 pro var. *J. Jacquinianae* (saltem pro p.) — Humile, saepe condensatum; rhizomate plerumque indurato; caulibus saepissime simplicibus, monocephalis; foliorum laciniis oblongis suborbiculatisve, terminali majore, subrotunda vel lanceolato-elongata. — Formis innumeris ad typum transit. — Exsicc.: Heldr. herb. norm. n. 273; Orph. fl. gr. n. 371 (f. *transiens*); Dörfl. fl. aeg. n. 53.

In collibus, herbidis regionis inferioris, in alpinam usque adscendens, varietas et praesertim formae intermediae typo multo vulgatiores — Epirus: mt. Nimercka (Bald.), jam extra ditionem nostram; Thessalia: mt. Ghavellu, Karava, pr. Malakasi, Kalabaka (Haussk.); Aetolia: mt. Tymphrestus (Neumayer); mt. Parnassus (Guicc.); Attica: mt. Pateras, pr. Eleusis, Athenas, in colle Philopappo, pr. Kephissia, Tatoi, mt. Parnes, Pentelicon, Hymettus, pr. Trakones, Phaleron, Piraeus, Megara (Heldr.); Achaia: mt. Olenos, Panachaicon, in collibus arenosis pr. Patras, mt. Chelmos (Hal.), Kyllene (Orph.); Arcadia: pr. Leontari, Zulatika (Nied.); Argolis: pr. Nauplia (Orph.), Torniki (Heldr.); Laconia: mt. Taygetos (Sart.); Messenia: a Pylos usque Arcadia (Chaub.); Cycladum insula Tenos (Weiss), Syra (Orph.). — Mart. Jul. ♃.

2. **P. laciniatum** L. sp. p. 791; Willd. sp. III. p. 1506; Bois. fl. or. III. p. 757 (*Scorzonera*); DC. fl. fr. IV. p. 62; Link in Linnaea IX. p. 582; Weiss in z. b. G. 1869 p. 45. — *S. octangularis* Willd. l. c. — Icon: Rchb. t. 34.

Bienne, glabrescens vel canescens; collo fasciculis sterilibus nullis; caulibus adscendentibus vel procumbentibus, ramosis, minute striatis; foliis pinnatipartitis, laciniis linearibus vel lanceolatis, terminali majore, involucro ligulis flavis duplo breviore, acheniis glabris, pappum aequantibus. — Polymorpha, uti praecedens.

β. **calcitrapifolia** Vahl symb. II. p. 87; Urv. enum. p. 99; Friedr. Reise p. 275; (*Scorzonera*) pro sp.; Bois. fl. or. III. p. 757 pro var. *S. laciniatae*. — *S. resedifolia* Retz. obs. III. p. 42; Ch. et B. fl. pelop. p. 54; Fraas fl. class. p. 197; non L. — *P. calcitrapifolium* DC. fl. fr. V. p. 455; Haussk. symb. p. 139. — Foliorum laciniae lanceolatae, ellipticae, oblongae vel obovatae.

In herbidis regionis inferioris. Boeotia: pr. Oropos (Bois.); Attica: pr. Eleusis (Haussk.), Athenas, Megara (Bois.); Euboea (Fraas); Acrocorinthus (Haussk.); Argolis pr. Nauplia (Link), insula Poros (Friedr.); Cycladum insula Syra, Tenos (Weiss), Melos (Urv.); Laconia et Messenia (Chaub.); sed loca nonnulla probabiliter ad speciem praecedentem spectant. — Maio, Jul. ⊙. N. v.

20. Tribus **CHONDRILLEAE** Koch. syn. p. 427.

84. **Taraxacum** Hall. hist. stirp. helv. I. p. 23.

a. Folia subcoriacea, rigidula, laciniarum dentibus mucrone albo cartilagineo terminatis.

1. **T. Haussknechtii** Uechtr. in Nym. consp. suppl. II. p. 190 solum nomen; Haussk. symb. p. 132. — *Leontodon serotinus* S. et S. pr. II. p. 129, Fl. gr. VIII. p. 71 t. 796, non W. et K. — *L. taraxacum v. laevigatum* Bois. fl. or. III. p. 789 p. p. — *T. gymnanthum* Hal. in ö. b. Z. 1890 p. 39, non (Link). — *T. leptocephalum* Hal. in ö. b. Z. 1892 p. 371, non Rchb. — Forsan huc: *T. vulgare v. pindicolum* Bald. riv. coll. bot. Alb. 1896 p. 73. — Exsicc.: Sint. it. or. a. 1889 n. 1828; Sint. et Bornm. it. turc. n. 1333.

Rhizomate crasso, saepe pluricipiti, collo inter folia lanuginoso; foliis synanthiis, rosulatis, terrae adpressis, glaucovirentibus, araneoso-puberulis, demum glabrescentibus, oblongis, vel oblongo-lanceolatis, runcinato-pinnatifidis, laciniis, minoribus interjectis, triangularibus; scapis floccoso-araneosis, demum glabrescentibus, monocephalis; involucri phyllis glabris vel ciliolatis, lividis, externis triangulari-lanceolatis, subadpressis, internis 2—3plo longioribus, lineari-lanceolatis; ligulis pallide flavis; acheniis pallidis, superne muricatis, in rostrum subaequilongum vel eis dimidio longius abeuntibus. — Ex affinitate *T. serotini* (W. et K.) Poir., cujus folia sunt majora, subintegra vel runcinata, sed tunc laciniis multo latioribus, scapi crassiores, capitula majora, achenia longiora. — Planta nostra cum icone Sibthorpii optime congruit.

In arenosis, herbidis regionis inferioris et montanae. Epirus: mt. Smolika (Bald.); Thessalia: pr. Malakasi in Pindo tymphaeo (Haussk.), pr. Litochori et ad Hagios Dionysios in Olympo (Sint.). — Jul. Sept. ♃.

b. Folia tenera, laciniarum dentibus mucrone molli terminatis.

α. Folia synanthia.

2. **T. officinale** Wigg. prim. fl. Holsat. p. 56; Bois. fl. or. III. p. 787; Form. in D. bot. Monat. 1890 p. 15. — *Leontodon taraxacum* L. sp. p. 798; S. et S. pr. II. p. 129; Dallap. prosp. p. 108; Ch. et B. exp. p. 232, Fl. pelop. p. 54; Fraas fl. class. p. 201. — *T. vulgare* Lam. fl. fr. II. p. 113. —

Glabrum vel subglabrum; rhizomate crasso, saepe pluricipiti; foliis rosulatis, oblongis vel oblongo-lanceolatis, runcinato-pinnatifidis, laciniis, denticulatis, interdum minoribus interjectis, triangularibus vel triangulari-lanceolatis; scapis monocephalis; involucri phyllis virescentibus, ecorniculatis, externis linearibus vel ovatis, adpressis vel reflexis, internis linearibus 2—3plo longioribus; ligulis flavis; acheniis pallide olivaceis, superne muricatis, in rostrum eis longius abeuntibas. — Stirps polymorpha.

α. **typicum.** — Folia profunde lobata vel partita; involucri phylla omnia linearia vel exteriora oblonga vel ovato-oblonga, deflexa; flores plerumque aurei. — Exsicc.: Sint. it. thessal. n. 182.

β. **alpinum** Hoppe in Sturm Heft 41 (*Leontodon*) pro sp.; Koch. syn. p. 428; Bois. fl. or. III. p. 788; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 49. — *T. officinale v. alpestre* Ung. Reise p. 125. — *T. vulgare v. alpinum* Bald. riv. coll. bot. alb. 1896 p. 73. — *T. Steveni* Hal. Beitr. fl. Epir. p. 29, quod phyllis corniculatis discedit. — *T. vulgare v. alpinum et Steveni* Bald. riv. coll. bot. Alb. 1896 p. 73 et 74. — Scapi humiles 5—20 cm. alti; involucri phylla sicca saepe nigrescentia, externa ovata, patentia vel deflexa. — Intermediis ad typum transit. — Exsicc.: Bald. it. alb. epir. IV. n. 246.

γ. **paludosum** Scop. fl. carn. ed. 2, II. p. 100 t. 48 (*Hedypnois*) pro sp. — *Leontondon palustre* Huds. fl. angl. ed. 2 p. 339. — *L. lividus* W. et K. pl. rar. hung. II. p. 120, t. 115; Friedr. Reise p. 263. — *T. palustre* DC. fl. fr. IV. p. 45; Haussk. symb. p. 133. — Folia anguste oblongo-lanceolata vel lanceolato-linearia, subintegra, sinuata, rarius pinnatifida; involucri phylla omnia adpressa, externa late ovata. — Exsicc.: N. v.

In pascuis regionis montanae in alpinam adscendens. Epirus: mt. Nimercka, Gamila, Smolika (Bald.), Peristeri, Tsumerka (Hal.); Thessalia: mt. Tringia (Hartl), pr. Kalabaka (Sint.), mt. Olympus, Pelion (Heldr.), Ossa (Form.); Aetolia: mt. Korax (Tunt.); Attica: mt. Parnes, Hymettus (Orph.); Euboea: mt. Dirphys (Heldr.); Argolis (Fraas); Achaia: mt. Panachaicon (Hal.), Kyllene (Heldr.); mt. Taygetos (Chaub.); Arcadia (Sibth.); Creta: mt. Sphaciotici (Bois.); Cephalonia: mt. Aenos (Ung.); — *γ.* in humidis mt. Ghavellu in Pindo (Haussk.); Corcyra (Friedr.). — Febr. in alpinis usque Aug. ♃.

3. **T. laevigatum** Willd. sp. pl. III. p. 1546; S. et S. pr. II. p. 129; (*Leontodon*); DC. pr. VII. p. 146; Raul. cret. p. 798 *v. polycephala*; Hal. Beitr. Fl. Epir. p. 29, Beitr. fl. thessal. p. 15; Haussk. symb. p. 133; Heldr. fl. Aegina p. 306 *f. stenolobum* (f. scapis decumbentibus foliis brevioribus, foliorum laciniis crebris anguste linearilanceolatis), chlor. Mykon. p. 246. — *L. taraxacoides* Hoppe in Sturm Heft 41; Friedr. Reise p. 267. — Exsicc.: Heldr. herb. norm. n. 951; Sint. it thessal. n. 735. —

Glabrum; rhizomate crasso, saepe pluricipiti; foliis rosulatis, oblongis vel oblongo-lanceolatis, runcinato-pinnatifidis, laciniis denticulatis, saepius minoribus interjectis, triangularibus vel triangulari-lanceolatis, acuminatis; scapis monocephalis; involucri phyllis virescentibus, apice corniculatis, externis lanceolatis, patulis vel reflexis, intimis linearibus 2—3 plo longioribus; ligulis pallide flavis; acheniis griseis, lateritiis vel purpurascentibus (*T. erythrospermum* Andrz. in Bess. en. volh. p. 75), superne muricatis, in rostrum eis longius abeuntibus. — Differt a praecedente statura humiliore, capitulis minoribus; involucri phyllis cornicu-

latis et floribus pallidis. — Species ceteroqui, quoad foliorum formam scaporum longitudinem, acheniorum colorem etc. uti istud variabile.

β. **scolopendrinum** Heldr. in Nym. syll. suppl. II. p. 190 pro sp., nomen solum. — Folia lanceolata, laciniis integerrimis, retrorsum versis, omnibus aequalibus, minoribus non interjectis. — Forma singularis. — Exsicc.: Heldr. in Baen. herb. europ. n. 4216.

In herbidis regionis inferioris in alpinam usque adscendens. Epirus: mt. Tsumerka, Peristeri (Hal.); Thessalia: mt. Tringia (Hartl.), Karava (Haussk.), pr. Pheras (Heldr.); Attica: pr. Athenas (Friedr.), mt. Parnes, pr. Menidi, Amarysia, Kephissia, mt. Pentelicon, Hymettus, ad promontorium Colias, pr. Ergastiria Laurii, insula Aegina (Heldr.); Petalium insula Megalonisi et Xeronisi (Holzm.); Cycladum insula Melos (Leon.); Achaia: mt. Chelmos pr. Sudena (Hal.); Laconia (Sibth.); Creta: ad Askyphos, Niato (Raul.). — Mart. Apr., in alpinis usque Aug. ♃.

4. **T. minimum** Brig. in Guss. syn. II. p. 397 (*Leontodon*); Tod. fl. sic. exs. n. 688; Haussk. symb. p. 136; Heldr. fl. Aegina p. 306. — *T. humifusum* Heldr. et Sart. in sched. sec. Haussk. l. c. — *T. gymnanthum* Heldr. herb. norm. n. 853 b, forma foliis synanthiis, non Link.

Differt a praecedente foliis primis oblongo-obovatis, denticulatis, reliquis runcinato-pinnatifidis, laciniis ovatis integris denticulatisve acutiusculis, involucri phyllis externis ovatis, adpressis et praesertim inflorescentia autumnali.

In montosis. Achaia: mt. Omplo pr. Patras (Heldr.); Attica: ad radices mt. Hymettus versus Caesariani, mt. Parnes, pr. Daphni (Heldr.), insula Aegina (Tunt.); Cycladum insula Keos (Tunt.); et probabiliter alibi. — Oct. Nov. ♃.

β. Folia hysteranthia.

5. **T. gymnanthum** Link in Linn. IX. p. 582 (*Leontodon*); DC. pr. VII. p. 145; Clem. sert. p. 53; Bois. fl. or. III. p. 788; Heldr. chlor. Mykon. p. 246, prosth. chlor. Thera p. 3. — Exsicc.: Orph. fl. gr. n. 87; Heldr. herb. norm. n. 853 a (mixtum cum T. minimo).

Rhizomate crasso, saepe pluricipiti, collo inter folia lanuginoso; foliis rosulatis, subtus araneosis, demum glabriusculis, oblongo-lanceolatis, in lacinias triangulares vel ovatas, acutas vel obtusas, subintegras sublyrato-pinnatipartitis; scapis araneosis, monocephalis; involucri glabri, lividi, phyllis adpressis, externis ovato-oblongis, saepe corniculatis, intimis linearibus 2—3 plo longioribus; ligulis pallide flavis, extus rubentibus; acheniis fuscidulis, superne muricatis, in rostrum eis longius abeuntibus.

In collibus siccis, locis aridis, ad vias regionis inferioris. Attica: pr. Athenas (Orph.), Daphni, Kephissia, Tatoi (Heldr.), ad radices Hymetti pr. Caesariani (Clem.); Argolis: ad ripas Erasini pr. Argos (Heldr.), pr. Nauplia, Poros (Link); Cycladum insula: Naxos (Leon.), Delos, Rhenea (Tunt.), Thera (Heldr.). — Sept. Nov. ♃.

85. Chondrilla L. gen. n. 910.

1. **C. ramosissima** S. et S. pr. II p. 128, Fl. gr. VIII p. 70 t. 795; Urv. en. p. 100; Link in Linnaea IX p. 582; Ch. et B. fl. pelop. p. 54; Fraas fl. class. p. 199; Clem. sert. p. 53; Raul cret. p. 798; Bois. fl. or. III p. 791; Haussk. symb. p. 132; Heldr. fl. Aegina p. 306; Form. in D. bot. Mon. 1898 p. 77. — *Prenanthes attica* Spreng. syst. III p. 654. — Exsicc.: Heldr. herb. norm. n. 436 et 1450; Orph. fl. gr. n. 757.

Caule divaricato-ramosissimo, angulato-sulcato, cano, setulis rigidis scabro; foliis radicalibus et inferioribus cito evanidis, glabris, oblongo-spathulatis, runcinato-dentatis, caulinis minimis, oblongo-linearibus, spinuloso-ciliatis, cito emarcidis; capitulis lateralibus, sessilibus, 2—5 glomeratis vel solitariis, paucifloris; involucri cylindrici, cani, phyllis extimis minimis, squamaeformibus, ceteris linearibus, dorso setosis; ligulis flavis; acheniis a medio sursum aculeolatis, in rostrum tenuissimum eis sesquilongius abeuntibus.

In campis, arvis, locis incultis regionis inferioris. Boeotia (Fraas); vulgatissima circa Athenas, nec non in planitia Attica tota; insula Aegina (Heldr.); in Peloponneso (Link) et Creta (Raul.), sine loci specialis indicatione. — Jun. Sept. ⊙

2. **C. juncea** L. sp. p. 796; S. et S. pr. II p. 127; Sieb. avis p. 4; Friedr. Reise p. 273 et 284; Weiss in z. b. G. 1869 p. 45; Bois. fl. or. III p. 792; Heldr. fl. cephal. p. 49, chlor. Thera p. 15; Form. in D. bot. Monat. 1890 p. 15, in Ver. Brünn 1895 p. 25, 1896 p. 42, 1897 p. 31; Bald. in nuov. giorn. bot. 1894 p. 98; Haussk. symb. p. 131; Hal. in ö. b. Z. 1895 p. 460. — Icon: Fl. dan. t. 1652. — Exsicc.: Sint. et Bornm. it. turc. n. 1336; Sint. it. thessal. n. 1286.

Caule longe et virgatim ramosissimo, teretiusculo, glaucescente, inferne setoso-hispido, superne glabro; foliis radicalibus cito evanidis, glabris, oblongo-spathulatis, runcinato-dentatis, caulinis parvis, lanceolatis vel linearibus, integris vel denticulatis; capitulis lateralibus et terminalibus, solitariis vel 2—5 nis, subsessilibus, paucifloris; involucri cylindrici, glaucescentis, phyllis extimis minimis, squamaeformibus, ceteris linearibus, minutissime puberulis; ligulis flavis; acheniis superne muriculatis, in rostrum tenuissimum eis sesquilongius abeuntibus.

β. **acantholepis** Bois. diagn. XI p. 48 pro sp.; Bois fl. or. III p. 792. — Involucri phylla secus nervum setoso-aculeata. — Exsicc.: N. v.

In vineis, arenosis, ad vias regionis inferioris. Epirus: pr. Janina, Han Balduma, mt. Prosgoli (Form.); Thessalia: frequens a Pindo usque ad mare Aegaeum; Aetolia: pr. Mesolongion (Nied.); Attica: pr. Tatoi, Kephissia (Heldr.); Achaia: pr. Patras (Reinh.); Argolis: pr. Nauplia (Sart.), Poros (Friedr.); Cyclades (Sibth.): Andros (Sart.), Naxos (Philoti), Thera (Heldr.); Creta: pr. Canea (Weiss); Cephalonia: pr. Pessada (Heldr.); Corcyra (Bald.): pr. urbem, Kanali, Potamo, Kontokali (Form.). β. Attica: pr. Kephissia (Heldr.). — Jul. Sept. ⊙

Obs. *C. capitata* Sieb. in Flora V, 2 p. 639; sec. DC. pr. VII p. 180 = *Lomatolepis glomerata* Cass., in Creta erronee indicatur.

21 Tribus. **CREPIDEAE** Bois. fl. or. III p. 712.

86. Sonchus L. gen. n. 908.

a. Perennes.

1. **S. palustris** L. sp. p. 793; S. et S. pr. II p. 124; Ch. et B. exp. p. 231, Fl. pelop. p. 54; Bois. fl. or. III p. 798. — Icon: Fl. dan. t. 1109.

Rhizomate crasso, subtuberoso, non repente; caule elato, 1—2 m. alto, crasso, apice fastigiatim ramoso, superne glanduloso-piloso, ceterum glabro; foliis glabris, spinuloso-denticulatis, inferioribus in lacinias 1—2 retrorsas, late lanceolatas runcinato-pinnatifidis, mediis lanceolatis, sagittato-amplexicaulibus, summis lineari-lanceolatis, semiamplexicaulibus; capitulis numerosis, cymosis; involucri nigricantis, dense glanduloso-pilosi, phyllis lanceolato-linearibus; acheniis testaceis, oblongo-linearibus, subcompressis, utrinque crasse 5 costatis, transverse tenuissime rugulosis.

In Elidis paludibus (Sibth.); postea a nemini lectum. — Jul. Sept. ♃ N. v.

2. **S. maritimus** L. sp. ed. 2 p. 1116; S. et S. pr. II p. 125; Friedr. Reise p. 275, Raul. cret. p. 800. — Icon: Rchb. t. 62.

Rhizomate tenui, repente; caule erecto, 30—100 cm. alto, simplici, vel apice parce ramoso, glabro; foliis glabris, longe et anguste lanceolato-linearibus, rarius parce sinuato-denticulatis, infimis in petiolum vaginantem attenuatis, superioribus breviter auriculato-amplexicaulibus; capitulis paucis, solitariis vel cymosis; involucri nigricantis, glabri, phyllis extimis oblongis, intimis oblongo-lanceolatis; acheniis testaceis, linearibus, subcompressis, utrinque tricostatis, transverse tenuissime rugulosis.

In humidis regionis litoralis. Insula Poros (Friedr.); Creta (Sibth.); a recentioribus ut videtur non lectum. Jul. Aug. ♃ N. v.

b. Annui vel biennes, raro (f. perennis *S. tenerrimi*) perennes sed tunc folia fere omnia petiolata, pinnatipartita.

α. Achenia transverse rugulosa.

3. **S. tenerrimus** L. sp. p. 794; S. et S. pr. II p. 125, Fl. gr. t. 790; Friedr. Reise p. 270 et 283, cum v. *spinulosa*; Clem. sert. p. 49 v. *muriculatus* (anthodiis praesertim inferne muriculatis; Ung. Reise p. 125; Raul. cret. p. 800; Bois. fl. or. III p. 797, suppl. p. 321; Hal. in z. b. G. 1888 p. 763; Haussk. symb. p. 132; Heldr. fl. Aegina p. 306, chlor. Thera p. 15. — *S. lacerus* Sieb. avis rem. p. 5, sec. Raul. cret. p. 800. non Willd. sp. III p. 1503. —

Biennis vel perennans; radice fusiformi; caule erecto, glabro, ramoso; foliis glabris, fere omnibus petiolatis, pinnatisectis vel pinnatipartitis,

87. Prenanthes L. gen. n. 911.

1. **P. purpurea** L. sp. p. 797; Bois. fl. or. III. p. 803. — Icon: Jacq. fl. austr. IV. t. 317. — Exsicc.: Orph. herb. a. 1857.

Glabra; caule erecto, ramosissime et ample paniculato; foliis inferioribus ovatis oblongisve, dentatis vel sinuatis, in petiolum alatum contractis, superioribus a basi rotundata auriculata, oblongis vel lanceolatis, integris; capitulis cernuis, tenuiter cylindricis; acheniis oblongo-linearibus, subcompressis, striatis, pappo albo brevioribus.

In regione silvatica mt. Olympus Thessaliae (Orph.). — Jul. Aug. ♃.

88. Lactuca L. gen. n. 909.

1. Sectio. *Scariola* DC. pr. VII. p. 133. — Capitula pluriflora, involucro imbricato; achenia abrupte rostrata.

a. Achenia 1 vel 3 nervia.

1. **L. cretica** Desf. ann. mus. XI. p. 160, t. 19, cor. Tourn. 44 t. 34; Urv. en. p. 100; Ch. et B. exp. p. 232, Fl. pelop. p. 54; Weiss in z. b. G. 1869 p. 45; Raul. cret. p. 798; Bois. fl. or. III. p. 305; Haussk. symb. p. 132; Heldr. fl. Aegina p. 306, chlor. Thera p. 15. — *L. leucophaea* S. et S. pr. II. p. 127, Fl. gr. VIII. p. 69 t. 794. — *L. sonchifolia* Sieb. avis rem. p. 5; Raul. cret. p. 798; non Willd. sp. III. p. 1530, quae flosculis coeruleo-purpurascentibus differt. — Exsicc.: Heldr. pl. gr. n. 3090, herb. norm. n. 1048; Sint. it. thessal. n. 538.

Glabra, glaucescens; rhizomate crasso, subtuberoso; caule crasso, erecto, fistuloso, simplici, in paniculam terminalem abeunte; foliis oblongo-lanceoatis, in lacinias ovatas vel oblongas, dentatas, runcinato-pinnatifidis, infimis in petiolum attenuatis, ceteris cordato-auriculatis, non decurrentibus, floralibus squamaeformibus, lanceolatis, integris; pedunculis patentibus, squamulosis, brevibus, capitulis solitariis magnis, 1 cm latis, ovato-cylindricis, multifloris; involucri phyllis nigricantibus, ab infimis ovatis ad superiora late linearia sensim auctis; flosculis luteis, acheniis nigris, ovatis, valde compressis, abrupte mucronulatis, latiuscule marginatis, subtrinerviis, minutissime rugulosis, rostro albo eis parum usque 3 plo longiore superatis.

In collibus, olivetis, fruticetis regionis inferioris et montanae. Thessalia: pr. Kalabaka (Sint.); Phocis: mt. Parnassus pr. Pagna (Guicc.); Attica: mt. Cithaeron, pr. Eleusis, mt. Parnes, Pentelicon, Hymettus (Heldr.), Lycabettus (Orph.), Phaleron (Haussk.), Pharmacusarum insula Lero et Mikrakyra, pr. Leopesi, Laurion; insula Aegina, in isthmo Corinthiaco pr. Lutraki (Heldr.); Argolis: in peninsula Methana (Haussk.), in collibus litoris oppido Poros oppositis (Heldr.); Messenia: mt. Ithome (Heldr.), pr. Methone, Kardamyle, Mavrovuni (Chaub.); pr. Megali-Anastasova distr. Alagoniae (Zahn.); Cycladum insula Syra (Weiss), Melos, Thera (Urv.); Crete (Tourn.). — Apr. Jun. ♃.

2. **L. graeca** Bois. fl. or. III, p. 812; Heldr. chlor. Parn. p. 22. — *L. intricata* Haussk. symb. p. 132, an Bois. diagn. IV. p. 27, quae sec. auctorem acheniis minoribus, minus acutis, non marginatis, subtrinerviis, sed meo sensu vix specifice differt. — *L. tenerrima* Bald. riv. coll. bot. Alb. 1895 p. 641, 1896 p. 74, non Pourr. act. Toul. III. p. 321, quae praesertim acheniis nigris obovatis a *L. graeca* specifice discedit. — Exsicc.: Heldr. fl. gr. a. 1855, herb. n. 1488; Orph. fl. gr. n. 766; Haussk. it. gr. n. 1885; Bald. it. alb. alt. n. 94.

Glaucescens; rhizomate crasso, multicipiti; caulibus gracilibus, erectis, fistulosis, glabris, a basi divaricatim ramosis, ramis virgatis, monocephalis; foliis subtus crispule puberulis, radicalibus petiolo basi dilatato suffultis, oblongis, sublyratis vel pinnatipartitis, laciniis linearibus, integris dentatisve, caulinis paucis, minutis, linearibus, integris, basi auriculatis, non decurrentibus; pedunculis elongatis; capitulis solitariis, mediocribus, 3—4 mm. latis, cylindricis, plurifloris; involucri glabri phyllis caesiis, ab infimis ovatis ad superiora linearia, auctis; flosculis coeruleis, acheniis pallide rufescentibus, elliptico-linearibus, compressis, apice acutatis, tumidulo marginatis, subtrinerviis, minutissime rugulosis, rostro albo, eis aequilongo vel breviore superatis.

In saxosis, glareosis regionis mediae et superioris. Epirus: ad radices mt. Peristeri supra Chaliki (Haussk.), mt. Konitza, nec non extra ditionem nostram in mt. Cika in mt. Acroceraunicis (Bald.); Phocis: mt. Parnassus in summo jugo Mavrolithari dicto (Guicc.); Thessalia: mt. Olympus supra Hagios Dionysios (Heldr.). — Jul. Aug. ♃.

b. Achenia 5—9 nervia.

α. Caulis fistulosus, viridis; rostrum acheniorum nigrum.

3. **L. Chaixi** Vill. prosp. p. 33, hist. pl. Dauph. III. p. 155 t. 32. — *L. sagittata* W. et K. pl. rar. hung. I. p. 1 t. 1; Form. in Ver. Brünn 1895 p. 25. —

Glabra, radice napiformi; caule erecto, simplici, superne paniculato-corymboso; foliis radicalibus in petiolum attenuatis, sinuato-dentatis, caulinis a basi acute sagittata oblongis, denticulatis, non decurrentibus summis ovato-lanceolatis, integris; capitulis cylindricis, 8—13 floris; involucri phyllis ab externis ovatis, ad interna linearia sensim auctis; flosculis pallide luteis; acheniis oblongis, nigris, compressis, utrinque 5striatis, in rostrum concolor eis dimidio brevius attenuatis.

In nemorosis. Epirus: pr. Safikbi et Han Balduma (Form.). — Jun. Aug. ⊙. N. v.

β. Caulis solidus, albicans; rostrum acheniorum album.

× Folia ovali-oblonga, mucronato-denticulata, runcinato-pinnatifida vel indivisa; capitula paniculam amplam formantia; rostrum achenio aequilongum vel brevius.

4. **L. virosa** L. sp. p. 795 excl. β. γ. δ.; L. sp. pl. ed. 2. p, 1119; Form. in D. bot. Mon. 1890 p. 15, in Ver. Brünn 1895 p. 25. 1196, p. 42, 1897 p. 31. — Icon. Rchb. t. 71.

Caule elato, saepe ramoso, inferne interdum parce aculeolato; foliis horizontalibus, glauco-virentibus, subtus ad nervum venasque aculeolato-setosis, radicalibus obovatis, petiolatis, caulinis a basi sagittata subcordata ovato-oblongis, indivisis vel sinuatis, rarius subruncinato-lobatis, non decurrentibus; capitulis cylindricis; involucri glabri phyllis obtusiusculis, sensim auctis; flosculis sulfureis; acheniis ellipticis, nigris, compressis, utrinque 5—7 costatis, glabris, apice margineque scabris, in rostrum eis subaequilongum attenuatis.

In rupestribus, fruticetis regionis inferioris et montanae. Thessalia: pr. Trikala, Karditza, Povelci, Larissa, Velestinos, Lechonia, Volo (Form.); Corcyra: pr. urbem et pr. Alipu (Form.). — Jul. Sept. ⊙ N. v.

5. **L. scariola** L. amoen. acad. IV p. 328 (*L. serriola*); L. sp. ed. 2 p. 1119; S. et S. pr. II p. 126; Fraas fl. class. p. 199; Clem. sert. p. 53; Bois. fl. or. III p. 809; Heldr. fl. cephal. p. 49; Form. in D. bot. Monat. 1890 p. 15, in Ver. Brünn 1895 p. 25, 1896 p. 41, 1897 p. 31; Haussk. symb. p. 132; Bald. viagg. cret. p. 71. — Icon: Rchb. t. 70.

Caule elato, superne ramoso, inferne interdum aculeolato; foliis erectis, glaucis, subtus ad nervum medium aculeolatis vel inermibus, ovali-oblongis, runcinato-pinnatifidis, caulinis basi sagittato-amplexicaulibus, non decurrentibus; capitulis cylindricis; involucri glabri phyllis obtusiusculis, sensim auctis; flosculis citrinis; acheniis ellipticis, spadiceis, compressis, utrinque 7—9 costatis, apice setuloso-puberulis, in rostrum eis aequilongum attenuatis. — Foliis erectis, plerumque runcinato-pinnatifidis, acheniis apice setuloso-puberulis a praecedente diversa.

β. **coriacea** Schultz in Linnaea XV. p. 725, in Flora XXV. p. 174; Fraas fl. class. p. 200. — *L. albicaulis* Bois. mss. ex Fl. or. III. p. 809. — Caulibus usque ad inflorescentiam foliisque crebre aculeolatis. — Exsicc.: Bald. it. cret. n. 129.

γ. **angustana** All. fl. ped. II. p. 224 t. 52; Friedr. Reise p. 273; pro sp. — Foliis omnibus indivisis. —

In cultis, dumetis, ad sepes regionis inferioris et montanae. Epirus: pr. Janina, Katschka, mt. Micikeli et Prosgoli (Form.); Thessalia: pr. Koturi, Said Pascha, Kalabaka, Trikala, mt. Chassia, Demirli, Hadsi Obasi, Larissa, Tyrnavos, Tafilvris, Rapsani, Pharsalus, Velestinos, Volo, Stylida, Megalivris, Lamia (Form.); Euboea: pr. Kurbatzi (Willd.); Attica: pr. Tatoi (Haussk.), in valle Cephissi (Heldr.), in Acropoli Athenarum (Clem.); Elis: pr. Olympia (Nied.); Argolis (Fraas); Cycladum insula Keos (Heldr.), Kythnos (Tunt.); Creta: pr. Suda (Bald.); Cephalonia: pr. Argostoli, Omala (Heldr.); Corcyra: pr. Kanali (Form.); — *γ.* Argolis: pr. Poros (Friedr.). — Typum et var. *γ.* non vidi. — Jul. Sept. ⊙

Obs. *L. sativa* L. sp. p. 795; Dallap. prosp. p. 107; Marg. et R. fl. Zante p. 85; Friedr. Reise p. 275; Fraas fl. class. p. 199;

Heldr. Nutzpfl. p. 28, Fl. cephal. p. 48, chlor. Thera p. 29. — Colitur ad usum culinarem.

×× Folia elongato-linearia, integerrima, infima tantum runcinato-pinnatifida (laciniis integris vel denticulatis); capitula racemoso-spicata; rostrum achenio aequilongum vel duplo longius.

6. **L. saligna** L. sp. p. 796; Marg. et R. fl. Zante p. 69; Friedr. Reise p. 284; Schultz in Flora XXV Beibl. p. 160; Bois. fl. or. III p. 810; Heldr. fl. cephal. p. 49, Fl. Aegina p. 306, chlor. Mykon. p. 246; Form. in D. bot. Monat. p. 15, in Ver. Brünn 1896 p. 41, 1897 p. 31; Haussk. symb. p. 132; Bald. viagg. Creta p. 72. — Huc vel ad praecedentem spectat: *L. Tommasiniana* Schultz in Flora XXV Beibl. p. 160, ex Atticae locis humidis a Fraas lecta („foliis carina laevibus, caulinis oblongo-lanceolatis, omnibus runcinatis, ramis linearibus subintegris"). — Icon: Jacq. fl. austr. III t. 250. — Exsicc.: Sint. it. thessal. n. 1071.

Caule erecto, glabro, simplici vel ramoso, ramis elongatis, virgatis; foliis inermibus vel ad nervum aculeolato-setosis, infimis in petiolum attenuatis, caulinis basi sagittatis, non decurrentibus; capitulis cylindricis, brevissime pedicellatis, fasciculatis vel solitariis; involucri glabri phyllis externis ovatis, intimis linearibus; flosculis luteis; acheniis brunneis, obovato-oblongis, utrinque 5—7 striatis, apice scabridis.

In cultis, ruderatis, ad sepes regionis inferioris et submontanae. Thessalia: pr. Velitsena, Malakasi, Han Kriavris in valle Penei, Kalabaka, Karditza, Trikala, Pharsalus, Demirli, Plessidi, Lechonia, Volo (Form.); Eurytania: pr. Mikrochorio (Heldr.); Attica: pr. Tatoi (Haussk.), mt. Hymettus (Hal.), pr. Caesariani, Heracleon, Phaleron, insula Aegina (Heldr.); Argolis: pr. Nauplia (Friedr.); Laconia: mt. Malevo (Orph.); Cycladum insula Keos, Rhenea (Tunt.); Creta: mt. Ida (Bald.); Zante: pr. Agria (Marg.); Cephalonia: pr. Same (Heldr.). — Jul. Oct. ⊙.

2. Sectio. *Phaenixopus* Cass. dict. 39 p. 391. — Capitula 5flora, involucro imbricato; achenia in rostrum sensim attenuata.

a. Folia caulina non decurrentia.

7. **L. acanthifolia** Willd. sp. III. p. 1542; Sieb. avis p. 5; (*Prenanthes*); Bois. fl. or. III. p. 818. — *Phaenixopus acanthifolius* Cass. dict. 48 p. 426. — *Phaenopus acanthifolius* DC. pr. VII. p. 176; Raul. cret. p. 799. — ? *Prenanthes aconitifolia* Sieb. in Flora I. p. 271. —

Caule erecto, simplici?, superne nudo, densiuscule spicato; foliis amplis, inferne secus costam latam arachnoideis, ceterum glabris, inferioribus late oblongo-spathulatis, in petiolum brevem attenuatis, grosse lobatis, basi sublyratis, caulinis amplexicaulibus pinnatipartitis, laciniis oblongo-triangularibus, dentatis; capitulis sessilibus, cylindricis, in fasciculos sessiles congestis, spicam longam formantibus; involucri phyllis 5—6, inferioribus ovatis, internis linearibus; flosculis?; acheniis lineari-

lanceolatis, compressis, scabridulis, utrinque 6—7 striatis, in rostrum eis subdimidio brevius sensim attenuatis. — Species incomplete nota; folia fere *Acanthi mollis*, pedalia, flores magnitudine *Prenanthis purpureae*. —

Creta (Tourn. cor. 35): in collibus ad promontorium Meleka (Sieb.). — Febr. N. v.

8. **L. amorgina** Heldr. et Orph. in ö. b. Z. 1898 p. 184 nomen solum; Hal. in z. b. G. 1899 p. 188. — Exsicc.: Heldr. pl. fl. hellen. a. 1897.

Caule erecto, simplici, fistuloso, collo lanato-villoso excepto glabro, superne nudo; foliis glaberrimis, glaucescentibus, oblongis, inferioribus grosse lobatis, in petiolum brevem attenuatis, ceteris integris, acutis, appendicula brevi semiamplexicaulibus; capitulis sessilibus, cylindricis, in fasciculos sessiles congestis, spicam longam, superne densiusculam formantibus; involucri phyllis 5—7, apice lanatulis, externis ovatis, internis linearibus; flosculis luteis; acheniis pallidis, lineari-lanceolatis, compressis, scabridulis, utrinque 5—7 costatis, in rostrum eis subdimidio brevius sensim attenuatis. — Ut videtur praecedenti maxime affinis et ab ea foliis glaberrimis, caulinis sagittatis forsan vix specifice distincta. Ceteroqui ambae ulterius observandae.

In Cycladum insula Amorgos, in mt. Profetae Elias ab autoribus a. 1861 detecta et nuper a Leonis lecta. — Jul. ⊙. ?

b. Folia caulina longe decurrentia.

9. **L. viminea** L. sp. p. 797; S. et S. pr. II. p. 128 (*Prenanthes*); Presl fl. cech. 160; Bois. fl. or. III. p. 818; Heldr. fl. cephal. p. 49, chlor. Parn. p. 22; Form. in D. bot. Monat. 1890 p. 15, in Ver. Brünn 1895 p. 25, 1896 p. 41, 1897 p. 30; Haussk. symb. p. 132. — *Prenanthes ramosissima* All. fl. ped. I. p. 226 t. 33; Ch. et B. fl. pelop. p. 54. — *Phaenopus vimineus* DC. pr. VII. p. 176; Raul. cret. p. 799. — Exsicc.: Sint. et Bornm. it. turc. n. 1337; Sint. it. thessal. n. 1123; Dörfl. fl. gr. n. 207.

Glabra, caulibus erectis, solidis, a basi vel a medio in ramos virgatos, racemoso-spicatos abeuntibus; foliis in lacinias lineares, saepe dentatas pinnatipartitis, infimis breviter petiolatis, ceteris appendicula adnata longe decurrentibus, summis linearibus integris; capitulis cylindricis, brevissime pedicellatis; iuvolucri phyllis externis ovato-oblongis, internis linearibus; flosculis luteis; acheniis lineari-lanceolatis, nigris, scabridulis, utrinque 7—9 costatis, in rostrum concolor, aequilongum sensim attenuatis. — Foliis decurrentibus et acheniis omnino nigris ab affinibus statim diagnoscenda.

In saxosis, fruticetis, regionis montanae in alpinam usque adscendens. Epirus: pr. Janina, Hagios Paraskevi (Form.); Thessalia: pr. Chaliki, Malakasi, Koturi, mt. Ghavellu, Agrapha, Chassia, pr. Patsios et Rapsani in Olympo (Form.), pr. Litochori (Sint.), mt. Pelion (Haussk.), Othrys (For[illegible] [illegible]nt. Parnassus (Heldr.); Attica: mt. Parnes, Hymettus (Heldr.); [illegible] Kyllene (Heldr.); Laconia: mt. Malevo (Orph.), [illegible]ia: pr. Kalamata (Gittard); Creta: pr. Drakona,

Omalos in mt. Sphacioticis (Raul.); Cephalonia: mt. Aenos, Phalaris (Heldr.). — Jul. Aug. ⊙.

3. Sectio. *Mycelis* DC. pr. VII. p. 139. — Capitula 5 flora, involucro squamis minutis calyculato; achenia in rostrum brevem attenuata.

10. **L. muralis** L. sp. p. 797; S. et S. pr. II. p. 128; Ch. et B. exp. p. 232, Fl. pelop. p. 54; Gaertn. fr. II. t. 158; Bois. fl. or. III. p. 817; Heldr. chlor. Parn. p. 22; Hal. Beitr. Fl. Epir. p. 29, Beitr. fl. thessal. p. 15, in ö. b. Z. 1897 p. 285; Haussk. symb. p. 132; Form. in Ver. Brünn 1896 p. 41, 1897 p. 30. — Exsicc.: Sint. it. thessal. n. 729.

Glabra; caule erecto, fistuloso, superne laxe panicato; foliis in segmenta ovato-rhomboidea, angulato-dentata, terminali multo majore, lyrato-pinnatisectis, inferioribus petiolatis, ceteris sagittato-amplexicaulibus, summis lanceolatis, saepius indivisis; capitulis cylindricis, involucri phyllis externis minimis, calyculiformibus, intimis lanceolatis; flosculis sulfureis; acheniis nigris, ellipticis, compressis, utrinque 5 costatis, in rostrum albidum eis 2—3 plo brevius attenuatis.

In silvaticis regionis montanae et subalpinae. Epirus: pr. Kalentini ad septentrionem urbis Arta, pr. Vulgarelion in mt. Tsumerka, pr. Mazuki (Hal.); Thessalia: pr. Chaliki (Sint.), mt. Oxya (Hal.), Zygos (Haussk.), Chassia (Form.), Olympus, Oeta (Heldr.); mt. Parnassus, mt. Telethrion et Dirphys Euboeae (Heldr.); Peloponnesus: mt. Olenos, Kyllene (Heldr.), Malevo (Orph.). — Jul. Aug. ♃.

89. Zacintha Gaertn. fr. II. p. 358.

1. **Z. verrucosa** Gaertn. l. c.; S. et S. pr. II. p. 145; Sieb. avis p. 5, rem. p. 5; Marg. et R. fl. Zante p. 63; Weiss in z. b. G. 1869 p. 45; Raul. cret. p. 800; Bois. fl. or. III. p. 830; Spreitz. in z. b. G. 1897 p. 713; Heldr. fl. cephal. p. 49, Fl. Aegina p. 307; Form. in D. bot. Monat. 1890 p. 16; Hal. Beitr. fl. Epir. p. 30, in z. b. G. 1899 p. 190; Haussk. symb. p. 138; Bald. viagg. Creta p. 71. — *Lapsana zacintha* L. sp. p. 811; Ch. et B. exp. p. 238, Fl. pelop. p. 55. — Icon: Fl. gr. t. 820. — Exsicc.: Rev. pl. cret. n. 99; Dörfl. pl. cret. n. 13.

Parce setulosa, plerumque multicaulis; caulibus rigidis, divaricato-ramosis, subnudis, fructiferis incrassatis; foliis fere omnibus radicalibus, lyrato- vel runcinato-pinnatifidis dentatisve, caulinis paucis, sagittato-amplexicaulibus; capitulis laxe cymosis, aliis terminalibus, aliis in bifurcationibus subsessilibus; flosculis flavis; acheniis pallidis, pappo albo plus duplo longioribus.

In collibus, cultis, ruderatis regionis inferioris et submontanae. Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.); Thessalia: ad monasterium Meteora, pr. Pharsalum, Aivali (Haussk.), Kapurna (Form.); Sporadum insula Scopelos (Philippides); Attica: pr. Marathon, in valle Cephissi pr. Chelidonu, Kephissia, pr. Podoniphti, Tatoi, mt.

Pentelicon, pr. Oropos (Heldr.), mt. Geraka, Turcovuni, pr. Laurion (Haussk.); insula Aegina (Heldr.), Andros (Sart.); pr. Corinthum (Heldr.); Argolis (Chaub.); Elis: pr. Chlemutzi (Heldr.); Messenia: in collibus litoris insulae Prodano oppositis (Chaub.), pr. Kalamata (Zahn); Creta: mt. Hagios Ilias pr. Kissamos (Bald.), pr. Canea (Weiss), Akroteri, Suda, Candia, Guves, Avdu, Potamies, Istovai, in promontorio Sidero (Raul.), pr. Males distr. Hierapetra (Leon.); Zante (Sibth.); Cephalonia: pr. Drapano, Katelios, Piscardo, Omala (Heldr.); Ithaca: mt. Skino (Spreitz.); Corcyra (Bickn.). — Maio. Jul. ⊙

90. Picridium Desf. fl. atl. II p. 220.

a. Achenia conformia, omnia valde rugoso-tuberculata.

1. **P. tingitanum** L. sp. p. 791 (*Scorzonera*); Desf. l. c.; Bois. fl. or. III p. 827; Heldr. chlor. Mykon. p. 346. — *P. hispanicum* Poir. dict. 40 p. 197. — Icon: Fl. gr. t. 792.

Glabrum; caulibus erectis vel adscendentibus, simplicibus vel ramosis, inferne foliisque papillis minutis albis plus minus crebre obsitis; foliis glaucis, spinuloso-denticulatis, pinnatifidis integrisve, caulinis basi cordato-amplexicaulibus; capitulis terminalibus, pedunculo apice tubaeformi-incrassato suffultis; involucri phyllis exterioribus squarrosis, ovatis, interioribus lineari-lanceolatis, omnibus late albo-marginatis; ligulis aureis, extus purpurascentibus; acheniis tetragonis, quadrisulcatis, valde rugoso-tuberculatis, demum fuscescentibus, pappo duplo brevioribus.

In cultis et arenosis Atticae (Sprun.) et in Cycladum insula Mykonos (Heldr.). — Apr. Jun. ⊙ sec. Bois., ♃ sec. DC., Willk. et Lge. — N. v.

b. Achenia difformia, exteriora valde tuberculato-rugosa, intima sublaevia.

2. **P. intermedium** Schultz ap. Webb. phytogr. Canar. II p. 451; Bois. fl. or. III p. 828. — *Sonchus picroides* S. et S. pr. II p. 126, Fl. gr. VIII p. 68 t. 793; Sieb. avis p. 4; Ch. et B. exp. p. 231, Fl. pelop. p. 54; non (L.) Lam.. —

Annuum, glabrum; caulibus erectis, simplicibus vel ramosis, foliisque laevibus; foliis laete viridibus vel glaucescentibus, argute et inaequaliter spinuloso-denticulatis, radicalibus obovato-oblongis vel repando-lobatis, superioribus a basi cordato-auriculata lanceolatis; capitulis terminalibus, pedunculo apice incrassato suffulis; involucri phyllis adpressis, anguste albo-marginatis, exterioribus ovatis, cordato-auriculatis, interioribus oblongo-linearibus; ligulis dilute aureis, concoloribus; acheniis externis ovato-oblongis, sulcatis, fuscis, valde tuberculato-rugosis, intimis prismaticis, albidis, sublaevibus. — A sequente radice annua, foliis subintegris, ligulis concoloribus distinctum.

In arenosis, cultis, herbidis regionis inferioris Achaia, Laconia (Sibth.): ad promontorium Malea, pr. Chimova, Kardamyle (Chaub.); Archinolagus (Sibth.); Creta (Sieb.). — Jan. Maio. ⊙ N. v.

3. **P. picroides** L. sp. p. 792 (*Scorzonera*); Hal. Beitr. fl. Epir. p. 30, in ö. b. Z. 1899 p. 25; Heldr. fl. Aegina p. 307, in ö. b. Z. 1898 p. 184, chlor. Thera p. 16, chlor. Mycon. p. 246. — P. *vulgare* Desf. fl. atl. II p. 221; Friedr. Reise p. 263, 265, 270 et 278; Clem. sert. p. 51; Weiss in z. b. G. 1869 p. 45; Raul. cret. p. 800; Bois. fl. or. III p. 828; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 49; Form. in D. bot. Monat. 1890 p. 15; Haussk. symb. p. 138. — *Sonchus chondrilloides* S. et S. pr. II p. 125, Fl. gr. VIII p. 67 t. 791. — *Picridium ligulatum* DC. pr. VII p. 182 quoad plantam Zacynthiam; Marg. et R. fl. Zante p. 63. — *Zollikoferia chondrilloides* Marg. et R. l. c. — ? *Scorzonera orientalis* Dallap. prosp. p. 107, non L. . — *P. perenne* Fraas fl. class. p. 197. — Exsicc.: Heldr. herb. dimorph. n. 35; Baen. herb. europ. n. 9304; Dörfl. fl. aeg. n. 71 et 440, pl. cret. n. 128.

Perenne, glabrum; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis glaucescentibus, plerumque denticulatis, interdum parce albo-papillosis, inferioribus sinuatis vel pinnatim lobatis partitisve, superioribus a basi cordato-amplexicauli lanceolatis; capitulis terminalibus, pedunculo apice incrassato suffultis; involucri phyllis adpressis, anguste albo-marginatis, exterioribus ovatis, cordato-auriculatis, interioribus oblongo-linearibus; ligulis flavis, subtus lividis; acheniis externis oblongis, sulcatis, olivaceis, valde tuberculato-rugosis, intimis prismaticis, albis, sublaevibus.

In arenosis, herbidis, rupestribus regionis inferioris, per omnem Graeciam. — Oct. Jul. ♃

91. Microrrhynchus Less. syn. p. 139.

1. **M. nudicaulis** L. mant. p. 273 (*Chondrilla*); Less. l. c.; Raul. cret. p. 800. — *Crepis nudicaulis* Sieb. avis p. 5, in Spreng. syst. III p. 634. — *Sonchus divaricatus* Desf. ann. mus. II p. 212 t. 46; Sieb. in Flora V, 2, p. 639. — Exsicc.: Samar. del. pl. Aeg. inf. cur. Heldr. n. 3050 (Alexandria).

Glaber; caulibus erectis vel decumbentibus, divaricatim ramosis; foliis radicalibus rosulatis, runcinatim partitis, laciniis numerosis, cartilagineo-denticulatis, caulinis paucis, minutis, pectinato-dentatis integrisve; capitulis cylindricis, pedunculo brevi dense squamato-suffultis; involucri phyllis late albo-marginatis, exterioribus ovatis, interioribus linearibus; ligulis flavis; acheniis anguste 5—6 sulcatis, nigricantibus, pappo 4 plo brevioribus.

In collibus, arenosis. Creta: pr. Vrises ad radices mt. Kedros, pr. Avdu, inter Prina et Hierapetra (Raul.). — Apr. Maio. ♃ N. v.

92. Cephalorrynchus Bois. diagn. IV p. 28.

1. **C. glandulosus** Bois. l. c.; Haussk. symb. p. 55. — Icon: Rouy ill. t. 162. —

[illegible] foliis glabris [illegible] basi sensim [illegible] basi sessili late [illegible] capitulis cylindricis [illegible] acuminatis [illegible] achenis elliptico- [illegible] 2—3 plo breviore [illegible]

[illegible] Meteora supra Kalabaka [illegible]

[illegible]

[illegible]

[illegible]

β. Annuae vel biennes.

× Involucri phylla glaberrima, externa minima, multo breviora, adpressa, interna fructifera indurata, nervo incrassato.

15. C. pulchra L.

×× Involucri phylla pubescentia, externa 2—3plo breviora, laxiuscula, interna fructifera immutata.

○ Involucri phylla interna facie interna hirta.

16. C. biennis L. **17. C. tectorum** L

○○ Involucri phylla interna facie interna glabra.

18. C. virens L. **19. C. neglecta** L.

b. Rhizoma ramos filiformes tuberiferos edens, collo stoloniferum.

20. C. bulbosa L.

2. Sectio. *Barkhausia* Moench meth. p. 537. — Achenia exalata, omnia vel centralia manifeste rostrata.

a. Involucri phylla externa lanceolata vel oblongo-lanceolata.

α. Ligulae roseae.

21. C. rubra L.

β. Ligulae luteae.

22. C. foetida L. **23. C. setosa** Hall.

b. Involucri phylla externa ampla, ovata, concava, omnino scariosa.

24. C. vesicaria L.

3. Sectio. *Gatyona* Cass. dict. 18 p. 184. — Achenia apice attenuata vel rostrata, radii intus carinata vel alata.

a. Achenia apice breviter attenuata.

α. Pedunculi demum vix incrassati; involucri fructiferi ovati, apice non contracti, phylla interna obtuse carinata, achenia marginalia intus carinata non includentia.

25. C. multiflora S. et S.

β. Pedunculi demum evidenter incrassati; involucri fructiferi globosi, apice contracti, phylla interna basi valde incrassata, achenia marginalia intus carinato-alata includentia.

26. C. Dioscoridis L. **27. C. tubaeformis** Hal.

b. Achenia interna in rostrum eis 2—3plo longius abeuntia.

28. C. aspera L.

1. Sectio. *Eucrepis* Bois. fl. or. III. p. 831. — Achenia exalata, apice plus minus attenuata, sed non rostrata.

a. Estolonosa, non tuberifera.

α. Perennes.

× Caulis foliatus (foliis interdum minimis), normaliter pleiocephalus.

○ Involucri phylla eglandulosa, praeter indumentum farinosum, dorso pilis crebris longis nigricantibus ciliata.

1. **C. geracioides** Haussk. symb. p. 135. — Exsicc.: Sint. it. thessal. n. 912.

Rhizomate obliquo, fusiformi; caule elato, paucifoliato, sparse piloso, a medio in ramos paucos, elongatos, monocephalos diviso; foliis obscure

viridibus, subtus ad nervum pilosis, inferioribus breviter petiolatis, oblongis, lyrato-pinnatipartitis, segmentis oblongis, denticulatis, terminali multo majore, irregulariter sinuato-dentato, foliis superioribus 1—2, ovatis vel ovato-lanceolatis, denticulatis, basi cordata amplexicaulibus, summis squamaeformibus; pedunculis apice incrassatis, capitulis majusculis; involucri phyllis externis triangulari-lanceolatis, dimidio brevioribus, internis lineari-lanceolatis; receptaculo glabro; ligulis flavis; acheniis brunneis, tenuiter striatis, apice vix attenuatis. — Ex affinitate *C. sibiricae* L. et *C. blattarioidis* (L.) Vill., ab ambabus praesertim foliis inferioribus lyrato-pinnatipartitis distincta.

In subhumidis jugi Zygos in Pindo tymphaeo (Haussk.). — Jun. Jul. ♃.

⊙⊙ Involucri phylla plus minus farinoso-tomentella, interdum glandulosa, pilis longis nigricantibus nullis.

. Folia radicalia obovato-oblonga, oblonga vel ovato-lanceolata, varie partita vel dentata.

; Rhizoma fusiforme.

, Flosculi lutei.

.— Folia glandulosa.

2. **C. Baldaccii** Hal. in z. b. G. 1892 p. 577. — Exsicc.: Bald. it. alb. 1892 n. 209, it. alb. alt. a. 1894 n. 144, it. alb. epir. IV. a. 1896 n. 183.

Caule erecto, paucifoliato, glanduloso, infra medium longe ramoso, rarius subsimplici; foliis viridibus, utrinque pilis glanduliferis obsitis, radicalibus in petiolum longum attenuatis, oblongis vel oblongo-lanceolatis, lyrato-runcinatis, segmentis irregulariter dentatis, lateralibus parvis, triangularibus, terminali maximo, foliis caulinis minoribus, inferioribus radicalibus similibus, basi amplexicauli sessilibus, summis diminutis, lanceolatis subintegris; pedunculis crassiusculis; capitulis majusculis; involucri phyllis lineari-lanceolatis, densissime glandulosis, externis 2—3plo brevioribus; receptaculo glabro; acheniis brunneis, striatis, scabridulis, apice subattenuatis. — Caulis 20—30 cm. altus, folia 12—20 cm. longa, 3—5 cm. lata.

In praeruptis regionis mediae et superioris. Epirus: mt. Konitza, nec non extra ditionem nostram in mt. Cika in mt. Acroceraunicis et in mt. Tomor Maja Albaniae (Bald.). — Jul. Aug. ♃.

3. **C. Raulini** Bois. diagn. XI. p. 58, Fl. or. III. p. 836; Raul. cret. p. 799; Bald. viagg. Cret. p. 72. — Exsicc.: Bald. it. cret. n. 77.

Caule erecto, subnudo, glanduloso, simplici vel furcato-dicephalo; foliis viridibus, utrinque pilis glanduliferis obsitis, radicalibus oblongis, in petiolum attenuatis, repando-denticulatis, caulinis ad 1—2 squamulas lineares reductis; pedunculis tenuibus; capitulis mediocribus; involucri phyllis lineari-lanceolatis, densissime glandulosis, externis 3 plo brevioribus; receptaculo glabro; acheniis brunneis, striatis, scabridulis, apice subattenuatis. — Caulis 5 cm. altus, folia 3 cm. longe, 1 cm. lata.

In regione alpina, ad nives. Creta: mt. Volakia distr. Sphakia (Bald.), mt. Psiloriti (Raul.). — Maio, Jul. ♃.

:= Folia eglandulosa.

— Achenia scabra; folia tenera, dentata.

4. **C. auriculaefolia** Sieb. avis p. 5, rem. p. 5, in Spreng. syst. III. p. 634; Raul. cret. p. 799 cum var. *minor*; Bois. fl. or. III. p. 834. — Exsicc.: Heldr. pl. cret. n. 1435; Bald. it. cret. alt. n. 233.

Caule erecto, subnudo, collo lanato, ceterum glabro vel parce glanduloso, superne ramoso; foliis radicalibus oblongis vel obovato-oblongis, acute dentatis, glabriusculis, in petiolum brevem attenuatis, caulinis paucis, diminutis, lanceolatis linearibusve, ad dichotomias sessilibus, basi lanatis; pedunculis vix incrassatis; capitulis mediocribus; involucri farinosi, interdum parce glandulosi, phyllis lanceolatis, externis duplo brevioribus; receptaculo setuloso; acheniis brunneis, apice sensim attenuatis. — Caulis 20—30 cm. altus, folia radicalia 10—15 cm. longa, 3—4 cm. lata, tenerrima.

In rupestribus regionis inferioris et montanae. Creta: mt. Sphaciotici (Raul.), mt. Lazaro in mt. Lassiti (Bald.), pr. Kritza distr. Mirobello (Heldr.). — Apr. Jun. ♃.

= Achenia laevia; folia radicalia runcinata; plantae omnino eglandulosae.

5. **C. turcica** Deg. et Bald. in ö. b. Z. 1896 p. 417; Bald. riv. coll. bot. Alb. 1895 p. 53, 1896 p. 74. — Exsicc.: Bald. it. alb. epir. IV. n. 182.

Caule elato, floccoso-lanuginoso, demum glabrescente, inferne folioso, superne subnudo, a medio ramoso polycephalo; foliis utrinque tenuiter lanuginosis, oblongo-lanceolatis, acute runcinato-dentatis, inferioribus petiolatis, superioribus basi semiamplexicauli, acute auriculata sessilibus, summis lineari-lanceolatis, subintegris; pedunculis crassiusculis; capitulis mediocribus; involucri dense niveo-tomentosi phyllis lanceolatis, externis triplo brevioribus; receptaculo glabro; ligulis flavis concoloribus; acheniis brunneis, striatis, apice vix attenuatis. — Ex affinitate *C. lacerae* Ten. et *C. rigidae* W. et K., a quibus foliorum forma et praesertim indumento involucri niveo-tomentoso distinguitur.

In rupestribus montanis. Epirus: mt. Profeta Elias et Kuruna supra Vonicko distr. Ljaskovik, mt. Vratedon supra Cepelovon distr. Zagorion, ad radices mt. Peristeri pr. Kalarrytes (Bald.). — Jun. Jul. ♃.

6. **C. divaricata** Bois. et Heldr. diagn. VII. p. 13, Fl. or. III. p. 836. — Exsicc.: Heldr. herb. norm. n. 1453.

Caule erecto, a basi divaricatim ramoso, glabro vel inferne parce lanuginoso; foliis viridibus, parce breviterque papilloso-asperulis, radicalibus lanceolatis, in petiolum brevem attenuatis, in lobos triangulares versus basin diminutos pinnatipartitis runcinatis, caulinis ad dichotomias minutis; pedunculis rigidis, crassiusculis; capitulis mediocribus; involucri tomentelli phyllis lanceolatis, externis subtriplo brevioribus, medio dorso

nigricantibus; receptaculo glabro; ligulis flavis, extus saepe rubentibus; acheniis brunneis, striatis, apice vix attenuatis. — Caulis 15—20 cm. altus, folia 5—10 cm. longa, 10—15 mm. lata.

Ad rupes regionis alpinae mt. Taygetos Laconiae loco unico Kakochioni dicto (Heldr.). — Jun. Aug. ♃.

7. **C. Sibthorpiana** Bois. et Heldr. diagn. XI. p. 56, Fl. or. III. p. 836; Raul. cret. p. 799; Spreitz. in z. b. G. 1890 p. 296; Bald. viagg. Creta p. 72. — *Hieracium foetidum* S. et S. pr. II. p. 134, Fl. gr. t. 799, non Willd. — Exsicc.: Heldr. pl. cret. n. 1637; Bald. it. cret. alt. n. 236.

Differt a praecedente, cujus forsan forma macra, caule nano, 3—10 cm. alto, parce breviterque ramoso, 1—4 cephalo et indumento foliorum et phyllorum densiori.

In rupestribus regionis mediae et superioris. Creta (Sibth.): mt. Stravopodia, Theodori (Raul.) et Hagion Pneuma (Bald.) in mt. Sphacioticis, mt. Ida (Heldr.). — Jun. Aug. ♃.

,, Flosculi rosei.

8. **C. incana** S. et S. pr. II. p. 136, Fl. gr. IX. p. 1 t. 802; Bois. fl. or. III. p. 835; Heldr. chlor. Parn. p. 22. — *C. albida* Ch. et B. fl. pelop. p. 54, non All. — Exsicc.: Heldr. herb. norm. n. 333 et 1657, Orph. fl. gr. n. 477.

Tota pube arachnoidea, pilis glandulosis immixtis cana; caulibus pumilis, erectis vel adscendentibus, a basi divaricatim semel vel pluries dichotomis; foliis radicalibus in petiolum brevem attenuatis, oblongo-lanceolatis, runcinato-pinnatipartitis vel sublyratis, caulinis sessilibus, amplexicaulibus, pedunculis rigidis, crassiusculis, dense glandulosis; capitulis mediocribus; involucri phyllis linearibus, externis paucis brevibus; receptaculo glabro; acheniis brunneis, striatis, apice sensim attenuatis. — Floribus pulchre roseis insignis.

Ad rupes regionis mediae et alpinae. Phocis: mt. Parnassus ad Gurna (Heldr.); Attica: mt. Parnes (Sart.); Euboea (Sibth.): mt. Dirphys (Heldr.); Elis: mt. Olenos (Heldr.); Achaia: mt. Clocos (Orph.); Laconia: mt. Malevo (Orph.); Messenia: pr. Andrusa (Chaub.). — Jun. Jul. ♃.

;; Rhizoma praemorsum, longe et dense fibrosam.

,, Involucrum pedunculisque glandulosum.

9. **C. Fraasii** Schultz in Flora 1842 p. 173; Haussk. symb. p. 135. — *C. Sieberi* Bois. diagn. XI. p. 53, Fl. or. III. p. 844; Raul. cret. p. 799; Spreitz. in z. b. G. 1877 p. 713, 1887 p. 663; Heldr. fl. cephal. p. 49, Hal. Beitr. fl. Achaia p. 25. — *C. interrupta* Sieb. avis p. 5, rem. p. 5, in Flora I. p. 272, DC. pr. VII. p. 161; Marg. et R. fl. Zante p. 63; non S. et S. — Exsicc.: Orph. fl. gr. n. 764; Heldr. herb. norm. n. 236 et 1050; Rev. pl. cret. n. 256 (specimina alia ad typum spectant, alia transitum ad var. praebent).

Colle dense rufo-lanato; caulibus elatis, glabris pilosulisve, superne laxe corymboso-paniculatis, glandulosis; foliis teneris, laete virentibus, supra glabrescentibus, subtus parce setulosis, radicalibus petiolatis, interrupte lyrato-pinnatipartitis, segmentis lateralibus utrinque 5—7, ab infimis triangularibus ad superiora ovata vel ovato-triangularia, obtusa vel acutiuscula, margine saepius integra sensim auctis, terminali majori, basi truncato vel subcordato, obsolete dentato, folio caulino ad ramum inferiorem sito, sublyrato vel lobato, basi amplexicauli, ceteris minimis, linearibus integris; pedunculis gracilibus, glandulosis; capitulis mediocribus; involucri glandulosi phyllis externis triangularibus, 4plo brevioribus, interioribus lanceolatis, ligulis flavis; acheniis striatis, apice attenuatis.

β. **Mungierii** Bois. et Heldr. diagn. XI. p. 55; Raul. cret. p. 799; pro sp.; Bois. fl. or. III. p. 845. — Caulis nanus, foliis vix sesquilongior, saepe a basi dichotomus, oligocephalus; indumentum saepe copiosius, interdum inferne et ad folia glandulosum. — Exsicc.: Heldr. pl. cret. a. 1846; Bald. it. cret. n. 78, it. cret. alt. n. 333.

In dumosis, silvaticis, rupestribus humidis regionis montanae et subalpinae. Thessalia: mt. Pelion pr. Zagora, Milies (Heldr.); Attica: mt. Pateras, Parnes, Pentelicon, Hymettus (Heldr.), pr. Kamariza (Haussk.); Achaia mt. Olenos, Panachaicon (Hal.); Argolis pr. Nauplia (Raul.); Laconia: mt. Malevo (Orph.), mt. Taygetos pr. Anavryti (Heldr.); Creta: ad promontorium Meleka, Hagios Joannes Kaimenos (Raul.), Omalos (Rev.); Zante (Marg.); Cephalonia: pr. Argostoli (Heldr.); Corcyra: mt. San Deca, pr. Lacones, mt. Pantocrator pr. Barbati (Spreitz.); — *β.* Creta: ad Hellinoseli in mt. Sphacioticis, mt. Psiloriti supra Voriza (Raul.). — Apr. Jun. ♃.

10. **C. smyrnaea** DC. pr. VII. p. 170; Haussk. symb. p. 135. — Exsicc.: Pichl. pl. fl. rum. et bithyn. a. 1874.

Differt ab affini *C. Fraasii* colle glabro, foliorum segmentis paucis, acutis, sinuato-dentatis, segmento terminali multo minoribus, pedunculis glandulosissimis, crassioribus, involucri phyllis externis brevissimis.

In silvis deustis mt. Pentelicon Atticae (Hausk.). — Maio, Jul. ♃. N. v.

,, Involucrum pedunculisque eglandulosum.

11. **C. Reuteriana** Bois. diagn. XI. p. 55. — Exsicc.: Grimb. pl. corc. a. 1897.

Collo glabro; caulibus elatis, inferne hirtulis, ceterum glabris, a medio stricte longeque ramosis; foliis teneris, supra glabrescentibus, subtus ad nervum medium crebrius setulosis, radicalibus petiolatis, runcinato-lyratis, segmentis saepius integris, lateralibus utrinque 3—5, oblongo-triangularibus, retrorsis, acutissimis, terminali basi truncato vel cordato triangulari, aliis 2—3plo majori, foliis caulinis 1—2, lineari-lanceolatis, integris; pedunculis gracilibus, glabris vel parce araneosis; capitulis mediocribus; involucri glabri vel parce hirtuli phyllis externis triangu-

laribus, 4 plo brevioribus, interioribus linearibus, fructiferis ad carinam incrassatis; ligulis flavis; acheniis striatis apice vix attenuatis. — Specimina nostra cum asiaticis bene congruunt.

In collibus, fruticetis pr. Potamos Corcyrae (Grieseb.). — Apr. Maio. ♃.

.. Folia radicalia linearia, graminea, integra.

12. **C. crocifolia** Bois. et Heldr. diagn. VII. p. 14, Fl. or. III. p. 837. — Exsicc.: Heldr. herb. norm. n. 1452.

Glabra, rhizomate fusiformi, verticali, collo squamoso; caulibus erectis, tenuiter virgatis, simplicibus vel a parte inferiori parce ramosis, 1—4 cephalis; foliis glauco-caesiis, radicalibus numerosis, linearibus, infimis subspathulatis obtusiusculis, ceteris acutis, caulinis paucis, minoribus; pedunculis apice subincrassatis; capitulis sat parvis; involucri farinoso-puberuli phyllis dorso nigricantibus, infimis brevissimis subpatulis, ceteris linearibus; ligulis flavis; acheniis pallidis, tenuissime striatis, apice paululum attenuatis. — Habitu *Hieracio porrifolio* L. simillima.

In regione alpina mt. Taygetos Laconiae loco unico Megala Zonaria et Chupata dicto (Heldr.). — Jun. Aug. ♃.

×× Caulis nanus, aphyllus vel subnudus, scapiformis, mono- vel dicephalus.

13. **C. bithynica** Bois. diagn. IV. p. 29, Fl. or. III. p. 29. — Exsicc.: Heldr. herb. n. 2484.

Rhizomate praemorso, fibrosissimo; foliis radicalibus, pube brevi papillari subglanduloso hirtis, obovatis, runcinato-lyratis, lobis triangularibus, deorsum decrescentibus, terminali maximo, rotundato; scapis adscendentibus, flexuosis, foliis parum longioribus, nudis, monocephalis vel furcatis, foliolo lineari-instructis, dicephalis, superne praesertim glanduloso-setosis; capitulis mediocribus; involucri nigricantis, parce farinoso-tomentelli, pilisque sparsis longioribus obsiti, phyllis lanceolatis, externis dimidio brevioribus; acheniis striatis, apice vix attenuatis. — Differt a *C. Raulini* radice praemorsa, foliorum forma et indumento minus glanduloso.

In herbidis regionis alpinae mt. Olympus (Heldr.). — Jul. Aug. ♃

14. **C. Columnae** Ten. syll. p. 398 *(Hieracium)*; Froel. in DC. pr. VII. p. 167; Bois. fl. or. III. p. 841, suppl. p. 324; Hal. Beitr. fl. thessal. p. 15, in ö. b. Z. 1897 p. 285; Haussk. symb. p. 136; Bald. riv. coll. bot. Alb. 1895 p. 53. — *C. Sartoriana* Bois. et Heldr. diagn. ser. 2, III. p. 100. — Huc forsan: *Hieracium alpestre* S. et S. pr. II. p. 132, non Jacq. — Exsicc.: Bald. it. alb. epir. III. n. 169.

Rhizomate praemorso, fibroso; foliis radicalibus, expansis, glabriusculis, lanceolatis, runcinato-dentatis vel pinnatipartitis, laciniis triangularibus irregularibus, terminali saepius majori; scapis decumbentibus, foliis 2—3 plo longioribus, nudis vel squamulis setaceis paucis obsitis,

monocephalis, glabris puberulisve; capitulis sat parvis; involucri livescentis, puberuli phyllis lanceolatis, externis triplo brevioribus; acheniis striatis, apice vix attenuatis. — Differt a praecedente rhizomate non fibrosissimo, foliis glabris, non obovato-lyratis, capitulo tenuiori, involucro non setuloso.

In pascuis alpinis. Epirus: in jugo Kakardista mt. Tsumerka (Bald.), mt. Peristeri (Hartl.); Thessalia: mt. Tringia (Hartl.), Zygos, Karava (Haussk.); mt. Tymphrestus Eurytaniae, mt. Oeta Phthiotidis (Heldr.); mt. Korax (Tuntas); Achaia: mt. Olenos, Kyllene (Heldr.). — Jul. Aug. ♃.

β. Annuae vel biennes.

× Involucri phylla glaberrima, externa minima, multo breviora, adpressa, interna fructifera indurata, nervo incrassato.

15. **C. pulchra** L. sp. p. 806. — Icon: Rchb. t. 80.

Caule erecto, inferne pubescenti, saepe glanduloso, superne glabro, laxe paniculato-corymboso; foliis puberulis, interdum glandulosis, inferioribus oblongo-spathulatis, in petiolum attenuatis, dentatis vel runcinatis, superioribus ovato-lanceolatis lanceolatisve, dentatis vel integris, basi truncata vel subsagittata sessilibus; capitulis cylindraceis; receptaculo glabro, ligulis flavis; acheniis linearibus, obsolete striatis, apice vix attenuatis, marginalibus scabriusculis, interioribus laevibus.

β. **adenoclada** Haussk. symb. p. 136. — Caulis ad inflorescentiam usque dense glandulosus, pedunculi parte superiore tantum glabri, folia utrinque glandulosa, achenia etiam interiora scabriuscula. — Exsicc.: Sint. it. thessal. n. 408.

In dumetis saxosis, vineis Thessaliae: pr. monasterium Meteora (Haussk.) et Kalabaka (Sint.), ubi var. tantum lecta fuit. — Maio, Jun. ⊙.

×× Involucri phylla pubescentia, externa 2 – 3 plo breviora, laxiuscula, interna fructifera immutata.

○ Involucri phylla interna facie interna hirta.

16. **C. biennis** L. sp. p. 807; Friedr. Reise p. 283; Ung. Reise p. 125. — Icon: Fl. dan. t. 1997.

Pilis brevibus asperula; caule erecto, corymboso; foliis oblongis, runcinatis, sinuato-dentatis vel subintegris, margine non revolutis, inferioribus in petiolum attenuatis, superioribus basi rotundata, auriculata vel subsagittata sessilibus; capitulis majusculis; involucri adpresse canescentis phyllis internis facie interna sericeo-pubescentibus; receptaculo villoso; ligulis luteis; stylis flavis; acheniis striatis, superne subattenuatis.

In herbidis regionis inferioris. Indicatur in: Corcyra (Ung.) et inter Pyrgos et Arcadia Elidis (Friedr.); species pro Graecia dubia. — Maio, Jul. ⊙ N. v.

17. **C. tectorum** L. sp. p. 807; Ch. et B. exp. p. 236, Fl. pelop. p. 55. — Icon: Curt. fl. lond. V. t. 55.

Glabra vel adpresse puberula; caule erecto, corymboso; foliis inferioribus oblongis vel lanceolatis, sinuato-dentatis vel runcinatis, in petiolum attenuatis, superioribus linearibus, integris, margine revolutis, basi sagittata sessilibus; capitulis mediocribus; involucri cano-pubescentis phyllis internis facie interna adpresse hirtis; receptaculo breviter setuloso; ligulis flavis; stylis siccis nigricantibus; acheniis striatis, apice valde attenuato contractis.

Ad radices mt. Taygetos pr. Andruvista (Chaub.); species pro Graecia dubia. — Maio, Jun. ⊙ N. v.

○○ Involucri phylla interna facie interna glabra.

18. **C. virens** L. sp. ed. 2 p. 1134; Ch. et B. exp. p. 235, Fl. pelop. p. 55, var. *α. arvensis, β. silvatica et γ. bellidifolia*; Friedr. Reise p. 277. — *C. polymorpha* Wallr. sched. crit. p. 426; Clem. sert. p. 53 v. *humilis* DC. pr. VII p. 162. — Icon: Rchb. t. 90.

Glabriuscula; caule erecto, laxe corymboso; foliis margine non revolutis, inferioribus oblongis lanceolatisve, sinuato-dentatis vel runcinato-pinnatipartitis, in petiolum attenuatis, superioribus lanceolatis linearibusve, integris vel inferne inciso-dentatis, basi sagittata sessilibus; capitulis parvis, erectis; involucri adpresse puberuli, interdum glandulosi phyllis externis brevissimis, internis lanceolato-linearibus; receptaculo glabro; ligulis luteis; stylis flavis vel siccis virescentibus; acheniis striatis, apice breviter attenuatis.

In herbidis, arvis. Indicatur in: Attica: mt. Pentelicon (Clem.); Elis (Ch. et B. fl. pelop.); Messenia: pr. Pylos, Chimova, Messene (Ch. et B. exp.); Cycladum insula Syra (Orph. ex Bois. l. c.); loca nonnulla forsan ad *C. neglectam* spectant. — Jun. Jul. ⊙ N. v.

19. **C. neglecta** L. mant. p. 107; S. et S. pr. II p. 137, Fl. gr. t. 805; Ch. et B. fl. pelop. p. 55; Bois. fl. or. III p. 849; Spreitz. in z. b. G. 1877 p. 713; Heldr. fl. cephal. p. 49. Fl. Aegina p. 307, chlor. Thera p. 16; Form. in D. bot. Monat. 1890 p. 14; Hal. Beitr. fl. Epir. p. 29; Haussk. symb. p. 137. — *C. stricta* Scop. fl. carn. ed. 2 II p. 99 t. 47. — *C. cernua* Ten. fl. neap. pr. p. 47, Fl. nap. t. 183, annot. fl. gr. p. 15. — *C. cretica* Bois. diagn. XI p. 53, Fl. or. III p. 849, suppl. p. 325; Raul. cret. p. 799; quae ex descriptione et speciminibus a Reverchon lectis, ab auctore ipso sub hoc nomine enumeratis a *C. neglecta* nullo modo discedit. — Exsicc.: Heldr. herb. norm. n. 520, 952 et 1353; Orph. fl. gr. n. 1083; Rev. pl. cret. n. 92; Sint. it. thessal. n. 246. —

Parce setuloso-hirta; caulibus gracilibus, adscendentibus vel erectis, saepe numerosis, a basi ramosis, superne laxe cymosis; foliis margine non revolutis; [illegible]ibus obovato-oblongis vel oblongo-lanceolatis, den- [illegible] iolum attenuatis, superioribus lanceolato- [illegible] nectinato-pinnatifidis, basi sagittata sessili- [illegible] hesin nutantibus; involucri glabriusculi

vel puberuli phyllis externis bevissimis, internis lanceolato-linearibus; receptaculo glabro; ligulis luteis; stylis flavis; acheniis striatis, apice sensim attenuatis. — Caules usque 25 cm. alti, ramosi, occurunt tamen specimina 5 cm. alta, mono- vel dicephala. — Stirps acheniis subrostratis transitum praebet ad sectionem sequentem, habitu toto attamen a speciebus praecedentibus non sejungenda. — Differt a *C. virenti* capitulis minoribus, ante anthesin nutantibus et acheniis subrostratis.

β. **fuliginosa** S. et S. pr. II p. 138; Urv. enum p. 102; DC. pr. VII p. 161 cum *β. adscendens* (forma caule adscendente); Marg. et R. fl. Zante p. 63; Ung. Reise p. 125; Weiss in z. b. G. 1869 p. 45; pro sp.; Ch. et B. fl. pelop. p. 55; Raul. cret. p. 799. — Huc forsan: *C. leontodontoides* Gelmi in bull. soc. bot. ital. 1889 p. 450, vix All. — Involucrum plus minus dense setulosum. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889; Baen. herb. europ. n. 9244.

In collibus herbidis, ad vias, in pascuis regionis inferioris totius Graeciae, in superiorem usque adscendens. — Mart. Jul. ⊙

b. Rhizoma ramos filiformes tuberiferos edens, collo stoloniferum.

20. **C. bulbosa** L. sp. p. 798; Ch. et B. exp. p. 232, Fl. pelop. p. 54; Tausch in Flora XI p. 78; Weiss in z. b. G. 1869 p. 45; Bois. fl. or. III p. 832; Heldr. fl. cephal. p. 49, Fl. Aegina p. 307; Gelmi in bull. soc. bot. it. 1889 p. 450; Haussk. symb. p. 136; Hal. in ö. b. Z. 1897 p. 96. — *Hieracium bulbosum* Willd. sp. III p. 1562; S. et S. pr. II p. 133, Fl. gr. t. 798; Sieb. avis rem. p. 5. — *Aetheorrhiza bulbosa* Cass. dict. sc. nat. 48 p. 425; Marg. et. R. fl. Zante p. 63; Raul. cret. p. 798. — Exsicc.: Heldr. et Hal. fl. aegaea a. 1889, fl. Sporad. a. 1896.

Caule erecto, simplici, monocephalo, ima basi paucifolio, ceterum nudo, apice glanduloso; foliis glabris, oblongo-lanceolatis, dentatis, in petiolum attenuatis; capitulo mediocri; involucri basi glandulosi phyllis adpressis, lineari-lanceolatis, externis dimidio brevioribus; ligulis flavis; acheniis testaceis, paucistriatis, fusiformi-subtetragonis.

In arenosis, olivetis, rupestribus maritimis. Thessalia: pr. Volo (Heldr.); Sporadum insula Scopelos (Leon.); Attica: pr. Laurion (Heldr.); Phaleron, Pikermi, mt. Hymettus, insula Salamis, Lero, Aegina et in scopulo adjacente Angistri (Heldr.); Corinthus (Haussk.); Argolis (Sprun.): pr. Nauplia (Haussk.); Cycladum insula: Keos, Kythnos (Tunt.), Syra (Weiss), Melos (Leon.); Laconia (Sibth.); Messenia: pr. Pylos (Chaub.); Creta: inter Perama et Damasta (Raul.); Zante (Sibth.); Cephalonia: pr. Drapano (Heldr.); Corcyra (Gelmi). — Febr. Jun. ♃.

2. Sectio. *Barkhausia* Moench meth. p. 537. — Achenia exalata, omnia vel centralia manifeste rostrata.

a. Involucri phylla externa lanceolata vel oblongo-lanceolata.

α. Ligulae roseae.

21. **C. rubra** L. sp. p. 806; S. et S. pr. p. 136, Fl. gr. t. 805; Ch. et B. exp. p. 234, Fl. pelop. p. 54; Bois. fl. or. III p. 855; Spreitz. in z. b. G. 1877 p. 713, 1887 p. 663; Heldr. fl. cephal. p. 50; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Aetol. p. 8, Beitr. fl. Achaia p. 25, in ö. b. Z. 1895 p. 460; Haussk. symb. p. 136; Bald. riv. coll. bot. Alb. 1896 p. 74. — *Barkhausia rubra* Moench meth. p. 537; Marg. et R. fl. Zante p. 62; Friedr. Reise p. 282; Raul. cret. p. 798. — *C. incarnata* Vis. stirp. dalm. sp. p. 17; Friedr. Reise p. 283; non (Wulf.) Tausch. — *B. purpurea* Ung. Reise p. 125 et Heldr. fl. cephal. p. 50, ex speciminibus in herb. mus. palat. vindob. asservatis; non (Biv.) Guss. . — Exsicc.: Baen. herb. europ. n. 9245; Sint. it. thessal. n. 82; Dörfl. fl. gr. n. 408.

Parce puberula; caulibus solitariis, erectis aut numerosis adscendentibus, simplicibus vel parce ramosis, superne nudis; foliis plerisque radicalibus, in petiolum attenuatis, oblongis, runcinato-dentatis vel pinnatipartitis, rarius indivisis, acute dentatis; pedunculis elongatis, monocephalis; capitulis majusculis, ante anthesin cernuis; involucri pallidi phyllis ab externis subpatulis, glabrescentibus, ovato-oblongis, ad interna oblongo-lanceolata, glanduloso-hispida sensim auctis; acheniis tuberculato-striatis, externis breviter attenuatis, ceteris in rostrum eis longius abeuntibus.

β. **integrifolia.** — Foliis indivisis, oblongo-spathulatis, acute dentatis. — Exsicc.: Baen. herb. europ. n. 9245 b; Reis. fl. gr. n. 1898.

In collinis, ad vias agrestes, in rupestribus regionis inferioris et subalpinae. Epirus: mt. Cuka distr. Janina (Bald.); Thessalia: pr. (Kalabaka (Sint.), Pharsalus, Aivali, Ormanmagula (Haussk.), Volo (Heldr.); Acarnania: pr. Vonitza (Lacon); Aetolia: mt. Arapokephalo (Heldr.), Aetolikon (Reis.), Mesolongion (Nied.), mt. Taphiassos ad sinum Patranum (Hal.); Eurytania: pr. Karpenisi (Samarit.); Attica: mt. Cithaeron (Guicc.), mt. Parnes pr. Tatoi, Megalivrysi et Menidi (Heldr.); Petalium insula Xeronisi (Holzm.); Achaia: mt. Olenos (Hal.), pr. Kalavryta (Leon.), Trikala (Orph.); Argolis: mt. Palamidi pr. Nauplia (Haussk.); Messenia: mt. Ithome (Heldr.), pr. Arcadia, Pylos, insula Sapienza (Chaub.); Laconia (Sibth.); Creta (Sibth.): pr. Malaxa, Prosnero, Askyphos, Kritza, Prina (Raul.); Zante (Marg.); Cephalonia: pr. Argostoli (Spreitz.), mt. Aenos pr. Omala (Heldr.); Ithaca: pr. Vathy (Spreitz.); Corcyra: pr. San Deca (Spreitz.), Garuna (Baen.). — β. ut videtur multo rarior: Zante (Reiser); Corcyra: pr. San Deca (Baen.) — Apr. Jun. ⊙

Obs. Habeo *Crepidem* rubrifloram e Corcyra (Sagb.) et Aetolikon Aetoliae (Reis.), habitu *C. bienni* simillimam, caulibus crassibus, fistulosis, elatis, 30—60 cm. altis, longe stricteque ramosis, foliatis, foliis runcinatis vel acute dentatis, capitulis eis *C. rubrae* similibus, sed acheniis nondum evolutis, quamombrem dijudicare nequeo, an huic sectioni vel priori adnumeranda sit. Ulterius observandum anne forte speciem pro-

priam (*C. corcyrensis*) sistet. — Huc forsan spectat *C. purpurea* Gelmi in bull. soc. bot. ital. 1889 p. 450 e Corcyra pr. Signes.

β. Ligulae luteae.

22. **C. foetida** L. sp. p. 807; Ch. et B. fl. pelop. p. 55; Weiss in z. b. G. 1869 p. 45; Bois. fl. or. III. p. 851; Heldr. fl. cephal. p. 50, Fl. Aegina p. 307, chlor. Thera p. 16; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1895 p. 25; Hal. Beitr. fl. Achaia p. 25. — *Barkhausia foetida* DC. fl. fr. IV. p. 42; Raul. cret. p. 798. — Icon: Rchb. t. 83.

Pubescens vel hirsuta, saepe glandulosa; caulibus solitariis, erectis vel numerosis, adscendentibus, simplicibus vel ramosis; foliis inferioribus in petiolum attenuatis, runcinato-pinnatifidis, interdum sublyratis, rarius sinuato vel repando-dentatis, caulinis auriculato-amplexicaulibus, inciso-dentatis, summis integris; capitulis solitariis cymosisve, majusculis, ante anthesin cernuis; involucro cano-tomentello et plus minus longe et crebre setuloso, simulque saepissime glanduloso, phyllis lanceolatis, adpressis, internis triplo longioribus, demum ad carinam induratis; acheniis subtiliter striatis, externis breviter attenuatis, ceteris in rostrum eis longius abeuntibus. — Stirps valde polymorpha.

α. **typica.** — Annua, caule erecto, solitario, infra medium vel superne ramoso; involucro cano-tomentello, pilisque glandulosis brevibus obsito. — Exsicc.: N. v.

β. **interrupta** S. et S. pr. fl. gr. II. p. 137, Fl. gr. IX. p. 2 t. 803 pro sp. — *C. glandulosa v. interrupta* Haussk. symb. p. 136. — *C. glandulosa v. acutiloba* Haussk. l. c. p. 137 (f. foliis profunde pinnatifidis, laciniis angustis, longe acutatis). — *C. foetida v. lagoserioides* Bald. viagg. Creta p. 72. — Annua, caulibus solitariis vel pluribus, adscendentibus, humilibus, a collo ramosis; involucro cano-tomentello pilisque longioribus et brevioribus, glandulosis obsito. — Exsicc.: Orph. fl. gr. n. 342; Sint. et Bornm. it. turc. n. 1339; Bald. it. cret. n. 79.

γ. **glandulosa** Guss. ind. sem. hort. Bocc. 1825 p. 4, pro sp.; Bisch. Beitr. p. 252. — *C. foetida v. rhoeadifolia* Bois. fl. or. III. p. 851, saltem quoad loca graeca; non *Barkhausia rhoeadifolia* M. a. B. fl. taur. cauc. II. p. 259, quae involucro setis longis rigidis, plerisque eglandulosis hispido differt. — *C. glandulosa v. simplex* Haussk. symb. p. 136; Form. in Ver. Brünn 1897 p. 30. — Annua, caule erecto, solitario, infra medium vel superne ramoso; involucro pilis longis flavescentibus glanduliferis dense hispido. — Exsicc.: Heldr. herb. norm. n. 310; Dörfl. fl. aeg. n. 17.

δ. **radicata** S. et S. pr. II. p. 136, Fl. gr. VIII. p. 74 t. 800. — *C. foetida v. maritima* Bois. fl. or. III. p. 851. — *C. glandulosa v. maritima* Haussk. symb. p. 137. — Perennans, caulibus humilibus, adscendentibus, parce ramosis vel monocephalis; foliis glabres-

centibus; involucro pilis flavescentibus longis, brevioribusque glandulosis intermixtis obsito. — Exsicc.: N. v.

ε. **zacynthia** Marg. et R. in DC. pr. VII. p. 158, Fl. Zante p. 62 (*Barkhausia*) pro sp. — *C. glandulosa v. eglandulosa* Haussk. symb. p. 136. — Annua, caule a basi ramosissimo, ramis elongatis subnudis; foliis subdentatis, incisis; involucro longe hispido, eglanduloso. — Exsicc.: N. v.

In ruderatis, cultis, glareosis regionis inferioris et montanae. Typum e Graecia non vidi et loca indicata probabiliter omnia ad varietatem β. et γ. spectant, quae per omnem territorium divulgatae videntur; — δ. indicatur tantum in arenosis maritimis Thessaliae pr. Caterina (Orph.) et ad Ergastiria pr. Laurion Atticae (Haussk.); — ε. in Zacyntho pr. Pisinonda et Litakia (Marg.) et in mt. Palamidi supra Naupliam Argolidis (Haussk.). — Maio, Jul. ⊙ vel ⊙.

23. **C. setosa** Hall. in Roem. arch. 1797, I. 2 p. 1; Bois. fl. or. III. p. 853; Hal. Beitr. fl. Epir. p. 29; Haussk. symb. p. 137. — *C. hispida* W. et K. pl. rar. hung. I. p. 42. t. 43; Ch. et B. exp. p. 234. Fl. pelop. p. 54. — *Barkhausia setosa* DC. fl. fr. IV. p. 44. — *B. hispida* Link. en. hort. berol. II. p. 290; Friedr. Reise p. 286. — Exsicc.: Sint. it. thessal. n. 245 et 1075.

Caule erecto, ramosissimo, foliisque plus minus setuloso; foliis inferioribus runcinato-dentatis vel pinnatifidis, in petiolum attenuatis, superioribus basi sagittata, saepe inciso-dentata, sessilibus; capitulis parvis, laxe cymosis, semper erectis; involucri phyllis valde hispido-spinulosis, demum ad carinam induratis, externis patulis vel reflexis, 2—3 plo brevioribus; acheniis striato-tuberculatis, omnibus rostro eis vix breviore superatis.

In pratis, herbidis, arvis regionis inferioris et montanae. Epirus: pr. Kalentini, Vulgarelion (Hal.); Thessalia: frequens in Pindo dolopico et in planitie (Haussk.); Aetolia: pr. Mesolongion (Nieder); Attica: pr. Athenas (Sart.), in valle Cephissi (Orph.) pr. Podoniphti (Heldr.); Arcadia: ad lacum Pheneon (Heldr.); Laconia: mt. Malevo (Orph.); Elis: pr. Olympia (Nied.); Messenia: pr. Methone, Kalamata (Chaub.). — Maio, Aug. ⊙.

b. Involucri phylla externa ampla, ovata, concava, omnino scariosa.

24. **C. vesicaria** L. sp. p. 805; S. et S. pr. II. p. 136; Pieri corc. fl. p. 114; Sieb. avis p. 5. rem. p. 5; Ch. et B. fl. pelop. p. 54; Bois. fl. or. III. p. 853; Haussk. symb. p. 137. — *Barkhausia vesicaria* Spreng. syst. III. p. 652; Raul. cret. p. 798. — Icon: Rchb. t. 86. — Exsicc.: Rev. pl. cret. n. 95; Sint. it. thessal. n. 243; Bald. it. cret. alt. n. 88; Dörfl. pl. cret. n. 12.

Parce setuloso-pubescens; caule erecto, superne ramoso, corymboso; foliis inferioribus oblongo-spathulatis, in petiolum attenuatis, dentatis, runcinato-pinnatifidis subiyratisve, caulinis lanceolatis, basi sessili auri-

culatis; pedunculis basi bractea scariosa, ovato-oblonga, cucullata suffultis; capitulis mediocribus, semper erectis; involucri phyllis externis glabris, calyculum formantibus, dimidio brevioribus, interioribus linearibus, hirtis vel villosis, ligulis flavis, acheniis striatis, marginalibus apice attenuatis, interioribus in rostrum eis longius abeuntibus.

β. **scariosa** Willd. sp. III. p. 1595, pro sp.; Vis. fl. dalm. II. p. 117; Weiss in z. b. G. 1869 p. 45. — Calyculum involucrum subaequans. — Exsicc.: N. v.

In herbidis, agrorum marginibus regionis inferioris et montanae. Thessalia: ad Metochi pr. Kalabaka (Sint.); Attica: ad radices mt. Pentelicon pr. Pikermi, pr. Chelidomi, Menidi, mt. Pateras (Heldr.), pr. Kamariza Laurii (Haussk.); Achaia: ad Gerokomion pr. Patras (Heldr.); Messenia: inter Armyros et Kalamata (Gittard); Creta: pr. Canea, Epanochorio, mt. Karadagh pr. Candia (Raul.), pr. Stavraka distr. Malevisi (Bald.), pr. Males distr. Hierapetra (Leon.); Corcyra (Pieri); — β. pr. Canea Cretae (Weiss). — Maio, Jun, ⊙.

Obs. *C. bursifolia* L. sp. p. 805; S. et S. pr. II. p. 135; Ch. et B. exp. p. 234, Fl. pelop. p. 54. — Probabiliter erronee indicatur in Laconia (Sibth.) et pr. Navarin Messeniae (Chaub.).

3. Sectio. *Gatyona* Cass. dist. 18, p. 184 sensu ampl.; Bois. fl. or. III. p. 832. — Achenia apice attenuata vel rostrata, radii intus carinata vel alata.

a. Achenia apice breviter attenuata.

α. Pedunculi demum vix incrassati; involucri fructiferi, ovati, apice non contracti, phylla interna obtuse carinata, achenia marginalia intus carinata non includentia.

25. **C. multiflora** S. et S. pr. II. p. 138, Fl. gr. IX. p. 4 t. 806; Bois. fl. or. III. p. 855; Hal. in ö. b. Z. 1895 p. 460, 1897 p. 96, Heldr. chlor. Thera p. 16. — *Endoptera dichotoma* Bois. et Bal. diagn. ser. 2 III. p. 98; Weiss in z. b. G. 1869 p. 45; Heldr. Gartenfl. 1873 p. 323, chlor. Mykon. p. 246; Haussk. symb. p. 138. — Exsicc.: Heldr. herb. fl. hellen. n. 26.

Caule erecto, solitario, glabro vel parce pilosulo, interdum glanduloso, paulo supra basin dichotome ramoso; foliis glabris, inferioribus oblongo-spathulatis, dentatis vel runcinato-pinnatifidis, breviter petiolatis, mediis oblongis, acute auriculato-amplexicaulibus, summis lineari-lanceolatis, subintegris; capitulis parvis; involucri tomentelli simulque parce glandulosi phyllis externis brevissimis, lineari-setaceis, interioribus linearibus, demum navicularibus, carinatis; ligulis luteis, involucrum parum superantibus; acheniis leviter incurvis, 5 costatis, apice breviter attenuatis; pappo caduco, involucrum vix superante. — Praeter notas indicatas, capitulis multo minoribus, eis *C. virentis* aeque magnis, a sequente diversissima.

In saxosis, collibus apricis, maritimis. Attica: pr. Laurion, Sunium, Buliasmene, in scopulo Rhaphti, Pharmacusarum insula Lero et Mikra

Kyra (Heldr.); Argolis: pr. Nauplia (Haussk.), Vromolimni, in scopulo Angistri (Heldr.); Sporadum insula Jura (Leon.); insula Pontikonisi ad septentrionem Euboeae (Wild); Cycladum insula: Tenos, Syra (Weiss), Keos, Kythnos (Tunt.), Mykonos (Sart.), Thera (Heldr.); Creta pr. Canea (Weiss). — Apr. Jun. ⊙.

β. Pedunculi demum evidenter incrassati; involucri fructiferi, globosi apice contracti, phylla intima basi valde incrassata, achenia marginalia intus carinato-alata includentia.

26. C. **Dioscoridis** L. sp. ed. 2 p. 1133; S. et S. pr. II. p. 139, Fl. gr. t. 808; Urv. enum. p. 102; Ch. et B. exp. p. 235, Fl. pelop. p. 54; Bois. fl. or. III. p. 856; Heldr. fl. cephal. p. 50, chlor. Thera p. 16; Hal. Beitr. fl. Epir. p. 29, Beitr. fl. Achaia p. 25, in ö. b. Z. 1895 p. 460, in z. b. G. 1899 p. 190. — *C. globifera* Hall. in naturw. Anz. 1818 p. 91. — *C. croatica* Horn. hort. hafn. suppl. p. 90. — *Gatyona globulifera* Cass. dict. 18 p. 184. — *G. Dioscoridis* Rchb. fl. germ. exc. p. 254. — *Endoptera Dioscoridis* DC. pr. VII. p. 172; Raul. cret. p. 800; Spreitz. in z. b. G. 1877 p. 663; Haussk. symb. p. 137. — Exsicc.: Orph. fl. gr. n. 765; Heldr. herb. norm. n. 242 et 1354.

Parce papilloso-puberula vel glabra; caule erecto, solitario, superne vel a basi dichotome ramoso, ramis subdivaricatis, raro simplici, monocephalo; foliis inferioribus obovatis vel oblongis, runcinato-dentatis vel lyratis, in petiolum attenuatis, mediis lanceolatis, basi laciniata sagittata sessilibus, summis linearibus integris; pedunculis glabris vel parce glandulosis, fructiferis parum incrassatis; capitulis mediocribus; involucri tomentelli, interdum parce glandulosi, phyllis externis brevissimis, linearisetaceis, interioribus lanceolatis, demum incurvatis, basi valde incrassatis; ligulis luteis, involucro subduplo longioribus; acheniis incurvis, 5 costatis, apice breviter attenuatis; pappo caduco, involucrum parum superante.

In saxosis regionis inferioris et submontanae. Epirus: pr. Kalentini, Mazuki (Hal.); Thessalia: pr. Klinovo (Haussk.); Aetolia: pr. Guritza (Reis.), Mesolongion (Nied.), mt. Taphiassos (Hal.); Attica: insula Salamis, mt. Lycabettus, Hymettus, Parnes (Heldr.), pr. Geraka, Laurion (Haussk.); Acrocorinthus (Orph.); Argolis: in peninsula Methana (Haussk.), pr. Nauplia (Orph.), Astros (Bois.); Achaia: pr. Trikala (Heldr.), mt. Panachaicon, pr. Lopesi (Hal.); Messenia: pr. Pylos (Chaub.), Kalamata (Zahn); Laconia: pr. Androvista (Heldr.); Cythaera: pr. San Nicolo (Spreitz.); Creta (Raul.); Cycladum insula Camini pr. Thera (Urv.); Zante (Mazz.); Cephalonia: pr. Kastro (Heldr.). — Apr. Jun. ⊙

27. C. **tubaeformis**. — Exsicc.: Hal. it. gr. secund. a. 1893.

Glabra; caulibus numerosis, erectis vel adscendentibus, gracilibus, simplicibus monocephalis vel inferne furcatis dicephalis, superne nudis, valde tubuloso-incrassatis; foliis fere omnibus radicalibus, anguste lanceolatis, dentatis, in petiolum attenuatis, caulinis paucis, valde diminutis, anguste linearibus, integris vel basi laciniatis; capitulis parvulis; in-

volucri tomentelli, eglandulosi, phyllis externis lineari-setaceis, interioribus lanceolatis, demum incurvatis, basi valde incrassatis; ligulis ignotis; acheniis incurvis, 5 costatis, apice breviter attenuatis; pappo caduco, involucro breviore. — Differt a praecedente, cui affinis, caulibus numerosis simplicibus vel tantum semel furcatis, valde elongatis, subnudis, superne tubaeformibus, capitulis minoribus, pappo breviore et toto habitu, eo *Arnoseridis minimae* simili.

In rupestribus calcareis mt. Taphiassos ad sinum Patranum loco Kakiskala dicto Aetoliae (Hal.). — Maio. ⊙

b. Achenia interna in rostrum eis 2—3 plo longius abeuntia.

28. **C. aspera** L. sp. ed. 2 p. 1132; S. et S. pr. II p. 137, Fl. gr. t. 804; Bois. fl. or. III p. 857. — *Nemauchenes aculeata* Cass. dict. 34 p. 362. — *Endoptera aspera* DC. pr. VII p. 178; Raul cret. p. 800. —

Caulibus a medio dichotome ramosis, laxe cymoso-corymbosis, setis rigidis aculeiformibus hispidis; foliis margine asperis vel setoso-ciliatis, radicalibus oblongo-spathulatis, in petiolum brevem attenuatis, acute multidentatis, caulinis oblongis, laciniato-dentatis, auriculato-amplexicaulibus, summis linearibus, integris; capitulis parvis, centralibus et saepe lateralibus cymarum subsessilibus; pedunculis fructiferis crassiusculis; involucri fructiferi phyllis externis ovatis, scariosis, parvis, deciduis, internis tandem induratis, tomentellis, navicularibus, dorso incrassato aculeatis; acheniis marginalibus intus carinato-alatis, apice acutatis, internis, rostratis.

In graminosis Cretae (Sibth.); a recentioribus non lecta. — Maio, Jun. ⊙

94. Hieracium L. gen. n. 913.

Dispositio specierum:

1. Sectio. *Pilosella* Tausch in Flora 1828 Ergbl. I. p. 50. — Achenia ad summum 1½ mm. longa, nigra, costis apice in dentem brevem procurrentibus. — Rhizoma saepe stoloniferum; folia omnia vel plurima radicalia, rosulata; caulis scapiformis, aphyllus vel oligophyllus; innovatio per stolones, raro per rosulas.

a. Caulis monocephalus.

1. H. macranthum Ten.

b. Caulis pleiocephalus.

α. Caulis humilis, aphyllus vel monophyllus, apice 2—5cephalus.

2. H. auricula L.

β. Caulis elatior, foliatus, polycephalus, corymbosus.

× Folia elliptica, oblonga vel lanceolata, viridia, mollia, apice (praesertim superiora) saepius glandulosa; involucrum dense setosum griseum.

3. H. cymosum L.

×× Folia linearia vel lanceolata, glauca, apice eglandulosa; involucrum glabrum vel breviter puberulum et glanduliferum.

4. H. florentimum All. **5. H. Bauhini** Schult.

2. Sectio. *Euhieracium* Scheele in Linn. XVI. p. 652. — Achenia minima 3 mm. longa, pallida, fusca vel nigra, costis apice plica annuliforme connexis. — Rhizoma rarissime stoloniferum; folia rarius rosulata; caulis saepissime pleiophyllus; innovatio per rosulas vel gemmas, rarissime per soboles.

1. Subsectio. *Graeca.* — Rhizoma repens, soboles (stolones subterraneos) edens; folia glauca, glabra vel villosa, radicalia sub anthesi adhuc vigentia; caulis scapiformis, 1—3phyllus, monocephalus, rarius furcatus dicephalus; involucrum irregulariter imbricatum, phyllis acutis vel acuminatis; ligulae apice glabrae vel ciliatae; pappus albus.

6. H. Naegelianum Panc. **7. H. graecum** Bois. et Heldr.

2. Subsectio. *Andryaloidea* Monn. ess. p. 51. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia cinerea, pilis distincte plumosis villosa vel tomentosa, radicalia saepius sub anthesi adhuc vigentia; caulis plerumque pleiocephalus, foliatus; involucrum irregulariter imbricatum, phyllis acutis vel acuminatis; ligulae (in nostris) apice glabrae; pappus rufescenti-albus.

a. Achenia fusco-atra.

8. H. Parnassi Fr.

b. Achenia testacea vel brunnea.

α. Caulis praeter pubem stellatam pilis longis patulis plus minus crebre obsitus.

× Involucrum dense villosum.

○ Ligulae glabrae.

. Folia rosulata, sub anthesi vigentia.

9. H. aetolicum Hal. **10. H. Heldreichii** Bois.

.. Folia infima sub anthesi emarcida.

11. H. euboeum Hal.

○○ Ligulae extus pilosae.

12. H. pannosum Bois. **13. H. Gaudryi** Bois. et Orph.

×× Involucrum pilis stellatis minutis cano-tomentellum.

14. H. thessalum Form.

β. Caulis glaber.

15. H. oligadenum Belli.

3. Subsectio. *Glutinosa* Koch syn. p. 459. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia viridia, glanduloso-viscida, radicalia sub anthesi saepe adhuc vigentia; caulis scapiformis vel plus minusve foliatus, mono-vel polycephalus; involucrum imbricatum, phyllis acutis; ligulae apice glabrae, pappus rufescenti-albus.

16. H. scapigerum Bois. Orph. et Heldr.

4. Subsectio. *Pulmonaroidea* Koch syn. p. 454. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia viridia vel glaucescentia, nunquam tomentosa vel glandulosa, radicalia sub anthesi adhuc vigentia; caulis scapiformis vel plus minusve foliatus, plerumque pleiocephalus; involucrum irregulariter imbricatum, phyllis acutis; ligulae apice glabrae; pappus rufescenti-albus.

a. Folia pleraque lanceolata vel oblongo-lanceolata, basi longe attenuata.

17. H. Sartorianum Bois. et Heldr.

b. Folia pleraque ovata vel oblonga, basi subcordata vel rotundata, rarius attenuata.

18. H. incisum Hoppe. **19. H. murorum** L. **20. H. abietinum** Bois. et Heldr.

5. Subsectio. *Italica* Fr. epicr. p. 107. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia viridia, nunquam tomentosa vel glandulosa, radicalia

sub anthesi saepe jam emarcida; caulis foliatus; inflorescentia racemosa; involucrum imbricatum, phyllis obtusis; ligulae apice glabrae; pappus rufescenti-albus.

a. Caulis stricte ramosus, rarius simplex, ramis tenuibus, saepe virgatis, elongatis, glabris vel parce pilosis.

21. H. bracteolatum S. et S. **22. H. Koracis** Bois.

b. Caulis simplex, crassiusculus, foliatus, stricte racemosus, patule pilosus.

23. H. racemosum W. et K. **24. H. crinitum** S. et S.

1. Sectio. *Pilosella* Tausch in Flora 1828 Ergbl. I. p. 50. — Achenia ad summum 2 ½ mm. longa, nigra, costis apice in dentem brevem procurrentibus. — Rhizoma saepe stoloniferum; folia omnia vel plurima radicalia, rosulata; caulis scapiformis, aphyllus vel oligophyllus; innovatio per stolones, raro per rosulas.

a. Caulis monocephalus.

1. **H. macranthum** Ten. fl. nap. IV. p. 114 pro var. *H. pilosellae*; Ten. fl. nap. V. p. 190 t. 184; Bois. fl. or. III. p. 860; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 4, chlor. Parn. p. 22; Hal. in z. b. G. 1888 p. 759, Hal. Beitr. fl. Epir. p. 29, in ö. b. Z. 1897 p. 285; Haussk. symb. p. 135; Bald. riv. coll. bot. Alb. 1895 p. 54, 1896 p. 75. — *H. pilosella* S. et S. pr. II. p. 133, quoad locum peloponnesiacum; Ch. et B. exp. p. 233, Fl. pelop. p. 54; Form. in D. bot. Monat. 1890 p. 15, in Ver. Brünn 1896 p. 39, 1897 p. 30; non L. — Exsicc.: Orph. fl. gr. n. 79; Heldr. herb. norm. n. 772, it. Thessal. n. 9; Bald. it. alb. epir. III. n. 176.

Rhizomate repente, stolones breves edente; caule erecto vel adscendente, aphyllo, pube brevissimo obducto, praesertim inferne plus minus setuloso et insuper superne saepe breviter glanduloso; foliis radicalibus, obovato-lanceolatis lanceolatisve, in petiolum attenuatis, subtus cano-tomentosis, utrinque setoso-pilosis; capitulis majusculis; involucri adpresse cano-tomentelli pilisque brevissimis nigris, saepe glandulosis conspersi vel simul pilis longis albidis hispidi, phyllis ab externis ovatis vel oblongis, obtusis ad interna lanceolata auctis; ligulis flavis, marginalibus extus saepe rubellis. — Subspecies *H. pilosellae* L. sp. p. 800, quod differt stolonibus elongatis, phyllis externis lanceolatis acutis, internis linearibus. — Variat caulibus 2—3 cm. usque 25 cm. longis, capitulis parvulis (*v. obscurius* Naeg. et. P. Pilos. p. 125) vel majusculis, involucro adpresse tomentello pilisque brevibus nigris glandulosis (*v. glandulosum* Naeg. et. P. l. c.) vel eglandulosis (*v. eglandulosum* Naeg. et P. l. c.) consperso et pilis albidis elongatis nullis vel (*v. anatolicum* Naeg. et P. l. c. p. 126) plus minus crebris obducto.

In pascuis regionis mediae et superioris fere omnium montium altiorum. Epirus: mt. Profetae Elias distr. Ljaskovik, mt. Micikeli, Peristeri (Bald.), Tsumerka (Hal.); Thessalia: in Pindo tymphaeo pr. Malakasi (Heldr.), in oropedio Neuropolis (Haussk.), mt. Giseltepe, Dokimi, Chassia (Form.), mt. Olympus, Pelion (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Euboea: mt. Ocha (Sart.),

Dirphys (Heldr.); Attica: mt. Parnes (Heldr.); Peloponnesus (Sibth.): mt. Malevo (Orph.), Kyllene, Taygetos (Heldr.). — Jun. Aug. ♃.

b. Caulis pleiocephalus.

α. Caulis humilis, aphyllus vel monophyllus, apice 2—5 cephalus.

2. **H. auricula** L. sp. p. 800; S. et S. pr. II. p. 133. — Icon: Fl. dan. t. 1111. —

Rhizomate repente, stolonifero; caule erecto, superne puberulo pilisque brevibus nigricantibus, saepius glanduliferis obsito; foliis radicalibus, glaucis, lanceolato-spathulatis, glabris, basin versus ciliatis, inferioribus obtusis; pedunculis brevibus; involucri phyllis nigricantibus, dorso breviter setoso-glandulosis, lineari-lanceolatis, obtusis; ligulis flavis.

In montosis (Sibth.), sine loco speciali; a recentioribus non lectum. — Maio, Jul. ♃. N. v.

β. Caulis elatior, oligophyllus, polycephalus, corymbosus.

× Folia elliptica, oblonga vel lanceolata, viridia, mollia, apice (praesertim superiora) saepius glandulosa; involucrum dense setosum, griseum.

3. **H. cymosum** L. sp. ed. 2, p. 1126; Bald. riv. coll. bot. Alb. 1895 p. 54, 1896 p. 75. —

Rhizomate truncato, obliquo, estolonoso; caule erecto, 1—4 phyllo, stellatim puberulo, setisque patulis saltem inferne plus minus dense obsito, superne saepe glandulifero, in cymam umbelliformem laxiusculam vel compactam abeunti; foliis oblongo-lanceolatis, setosis, pubeque stellata minuta adspersis, infimis obtusis, ceteris acutis, caulinis diminutis; capitulis parvis; involucri dense setosi, grisei, phyllis acutis; ligulis flavis.

α. **typicum.** — Caulis saepius 3 phyllus; setae caulis diametrum aequantes vel breviores; inflorescentia laxiuscula, eglandulosa vel involucri phylla apice pedunculisque uti caulis superne parce breviterque glandulosa. — Exsicc.: Sint. it. thessal. n. 737.

β. **sabinum** Seb. rom. pl. fasc. I. p. 8 t. 1; Seb. et Maur. fl. rom. pr. p. 270, t. 6; Bois. fl. or. III. p. 863 p. p.; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Epir. p. 29, Beitr. fl. Achaia p. 25; Heldr. chlor. Parn. p. 22; Haussk. symb. p. 135; Bald. riv. coll. bot. Alb. 1895 p. 54; Form. in Ver. Brünn 1896 p. 39; pro sp.; Naeg. et Pet. pilosell. p. 407. — Caulis saepissime 2 phyllus; setae caulis diametro saepius longiores; inflorescentia congesta; involucri phylla apice parce, pedunculi et caulis superne crebrius glandulosa. — Exsicc.: Bald. it. alb. epir. III. n. 174.

γ. **Heldreichianum** Naeg. et Pet. pilosell. p. 408. — *H. sabinum v. Heldreichianum* Hal. in z. b. G. 1888 p. 759. — *H. Heldreichianum* Haussk. symb. p. 135. — Caulis saepissime 2 phyllus; setae caulis diametro saepius longiores; inflorescentia laxiuscula vel congesta; involucri phylla apice parce glandulosa, caulis et pedunculi omnino

eglandulosi. — Intermediis cum β. conjuncta est. — Exsicc.: Heldr. herb. n. 2729; Orph. herb. n. 2741; Bald. it. alb. epir. III. n. 175; Dörfl. fl. gr. n. 289.

In herbidis regionis subalpinae et alpinae. Thessalia: in valle Negerli pr. Chaliki (Sint.); — β. et γ. Epirus: mt. Gamila (Bald.), Tsumerka, Peristeri (Hal.); Thessalia: mt. Zygos, mt. Karava, in oropedio Neuropolis (Haussk.), mt. Oeta (Heldr.); Aetolia: mt. Tymphrestus (Samar.), Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Attica: mt. Parnes (Heldr.); Achaia: mt. Olenos, Panachaicon (Hal.). — Jun. Aug. ♃.

1 × 3. **H. macranthum × Heldreichianum.** — *H. Halacsyi* Heldr. herb. norm. n. 1656. — Rhizomate praemorso, estolonoso; caule furcato, oligocephalo, stellatim puberulo, pilisque patulis parce obsito, eglanduloso, foliis paucis squamiformibus munito; foliis radicalibus lanceolatis, subtus cano-tomentosis, utrinque setoso-pilosis; capitulis mediocribus; involucri phyllis cano-tomentellis, pilisque brevissimis nigris eglandulosis simulque longis albis hispidis; ligulis extus rubellis. — Hybrida manifeste inter parentes intermedia, habitu *H. cano (cymosum × pilosella)* Naeg. et Pet. pilos. p. 425 simile, a *H. macrantho* praesertim caulibus furcatis, capitulis dimidio minoribus, a *H. Heldreichiano* caulibus humilibus, furcatis, oligocephalis, nec corymbosis, foliis subtus canis, capitulis multo majoribus, ligulis extus rubellis discedens. — In regione abietina mt. Parnes Atticae (Heldr.). — Jun. Jul. ♃.

3 × 5. **H. sabinum × Bauhini.** — (*H. panachaicum.*) — Habitu magis *H. Bauhini* accedens, differt tamen ab ea foliis latioribus, mollioribus, vix glaucescentibus, inflorescentia congesta et praesertim involucro dense longeque setoso, griseo, a *H. sabino* rhizomate longe stolonifero. — In herbidis regionis mt. Panachaicon Achaiae (Hal.). —

Obs. Hybridam aliam, *H. Heldreichiannm × praealtum* memorat sine descriptione Haussknecht in symb. p. 135 ex oropedio Neuropolis Thessaliae.

×× Folia linearia vel lanceolata, glauca, apice eglandulosa; involucrum glabrum vel breviter puberulum et glanduliferum.

4. **H. florentinum** All. fl. ped. I. p. 213; Hal. in ö. b. Z. 1890 p. 39; Haussk. symb. p. 135; Form. in Ver. Brünn 1896 p. 39, 1897 p. 30. — *H. piloselloides* Vill. hist. pl. Dauph. III. p. 100 t. 27. —

Rhizomate praemorso, estolonoso; caulibus gracilibus, glabris, a medio vel superne ramosis, 1—3 phyllis; foliis glabris vel basi petioloque ciliatis, foliis lineari-lanceolatis linearibusve, acutiusculis, radicalibus in petiolum attenuatis, caulinis basi angusta sessilibus; capitulis minutis, in corymbum laxum dispositis; involucri phyllis linearibus petiolisque glandulosis; ligulis flavis.

β. **hololeion** Naeg. et P. pilos. p. 524. — Folia caulina basi parum dilatata sessilibus; involucrum omnino glabrum. — Exsicc.: Sint. et Bornm. it. turc. n. 1342 et 1824.

γ. **praealtum** Vill. voy. bot. p. 62; Haussk. symb. p. 135; Bald. riv. coll. bot. Alb. 1896 p. 75; pro sp. — Caules crassiores, inflorescentia minus laxa, capitula majora, involucrum obscurum, breviter puberulum et glanduliferum. — Exsicc.: N. v.

In herbidis, fruticetis, glareosis regionis montanae et subalpinae. Indicatur in Thessalia: mt. Zygos (Haussk.), Mandra Hodza, mt. Mitritza in mt. Chassia (Form.); — β. ad Hagios Diomysios in Olympo (Sint.); — γ. Epirus: mt Olycika (Bald.); Thessalia: in oropedio Neuropolis, ad monasterium Korona (Haussk.); Attica: mt. Pentelicon (Haussk.). — Jul. Aug. ♃.

5. **H. Bauhini** Schult. obs. bot. p. 164; Bess. prim. fl. gal. II. p. 149; Hal. Beitr. fl. Epir. p. 29, Beitr. fl. Achaia p. 25. — *H. praealtum* Ch. et B. exp. p. 234, Fl. pelop. p, 54; Bois. fl. or. III. p. 862 cum β. *hispidissimum*; Bois. fl. or. suppl. p. 326; Heldr. chlor. Parn. p. 22, non Vill. — *H. praealtum v. Baulini* Griesb. revis. p. 13; Bald. riv. coll. bot. Alb. 1896 p. 75. – *H. atticum* Nym. consp. p. 452. — *H. magyaricum* Naeg. et Pet. pilos. p. 576, cum var. *laconicum* et subsp. *graecum* p. 579. — Huc quoque verosimiliter: *H. pannonicum subsp. leiocaulon* Naeg. et P. pilos. p. 759, quoad locum mt. Malevo, quod dicitur hybrida e *H. magyarico* et *H. echioide*, tametsi species ultima in omni Graecia desideratur. — Exsicc.: Heldr. herb. norm. n. 46, 1049, 1655 f. *subexstolonosum*; Orph. fl. gr. n. 759; Sint. it. thessal. n. 911.

Rhizomate praemorso, longe stolonifero; caulibus elongatis, glabriusculis vel puberulis, simulque setis longis patentibus plus minus dense hispidis, interdum glanduliferis, 1—4 phyllis, apice corymbosis; foliis lanceolatis oblongo-lanceolatisve, plus minus dense setoso-hispidis, radicalibus in petiolum attenuatis, caulinis basi angusta sessilibus; capitulis minutis in corymbum laxiusculum dispositis; involucri phyllis linearibus, petiolisque pube brevi stellata, setis simplicibus basi nigricantibus, et brevióribus glanduliferis obsitis. — Differt a praecedente praesertim rhizomate valde stolonifero.

In montosis, in regionem subalpinam adscendens per totam Graeciam continentalem; deesse videtur in insulis Ionicis et maris Aegaei, nec non in Creta. — Maio, Jul. ♃.

2. Sectio. *Euhieracium* Scheele in Linn. XVI. p. 652. — Achenia minime 3 mm. longa, pallida, fusca vel nigra, costis apice plica annuliformi connexis. — Rhizoma rarissime stoloniferum; folia rarius rosulata; caulis saepissime pleiophyllus; innovatio per rosulas vel gemmas, rarissime per soboles.

1. Subsectio. *Graeca*. — Rhizoma repens, soboles (stolones subterraneos) edens; folia glauca, glabra vel villosa, radicalia sub anthesi

adhuc vigentia; caulis scapiformis, 1—3 phyllus, monocephalus, rarius furcatus dicephalus; involucrum irregulariter imbricatum, phyllis acutis vel acuminatis; ligulae apice glabrae vel ciliatae; pappus albus (characteribus nonnullis ad sectionem *Pilosella*, aliis ad *Stenotheca* vergens, a priori acheniis, ab altera involucro non biseriali discedens).

6. **H. Naegelianum** Panc. elench. pl. vasc. Crnag. p. 57; Naeg. et P. archierac. p. 6; Bald. riv. coll. bot. Alb. 1895 p. 54. — *H. undulatum* Bois. fl. or. III. p. 867, suppl. p. 327; Hal. Beitr. fl. Epir. p. 29; non Ait. — Exsicc.: Heldr. it. gr. septentr. a. 1879; Hal. it. gr. sec. a. 1893; Bald. it. alb. epir. III. n. 173.

Glaucum; rhizomate repente, multicipite, soboles squamatos, apice demum rosulam ferentibus edente; caulibus gracilibus, glabris, monocephalis, foliis 1—3 minutis, anguste linearibus obsitis; foliis radicalibus rosulatis, glabris vel subtus ad nervum et margine parce setosis, infimis spathulatis, ceteris linearibus, margine undulatis; involucri phyllis nigricantibus, lineari-lanceolatis, acuminatis, parce setulosis et glandulosis; ligulis flavis; glabris; acheniis pallidis.

In saxosis herbidis regionis alpinae, rarissima. Epirus: in cacumine mt. Peristeri (Hal.); Thessalia: mt. Olympus, Oeta (Heldr.); Aetolia: mt. Tymphrestus, Korax (Heldr.). — Jul. Aug. ♃.

7. **H. graecum** Bois. et Heldr. diagn. ser. 2 III. p. 101; Fl. or, III. p. 866, suppl. p. 327; Heldr. chlor. Parn. p. 22. — Fortasse huc: *H. alpinum* S. et S. pr. III. p. 133, non L. — Exsicc.: Heldr. herb. norm. n. 723; Orph. herb. n. 2725.

Glaucum; rhizomate repente, collo squamoso; caulibus flexuosis, monocephalis, rarius supra basin furcatis, dicephalis, foliisque pilis longissimis flexuosis denticulatis villosis, foliis 1—3 minutis, linearibus obsitis; foliis radicalibus rosulatis, infimis oblongo-spathulatis, obtusis, ceteris late linearibus, margine interdum obsolete repandis; involucri phyllis lineari-lanceolatis, acutiusculis, pilis flexuosis, denticulatis, eglandulosis villosissimis; ligulis flavis, eximie ciliatis; acheniis . . . — Praecedenti affine et probabiliter etiam soboliferum, differt ab eo praeter indumentum villosum et ligulis ciliatis, capitulis multo majoribus.

In regione alpina mt. Parnassus loco Trypios vrachos (Held.) et Kedrozasteno (Orph.) dicto, dein in mt. Chelmos Achaiae ad fontem Stygis (Heldr.) et in mt. Taygetos (Zahn), rarissime. — Jul. Aug. ♃.

2. Subsectio. *Andryaloidea* Monn. ess. p. 51. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia cinerea, pilis distincte plumosis villosa vel tomentosa, radicalia saepius sub anthesi adhuc vigentia; caulis plerumque pleiocephalus, foliatus; involucrum irregulariter imbricatum, phyllis acutis vel acuminatis; ligulae (in nostris) apice glabrae; pappus rufescenti-albus.

a. Achenia fusco-atra.

8. **H. Parnassi** Fr. symb. p. 86, epicr. p. 76; Bois. fl. or. III p. 867; Heldr. chlor. Parn. p. 22; Freyn in bull. herb. Bois. III

p. 509. — *H. Heldreichii* Bois. diagn. ser. 2 III p. 102 p. p. quoad. plant. Parn. —

Pilis elongatis, flexuosis, remote plumosis tenuiter villosum; caulibus e collo 1—3, monocephalis vel furcato-dicephalis, nudis vel ad dichotomiam folio parvo lineari obsitis; foliis radicalibus sub anthesi vigentibus, rosulatis, obovatis, sinuato-dentatis, obtusis, in petiolum brevissimum attenuatis; capitulis mediocribus; involucri pilis simplicibus, basi nigricantibus, eglandulosis, hirsuti, phyllis acuminatis; ligulis flavis, extus pilosis. — Species mihi ignota.

In mt. Parnasso (Sprun.). — N. v.

b. Achenia testacea vel brunnea.

α. Caulis praeter pubem stellatam pilis longis patulis plus minus crebre obsitus.

× Involucrum dense villosum.

○ Ligulae glabrae.

• Folia rosulata, sub anthesi vigentia.

9. **H. aetolicum.** — *H. Gaudryi* Heldr. it. gr. septentr. a. 1879, non Bois. —

Pilis elongatis, flexuosis, plumoso-scabris tenuiter villosum; caulibus e collo 1—2, tenuibus, furcato 2—3 cephalis, ad dichotomiam folio parvo lanceolato obsitis, superne 1—2 squamulosis; foliis radicalibus rosulatis, oblongis oblongo-lanceolatisve, acutis, integris, basi attenuatis; capitulis mediocribus; involucri pilis patulis longis villosi, vix stellulato-puberuli, eglandulosi, phyllis acutis; ligulis flavis; acheniis immaturis brunneis. — Caulis 15—25 cm. altus, folia 7—9 cm. longa, 2 cm. lata. — Sequenti affine, ab eo caulibus crebrius pilosis, foliis lanceolatis, acutis, angustioribus, integris et involucro pilis stellulatis fere omnino destituto differt. *H. Gaudryi* caule elatiori crassiusculo, superne divaricatim et dichotome 2—4 cephalo, foliis remote denticulatis, ligulis extus pilosulis et indumento copiosiori discedit.

In rupibus regionis alpinae mt. Korax Aetoliae, rarissime (Heldr.). — Jul. Aug. ♃.

10. **H. Heldreichii** Bois. diagn. ser. 2 III p. 102 excl. plant. Parnass., Fl. or. III p. 868; Fr. epicr. p. 75; Hal. Beitr. fl. Thessal. p. 15; Freyn in bull. herb. Bois. III p. 655. — Exsicc.: Heldr. herb. norm. n. 1654.

Pilis elongatis, flexuosis, plumoso-scabris tenuiter villosum; caulibus e collo 1—3, tenuibus, monocephalis vel furcato 2—3 cephalis, foliis 1—3 minimis obsitis; foliis radicalibus rosulatis, obovatis vel obovato-oblongis, obtusis, integris vel inferne repando-denticulatis, subsessilibus; capitulis mediocribus; involucri dense stellato-puberuli, pilisque patulis longis villosi, eglandulosi, phyllis acutis; ligulis flavis; acheniis pallidis. — Ab *H. Parnassi*, facie simili, ligulis glabris et acheniis pallidis, a formis scaposis *H. pannosi* indumento adeo tenui, ut color viridis transluceat, foliis tenuiter membranaceis et ligulis glabris discedit.

Ad rupes regionis subalpinae. — Thessalia: mt. Oxya supra Chaliki (Hal.); Attica: mt. Parnes (Sart.); Arcadia: ad lacum Pheneon (Heldr.). — Jun. Jul. ♃.

•• Folia infima sub anthesi emarcida.

11. **H. euboeum** Hal. in z. b. G. 1899 p. 189. — Exsicc.: Heldr. pl. fl. hellen. a. 1895 et 1896.

Pilis elongatis, plumoso-scabris villosum, papillisque minutis asperulum; caule crasso, abbreviato, rigido, folioso, apice subito in pedunculos paucos, elongatos, monocephalos diviso; foliis oblongis, acutis, remote acuteque denticulatis, basi attenuatis; pedunculis praeter pubem stellatam papilloso-asperulis et parce patule pilosis, foliis squamaeformibus jubatis obsitis; capitulis mediocribus; involucri stellato-puberuli, pilisque patulis longis villosi, eglandulosi phyllis acutis; ligulis flavis; acheniis pallidis. — *H. pannoso* affine, ab eo foliis acute dentatis, capitulis duplo minoribus, ligulis glabris et praesertim indumento alio, asperulo, nec molli specifice differre videtur.

In mt. Dirphys Euboeae et ad summum cacumen mt. Pentelicon Atticae, rare (Heldr.). — Jul. Aug. ♃.

○○ Ligulae extus pilosae.

12. **H. pannosum** Bois. diagn. IV p. 32, Fl. or. III p. 869, suppl. p. 327; Hal. in z. b. G. 1888 p. 759, Beitr. fl. Thessal. p. 15, in ö. b. Z. 1897 p. 285; Heldr. chlor. Parn. p. 22; Form. in D. bot. Monat. 1890 p. 15, in Ver. Brünn 1896 p. 40; Bald. riv. coll. bot. Alb. 1895 p. 55; Freyn in bull. herb. Bois. III p. 508. — *Andryala lanata* S. et S. pr. II p. 141, non L.. — *H. chalcidicum* Hal. in ö. b. Z. 1890 p. 39, non Bois. et Heldr. — *H. dolopicum* Freyn. et Sint. in bull. herb. Bois. V p. 785 (f. foliis basin versus longius angustatis). — Exsicc.: Orph. herb. a. 1851; Heldr. it. gr. sept. a. 1879, herb. fl. hellen. a. 1896; Sint. et Bornm. it. turc. n. 1343; Sint. it. thessal. n. 1265; Bald. it. alb. epir. III n. 172.

Pilis flexuosis, plumosis, densis pannosum; caule nunc abbreviato, supra partem inferiorem, crebre foliosam ramoso, nudo, 3—5 cephalo, nunc elongato et remotiuscule foliato, vel interdum contracto ad rosulam reducto et monocephalo; foliis amplis, ovato- vel oblongo-spathulatis, obtusis vel acutis, basi sessili attenuatis, integris vel denticulato-repandis, superioribus semiamplexicaulibus; capitulis majusculis; involucri pilis longis flexuosis lanati, eglandulosi phyllis linearibus acutis; ligulis flavis; acheniis pallidis. — Stirps quoad caulis longitudinem, foliorum formam, indumentum valde polymorpha, variat nempe caule plus minus elongato vel abbreviato usque contracto brevissimo, phyllopodo, hypophyllopodo vel aphyllopodo, foliis obovatis usque lanceolatis, indumento breviori pannoso vel longo, jubato, argenteo-sericeo, pilis longius vel brevius plumosis; occurrit insuper flosculis monstrosis, tubulose deformatis.

β. **taygeteum** Bois. et Heldr. diagn. VII. p. 15; Haussk. symb. p. 135; Form. in Ver. Brünn 1897 p. 30; pro sp.; Bois. fl. or. III.

p. 869; Bald. riv. coll. bot. Alb. 1896 p. 75. — Indumentum longius, argenteo-sericeum, praesertim ad margines et costam foliorum et ad caules longe villosum. Occurrit quoque (*v. scapiferum* Bois. l. c. p. 16) caule brevissimo, foliis subrosulatis. — Intermediis ad typum transit. — Exsicc.: Heldr. herb. norm. sc. 47, n. 1451 (*f. scapifera*); Orph. fl. gr. n. 346; Bald. it. alb. epir. IV. n. 398.

γ. **Friwaldii** Rchb. ic. XIX. p. 94 t. 196; Raul. cret. p. 801; Bald. riv. coll. bot. Alb. 1895 p. 55; pro sp.; Bois. fl. or. suppl. p. 327. — *H. versutum* Friv. exs. sec. Fries epicc. p. 77; Raul. cret. p. 801. — Folia angustiora, minora, oblongo-lanceolata vel lanceolata, acuta. — Exsicc.: Bald. it. alb. epir. n. 170.

Ad rupes regionis abietinae et superioris. Epirus: pr. Kalarrytes et Syraku in mt. Peristeri (Bald.); Thessalia: mt. Oxya supra Chaliki (Hal.), pr. Sermeniko (Sint.), Malakasi, mt. Dokimi, pr. Velitsena, Vendista, Kastania in Pindo, pr. Patsios (Form.) et Hagios Dionysios (Sint.) in Olympo, mt. Oeta Phthiotidis (Heldr.); Aetolia: mt. Tymphrestus (Samar.), mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Peloponnesus: mt. Kyllene, Chelmos (Orph.), Olenos, Taygetos (Heldr.); — β. Epirus: mt. Gamila, Micikeli (Bald.); Thessalia: pr. Chaliki, mt. Zygos, Oxya (Haussk.), Ghavellu in Pindo, mt. Pyrgo in Othrys (Form.), Olympus (Heldr.); mt. Dirphys Euboeae, mt. Parnassus, Parnes Atticae, mt. Taygetos (Heldr.); — γ. Epirus: mt. Peristeri (Bald.); Creta (Frid.): mt. Drakona (Rev.). — Forma *monstrosa* flosculis tubulosis in mt. Kiona (Hal.). — Jul. Aug. ♃.

13. **H. Gaudryi** Bois. et Orph. diagn. ser. 2 III. p. 105, Fl. or. III. p. 869; Fr. epicr. p. 75; Heldr. chlor. Parn. p. 22; Naeg. et Pet. archier. p. 331, ubi autores hybridam autumant inter *H. pannosum* et *H. prenanthoides*, sed certissime erronee, nam species ultima in Graecia non provenit; Freyn in bull. herb. Bois. III. p. 511. — Exsicc.: Orph. fl. gr. n. 347.

Pilis elongatis, plumosis sericeo-villosum; caule elongato, foliato, superne divaricatim et dichotome 2—4 cephalo, rarius monocephalo; foliis ab infimis sub anthesi adhuc vigentibus, ad superiora sensim diminutis, acutis, inferioribus lingulato-oblongis, basi attenuatis, remote denticulatis, superioribus oblongis, basi rotundata semiamplexicaulibus, dichotomialibus ad squamas lineares reductis; capitulis mediocribus; involucri nigricantis, breviuscule hirsuti vel villosi, eglandulosi, phyllis lanceolatis, acutiusculis; ligulis flavis; acheniis pallidis. — Caulis 25—40 cm. altus, folia eis praecedentis angustiora, non ut in eo secus caulis partem inferiorem condensata, sed superne sparsa, diminuta, capitula minora, involucri phylla latiora, indumento breviori obducta; sed formae intermediae adesse videntur.

In regione superiori mt. Parnassus pr. Kedrozasteno (Orph.). — Jun. Aug. ♃.

×× Involucrum pilis stellatis minutis cano-tomentellum.

14. **H. thessalum** Form. in Ver. Brünn 1896 p. 40. — Exsicc.: Form. pl. thessal. a. 1895.

Caule erecto, foliato, pilis plumoso-scabris inferne dense superne sparsim obsito, a medio vel superne ramoso, oligocephalo; foliis sericeo-villosis, ab infimis, sub anthesi jam emarcidis, ad superiora sensim diminutis, acutis, inferioribus oblongo-lanceolatis, acute denticulatis, basi attenuatis, superioribus ovato-lanceolatis lanceolatisve, semiamplexicaulibus, summis ad squamas lineares reductis; capitulis mediocribus; involucri stellato-cani, eglandulosi phyllis lanceolatis, acutis; ligulis flavis, extus pilosis; acheniis pallidis. — Praecedenti affine, ab eo caule aphyllopodo, involucro brevissime tomentello, nec pilis elongatis hirsuto certe distinctum, mihi ex unico specimine male praeparato tantum notum.

Thessalia: pr. Tafilvris in mt. Olympo (Form.). — Jul. Aug. ♃.

β. Caulis glaber.

15. **H. oligadenum** Belli ap. Bald. riv. coll. bot. Alb. 1895 p. 55 pro forma *H. Waldsteinii* Tausch. — Exsicc.: Bald. it. alb. epir. III. n. 171.

Caule elato, nigricante, glabro, parte infima folioso, ceterum foliis minutis squamaeformibus munito, superne parce ramoso, 2—4 cephalo; foliis pilis plumosis flexuosis pannosis, oblongis, obtusis, integris vel repando-subdenticulatis, basi attenuatis, infimis sub anthesi adhuc vigentibus; capitulis mediocribus; involucri pedunculisque minute stellato-puberuli, eglandulosi, phyllis lanceolatis, acutis; ligulis flavis, extus pilosis; acheniis pallide fuscis. — *H. plumuloso* Kern in ö. b. Z. 1874 p. 170, dinarico, caule pilis crispulis et involucro praeter indumentum stellatum, pilis glanduliferis brevibus obsito, diverso, proximum. — *H. Waldsteinii* caulibus villosis, involucro hirsuto glandulifero, acheniis fusco-atris longius distat.

In rupestribus ad Jeranovuni in mt. Tsumerka in Epiro (Bald.). Jul. Aug. ♃.

3. Subsectio. *Glutinosa* Koch syn. p. 459. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia viridia, glanduloso-viscida, radicalia sub anthesi saepe adhuc vigentia; caulis scapiformis vel plus minusve foliatus, mono- vel polycephalus; involucrum imbricatum, phyllis acutis; ligulae apice glabrae, pappus rufescenti-albus.

16. **H. scapigerum** Bois. Orph. et Heldr. diagn. ser. 2, III. p. 103, Fl. or. III. p. 870, suppl. p. 328; Hal. in z. b. G. 1888 p. 759, in ö. b. Z. 1897 p. 285; Heldr. chlor. Parn. p. 22; Rouy ill. p. 39 t. 121. — Exsicc.: Orph. fl. gr. n. 345.

Caulibus erectis, 5—15 cm. altis, simplicibus et monocephalis vel furcato 2—3 cephalis, pilis densis glanduliferis et longioribus scabris hispidis, nudis vel 1—2 foliis minutis obsitis; foliis fere omnibus rosulatis, oblongo-lanceolatis, basi sessili attenuatis, grosse sinuato-dentatis, glandulosis; capitulis majusculis; involucri dense glandulosi pilisque

longioribus hispidi phyllis nigricantibus lanceolato-linearibus, acutis; ligulis flavis, extus pilosulis; acheniis pallidis.

In rupestribus regionis alpinae, rare. Phthiotis: mt. Oeta (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Orph.); Achaia: mt. Kyllene (Orph.). — Jun. Aug. ♃.

4. Subsectio. *Pulmonaroidea* Koch syn. p. 454. — Rhizoma cylindricum, induratum, nunquam soboliferum; folia viridia vel glaucescentia, nunquam tomentosa vel glandulosa, radicalia sub anthesi adhuc vigentia; caulis scapiformis vel plus minusve foliatus, plerumque pleiocephalus; involucrum irregulariter imbricatum, phyllis acutis; ligulae apice glabrae; pappus rufescenti-albus.

a. Folia pleraque lanceolata vel oblongo-lanceolata, basi longe attenuata.

17. **H. Sartorianum** Bois. et Heldr. diagn. VII. p. 15; Bois. fl. or. III. p. 870 probab. tantum pro p.; Heldr. chlor. Parn. p. 22; Haussk. symb. p. 134; Bald. riv. coll. bot. Alb. 1895 p. 15, 1896 p. 75. — Exsicc.: Hal. it. gr. a. 1888.

Glaucovirens; caule scapiformi, mono — rarius dicephalo, 1—2 squamis linearibus aucto, pilis albis, scabridulis, basi nigricantibus, diametro suo longioribus, aliquando evanidis, tomento raro brevissimo et interdum apicem versus pilis glandulosis rarissimis intermixtis hirsuto; foliis radicalibus rosulatis, lanceolatis oblongo-lanceolatisve, acutis, basi longe attenuatis, integris vel remote denticulatis, superne glabris, subtus et margine pilis albis longis patulis ciliatis; capitulis mediocribus; involucri phyllis anguste linearibus, pilis albis longis hirsutis, eglandulosis; ligulis flavis, extus pilosulis; acheniis nigris.

β. **Leithneri** Heldr. et Sart. in sched. pro sp.; Bois. fl. or. suppl. p. 328. — Huc spectare videtur: *H. eriobasis* Freyn. et Sint. in bull. herb. Bois. V. p. 787. — Folia radicalia sublatiora, exteriora interdum oblonga, obtusa, caulis superne glandulosus, capitula submajora. — Exsicc.: Orph. fl. gr. n. 1151, teste Heldr. ipso.

In rupibus regionis alpinae. Epirus: mt. Smolika, Tsumerka (Bald.), pr. Chaliki (Haussk.); mt. Olympus, Parnassus, Olenos, Taygetos (Heldr.); — β. mt. Parnassus (Orph.). — Jul. Aug ♃.

b. Folia pleraque ovata vel oblonga, basi subcordata vel rotundata, rarius attenuata.

18. **H. incisum** Hoppe in Sturm Heft n. 39; Koch. syn. p. 457. — *H. oligocephalum* Hal. Beitr. fl. Epir. p. 29, non Arv. Touv. — Huc probabiliter ut forma: *H. murorum* Sieb. avis rem. p. 5; Raul. cret. p. 800; = *H. murorum v. pilosissimum* Bois. fl. or. III. p. 873. — Exsicc.: Heldr. it. gr. sept. a. 1879; Hal. it. gr. sec. a. 1893. —

Caule erecto, 1—oligocephalo, nudo vel foliis 1—2 diminutis, linearibus, sessilibus obsito, inferne parce piloso, superne pedunculis involucroque canescentibus, hirsutis, saepe simulque glanduliferis; foliis radi-

calibus rosulatis, petiolatis, ovatis ovato-oblongisve, sinuato-dentatis, saepe purpureo-maculatis, subtus margineque hirsutis, basi rotundatis subcordatisve; capitulis mediocribus; involucri phyllis acuminatis; ligulis flavis, glabris; acheniis fusco-nigris. — Specimina graeca ab eis alpium non differunt.

In regione subalpina et alpina. Epirus: in herbidis lapidosis ad summum cacumen mt. Peristeri (Hal.); Phthiotis: in subalpinis mt. Oeta (Heldr.); nec non in Creta (Sieb.): ad Hellinoseli, Stravopodia, Hagion Pneuma, si. *H. murorum* Sieb. et Raul. huc. pertineat. — Jul. Aug. ♃.

19. **H. murorum** L. sp. p. 802 p. p.; Bois. fl. or. III. p. 871; Heldr. chlor. Parn. p. 22; Haussk. symb. p. 134. — *H. vulgatum* Hal. Beitr. fl. Thessal. p. 15, non Fr. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Hal. it. gr. sec. a. 1893.

Caule elato, apice corymboso, unifolio, inferne parce piloso, superne pedunculis involucroque canescentibus, hirsutis, glanduliferis; foliis radicalibus rosulatis, petiolatis, ovatis ovato-oblongisve, sinuato-dentatis, laete viridibus, subtus praesertim ad nervos hirsutis, basi subcordatis, rotundatis vel attenuatis; capitulis mediocribus; involucri phyllis acuminatis; ligulis flavis, glabris; acheniis fusco-nigris. — Praecedenti elatius, apice corymbosus, folia multo majora, subtus minus hirsuta, caulina radicalibus subsimilia.

In rupestribus, fagetis subalpinis. Thessalia: mt. Oxya supra Chaliki (Hal.), mt. Ghavellu, Zygos Haussk.); mt. Parnassus et Kyllene (Heldr.). — Jul. Aug. ♃.

20. **H. abietinum** Bois. et Heldr. diagn. ser. 2 III. p. 105 pro var. *H. vulgati*; Reut. in Bois. fl. or. III. p. 872; Heldr. chlor. Parn. p. 22; Haussk. symb. p. 134. — Exsicc.: Heldr. herb. n. 2731.

Caule tenui, 3—5 folio, apice corymboso, inferne parce piloso, superne pedunculis involucroque canescentibus, hirsutis, glanduliferis; foliis membranaceis, orbiculatis ovatisve, minute denticulatis, subtus ad nervos hirtis, basi cordatis rotundatisve, radicalibus longiuscule petiolatis, caulinis subsessilibus vel breviter petiolatis; capitulis parvis; involucri phyllis acutis; ligulis flavis, glabris; acheniis fusco-nigris. — A praecedentibus forma foliorum, caule foliato et capitulis minoribus discedit.

In faucibus umbrosis regionis abietinae mt. Parnassus ad Dipotamo (Heldr.) et in fagetis jugi Zygos in Pindo tymphaeo (Haussk.). — Jul. Aug. ♃.

5. Subsectio. *Italica* Fr. epicr. p. 107. — Rhizoma cylindricum, induratum, nunquam soboliferum, folia viridia, nunquam tomentosa vel glandulosa, radicalia sub anthesi saepe jam emarcida; caulis foliatus, inflorescentia racemosa; involucrum imbricatum, phyllis obtusis; ligulae apice glabrae, pappus rufescenti-albus.

α. Caulis stricte ramosus, rarius simplex, ramis tenuibus, saepe virgatis, elongatis, glabris vel parce pilosis.

21. **H. bracteolatum** S. et S. pr. II, p. 135; Bois. fl. or. III. p. 874; Heldr. chlor. Parn. p. 22; Freyn. in bull. herb. Bois. V. p. 788. — *H. Reinholdii* Heldr. et Sart. in Boiss. diagn. ser. 2 V. p. 117. — Exsicc.: Orph. herb. n. 44 (Athos).

Caule erecto, angulato, glabro vel pilis albis longis patulis obsito, simplici vel stricte et tenuiter ramoso; foliis parce hirtulo-asperulis, radicalibus et caulinis inferioribus, sub anthesi adhuc vigentibus, oblongo-lanceolatis, acutis, in petiolum patule pilosum attenuatis, grosse et remote dentatis vel basi pinnatifidis, foliis caulinis superioribus parvis, sessilibus, lanceolatis, obsolete 1—2 denticulatis, integrisve; capitulis tenuibus, paucifloris, pedunculo eis pluries longiori suffultis; involucri phyllis glabris vel medio dorso parce et minutissime hirtulis; ligulis flavis, glabris; acheniis pallidis.

β. **laxum** Bois. fl. or. III. p. 874; Haussk. symb. p. 134. — Caulis simplex; folia abbreviata; capitula 2—3, longe pedunculata. — Exsicc.: Heldr. herb. norm. n. 724.

γ. **pelium.** — *H. Koracis* Haussk. symb. p. 134, non Bois. — Caulis virgato-ramosus, rarius simplex, ramis elongatis, glabris, capitulis subsessilibus. — Exsicc.: Heldr. it. thessal. n. 67; Haussk. it. gr. a. 1885; Sint. it. thessal. n. 1283.

In nemorosis, fagetis regionis montanae et subalpinae. Phocis: mt. Parnassus pr. Rachova (Guicc.) sec. Boissier, sed specimina non vidi. — *β.* Thessalia: ad monasterium Korona et Meteora (Haussk.); Eurytania: pr. Laspi (Heldr.); — *γ.* Thessalia: mt. Pelion pr. Volo (Haussk.), Makrinitza (Heldr.), Zagora (Aphentulis); Attica: mt. Parnes (Heldr.). — Jun. Sept.

22. **H. Koracis** Bois. fl. or. suppl. p. 328. — Exsicc.: Heldr. it. gr. sept. a. 1879.

Caule erecto, angulato, parce et minute piloso, in paniculam laxam, strictam abeunte; foliis minute et adpresse setulosis, oblongis, obtusiusculis, grosse dentato-crenatis, radicalibus sub anthesi emarcidis et inferioribus in petiolum attenuatis, ceteris sessilibus, sensim diminutis, summis bracteaeformibus, integris; capitulis tenuibus, paucifloris, ad ramorum brevium paniculae apicem 2—5 confertis, pedunculis involucro brevioribus suffultis; involucri phyllis pedunculisque praeter glandulas sessiles pilis glanduliferis hirtis; ligulis flavis, glabris; acheniis pallidis. — Praecedenti affine, ab eo praeter inflorescentiam thyrsoideam, involucro glanduloso praesertim discedit.

In regione abietina et superiori mt. Korax Aetoliae (Heldr.). — Jul. Aug. ♃.

b. Caulis simplex, crassiusculus, foliatus, stricte racemosus, patule pilosus.

23. **H. racemosum** W. et K. pl. rar. hung. II p. 211 t. 193; Freyn in bull. herb. Bois. V p. 790. — Icon: Rchb. t. 178. — Exsicc.: Sint. it. or. a. 1889 p. 1823, it. thessal. n. 1197 b.

Caule erecto, simplici, stricte racemoso, praeter pubem minimam pilis patulis obducto; foliis praeter setas secus petiolos et subtus ad nervos sparsas glabriusculis, infimis sub anthesi saepe jam emarcidis, inferioribus rosulatis, ovatis vel oblongo-lanceolatis, acutis, denticulatis, in petiolium attenuatis, ceteris diminutis, sessilibus, ovato-oblongis; capitulis mediocribus, pedunculo brevissimo vel elongato suffultis; involucri pallidi phyllis glabriusculis vel parce setulosis, interdum parce glandulosis; ligulis flavis, glabris; acheniis fuscis.

In silvis montanis Thessaliae: pr. Sermeniko in Pindo et in Olympo (Sint.). — Aug. Sept.

21 × 23 **H. bracteolatum × racemosum.** — *H. sermenikense* Freyn et Sint. in bull. herb. Bois. V p. 790. — Probabiliter hybrida harum specierum; differt a *H. bracteolato* caule superne parce ramoso, patentim piloso breviterque puberulo, foliis inferioribus acute dentatis, involucro hirtulo et interdum parce glanduloso; a *H. racemoso* caule tenui virgato, capitulis minoribus, paucifloris, longe pedunculatis et acheniis pallidis. — In silvis montanis pr. Sermeniko in Pindo dolopico, nec non in mt. Pelion (Sint.). — Exsicc.: Sint. it. thessal. n. 1179.

24. **H. crinitum** S. et S. pr. II p. 134; Bois. fl. or. III p. 875; Heldr. chlor. Parn. p. 22; Form. in Ver. Brünn 1896 p. 40. — *H. eriopus* Bois. et Heldr. diagn. ser. 2 III p. 106. — Exsicc.: Sint. et Bornm. it. turc. n. 1346 et 1347.

Caule erecto, simplici, praeter pubem minimam pilis patulis crinitum; foliis ad petiolos et subtus ad costam crinitis, infimis sub anthesi jam emarcidis, inferioribus rosulatis, ovato-oblongis, acutis, acute dentatis, in petiolum attenuatis, ceteris sessilibus, saepe semiamplexicaulibus, cito decrescentibus; capitulis mediocribus, pedunculo brevi suffultis; involucri pallidi phyllis minute puberulis, parce breviterque setulosis, interdum parce glandulosis, ligulis flavis, glabris; acheniis pallidis. — Valde affine *H. racemoso*, a quo differt hispiditate et acheniis pallidis. Formae intermediae adesse videntur.

In montosis et silvaticis. Thessalia: mt. Olympus, mt. Pelion (Heldr.); mt. Parnassus (Heldr.); mt. Malevo Laconiae (Orph.). — Aug. Sept.

Obs. Species dubia, nomine tantum nota: *H. nudicaule* Sieb. avis rem. p. 5.

95. Andryala L. gen. n. 915.

1. **A. dentata** S. et S. pr. II. p. 140, Fl. gr. IX. p. 7 t. 811; Bois. fl. or. III. p. 879; Haussk. symb. p. 134; Heldr. chlor. Thera p. 16. — *A. integrifolia* Urv. enum. p. 103, non L. — Exsicc.: Heldr. herb. norm. n. 854.

Caule erecto, foliato, superne vel a parte inferiori ramoso, laxiuscule corymboso, breviter tomentello, glandulifero; foliis tenuiter tomentosis, infimis in petiolum brevem attenuatis, obovatis, obtusis,

repandis, ceteris oblongo-lanceolatis acutatis, dentatis vel pinnatifidis, sessilibus; capitulis parvis, globosis; involucro pilis glanduliferis hispidissimo, phyllis acuminatis basilaribus paucissimis vel nullis; ligulis flavis; acheniis nigris.

In cultis, vineis, arenosis, ad litora maris, rare. Argolis: pr. Vromolimni in peninsula Methana (Heldr.), insula Poros (Reinh.), Andros (Sart.), Melos (Sibth.), Ios, Thera, Therasia (Heldr.) et Kameni (Urv.). — Maio, Jul. ⊙.

LXIII. Ordo. **Ambrosiaceae** Cass. in bull. soc. phil. 1815 p. 173.

1. **Ambrosia** L. gen. n. 1057.

1. **A. maritima** L. sp. p. 988; Sieb. avis p. 5; Ch. et B. exp. p. 270, Fl. pelop. p. 63; Raul. cret. p. 780; Bois. fl. or. III. p. 252; Heldr. fl. cephal. p. 44, chlor. Thera p. 16; Bald. in nov. giorn. bot. 1894 p. 97; Form. in Ver. Brünn 1895 p. 25. — Icon: Rchb. XIX. t. 216. — Exsicc.: Reverch. pl. cret. n. 70.

Villoso-canescens; caule stricto, divaricatim ramoso, folioso; foliis petiolatis, in lacinias oblongas, obtusiuscule dentatas bipinnati-partitis; capitulis ad apicem ramorum dense spicatis; spicis basi femineis, superne masculis; involucro viridi, femineo cornubus rectis 4—5 echinato; floribus flavis.

In arenosis maritimis. Epirus: pr. Prevesa (Bald.); Cephalonia: pr. Drapano, Lixuri (Heldr.); Argolis, Maina (Chaub.); Messenia: pr. Gialova (Zahn); Cycladum insula Paros (Sart.), Andros, Naxos, Thera (Heldr.); Creta (Sieb.): pr. Platania (Rev.), Retymo (Raul.). — Jul. Sept. ⊙.

2. **Xanthium** L. gen. n. 1056.

1. **X. strumarium** L. sp. p. 987; S. et S. pr. II. p. 234; Ch. et B. exp. p. 270, Fl. pelop. p. 63; Friedr. Reise p. 273; Fraas fl. class. p. 216; Raul cret. p. 780; Bois. fl. or. III. p. 251; Heldr. fl. cephal. p. 44; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1895 p. 25, 1896 p. 39, 1897 p. 30; Haussk. symb. p. 143. — Icon: Fl. dan. t. 970.

Caule erecto, asperulo, ramoso, foliato, inermi; foliis asperis, longe petiolatis, cordato-ovatis, irregulariter lobato-dentatis; capitulis apice caulis et in axillis ramorum glomerato-racemosis, superioribus masculis; involucris fructiferis ovatis vel oblongis, breviter pubescentibus, spinis apice uncinatis obsitis, rostris binis superatis.

α. **typicum.** — Involucri apex spinis destitutus, rostra subdivergentia. — Exsicc.: Orph. herb. a. 1850. —

β. **antiquorum** Wallr. Beitr. bot. I. p. 229; Haussk. symb. p. 143; pro sp.; Bois. fl. or. III. p. 252; Form. in Ver. Brünn 1896

p. 39, 1897 p. 30. — Involucrum usque ad apicem spinis obsitum, rostra magis divergentia. — Exsicc.: Orph. fl. gr. n. 752; Sint. et Bornm. it. turc. n. 1323 b.

Ad vias, in ruderatis totae Graeciae, varietas ut videtur typo vulgatior. — Jun. Sept. ⊙.

2. **X. spinosum** L. sp. p. 987; Ch. et B. exp. p. 270, Fl. pelop. p. 63; Friedr. Reise p. 263; Weiss in z. b. G. 1869 p. 47; Bois. fl. or. III. p. 252; Heldr. fl. cephal. p. 44, in Sitzungsb. acad. Wiss. Berl. 1883 p. 8; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1895 p. 25, 1896 p. 39, 1897 p. 30; Hal. Beitr. fl. Aetol. p. 8; Haussk. symb. p. 143. — Icon: Rchb. t. 214. — Exsicc.: Heldr. pl. fl. hellen. a. 1879.

Caule erecto, puberulo, ramoso, foliato, spinis tripartitis, longis, flavicantibus armato; foliis subtus canis, superne praeter nervum medium canescentem viridibus, breviter petiolatis, oblongo-lanceolatis, cuneatis, indivisis vel trilobis, lobo medio multo longiore; capitulis masculis terminalibus; involucris femineis ad axillas foliorum subsolitariis, tandem cernuis, oblongis, puberulis, spinis apice uncinatis obsitis, rostro saepius unico recto superatis.

Ad vias, in ruderatis regionis inferioris et submontanae per totam ut videtur Graeciam. — Jul. Sept. ⊙.

LXIV. Ordo. **Campanulaceae** Juss. gen. p. 163.

Dispositio tribuum generumque:

1. Tribus. *Campanuleae* Don gard. dict. III. p. 732. — Capsula lateraliter poris dehiscens.

a. Corolla 5 loba vel 5 fida.

α. Antherae connatae; corolla campanulata, 5 loba; stylus pilosus; stigmata 3; capsula 3 locularis, poris 3, lateralibus ad basin dehiscens.

1. Symphyandra DC.

β. Antherae liberae.

× Corolla campanulata vel infundibuliformis, rarius tubulosa vel subrotata; stylus pilis collectoribus obsitus; stigmata 3—5; capsula turbinata vel obconica, raro subglobosa vel disciformis, 2—5 locularis, poris 3—5, lateralibus ad basin, ad medium vel sub apice dehiscens.

2. Campanula L.

×× Corolla rotata; stylus pilosus, stigmata 3; capsula linearis vel oblonga, prismatica, 3 locularis, poris 3, lateralibus, versus apicem sitis dehiscens.

3. Specularia Heist.

b. Corolla 5 partita.

α. Stylus glaber, stigmatibus in capitulum sphaericum connatis; capsula 3 locularis, poris 3, lateralibus, versus medium sitis dehiscens.

4. Petromarula DC.

β. Stylus pilosus, stigmatibus 2—3 liberis.

× Corolla profunde 5 partita; stigmata 3, filiformia; capsula 3 locularis, poris 3, lateralibus, apice, ad medium vel basi dehiscens.

5. Podanthum Don.

×× Corolla ultra medium 5 partita; stigmata 2—3, brevissima; capsula 2—3 locularis, poris 2—3, lateralibus basi dehiscens.

6. Diosphaera Buser*).

2. Tribus. *Wahlenbergieae* Endl. gen. pl. p. 514. — Capsula apice intra calycem dehiscens.

a. Corolla 5 fida, tubuloso-campanulata; antherae liberae; stigmata 2—3; capsula 2—3 locularis, superne intra calycem fragmentis irregularibus lacera.

7. Edrajanthus DC.

b. Corolla profunde 5 partita, laciniis lineari-lanceolatis; antherae basi connatae; stigmata 2; capsula 2 locularis, apice intra calycem foramine lato dehiscens.

8. Jasione L.

1. Tribus. **CAMPANULEAE** Don. gard. dict. III. p. 732.

1. Symphyandra DC. mon. camp. p. 365.

1. **S. cretica** DC. mon. camp. p. 366 t. 8; Raul. cret. p. 804; Bois. fl. or. III. p. 888; Spreitz. in z. b. G. 1890 p. 296. — *Campanula nutans* Sieb. avis p. 3, rem. p. 3, non Lam. — Exsicc.: Spreitz. it. cret. n. 75; Bald. it. cret. alt. n. 262.

Glabra; caule simplici, inferne parce foliato; foliis radicalibus et caulinis inferioribus amplis, longe petiolatis, ovato-cordatis, acutis, acute et grosse subduplicatim serratis, superioribus diminutis, breviter petiolatis subsessilibusve, floralibus anguste lanceolatis, serratis, pedunculo multo longioribus; floribus paucis, in racemum terminalem dispositis, brevissime pedunculatis, solitariis, nutantibus; calycis exappendiculati laciniis lanceolato-acuminatis, integris vel denticulatis, tubo obconico-hemisphaerico triplo longioribus, erectis; corolla infundibuliformi-campanulata, coerulea, glabra, breviter lobata, calycis laciniis duplo longiore. — Caulis 30—45 cm. altus, foliorum radicalium lamina 10—15 cm. longa, 8—10 cm. lata, corolla 4 cm. longa.

Ad rupes faucium Cretae (Sieb.): pr. Enneachoria, Hagia Rumeli, Samaria, Aradhena (Raul.). — Jun. Jul. ♃.

2. **S. Sporadum** Hal. in ö. b. Z. 1895 p. 461, 1897 p. 96. — Icon: Hal. in ö. b. Z. 1895 p. 461. — Exsicc.: Heldr. et Hal. fl. sporad. a. 1896.

Glabra; caule simplici, subnudo; foliis radicalibus parvis, longe petiolatis, repando-crenatis, obtusis, aliis reniformibus, aliis ovato-cordatis, caulinis minimis, squamaeformibus, sessilibus, floralibus pedunculo brevioribus; floribus paucis, in racemum terminalem dispositis, breviter

*) Genus affine *Trachelium* differt praeter alias notas (cf. Bus. in bull. herb. Bois. II. p. 511) corolla breviter 5 loba, filamentis glabris, nec hispido-villosulis, antheris inclusis vel vix exsertis, nec longe exsertis.

pedunculatis, solitariis, nutantibus; calycis exappendiculati laciniis lineari-setaceis, integerrimis, tubo depresso-conico 3—5 plo longioribus, demum patulis vel reflexis; corolla e basi angustata ventricoso-campanulata, coerulea, glabra, breviter lobata, calycis laciniis dimidio usque triplo longiore. — Caules tenues, 10—25 cm. alti, foliorum lamina 1—5 cm. longa, 1—4 cm. lata, saepe purpurascens, corolla 2 cm. longa, medio ventricosa. — Affinis *S. samothracicae* Deg. in ö. b. Z. 1891 p. 333 pro var. *S. creticae;* Hal. in ö. b. Z. 1895 p. 463; caulibus foliatis, foliis irregulariter serratis, racemo composito, calycis laciniis lanceolato-linearibus callosis diversae.

In rupestribus Sporadum insulae Jura — Gyaros — veterum (Reiser). — Maio, Jun. ♃.

2. Campanula L.

Dispositio specierum.

1. Sectio. *Medium* Bois. fl. or. III. p. 893. — Capsula poris vel valvulis prope basin sitis dehiscens.

1. Subsectio. *Quinqueloculares* Bois. l. c. — Stigmata 5, capsula 5 locularis calycis sinus appendiculati.

a. Folia radicalia ovata vel oblonga, crenato-dentata, petiolis non lobulatis.

α. Corolla ampla, tubo 2—3 cm. lato.

1. C. pelviformis Lam. **2. C. corymbosa** Desf.

β. Corolla mediocris, tubulosa, tubo 1 cm. lato.

3. C. tubulosa Lam. **4. C. saxatilis** L.

b. Folia inferiora vel saltem radicalia irregulariter laciniata vel lyrata, petiolis nempe lobulatis (cf. *C. rupestris γ gracilis*).

α. Corolla lata et breviter campanulata, ampla, antice 4—5 cm. lata.

5. C. laciniata L.

β. Corolla tubuloso-campanulata.

× Calycis laciniae tubo aequilongae; corolla majuscula; stylus exsertus.

6. C. Reiseri Hal.

×× Calycis laciniae tubo 3—4 plo longiores; corolla mediocris; stylus inclusus.

○ Corolla cylindrica, pulchre cyanea, lobis patentibus.

7. C. anchusiflora S. et S.

○○ Corolla pallide coerulea, a basi sensim ampliata, lobis erecto-patulis.

8. C. Celsii DC. **9. C. Andrewsii** DC. **10. C. rupestris** S. et S.

2. Subsectio. *Triloculares* Bois. l. c. — Stigmata 3, capsula 3 locularis.

a. Calycis sinus appendiculati.

α. Folia inferiora basi cordata.

11. C. incurva Auch.

β. Folia basi non cordata.

× Flores sessiles, fasciculati.

12. C. lingulata W. et K.

×× Flores pedunculati, non fasciculati.

⊙ Perennes, caespitosae.

. Caules remote foliati; corolla majuscula, 3 cm. longa, campanulata.

13. C. **oreadum** Bois. et Heldr. 14. C. **rupicola** Bois. et Spr.

.. Caules foliosi; corolla parva, 10—15 mm. longa, tubuloso-infundibuliformis.

15. C. **heterophylla** L. 16. C. **calaminthifolia** Lam.

○○ Annua.

17. C. **dichotoma** L.

b. Calycis sinus exappendiculati.

α. Perennes.

× Flores sessiles, in fasciculos axillares subsessiles et capitulum terminale, bracteis ovatis acuminalis arcte involucratum congesti; capsula erecta.

○ Radix cylindrica, subrepens, simplex vel ramosa, caulem solitarium erectum vel plures e basi adscendente erectos edens; folia sub anthesi saepius jam emarcida.

18. C. **glomerata** L. 19. C. **stenosiphon** Bois. et Heldr. 20. C. **foliosa** Ten.

○○ Radix fusiformis, verticalis, simplex, caulem centralem erectum et laterales flagelliformes, decumbentes vel arcuato-adscendentes edens; folia radicalia rosalata, sub anthesi adhuc vigentia.

21. C. **tymphaea** Haussk.

×× Flores pedicellati, solitarii vel fasciculati, in racemum simplicem vel compositum dispositi.

⊙ Caules erecti, elati crassiusculi.

. Corolla campanulata; capsula nutans.

22. C. **athoa** Bois. et Heldr. 23. C. **bononiensis** L.

.. Corolla subrotata; capsula erecta.

24. C. **versicolor** Andr.

○○ Caules humiles, tenues, saepius decumbentes.

. Folia difformia, radicalia reniformi-cordata, caulina lineari-lanceolata; capsula nutans.

25. C. **rotundifolia** L.

.. Folia subconformia; capsula erecta.

; Calyx papillis pellucidis dense obsitus, glaber.

26. C. **Hawkinsiana** Haussk. et Heldr.

;; Calyx laevis, glaber vel hirsutus.

, Folia cordato-orbiculata vel cordato-ovata.

27. C. **cephallenica** Feer. 28. C. **Sartorii** Bois. et Heldr.

,, Folia oblonga, basi attenuata.

29. C. **radicosa** Chaub. et Bory.

β. Annuae; capsula nutans.

× Corolla majuscula, calyce duplo longior.

30. C. **scutellata** Griseb. 31. C. **drabifolia** S. et S.

×× Corolla minuta, calyce aequilonga.

32. C. **erinus** L.

2. Sectio. *Rapunculus* Bois. fl. or. III. p. 895. — Capsula 3locularis, poris vel valvulis prope apicem, rarius versus medium sitis dehiscens.

a. Perennes.

α. Radix cylindrica.

33. C. **persicifolia** L. 34. C. **abietina** Griseb. et Sch.

β. Radix napiformis.

35. C. Spruneriana Hampe. **36. C. Sibthorpiana** Hal.

b. Biennes vel annuae.

α. Folia radicalia numerosissima, in rosulam densam congesta; calycis laciniae tubo aequilongae vel parum longiores.

37. C. aizoon Bois. et Sprun.

β. Folia radicalia pauca; calycis laciniae tubo saepius multo longiores.

× Corolla infundibuliformis vel campanulata.

○ Radix fusiformis, carnosa.

38. C. rapunculus L.

○○ Radix cylindrica, ramosa.

. Biennis; folia radicalia sub anthesi adhuc vigentia; panicula subcorymbosa; calycis laciniae tubo 2—3 plo longiores.

39. C. patula L.

.. Annuae; folia radicalia sub anthesi jam emarcida; panicula racemosa vel thyrsoidea; calycis laciniae tubo multoties longiores.

40. C. sparsa Friv. **41. C. sphaerothrix** Griseb.

×× Corolla breviter obconica, subrotata.

42. C. phrygia. **43. C. ramosissima** S. et S.

1. Sectio. *Medium* Bois. fl. or. III. p. 893. — Capsula poris vel valvulis prope basin sitis dehiscens.

1. Subsectio. *Quinqueloculares* Bois. l. c. — Stigmata 5, capsula 5 locularis; calycis sinus appendiculati.

a. Folia radicalia ovata vel oblonga, crenato-dentata, petiolis non lobulatis.

α. Corolla ampla, tubo 2—3 cm. lato.

1. **C. pelviformis** Lam. dict. I. p. 586; Desf. choix. pl. p. 41 t. 31; Sieb. avis p. 3, rem. p. 3, in Flora V. I. p. 15; DC. mon. camp. p. 223; Raul. cret. p. 803; Bois. fl. or. III. p. 900 p. p. — Hirsuta; caulibus adscendentibus, simplicibus vel ramulis paucis, unifloris auctis; foliis ovatis, acutis, denticulato-serratis, radicalibus petiolatis, caulinis subsessilibus; calycis setosi laciniis ovato-acuminatis, tubo triplo longioribus, appendicibus ovato-rotundatis, laciniis longitudine subaequalibus; corolla campanulata, ampla, calyce 2—2 ½ plo longiore, griseo-coerulea, glabriuscula; capsula . . . — Secundum Sieber in Flora l. c. forma tantum a capris mutilata *C. corymbosae,* corollis majoribus eis *C. medii* aequimagnis, a qua tamen ex descriptione et icone Desfontainesii, caulibus adscendentibus, appendicibus ovato-rotundatis, tubum obtegentibus, corolla ampliore, tubo 3 cm. lato, medio inflato specifice differre videtur.

In collinis Cretae (Tourn.): mt. Karadagh (Raul.). — Apr. Maio. ⊖ N. v.

2. **C. corymbosa** Desf. choix. pl. p. 40 t. 30; DC. mon. camp. p. 223; Raul. cret. p. 803. — Exsicc.: Heldr. pl. cret. n. 1424; Bald. it. cret. alt. n. 335.

Hirsuta; caulibus erectis, superne ramosis, corymbosis; foliis ovatis, acutis, acute denticulato-serratis, radicalibus petiolatis, caulinis subsessilibus; calycis setosi laciniis ovatis, longe acuminatis, tubo triplo longioribus, appendicibus ovato-rotundatis, laciniis longitudine dimidio brevioribus; corolla campanulato-tubulosa, calyce duplo longiore, coerulea, undique sed ad nervos longius setulosa; capsula hemisphaerica, appendicibus auctis, reticulatis occultata. — Caules 25—50 cm. alti, corollae tubus 20—25 mm. latus, medio non inflatus. — Secundum Boissier fl. or. III. p. 900 forma praecedentis floribus minoribus.

In collinis, locis umbrosis Cretae (Tourn.): ad Kaenuriokorio, Kritsa, ad ripas Sklavotiae pr. Piskokephalo (Raul.), pr. Mirabello (Heldr.), mt. Lazari in mt. Lassiti (Bald.). — Apr. Jul. ⊙.

β. Corolla mediocris, tubulosa, tubo 1 cm. lato.

3. **C. tubulosa** Lam. dict. I. p. 586; Desf. choix. pl. p. 42 t. 32; DC. mon. camp. p. 222; Raul. cret. p. 803; Bois. fl. or. III. p. 900. — *C. pelviformis β. micrantha* DC. mon. camp. p. 222; pr. V. p. 459. — Exsicc.: Sieb. herb. cret. sub *C. pelviformis*; Bald. it cret. alt. n. 37.

Caulibus e radice fusiformi divergentibus vel adscendentibus, brevibus, patule pilosis, 1—3 floris; foliis pilosis, radicalibus ovato-oblongis, crenato-dentatis, in petiolum attenuatis, caulinis ovato-lanceolatis, subsessilibus sessilibusve, serrato-dentatis; calycis setulosi, reticulati, laciniis ovato-lanceolatis, acutis, tubo duplo longioribus, appendicibus ovato-rotundatis, laciniis dimidio brevioribus; corolla violaceo-cinerea, pubescente, calyce duplo longiore; capsula hemisphaerica, appendicibus auctis, reticulatis omnino occultata. — Caules 3—15 cm. longi, corolla 20—25 mm. longa.

In rupium fissuris Cretae (Tourn.): inter Askyphos et Anopolis (Raul.), supra Kani Kastelli distr. Temenos (Bald.). — Maio, Jun. ⊙.

Obs. Praecedenti affinem, sed ab ea foliis argute subbiserratis, calycis laciniis tubo 3plo longioribus, appendicibus pilis densissimis tomentum album formantibus vestitis, corolla longiore antice ampliata, certe specifice diverisam, quod scio speciem adhuc non descriptam, *C. carpatha* nominandam, sistit planta a Pichler a. 1883 in insula Karpathos lecta et a Boissier in fl. or. suppl. p. 330, nec non a Rouy ill. pl. p. 13 t. 40 erronee ad *C. saxatilem* L., caulibus foliisque subglabris, foliis oblongo-spathulatis crenulatis, calyce adpresse brevissimeque canescente, appendicibus parvis tubo 2—3plo brevioribus, corolla velutina antice subangustata diversissimam, ducta.

4. **C. saxatilis** L. sp. p. 167; S. et S. pr. I. p. 139; DC. mon. camp. p. 232; Raul. cret. p. 803; Bois. fl. or. III. p. 898; Bald. viagg. Creta p. 73. — *C. tubulosa* Heldr. in ö. b. Z. 1880 p. 346; Spreitz. in z. b. G. 1887 p. 663; Forsyth in bull. herb. Bois. V. p. 399; non Lam. — Icon: Jaub. et Sp. ill. or. t. 392. — Exsicc.: Spreitz. it. jon. a. 1880; Bald. it. cret. n. 229; Bois. fl. gr. a. 1898.

Pallide virens, caulibus a rhizomate crasso, caespitoso erectis vel adscendentibus, fragilibus, glabris, simplicibus vel ramoso-racemosis, paucifloris; foliis glabris vel minutissime puberulis, crenulatis, radicalibus rosulatis, oblongo-spathulatis in petiolum attenuatis, caulinis sessilibus, oblongis lineari-oblongisve; calycis adpresse canescentis laciniis triangulari-lanceolatis, acutis, tubo 2—3 plo longioribus, appendicibus obtusis, laciniis multo brevioribus; corolla laete coerulea, velutina, calyce duplo longiore; capsula obconica, appendicibus non reticulatis non occultata. — Species pulchra, eximia; a praecedente caulium et foliorum glabrescentia, calyce adpresse canescente, appendicibus parvis, corolla laete coerulea velutina, antice angustata diversissima.

β. **Simonellii** Bald. viagg. Creta p. 73. — Caules suberosi, brevissimi, uniflori; folia puberula caulina approximata. — Exsicc.: N. v.

In rupium fissuris regionis inferioris, rarissime. Cythaera: in fauce Kalamo (Spreitz.), pr. Kapsali (Reiser); Creta (Sibth.): ad promontorium Maleka, cum varietate (Bald.). — Jun. Jul. ⊙ an ♃.

b. Folia inferiora vel saltem radicalia irregulariter laciniata vel lyrata, petiolis nempe lobulatis.

α. Corolla late et breviter campanulata, ampla, antice 4—5 cm. lata.

5. **C. laciniata** L. sp. p. 165; DC. mon. camp. p. 220; Bois. fl. or. III. p. 896. — Icon: Tourn. voy. p. 99; Jaub. et Sp. ill. or. t. 421, optima; Rouy ill. t. 65. — Exsicc.: Veneris in Orph. fl. gr. n. 1119.

Minute puberula, pallide virens; caulibus elatis, crassis, laxe racemoso-paniculatis; foliis inferioribus ovato-lanceolatis, in petiolum lobulatum decurrentibus, grosse laciniatis, laciniis ovatis, obtuse dentatis, foliis superioribus sursum diminutis ovatis, sessilibus, irregulariter dentatis; paniculae ramis brevibus, 2—4 floris; calycis laciniis triangulari-lanceolatis, tubo sublongioribus; appendicibus ovatis, obtusis, tubi longitudine; corolla pallide coerulea, puberula, calyce duplo longiore; capsula sphaeroidea, appendicibus occultata.

Ad rupes scopuli Cardiotissa pr. insulam Pholegondros Cycladum (Tourn.). — Apr. Majo. ⊙.

β Corolla tubuloso-campanulata.

× Calycis laciniae tubo aequilongae; corolla majuscula, stylus exsertus.

6. **C. Reiseri** Hal. in ö. b. Z. 1896 p. 14, 1897 p. 96. — Exsicc.: Heldr. et Hal. fl. Sporad. n. 1896,

Pubescens; caulibus adscendentibus vel decumbentibus, flexuosis, elongatis, laxe et subunilateraliter racemoso-spicatis, simplicibus vel parce breviterque ramulosis; foliis radicalibus ignotis, caulinis ovato- vel oblongo-spathulatis, crenato-dentatis, inferioribus in petiolum, parce lobulatum attenuatis, superioribus sursum minoribus, sessilibus; floribus bre-

viter pedunculatis, racemum elongatum, simplicem vel subcompositum formantibus; calycis dense setulosi laciniis triangulari-lanceolatis, tubo aequilongis, appendicibus ovatis, acutis, tubo aequilongis; corolla puberula, pallide coerulea, tubuloso-campanulata, antice ampliata, calyce 3—4 plo longiore. — Caules 25—45 cm. longi, crassiusculi, folia inferiora cum petiolo ad 9 cm. longa, corolla 4 cm. longa, tubus antice 15—20 cm. latus. — A sequentibus caulibus crassioribus, calycis laciniis brevibus, corolla manifeste majori, antice manifeste ampliata et stylo exserto discedit.

β. **Leonis** Hal. in ö. b. b. Z. 1897 p. 96. — Caulis et folia glabra; flores simpliciter racemosi. — Exsicc.: Heldr. et Hal. fl. Sporad. a. 1896.

In Sporadum insula Jura (Leon.). — Maio, Jun. ⊙.

XX Calycis laciniae tubo 3—4 plo longiores; corolla mediocris, stylus inclusus.

O Corolla cylindrica, pulchre cyanea, lobis patentibus.

7. **C. anchusiflora** S. et S. pr. I. p. 141, Fl. gr. III. p. 9 t. 212; DC. mon. camp. p. 221; Bois. fl. or. III. p. 897; Heldr. in bot. Centralbl. 1890 n. 46 p. 212; Hal. in ö. b. Z. 1896 p. 13. — Exsicc.: Heldr. herb. norm. n. 1056.

Breviter pubescens, subincana; caulibus adscendentibus vel erectis, ramosis, paniculatis, multifloris; foliis radicalibus lyratis, segmentis obtuse crenatis, terminali magno, ovato-cordato, lateralibus parvis, secus petiolum lobulatum decurrentibus, foliis caulinis diminutis, sessilibus, obovatis; floribus pedunculatis; calycis laciniis triangulari-lanceolatis, tubo duplo longioribus, appendicibus rotundatis, tubo multo brevioribus; corollae extus pubescentis tubo cylindrico, lobis ovatis vel ovato-oblongis, obtusiusculis, patentibus duplo longiore; capsula turbinata. — Sequenti simillima et ab ea in herbario vix discernenda, sed sec. Heldreich, qui plantam vivam observavit, corolla pulchre cyanea, apice vix ampliata, limbo horizontali, specifice distincta.

Ad muros, in rupium fissuris. Attica: ad muros coenobii Panagia Cliston in mt. Parnes (Heldr.); Argolis: insula Hydra (Sibth.); Cycladum insula Pholegandros (Sibth.). — Apr. Maio. ⊙.

OO Corolla pallide coerulea, a basi sensim ampliata, lobis erecto patulis.

8. **C. Celsii** DC. mon. camp. p. 217; Hal. in ö. b. Z. 1896 p. 11, 1897 p. 96. — *C. tomentosa* Vent. descr. pl. jard. Cels. p. 18 t. 18; Bois. fl. or. III. p. 897 p. p.; Haussk. p. 143 p. p.; Heldr. in bot. Centralbl. 1890 n. 46 p. 213, *α. typica* et *β. diffusa*, Fl. Aegina p. 308; non Lam. — *C. eriantha* Hampe in Flora XXV. p. 76. — Huc vel ad sequentes pertinere videtur: *C. lyrata* Link. in Linn. IX. p. 579; Ch. et B. fl. pelop. p. 14; non Lam. — Exsicc.: Heldr. herb. norm. n. 345 et 1057, herb. dimorph. n. 71.

Breviter pubescens, subincana; caulibus adscendentibus vel decumbentibus, flexuosis, elongatis, laxe et subunilateraliter racemoso-spicatis, simplicibus vel parce ramulosis; foliis radicalibus oblongo-spathulatis, in petiolum lobulatum decurrentibus, crenato-dentatis, laciniatis vel lyratis, segmento terminali oblongo, acutiusculo, lateralibus parvis, foliis caulinis diminutis, ovatis, infimis petiolatis, ceteris sessilibus; calycis laciniis ovato-lanceolatis, tubo 2—4 plo longioribus, appendicibus parvis, tubo saepius multo brevioribus; corollae extus pubescentis tubo antice parum ampliato, calyce 2—3 plo longiore; capsula turbinata.

Ad rupes regionis inferioris et montanae. Attica: mt. Lycabettus, Parnes, Pentelicon, Hymettns (Heldr.); in Petalium insulis, insula Aegina, in Peloponneso, insula Hydra, Cyclades (Heldr.); Sporadum insula Skopelos (Leon.). — Apr. Jun. ⊙.

9. **C. Andrewsii** DC. mon. camp. p. 220; Hal. in ö. b. Z. 1895 p. 463. — *C. laciniata* Andrews bot. repos. VI. t. 385; S. et S. pr. I. p. 141; ? Ch. et B. fl. pelop. p. 14; non L. — *C. tomentosa* Bois. fl. or. III. p. 897, p. p.; Haussk. symb. p. 143 p. p.; Heldr. in bot. Centralbl. 1890 p. 213, *γ. bracteosa*; non Lam. nec. Vent. — Exsicc.: Orph. fl. gr. n. 660 et 1081; Sint. it. thessal. n. 1; Dörfl. fl. gr. n. 192.

Molliter villosa, cinerascens; caulibus adscendentibus vel decumbentibus, flexuosis, racemoso-paniculatis; foliis radicalibus et caulinis inferioribus lyratis, segmento terminali maximo, orbiculato vel ovato, inciso-crenato, lateralibus parvis, incisis integrisve, secus petiolum lobulatum decurrentibus, foliis caulinis sensim diminutis, superioribus sessilibus, ovatis, crenato-dentatis; calycis laciniis triangulari-lanceolatis, tubo 3—4 plo longioribus, appendicibus ovatis, tubo aequilongis; corollae extus pubescentis tubo antice ampliato, calyce 2—3 plo longiore. — A praecedente meo sensu, indumento molliter villoso foliorum inferiorum segmento rotundato et corolla majore antice manifeste ampliato, specifice diversa.

In rupium fissuris regionis inferioris et montanae. Thessalia: pr. Pharsalus (Haussk.)?, ad lacum Karla (Reis.), pr. Volo (Sint.); mt. Parnassus (Heldr.), pr. Lebadea Boeotiae (Orph.); Acrocorinthus (Bois.)?; Argolis: pr. Nauplia (Orph.), peninsula Methana (Haussk.)?. — Apr. Jul. ⊙.

10. **C. rupestris** S. et S. pr. I. p. 142, Fl. gr. III. p. 10 t. 213; DC. mon. camp. p. 218; Ch. et B. exp. p. 71, Fl. pelop. p. 14; Friedr. Reise p. 278; Ung. Reise p. 125; Hal. in ö. b. Z. 1896 p. 13. — *C. lanuginosa* Willd. en. hort. berol. I. p. 213; cf. Vatke in Linnaea 1874 p. 705, non Lam. — *C. tomentosa β. brachyantha* Bois. fl. or. III. p. 898; Heldr. in bot. Centralbl. 1890 p. 213; Haussk. symb. p. 143. — *C. tomentosa* Hal. Beitr. fl. Achaia p. 26, non Lam. — Exsicc.: Heldr. pl. fl. hellen. a 1887, 1895, 1897; Pichl. pl. gr. a. 1876.

Incano-cinerea; caulibus decumbentibus, flexuosis, laxe et subunilateraliter racemoso-spicatis, simplicibus vel parce ramulosis, saepius abbreviatis; foliis radicalibus lyratis, segmento terminali multo majori, ovato, crenato, lateralibus minutis, secus petiolum decurrentibus, foliis caulinis sessilibus, ovatis vel spathulatis, crenatis; calycis laciniis triangulari-lanceolatis, tubo 3—4 plo, appendicibus ovatis, tubo aequilongis; corollae extus pubescentis, tubo antice parum ampliato, calyce saepius parum longiore. — A praecedente, quacum formis intermediis ut videtur conjuncta est, foliis radicalibus multo minoribus, segmentis lateralibus minimis, paucis, interdum fere nullis instructis, indumento saepius densiori, sed breviori, corollis manifeste minoribus antice parum ampliatis, calycis lacinias saepe parum superantibus discedit.

β. **virescens.** — Folia tenera, parva, parce puberula, virescentia, radicalia uno alterove lobulo tantum ad petiolum munita. — Exsicc.: Orph. herb. a. 1850; Pichl. pl. gr. a. 1876.

γ. **gracilis** Heldr. ap. Hal. in ö. b. Z. 1896 p. 13. — Caules graciles, simplices, folia minuta, cano-tomentosa, omnia indivisa, minute crenulata, radicalia in petiolum integrum attenuata. — Exsicc.: Heldr. pl. fl. hellen. an. 1880.

δ. **calycina** Heldr. in bot. Centralbl. 1890 p. 214, pro var. *C. tomentosae.* — Virescens, caulibus abbreviatis, calycis laciniis valde acuminatis, corollam aequantibus. — Exsicc.: N. v.

In rupestribus regionis montanae. Locris: ad Thermopylas (Heldr.); Euboea: pr. Steni (Pichl.), Kyme (Ung.); Boeotia (Sibth.): pr. Lebadia (Heldr.); Attica: mt. Hymettus (Heldr.), Pentelicon (Haussk.), Lycabettus (Ung.); Achaia: pr. Sudena, mt. Kastro pr. Kalavryta, in valle Voreikos, pr. Megaspilaeon (Hal.), mt. Kyllene pr. Trikala, pr. Diakophto (Heldr.); Arcadia: pr. Zulatika (Nied.); Messenia: pr. Phigalea (Chaub.); Laconia: pr. Selitza (Zahn), Gythion, mt. Taygetos, pr. Androvista (Chaub.); Cycladum insula Naxos (Chaub.). — β. mt. Malevo (Orph.); — γ. mt. Cithaeron Atticae (Heldr.); — δ. Euboea: pr. Kumi (Heldr.). — Maio, Jul. ⊙.

Obs. *C. stricta* S. et S. pr. I. p. 140, non Lab. = *C. lyrata* Lam. dict. I. p. 588, in ditione nostra non crescit.

2. Subsectio. *Triloculares* Bois. fl. or. III. p. 893. — Stigmata 3, capsula 3 locularis.

a. Calycis sinus appendiculati.

α. Folia inferiora basi cordata.

11. **C. incurva** Auch. in DC. pr. VII. p. 464. — *C. Leutweinii* Heldr. cat. sem. hort. Athen. a. 1860 p. 7, in Sitzungsb. acad. Wiss. Berlin 1883 p. 4; Bois. fl. or. III. p. 896. — Exsicc.: Aucher n. 1864; Heldr. it. thessal. n. 65. —

Pubescens; caulibus elongatis, laxe racemoso-paniculatis, rarius simplicibus; foliis ovatis ovato-oblongisve, obtuse crenatis, radicalibus et

inferioribus longiuscule petiolotis, basi cordatis, superioribus breviter petiolatis, basi truncatis, summis sessilibus; floribus magnis; calycis pilosiusculi laciniis late ovato-triangularibus, tubo 3plo longioribus, appendicibus ovatis, obtusis, tubo subaequilongis; corolla pallide coerulea, ample campanulata, calyce subtriplo longiore, extus ad nervos et ore setulosa, ceterum glabra; capsula hemisphaerica, appendicibus auctis, reticulatis occultata. — Caules 20—40 cm. alti, folia radicalia cum petiolo 10—15 cm. longi, corolla 4 cm. longa, 20—25 mm. lata. — Foliis cordatis ab omnibus affinibus distincta.

In dumetis, rupestribus regionis submontanae et montanae. Thessalia: mt. Olympus juxta sinum Thermaicum (Auch.), mt. Ossa pr. Tzaisi (Heldr.), mt. Pelion pr. Zagora (Aphentulis); Euboea septentrionalis: supra Orobias et in mt. Kavallari supra Agianako (Leutwein). — Jul. Aug. ⊙.

β. Folia basi non cordata.

× Flores sessiles, fasciculati.

12. **C. lingulata** W. et K. pl. rar. hung. I. p. 65 t. 64; Bois. fl. or. III. p. 926; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1896 p. 38. — Exsicc.: N. v.

Hispida; caulibus erectis, simplicibus; foliis subundulatis, repando-dentatis, inferioribus oblongo-spathulatis, in petiolum attenuatis, caulinis basi ovata sessilibus, oblongis vel lanceolatis; floribus 5—9, terminalibus, fasciculatis, foliis summis bracteatis; calycis laciniis oblongis, obtusis, appendicibus ovatis, tubo sublongioribus, laciniis vix brevioribus; corolla anguste tubulosa, limbo infundibuliformi, coerulea, glabra vel ad nervos setulosa, laciniis calycis 3—4plo longiore.

β. **cichoracea** S. et S. pr. I. p. 140, Fl. gr. III. p. 7 t. 209 pro sp.; Griseb. spic. II. p. 288; Bois. fl. or. III. p. 926; Haussk. symb. p. 143; Bald. riv. coll. bot. Alb. 1896 p. 77. — *C. capitata* Sims. in Curt. bot. mag. XXI. p. 811 t. 811. — Caulibus elatis, ramosis; fasciculis florum axillaribus et terminalibus. — Exsicc.: Sint. et Bornm. it turc. n. 1351 et 1352; Sint. it. thessal. n. 323. —

In saxosis apricis. Thessalia: mt. Dokimi in Pindo, pr. Patsios in Olympo (Form.). — β. Epirus: mt. Profeta Ilias et Maria distr. Ljaskovik (Bald.); Thessalia (Sibth.): pr. Tsungeri et Malakasi (Haussk.), Kastreiki ad Meteora (Sint.), mt. Olympus (Auch.) ad Hagios Dionysios et Litochori (Sint.). — Apr. Jul. ⊙.

×× Flores pedunculati, non fasciculati.

○ Perennes, caespitosae.

. Caules remote foliati; corolla majuscula, 3 cm. longa, campanulata.

13. **C. oreadum** Bois. et Heldr. diagn. ser. 2 III. p. 107, Fl. or. III. p. 908. — Exsicc.: Heldr. herb. n. 1851; Orph. fl. gr. n. 655.

Adpresse hirta, pumila; caulibus tenuibus; erectis vel adscendentibus, 1—5 floris; foliis integris, radicalibus oblongo-spathulatis, obtusis, in

petiolum attenuatis, caulinis lanceolatis, sessilibus, acutis; floribus longiuscule pedunculatis; calycis dense hirsuti laciniis lanceolatis, acutis, integris, tubo 3—4 plo longioribus, appendicibus parvis, tubo brevioribus; corolla coerulea, campanulata, glabriuscula, laciniis calycis multo longiore. — Rhizoma crassum, caules 5—8 cm. alti, corolla 3 cm. longa, 20—25 mm. lata. — Species pulchra, egregia.

In rupestribus regionis alpinae mt. Olympus prope cacumina (Heldr.). — Jul. Aug. ♃.

14. **C. rupicola** Bois. et Sprun. diagn. VII. p. 17, Fl. or. III. p. 908; Hal. in z. b. G. 1888 p. 759, in ö. b. Z. 1897 p. 286; Heldr. chlor. Parn. p. 22. — Exsicc.: Heldr. herb. norm. n. 224; Orph. fl. gr. n. 418; Dörfl. fl. gr. n. 185.

Parce pubescens; caulibus tenuibus, flexuosis, decumbentibus, 1—3 floris; foliis denticulatis, inferioribus ovatis vel orbiculatis, basi breviter cuneatis, longe petiolatis, superioribus ovatis vel oblongis, breviter petiolatis; floribus breviter vel longiuscule pedunculatis; calycis glabri, laciniis ovato-lanceolatis, obtusiusculis, ciliatis, saepe denticulatis, tubo 2—3 plo longioribus, appendicibus minutissimis; corolla coerulea, campanulata, glabra, laciniis calycis multo longiore. — Caules 5—15 cm. longi, corolla 3 cm. longa 2 cm. lata, pulcherrime coerulea.

In rupestribus regionis alpinae. Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Sprun.). — Jul. Aug. ♃.

.. Caulis foliosi; corolla parva, 10—15 mm. longa, tubuloso-infundibuliformis.

15. **C. heterophylla** L. sp. ed. 2 p. 240; S. et S. pr. I. p. 139, Fl. gr. III. p. 6 t. 208; Ch. et B. fl. pelop. p. 15; Bois. fl. or. III. p. 912, suppl. p. 331. — Exsicc.: Heldr. pl. Cyclad. a 1881.

Viridis, inferne glabrata, superne minutissime puberula; caulibus e rhizomate verticali, carnosulo decumbentibus, fragilibus, foliosis, subsimplicibus; foliis integris, crassiusculis, radicalibus oblongo-spathulatis, obtusis, in petiolum attenuatis, caulinis subsessilibus, orbiculatis; floribus parvis, axillaribus, breviter pedicellatis, inferioribus 2—3 nis vel omnibus solitariis, racemosis; calycis puberuli laciniis triangulari-lanceolatis, tubo subaequilongis, appendicibus brevissimis; corolla tubuloso-infundibuliformi, coerulea, pubescente, calyce subduplo longiore.

In rupibus insulae Amorgos pr. Panagia (Heldr.) et in scopulo adjacente Cheros (Tourn. voy. p. 95 cum t.); indicatur in Ant. ion. V. 206 quoque in mt. Pantocrator et pr. Caragallo in insula Corcyra, sed certe ex confusione. — Jun. Jul. ♃.

16. **C. calaminthifolia** Lam. dict. I. p. 585; Desf. choix pl. p. 37 t. 27; Bois. fl. or. III. p. 912, Heldr. in ö. b. Z. 1898 p. 184. chlor. Mykon p. 247. — Exsicc.: Heldr. herb. norm. n. 1454.

Tota pubescens, cinerea; caulibus e rhizomate verticali, carnosulo decumbentibus, fragilibus, foliosis, simplicibus vel breviter ramulosis; foliis crenato-dentatis, crassiusculis, radicalibus oblongo-spathulatis, obtusis, in petiolum attenuatis, caulinis subsessilibus, ovatis; floribus parvis,

axillaribus, breviter pedicellatis, ad apicem caulis subconfertis vel ad apicem ramulorum solitariis; calycis tomentelli laciniis triangulari-lanceolatis, tubo subaequilongis, appendicibus brevibus; corolla tubuloso-infundibuliformi, coerulea, tomentella, calyce duplo longiore. — Differt a praecedente foliis pubescentibus, cinereis, crenato-dentatis.

β. **Olivieri** DC. mon. camp. p. 233 t. 9, pro sp. — Folia caulina orbiculata vel ovato-orbiculata, saepius integra; flores ut videtur submajores. — Exsicc.: Orph. fl. gr. n. 1118. —

In rupium fissuris. Cycladum insula Naxos (Tourn.) pr. Apirantos (Heldr.); — β. ad muros templi Iovis insulae Delos (Oliv.), insula Pholegandros (Veneris). — Jun. Jul. ♃.

○○ Annua.

17. **C. dichotoma** L. amoen. IV. p. 306; S. et S. pr. I. p. 141, Fl. gr. t. 211; Ch. et B. fl. pelop. p. 15; Friedr. Reise p. 277; Bois. fl. or. III. p. 929. —

Setis patulis hispidula; caule erecto, dichotome ramoso; foliis oblongis, integris vel repando-denticulatis; floribus solitariis, terminalibus et alaribus, breviter pedicellatis; calycis laciniis lanceolatis, acuminatis, appendicibus acutis, calycis tubo longioribus; corolla coerulea, glabra, vel parce setulosa, campanulata, calyce sesquilongiore; capsula nutante. ♂: In collibus apricis. Attica: pr. Athenas (Sibth.); insula Poros (Friedr.); sed a recentioribus non lecta. — Jun. Jul. ⊙. N. v.

b. Calycis sinus exappendiculati.

α. Perennes.

× Flores sessiles, in fasciculos axillares subsessiles et capitulum terminale, bracteis ovatis acuminatis arcte involucratum congesti; capsula erecta.

⊙ Radix cylindrica, subrepens, simplex vel ramosa, caulem solitarium erectum vel plures e basi adscendente erectos edens; folia sub anthesi saepius jam emarcida.

18. **C. glomerata** L. sp. p. 166. — *C. foliosa* Form. in Ver. Brünn 1897 p. 28, non Ten. — Exsicc.: Form. pl. thessal. a. 1896.

Hirta vel glabrata; caule erecto vel basi adscendente, foliato; foliis minute crenatis, inferioribus a basi cordata vel rotundata ovato-lanceolatis, longe petiolatis, petiolis non alatis, foliis superioribus sessilibus, oblongo-lanceolatis, basi cordata semiamplexicaulibus; floribus in fasciculos axillares et terminales congestis; bracteis ovato-acuminatis, floribus brevioribus; calycis laciniis plus minusve pubescentibus, tubo 3—4plo longioribus; corolla infundibuliformi, violacea, hirta, calycis laciniis sesqui vel duplo longiore. — Corolla 20—25 mm. longa.

β. **Vlachavae** Bois. et Orph. in Orph. fl. gr. exs. n. 653. — Bracteae et calycis laciniae breviter setuloso-ciliatae; corolla magna, glabriuscula, circa 3 cm. longa.

In subalpinis Thessaliae, rarissime: mt. Ghavellu in Pindo (Form); — β. mt. Olympus pr. Mavros Longhos (Orph.). — Jul. Aug. ♃.

19. **C. stenosiphon** Bois. et Heldr. diagn. VII. p. 18. — *C. glomerata β. stenosiphon* Bois. fl. or. III. p. 928. — Exsicc.: Heldr. herb. norm. n. 1562.

Pilis albis substrigosis hispida; caulibus e collo pluribus, erectis vel adscendentibus, foliatis; foliis minute crenatis, inferioribus oblongis vel lanceolatis, anguste in petiolum attenuatis, superioribus sessilibus ovato-oblongis, basi cordata semiamplexicaulibus; floribus in fasciculos axillares 2—3 floros et terminales 4—7 floros congestis; bracteis late ovato-acuminatis, recurvis, floribus subbrevioribus; calycis laciniis margine et ad nervum medium albo-strigosis, tubo secus nervos strigoso, ceterum glabro, longioribus; corolla infundibuliformi, violacea, basi parce hispida, calycis laciniis duplo longiore. — Corolla circa 20 mm. longa. — Indumento strigoso-hispido, foliis radicalibus angustis in petiolum attenuatis, corolla minore, superne glabriuscula praecedenti discrepat.

β. **majoriflora**. — Bracteae floribus subduplo breviores; corolla magna, 25 mm. longa. — Forma parallela varietatis praecedentis, a qua indumento magis strigoso et praesertim foliis radicalibus angustioribus, in petiolum sensim attenuatis differt. — Exsicc.: Orph. fl. gr. n. 659.

In regione abietina. Peloponnesus: in faucibus Langada, pr. Anavryti et Neraidovuni in mt. Taygetos (Heldr.); — β. mt. Malevo pr. Castanitza (Orph.). — Jun. Jul. ♃.

20. **C. foliosa** Ten. fl. nap. I. p. 71 t. 18; Hal. Beitr. fl. Thessal. p. 15; Haussk. symb. p. 143. — Exsicc.: Sint. it. thessal. n. 903.

Pilosiuscula; caule erecto, foliato; foliis irregulariter biserratis, radicalibus a basi subcordata vel rotundata ovatis, caulinis inferioribus ovato-oblongis, radicalibusque longe petiolatis, petiolis alatis, foliis superioribus sessilibus; floribus in fasciculos terminales, multifloros congestis; bracteis ovato-acuminatis, flores subaequantibus vel subsuperantibus, margine et ad nervos albo-ciliatis, ceterum glabris; calycis laciniis albo-ciliatis, tubo glabro 3—4 plo longioribus; corolla infundibuliformi, violacea, glabra, calycis laciniis sesquilongiore. — Differt a praecedentibus foliis multo majoribus, irregulariter biserratis, petiolis alatis, bracteis majoribus, capitulo multo majori, multifloro.

In subalpinis, ad silvarum margines, in fagetis. Thessalia: mt. Zygos, pr. Chaliki (Haussk.) et mt. Oxya (Hal.) in Pindo tymphaeo. — Jul. Aug. ♃.

○○ Radix fusiformis, verticalis, simplex, caulem centralem erectum et laterales flagelliformes, decumbentes vel arcuato-adscendentes edens; folia radicalia rosulata, sub anthesi adhuc vigentia.

21. **C. tymphaea** Haussk. in Mitth. thür. bot. Ver. V. p. 87, symb. p. 143. — *C. flagellaris* Hal. Beitr. fl. Epir. p. 30 t. 3, Beitr.

fl. Thessal. p. 15; Form. in Ver. Brünn 1896 p. 38; non Humb. Bonpl. et Kunth nov. gen. Am. III. p. 301. — *C. glomerata* Bald. riv. coll. bot. Alb. 1895 p. 57, non L. — *C. parnassica* Bald. riv. coll. bot. Alb. 1896 p. 76, non Bois. et Spr. — Exsicc.: Hal. it. gr. II. a. 1893; Bald. it. alb. epir. III. n. 179, IV. n. 184; Sint. it. thessal. n. 667 et 901.

Glabrata vel parce setulosa, viridis; caulibus foliatis; foliis minute crenulatis, radicalibus rosulatis, oblongo-spathulatis, in petiolum attenuatis, caulinis oblongis vel ovato-lanceolatis, sessilibus, semiamplexicaulibus; floribus in fasciculos terminales 4—7 floros, interdum quoque axillares 2—3 floros congestis; bracteis ovatis, breviter acuminatis, saepe denticulatis, margine ciliatis; calycis laciniis ciliatis, tubo glabro sublongioribus; corolla infundibuliformi, violacea, glabra, calycis laciniis 2—3 plo longiore. — Caulis 10—35 cm. altus, laterales breviores tenuiores, saepe stoloniformes, folia radicalia 5 cm. longa, 1 cm. lata, corolla 15—20 mm. longa. — Species eximia, in Pindo endemica, a *C. glomerata* et affinibus, nec non a *Edrajantho parnassico*, quocum etiam ab auctoribus, incredibile dictu, confusa fuit, certe distinctissima. Formae a capris vel ovibus mutilatae caespitem depressum, caulibus abbreviatis, paucifloris formant et a planta intacta habitu valde differunt.

In pascuis subalpinis et alpinis. Epirus: mt. Gamila, Smolika Micikeli (Bald.), Peristeri, Tsumerka (Hal.); Thessalia: mt. Baba, Zygos (Haussk.), Oxya (Hal.), Dokimi, Mandra Hodza in Pindo (Form.). — Jul. Aug. ♃.

×× Flores pedicellati, solitarii vel fasciculati, in racemum simplicem vel compositum dispositi.

○ Caules erecti, elati, crassiusculi.

× Corolla campanulata; capsula nutans.

22. **C. athoa** Bois. et Heldr. diagn. ser. 2 III. p. 110; Hal. Beitr. fl. Thessal. p. 15, in ö. b. Z. 1897 p. 286. — *C. trachelium β. orientalis* Bois. fl. or. III. p. 922; Heldr. in Sitzungs. acad. Wiss. Berl. 1883 p. 4, chlor. Parn. p. 23; Form. in Ver. Brünn 1896 p. 38. -- *C. trachelium v. parviflora* Form. in D. bot. Monat. 1890 p. 13, in Ver. Brünn 1896 p. 38. — *C. trachelium v. balcanica* Form. in Ver. Brünn 1897 p. 29. — Huc probabiliter: *C. trachelium* Mazz. in ant. ion. V. p. 202; Haussk. symb. p. 143; Bald. riv. coll. bot. Alb. 1896 p. 76; vix L.; nec non *C. urticaefolia* Mazz. l. c.; non Turra, nec Schm. — Exsicc.: Heldr. herb. norm. n. 620, it. thessal. n. 49; Sint. it. thessal. n. 1014.

Tota pilis patulis strigosa; caule argute angulato, simplici vel ramuloso, foliato; foliis scabris, duplicato-serratis, inferioribus petiolatis, e basi cordata triangulari-elongatis, superioribus sessilibus, oblongis vel lanceolatis; floribus in axillis foliorum superiorum subsessilibus, fasciculatis; calycis tubo angulato, pilis patulis praesertim ad angulos strigosissimo, laciniis erectis, lanceolatis, strigosis, tubo duplo longioribus; corolla tubuloso-campanulata, glabra, violacea, calycis laciniis parum vel

duplo longiore. — Subspecies austro-orientalis *C. trachelii* L., sed strigosior, florum fasciculi subsessiles, corolla diminuta, 15—20 mm. tantum longa et angustior.

In umbrosis, silvaticis regionis montanae et subalpinae. Epirus: pr. Cepelovon distr. Zagorion (Bald.); Thessalia: ad monasterium Korona (Haussk.), pr. Han Uranaeos (Sint.) et Kukleus in valle Penei, pr. Sermenikon et mt. Ghavellu in Pindo, pr. Kerasia Sina, mt. Cuka, Phlambures et mt. Mitrica in mt. Chassia, mt. Ossa (Form.), Olympus (Orph.), Pelion (Heldr.); Euboea: mt. Telethrion (Heldr.); Eurytania: pr. Karpenisi (Samarit.), mt. Chelidoni pr. Mikrochorio (Heldr.); Aetolia: mt. Korax (Tunt.); mt. Parnassus (Heldr.); Peloponnesus: mt. Kyllene (Heldr.), Malevo (Orph.); Corcyra: mt. St. Stefana et pr. Sidari, si plantae Mazziarii supra enumeratae huc spectent. — Jul. Aug. ♃.

Obs. *C. rapunculoides* L. sp. p. 165 a Mazziari in ant. ion. V. p. 202 pr. Curamades in Corcyra certe ex confusione quadam indicatur.

23. **C. bononiensis** L. sp. p. 165; Bois. fl. or. III. p. 923; Form. in D. bot. Monat. 1890 p. 13. — Icon: Rchb. t. 237.

Canescens; caule tereti; simplici vel ramuloso, folioso; foliis inaequaliter crenatis vel serratis, superne adpresse pilosis, subtus canotomentellis, ab infimis cordato-oblongis, petiolatis, ad intermedia sessilia, semiamplexicaulia, ovata vel acuminata diminutis, summis linearibus; floribus 2—3 nis, solitariisque, breviter pedunculatis, racemum longum, spiciformem, simplicem vel subpaniculatum formantibus; calycis minutissime puberuli laciniis patentibus, breviter triangulari-lanceolatis, tubo sublongioribus; corolla infundibuliformi, glabra, coerulea, calycis laciniis multo longiore.

In dumosis montanis. Thessalia: ad Han Kriavris in valle Penei (Form.), ad radices mt. Olympus (Orph.). — Jul. Aug. ♃ N. v.

×× Corolla subrotata; capsula erecta.

24. **C. versicolor** Andrews bot. repos. t. 396; S. et S. pr. I. p. 138, Fl. gr. III. p. 5 t. 207; Ch. et B. exp. p. 87, Fl. pelop. p. 14; Mazz. in ant. ion. V. p. 200; Ten. annot. fl. gr. p. 8; Heldr. Nutzpfl. p. 28 et 79, Fl. cephal. p. 50, chlor. Parn. p. 23; Bois. fl. or. III. p. 914; Spreitz. in z. b. G. 1887 p. 664; Hal. in z. b. G. 1888 p. 759. —

Caulibus e rhizomate crasso, carnoso erectis, curvatis, crassiusculis, foliatis, in racemum compositum thyrsoideum abeuntibus; foliis subcoriaceis, crenatis vel serrulatis, inferioribus longe petiolatis, ovato-vel oblongo-cordatis, ceteris sursum brevius petiolatis, oblongis, basi attenuatis; florum fasciculis breviter pedunculatis, 3—7 floris; pedicellis calyce brevioribus; calycis laciniis anguste lineari-lanceolatis, patentibus vel reflexis, tubo 3plo longioribus; corolla pallide coerulea, fundo violacea, in lobos ovatos partita, calycis laciniis duplo longiore.

α. **typica.** — Omnino glabra. — Exsicc.: Heldr. herb. norm. n. 515, 1356 et 1455.

β. **thessala** Bois. fl. or. III. p. 915. — Folia argutius serrata, basi attenuata nec cordata; flores minores; calycis laciniae abbreviatae. Formis intermediis ad typum transit. Planta glabra vel (f. *tomentella* Hal. in ö. b. Z. 1892 p. 372) ad caules et folia undique adpresse puberula vel tomentella. — Exsicc.: Sint. et Bornm. it turc. n. 1353, 1354 et 1847, in Baen. herb. europ. n. 7628.

In rupibus regionis submontanae in subalpinam adscendens. Thessalia: pr. Sermeniko (Sint.); Eurytania: mt. Chelidoni (Heldr.), mt. Tymphrestus (Sprun.), mt. Vardusia supra Vustinitza (Tuntas), pr. Mesolonghion (Nied.); mt. Kiona supra Segditza (Hal.), mt. Parnassus (Heldr.); Achaia: mt. Olenos, Kyllene (Heldr.); Arcadia: mt. Menalus et Diaforti (Chaub.); Laconia: mt. Malevo (Heldr.), pr. Mistra, mt. Taygetos (Bois.); Cephalonia: ad Hagios Varvaros supra Drapano, inter Valeriano et Markopulo, pr. Steno (Heldr.); Leucas: ad promontorium Zuana (Spreitz.); Corcyra: insula Othonus (Mazz.); — *β.* Thessalia: pr. Kalabaka (Haussk.), mt. Olympus pr. Hagios Dionysios, Litochoron (Heldr.), in fauce Megarema (Sint.). — Jul. Sept. ♃.

○○ Caules humiles, tenues, saepius decumbentes.

. Folia difformia, radicalia reniformi-cordata, caulina lineari-lanceolata; capsula nutans.

25. **C. rotundifolia** L. sp. 163; S. et S. pr. I. p. 137; Ch. et B. exp. p. 70, Fl. pelop. p. 14; ? Mazz. in ant. ion. V. p. 200; Bois. fl. or. III. p. 919; Bald. riv. coll. bot. Alb. 1895 p. 56; Form. in Ver. Brünn 1896 p. 37, 1897 p. 29; Hal. in ö b. Z. 1897 p. 286; non L. — Exsicc.: Bald. it. alb. epir. III. n. 180; Sint. it. thessal. n. 1269.

Glabra; radice verticali, caudiculosa, caules floriferos, adscendentes rosulasque steriles edente; foliis rosularum longe petiolatis, orbiculatis, cordatis vel ovato-lanceolatis, denticulatis, caulinis lanceolatis linearibusque, sessilibus, integris; floribus erectis, longiuscule pedunculatis; calycis laciniis lineari-filiformibus, erecto-patulis, tubo subduplo longioribus; corolla campanulata, violacea, calyce triplo longiore. — Caulis plantae graecae humilis, 1—2 florus.

In saxosis regionis superioris. Epirus: mt. Peristeri (Bald.); Thessalia: mt. Kokkino Lithari pr. Sermeniko (Sint.), mt. Karava (Form.) in Pindo; Aetolia: mt. Korax (Tunt.); Euboea: mt. Hagios Elias (Sart.); Achaia: mt. Kyllene (Orph.); Laconia (Sibth.); *C. rotundifolia* indicatur quoque in insula Samatrachi ad septentrionem Corcyrae (Mazz.), sed dubito quin identica sit cum planta alpina e Graecia. — Jul. Aug. ♃.

.. Folia subconformia; capsula erecta.

; Calyx papillis pellucidis dense obsitus, glaber.

26. **C. Hawkinsiana** Haussk. et Heldr. in Mitth. thür. bot. Ver. V. p. 87, symb. p. 144; Form. in D. bot. Monat. 1890 p. 14, in Ver. Brünn 1896 p. 38; Bald. riv. coll. bot. Alb. 1896 p. 76; Hal. Beitr.

fl. Thessal. p. 15. — *C. Halácsyana* Bald. riv. crit. coll. bot. Alb. 1892 p. 56. — Exsicc.: Heldr. herb. norm. n. 856; Bald. it. alb. IV. n. 248; Sint. it. thessal. n. 622 et 622b. —

Radice verticali, caudiculosa; caulibus e collo numerosissimis, decumbentibus vel adscendentibus, flexuosis, laxe foliatis, simplicibus vel ramulosis, glabris vel rarius (*β. scabriuscula* Haussk. l. c.) inferne pilis papillaribus plus minus obsitis; foliis carnosulis, glabris, suborbiculatis, ovatis vel ovalibus, integris vel leviter crenulatis, inferioribus in petiolum brevem attenuatis, superioribus diminutis, subsessilibus; floribus solitariis, longiuscule pedunculatis; calycis glabri laciniis lanceolatis, erecto-patulis, laevibus, tubo papilloso subtriplo longioribus; corolla subrotata, glabra, intense coeruleo-violacea, in lobos ovatos partita, calyce 2—3plo longiore. — Species elegans, egregia, nulla cum alia comparanda. Caules 10—20cm. longi, folia caulina minuta, corolla 15—20 mm. longa, capsula 5—7 mm. longa.

In rupium fissuris et lapidosis regionis subalpinae, substratu siliceo-serpentino. Epirus: mt. Smolika (Bald.); Thessalia: mt. Said Pascha (Form.), pr. Malakasi, mt. Zygos, pr. Chaliki, Krania, mt. Baba (Haussk.), mt. Oxya (Hal.); occurrit quoque in mt. Tomor Maja Albaniae (Bald.). — Jun. Aug. ♃.

;; Calyx laevis, glaber vel hirsutus.

, Folia cordato-orbiculata vel cordato-ovata.

27. **C. cephallenica** Feer in journ. bot. XXVIII. p. 273 (1890). — *C. garganica* Bois. fl. or. III. p. 918; Heldr. fl. cephal. p. 50; non Ten — Exsicc.: Heldr. pl. cephal. a. 1867.

Pubescens; caulibus e rhizomate carnoso procumbentibus, foliatis, simplicibus vel ramulosis, racemosis; foliis cordato-ovatis, grosse crenato-dentatis, radicalibus petiolatis, caulinis diminutis, summis breviter petiolatis; floribus parvis, paucis, pedunculis arcuato-patulis subaequilongis, in racemum terminalem laxum dispositis; calycis laciniis lineari-lanceolatis, acutissimis, patentibus; corolla plus minus pubescente, scutellato-patula, coerulea, quinque-partita, duplo brevioribus; stylo exserto. — Affinis *C. garganica* Ten. differt floribus numerosis, majoribus, pedunculis 3—5plo longioribus patulis suffultis; calycis laciniis acutis; corolla 10 mm. alta, 15 mm. lata (nec 6—8 mm. alta et 10—12 mm. lata); staminum squamis cordato acuminatis, filamentis aequilongis (nec ovatis filamentis duplo fere longioribus); antheris longioribus; capsula 3—4 mm. (nec 2—2 1/2 mm.) longa et seminibus planiusculis, opacis, rugulosis (nec subovalibus, crassiusculis, nitidis).

Cephalonia: in rupibus regionis litoralis pr. Hagios Soter, pr. Athera et pr. coenobium Taphius (Heldr.), nec non in regione superiori mt. Aenos (Letourn.). — Jul. Aug. ♃.

28. **C. Sartorii** Bois. et Heldr. fl. or. III. p. 919. — Exsicc.: Heldr. herb. norm. n. 954.

Velutino-pubescens; caulibus e rhizomate carnosulo procumbentibus, flexuosis, foliatis, laxe racemosis; foliis caulinis minutis, orbiculari-

subcordatis, obtuse 5—7 crenatis, inferioribus petiolo aequilongo, superioribus breviore suffultis, foliis rosularum sterilium multo majoribus, reniformi-cordatis, multicrenatis, longe petiolatis; floribus parvis, paucis, pedunculis arcuato-patulis, calyci subaequilongis suffultis, in racemum laxum dispositis; calycis laciniis lanceolatis, acutis, patentibus, corolla velutina, albida vel coerulescente, infundibuliformi-campanulata, quinquelobata subduplo brevioribus; stylo exserto. — Caules 3—15 cm. longi, folia caulina 5—10 mm., rosularum sterilium 25 mm. lata, flores circa 10 mm. longa.

In stillicidiis et rupinis umbrosis ad coenobium Panagia Panachrantos, alt. 500 m. circiter, in insula Andros Cycladum (Sart.). — Jul. Aug. ♃.

„ Folia oblonga, basi attenuata.

29. **C. radicosa** Ch. et B. fl. pelop. p. 14 t. 7; Bois. fl. or. III. p. 915; Hal. in z. b. G. 1888 p. 760, in ö. b. Z. 1896 p. 286; Heldr. chlor. Parn. p. 23. — Exsicc.: Heldr. herb. norm. n. 223 et 1456; Orph. fl. gr. n. 99; Dörfl. fl. gr. n. 230 et 315.

Hirsuta, subcanescens; radice verticali, crassa; caulibus extrarosularibus, numerosis, decumbentibus, solo adpressis, foliatis, in racemum spiciformem, unilateralem abeuntibus, interdum uni vel paucifloris; foliis rosularum oblongo-spathulatis, obtusis, in petiolum attenuatis, denticulatis, caulinis diminutis, oblongis; floribus parvis, ad axillas solitariis, pedicello cernuo, calyce subbreviore suffultis; calycis laciniis lanceolatis, acutiusculis, tubo aequilongis; corolla violacea, glabriuscula, breviter obconica, in lobos oblongos patentes partita, calycis laciniis duplo longiore; capsula ignota. — Stirps singularis, caulibus 3—8 cm. longis, saepissime solo adpressis, foliis rosularum 25—40 mm. longis, floribus minutis, intense violaceis.

In pascuis alpinis. Aetolia: mt. Tymphrestus (Sprun.), Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Peloponnesus: mt. Kyllene pr. Malakadia (Orph.), Taygetos (Chaub.). — Jun. Aug. ♃.

β. Annuae; capsula nutans.

× Corolla majuscula, calyce duplo longior.

30. **C. scutellata** Griseb. spic. II. p. 282; Form. in D. bot. Mon. 1890 p. 14, in Ver. Brünn 1896 p. 38, 1897 p. 29; Haussk. symb. p. 145. — Exsicc.: Orph. fl. gr. n. 1080 (Macedonia); Sint. it. thessal. n. 1480.

Patule hispidula; caule erecto, a basi vel superne dichotome ramoso; foliis oblongo-lanceolatis, acutiusculis, integris vel repandis, sessilibus, ima sub anthesi emarcidis; floribus in dichotomiis et terminalibus, solitariis, breviter pedicellatis; calycis laciniis ovato-lanceolatis, acuminatis, corolla subduplo brevioribus, post anthesin auctis, stellatim patentibus; corolla subrotata, coerulea, glabra vel ad nervum parce setulosa, lobis ovatis. — Caulis 10—30 cm. altus, flores 15—25 mm. longi. — Habitu *C. dichotomae* et *C. ramosissimae* similis, a priore calyce exappendi-

culato, ab altera calyce poris basilaribus, nec apicularibus dehiscente diversa.

In rupestribus umbrosis, fruticetis regionis inferioris et montanae. Thessalia: pr. Malakasi, Uranaeos, Tsungeri, Meteora (Haussk.), Kalabaka Kastri, Vlachava (Form.). — Jun. Jul. ⊙.

31. C. **drabifolia** S. et. S. pr. I. p. 142, Fl. gr. III. p. 11 p. 215; Ch. et B. exp. p. 71, Fl. pelop. p. 15; Ung. Reise p. 125; Hal. Beitr. fl. Aetol. p. 8; Haussk. symb. p. 145; Form. in D. bot. Monat. 1898 p. 77. — *C. attica* Bois. et Heldr. diagn. XI. p. 67; Haussk. symb. p. 145. — *C. drabifolia β. major* Bois. fl. or. III. p. 933; Heldr. fl. cephal. p. 50. — Exsicc.: Heldr. herb. norm. n. 420 et 857; Orph. fl. gr. n. 249.

Patule hispidula; caulibus e collo pluribus vel solitariis, erectis vel diffusis, dichotome ramosis; foliis oblongis, obtusis vel acutiusculis, dentatis, radicalibus in petiolum brevem attenuatis, sub anthesi saepe jam emarcidis, ceteris sessilibus, saepe profundius dentatis vel trifidis; floribus in dichotomiis et terminalibus, solitariis, breviter pedicellatis; calycis laciniis lanceolatis, acutis, corolla duplo brevioribus, post anthesin auctis, stellatim-patentibus; corolla campanulata, coerulescente, glabra, lobis ovatis, violaceis. — Caules 5—25 cm. longi, flores 10—18 mm. longi. — A praecedente foliis dentatis et corolla campanulata diversissima. — *C. attica* ab auctoribus caulibus longioribus magis diffusis, foliis latioribus, floralibus profunde et acute trifidis, corolla 8 nec 4 lineas longa a *C. drabifolia* differre dicitur, sed omnes hae notae valde variabiles et inconstantes sunt, quamobrem *C. attica* meo sensu vix varietas speciei Sibthorpiani dicenda.

In arvis, rupestribus regionis inferioris. Thessalia: ad monasterium Korona, pr. Sophates, Karditza (Haussk.); Aetolia: pr. Mesolonghion (Nieder), ad radices mt. Chalkis (Hal.); Attica: frequens pr. Athenas (Sibth.), in colle Philopappo, Lycabetto, ad Phalerum, mt. Corydalus, pr. Eleusis, Buliasmene, mt. Hymettus (Heldr.); Acrocorinthus (Haussk.): Argolis: pr. Nauplia (Sprun.), Vromolimni (Haussk.); Laconia: pr. Androvista, Scutari, in Maina, mt. Ithome Messeniae (Chaub.); Cephalonia: pr. Argostoli (Heldr.). — Apr. Jun. ⊙.

×× Corolla minuta, calyce aequilonga.

32. C. **erinus** L. sp. p. 169; S. et S. pr. I. p. 142, Fl. gr. III. p. 10 t. 214; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 71, Fl. pelop. p. 15; Mazz. in ant. ion. V. p. 202; Friedr. Reise p. 283 et 286; Weiss in z. b. G. 1869 p. 49; Raul. cret. p. 804; Bois. fl. or. III. p. 932; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 145; Hal. in ö. b. Z. 1897 p. 96; Held. fl. Aegina p. 308. — Exsicc.: Sint. it. thessal. n. 504; Dörfl. fl. aeg. n. 1, Fl. cret. n. 38.

Hirta; caule erecto, dichotome ramoso; foliis dentatis, inferioribus obovatis, breviter petiolatis, superioribus oblongis, sessilibus, floralibus 3—5 lobis; floribus axillaribus et terminalibus, solitariis, subsessilibus;

calycis laciniis triangulari-lanceolatis, acutis, corolla aequilongis, post anthesin auctis, stellatim patentibus; corolla tubulosa, albida vel roseo-coerulescente, glabra, lobis brevibus, patentibus. — Corolla minuta insignis.

In saxosis, graminosis regionis inferioris et submontanae. Thessalia: pr. Guvelzi (Sint.); Attica: pr. Athenas (Sart.), Laurion, Acrocorinthus (Haussk.); insula Aegina: pr. Perivolia (Heldr.); Argolis (Sibth.); Laconia pr. Misitra (Friedr.); Sporadum insula Peristeri (Leon.); Cyclades (Chaub.): insula Amorgos (Leon.); Creta: pr. Canea, Retymo (Raul.), Males distr. Hierapetra (Leon.); Corcyra (Mazz.). — Apr. Jun. ⊙

2. Sectio. *Rapunculus* Bois. fl. or. III. p. 895. — Capsula 3 locularis, poris vel valvulis prope apicem, rarius versus medium sitis dehiscens.

a. Perennes.

α. Radix cylindrica.

33. **C. persicifolia** L. sp. p. 154; Bois. fl. or. III. p. 935; Form. in Ver. Brünn 1896 p. 39, 1897 p. 29. — Icon: Fl. gr. t. 205. — Exsicc.: Heldr. herb. n. 2367.

Radice crassiuscula, subrepente; caule erecto, glabro vel inferne hirtulo, foliato, paucifloro, rarius unifloro; foliis firmis, lucidis, glabris, integris vel minute denticulatis, radicalibus anguste spathulato-oblongis, in petiolum attenuatis, caulinis sessilibus, lineari-lanceolatis linearibusve, superioribus valde diminutis; pedunculis calyce brevioribus; calycis laciniis tubo glabro vel hispidulo triplo longioribus; corolla late campanulata, glabra, coeruleo-violacea, ad quartam partem lobata, calyce duplo longiore; capsula recta.

In umbrosis regionis montanae, rarissime. Thessalia: mt. Mitrica et pr. Phlambures in mt. Chassia (Form.), mt. Olympus (Heldr.). — Jun. Jul. ♃.

34. **C. abietina** Griseb. et Schenk in Wiegm. arch. XV. I. p. 333; Bald. riv. coll. bot. Alb. 1895 p. 57. — *C. patula v. pauciflora* Roch. pl. Banat. rar. p. 42 t. 6, non *C. pauciflora* Lam. nec Desf. — *C. Rochelii* Schur sert. fl. Trans p. 47, in ö. b. Z. 1861 p. 45. —

Radice tenui, repente, caudiculos numerosos, filiformes, subterraneos, steriles, apice foliosos, caulesque adscendentes, tenues, remote foliatos, pauci vel multifloros edente; foliis membranaceis, crenatis, glabris vel margine setulosis, radicalibus ovatis vel ellipticis, basi rotundatis, longe petiolatis, caulinis inferioribus oblongo-spathulatis, breviter petiolatis, ad superiora lanceolata, sessilia sensim decrescentibus; floribus longiuscule pedunculatis; calycis glabri laciniis lineari-lanceolatis, erecto-patulis vel reflexis, tubo 3—4 plo longioribus; corolla patule infundibuliformi, coerulea, glabra, ad medium quinquefida, calyce duplo longiore.

In saxosis et pratis in summo jugo mt. Peristeri et in regione silvatica mt. Olycika distr. Janina in Epiro (Bald.). — Jun. Aug. ♃. N. v.

β. Radix napiformis.

35. **C. Spruneriana** Hampe in Flora 1842 p. 76; Spreitz. in z. b. G. 1886 p. 664; Haussk. symb. p. 145. — *C. patula* S. et S. pr. I. p. 137; Ch. et B. exp. p. 70, Fl. pelop. p. 14; M. et R. fl. Zante p. 64; non L. — *C. spathulata* Friedr. Reise p. 272; Ung. Reise p. 125; Hal. in ö. b. Z. 1897 p. 96 et 286 pro p.; non W. et K., nec S. et S. — *C. Spruneri* Raul. cret. p. 804; Bois. fl. or. III. p. 937; Spreitz. in z. b. G. 1877 p. 714; Heldr. fl. cephal. p. 50. — Exsicc.: Heldr. herb. norm. n. 23 et 1058a; Sint. it. thessal. n. 207; Dörfl. fl. gr. n. 333.

Radice napiformi, globosa, ovata vel oblonga, caudiculos numerosos, nudos, procumbentes vel subterraneos, in caules erectos abeuntes, edente; caulibus saepissime elatis, glabris vel inferne hirtis, patule ramosis, rarius simplicibus, ramis unifloris; foliis glabris vel breviter hirtis, remote et obsolete crenatis, inferioribus ovatis vel oblongis, in petiolum attenuatis, ceteris lanceolatis, elongatis, acuminatis, sessilibus; calycis glabri laciniis lanceolato-subulatis, basin versus interdum remote et obsolete denticulatis, alabastro dimidio usque duplo longioribus, corollam subaequantibus; corolla coerulea, infundibuliformi, ad medium in lobos oblongos, acutos partita; capsula obconica, laciniis calycis rectis, ea longioribus coronata. — Caules 15—50 cm. alti, corolla spectabilis, 25—30 mm. longa.

β. **hirsuta** Heldr. in Pichl. pl. gr. exs. a. 1876. — Caules simplices, inferne foliisque dense hirti.

In dumosis regionis sempervirentis et montanae. **Thessalia**: pr. Kalabaka (Sint.), mt. Pelion pr. Kissos et Milies (Heldr.); Sporadum insula Skopelos (Leon.); Euboea: mt. Telethrion (Heldr.); Aetolia: mt. Korax pr. Musinitza (Tunt.); mt. Parnassus (Sprun.); Attica: pr. Marathon, mt. Pentelicon (Heldr.); Acrocorinthus (Orph.); Argolis: pr. Vromolimni (Friedr.), insula Poros (Wied.), in olivetis insulae Poros oppositis (Heldr.), mt. Artemision supra Argos (Kalondzis), Port Tolon, mt. Palamidi pr. Nauplia (Haussk.); Achaia: mt. Kyllene (Heldr.); Messenia: mt. Diaforti, pr. Methone, Corone (Chaub.); Laconia: pr. Mistra (Bois.), mt. Taygetos (Chaub.); Creta: pr. Kaenuriokorio, in eparchia Sitia (Raul.); Zante (Marg.); Cephalonia: in oropedio Omala, pr. Lixuri, Dallaportata, Vorikes (Heldr.); Ithaca (Ung.): pr. Vathy (Spreitz.); Corcyra: pr. San Deca (Spreitz.). — β. in castanetis pr. Steni Euboeae (Pichl.). — Apr. Jun. ♃.

36. **C. Sibthorpiana.** — *C. spathulata* S. et S. pr. I. p. 137, Fl. gr. III. p. 2 t. 203 (specimen gracile, caule solitario, sine tubere); Hal. in z. b. G. 1888 p. 760; Beitr. fl. Epir. p. 31, Beitr. fl. thessal. p. 15, Beitr. fl. Achaia p. 26, in ö. b. Z. 1897 p. 286 p. p.; Heldr. chlor. Parn. p. 23; Form. in Ver. Brünn 1896 p. 39; non W. et K.

— *C. Spruneri β. alpina* Bois. fl. or. III. p. 937; Haussk. symb. p. 145; Form. in Ver. Brünn p. 30, non *C. alpina* Jacq. — Exsicc.: Heldr. herb. norm. n. 1058 b p. p.; Sint. it. thessal. 668 et 900.

Radice napiformi, globosa, ovata vel oblonga, (rarissime vix incrassata, cylindrica), caudiculos numerosos, nudos procumbentes vel subterraneos, in caules erectos abeuntes, edente; caulibus glabris vel inferne hirtis, simplicibus, unifloris, rarius ramulo auctis; foliis glabris vel breviter hirtis, remote et obsolete crenatis, inferioribus spathulatis, obovatis vel rotundatis, in petiolum attenuatis, ceteris oblongis vel breviter oblongo-lanceolatis, acutis, sessilibus, summis lanceolatis; calycis glabri laciniis lanceolatis, basin versus interdum remote et obsolete denticulatis, alabastro brevioribus vel sublongioribus, corolla duplo vel triente brevioribus; corolla coerulea, infundibuliformi, ad medium in lobos oblongos, acutos partita; capsula obconica, laciniis calycis rectis, ea longioribus coronata. — Caulis 6—25 cm. alti, corolla 12—25 mm. longa. — Differt a praecedente caulibus humilioribus, foliis caulinis abbreviatis, latioribus brevioribusque, calycis laciniis brevioribus et corolla minore.

β. **filicaulis.** — Caules filiformes, flaccidi, flexuosi, paucifoliati, simplicissimi, uniflori; folia membranacea, inferiora et media petiolata, ovata vel elliptica, capsula breviter obconica. — Ex paucis speciminibus incompletis, radice destitutis mihi tantum nota et ulterius observanda, anne speciem propriam sistet. — Exsicc.: Bald. it. cret. n. 336.

In pascuis et rupestribus regionis subalpinae et alpinae. Epirus: mt. Tsumerka, Peristeri (Hal.); Thessalia: pr. Chaliki, Malakasi (Sint.), mt. Zygos, Karava, Ghavellu (Haussk.), Dokimi, Mandra Hodza (Form.), mt. Oxya (Hal.), Tringia (Hartl); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Sibth.), Cithaeron (Heldr.); Achaia: mt. Olenos, Panachaicon, Chelmos (Hal.), Kyllene (Heldr.); Laconia: mt. Malevo (Leon.); — *β.* in lapidosis mobilibus mt. Lazaro in mt. Lassiti Cretae (Bald.). — Jun. Aug. ♃

b. Biennes vel annuae.

α. Folia radicalia numerosissima, in rosulam densam congesta; calycis laciniae tubo aequilongae vel parum longiores.

37. **C. aizoon** Bois. et Sprun. diagn. IV. p. 34; Hal. in z. b. G. 1888 p. 760; Heldr. chlor. Parn. p. 23. — Exsicc.: Heldr. herb. norm. n. 22; Orph. fl. gr. n. 417.

Glabra; radice crassa, cylindrica; caule crasso, sulcato, parce foliato, fere a basi thyrsoideo-paniculato, ramis numerosis, rigidis, erecto-patulis; foliis rosularum lingulatis, ex apice spathulato, obtuso, saepe mucronulato, sensim attenuatis, margine cartilagineis, integris vel crenatis, laevibus vel sub lente spinuloso-ciliatis, caulinis parvis, triangularilanceolatis, acutis, sessilibus; floribus secus ramos strictis, solitariis, pedicello calyce breviori suffultis, terminalibus saepius longius pedicellatis; calycis glabri laciniis triangularibus; corolla coerulea, campanulata, breviter lobata, calyce 4 plo longiore; capsula erecta, globosa,

valvis supra medium sitis dehiscente. — Species egregia, nulla cum alia comparanda, caule crasso 15—50 cm. alto, floribus circa 15 mm. longis; in mt. Kiona observavi specimina robusta, pulcherrima, floribus numerosissimis.

In rupibus regionis abietinae et superioris mt. Kiona (Hal.) et Parnassus (Sprun.), rarissime. — Jun. Aug. ⊙

β. Folia radicalia pauca; calycis laciniae tubo saepius multo longiores.

× Corolla infundibuliformis vel campanulata.

○ Radix fusiformis, carnosa.

38. **C. rapunculus** L. sp. p. 164; Pieri corc. fl. p. 30; Mazz. in ant. ion. V. p. 200; Form. in D. bot. Monat. 1890 p. 14. — *C. rapunculus v. Lambertiana* Haussk. symb. p. 145, f. calycis laciniis denticulatis; an *C. Lambertiana* DC. mon. camp. p. 327. — Icon: Fl. dan. t. 1326.

Glabra vel pubescens; caule erecto, inferne foliato, in racemum angustum, elongatum simplicem vel compositum, basi ramulis erectis auctum abeunti; foliis obtuse crenatis, radicalibus sub anthesi jam emarcidis, in petiolum decurrentibus, superioribus lanceolatis vel lineari-lanceolatis, sessilibus; floribus solitariis, rarius subternis, flore intermedio longius pedunculato; calycis laciniis lanceolato-subulatis, integris vel inferne parce et minute denticulatis, tubo obconico 2—3 plo longioribus; corolla infundibuliformi, coeruleo-violacea, ad tertiam partem in lobos oblongo-lanceolatos fissa; capsula obconica, erecta. — Caulis strictus, 45—80 cm. altus, corolla 20—25 mm. longa.

In nemorosis regionis inferioris et montanae. Thessalia: pr. Volo, Plessidi (Form.), Karditza, Palaeokastron, ad monasterium Korona, in oropedio Neuropolis (Haussk.); Corcyra (Pieri): pr. Calefactiones (Mazz.). — Maio, Jun. ⊙ N. v.

○○ Radix cylindrica, ramosa.

. Biennis; folia radicalia sub anthesi saepius adhuc vigentia: panicula subcorymbosa; calycis laciniae tubo 2—3 plo longiores.

39. **C. patula** L. sp. p. 163; Haussk. symb. p. 145; ? Form. in Ver. Brünn 1897 p. 29 subsp. *chassia*, quae a typo pedunculis multo longioribus, arcuato-adscendentibus, calycis laciniis tubo plus 3 plo longioribus, corollae lobis triangulari-ovatis, subobtusis, differre dicitur. — Icon: Fl. dan. t. 373. — Exsicc.: Sint. it. thessal. n. 902.

Glabra vel pubescens, caulibus solitariis vel pluribus, erectis, foliatis, patule ramosis, ramis paniculatis; foliis crenatis vel subintegris, radicalibus rosulatis obovato vel oblongo-lanceolatis, in petiolum decurrentibus, caulinis oblongo-lanceolatis lanceolatisve, sessilibus; floribus solitariis, longiuscule pedunculatis; calycis laciniis lineari-lanceolatis, integris vel inferne parce et minute denticulatis, tubo obconico 2—4 plo longioribus; corolla campanulata, lilacina, ad medium in lobos oblongo-

lanceolatos fissa; capsula ovoideo-cylindrica. — Caulis 30—50 cm altus, corolla 20—25 mm. longa.

In nemorosis Thessaliae: in jugo Zygos, supra Malakasi (Haussk.), pr. Hapdominta Aderfia et Phlamburo in mt. Oxya, pr. Phlambures et mt. Mitrica in mt. Chassia (Form.). — Maio, Jun. ⊙

.. Annuae, folia radicalia sub anthesi jam emarcida; panicula racemosa vel thyrsoidea; calycis laciniae tubo multoties longiores.

40. **C. sparsa** Friv. in magy. tud. társ. évk. 1840 p. 291. — *C. expansa* Friv. in Flora 1836 II. p. 434; Bois. fl. or. III. p. 941; non Rud., in mem. acad. St. Petersb. IV. p. 340. — *C. Frivaldszkyi* Steud. nom. bot. ed. 2 I. p. 267. — Exsicc.: Orph. fl. gr. n. 657.

Caulibus elatis, solitariis vel pluribus, foliatis, angulato-subalatis, praesertim ad angulos patule pilosis, saepius jam infra medium in paniculam diffusam, multifloram, rarius in racemum compositum abeuntibus, ramis inferioribus elongatis, superioribus sursum brevioribus; foliis crenatis, pilosulis, radicalibus oblongo-spathulatis, in petiolum decurrentibus, ceteris oblongo-lanceolatis, acutis, basi lata semiamplexicauli sessilibus, floralibus lineari-lanceolatis; floribus magnis, solitariis vel subternis, flore intermedio longius pedunculato; calycis laevis laciniis lineari-lanceolatis, erecto-patulis, inferne saepius parce et minute denticulatis, corollam subaequantibus; corolla infundibuliformi-campanulata, lilacina, ad tertiam partem in lobos oblongo-lanceolatos fissa; capsula obconica. — Caulis 50—100 cm. altus, folia media 5—6 cm. longa, basi 2 cm. lata, corolla 3 cm. longa. — Specimina originalia non vidi, sed si planta Frivaldszkyana cum planta ex Olympo revera identica sit, speciem certe propriam, a sequente distinctissimam sistit.

In regione silvatica mt. Olympus supra Hagios Dionysios, rare (Orph.). — Jul. Aug. ⊙

41. **C. sphaerothrix** Grieseb. Reise Rumal. II. p. 263, sol. nom., Spicil. fl. rum. II. p. 280; Haussk. symb. p. 145. — *C. Welandii* Heuff. in ö. b. Z. 1857, p. 118, in z. b. G. 1858 p. 154; Celak. in ö. b. Z. 1871 p. 6. — *C. expansa v. sphaerothrix* Bois. fl. or. III. p. 941. — Exsicc.: Sint. it. thessal. n. 1221.

Caulibus erectis, solitariis vel pluribus, striato-sulcatis, breviter pilosis, a medio vel superne in paniculam diffusam, multifloram abeuntibus, ramis tenuibus, sursum decrescentibus; foliis crenatis, pilosulis, radicalibus oblongo-spathulatis, in petiolum decurrentibus, ceteris ovato- vel oblongo-lanceolatis, acutis, basi lata, semiamplexicauli sessilibus, floralibus lineari-setaceis; floribus parvis, solitariis, pedunculis filiformibus elongatis suffultis; calycis tubo glandulis sphaericis plus minus exasperato, praeterea laevi, laciniis lineari-setaceis, patentibus, inferne saepius parce et minute denticulatis, corollam subaequantibus, cum fructu etiam excrescentibus; corolla infundibuliformi-campanulata, lilacina, ad tertiam partem in lobos oblongo-lanceolatos fissa; capsula obconica. — Differt a praecedente statura graciliori, ramis tenuioribus, pedunculis filiformibus,

calycis laciniis setaceis, patentibus et praesertim floribus duplo minoribus, 12—16 mm. longis. Affinis quoque *C. patulae*, a qua radice annua, caulibus acutius angulatis, magis asperis, foliis radicalibus sub anthesi emarcidis, caulinis magis approximatis, inflorescentia thyrsoidea, calycis laciniis setaceis, longioribus et corollae minoris tubo angustiore, discedit.

β. **oxya.** — *C. expansa* Hal. Beitr. fl. Thessal. p. 16, non Friv — Folia caulina majora, obtusa, inflorescentiae rami firmiores, calycis laciniae latiores, erecti, capsula ovoidea. — An species propria. — Exsicc.: Hal. it. gr. II. a. 1893.

In collibus dumosis et silvaticis regionis montanae et subalpinae Thessaliae: pr. Aivali (Haussk.), pr. Pezula in oropedio Neuropolis (Heldr.); — *β.* in silvis caeduis mt. Oxya supra Chaliki (Hal.). — Jun. Jul. ⊙.

×× Corolla breviter obconica, subrotata.

42. **C. phrygia** Jaub. et Sp. ill. or. III. t. 233; Haussk. symb. p. 146. — Heldr. it. thessal. a. 1885.

Puberula; caule tenui, a basi ramuloso, flexuoso; foliis parvis, inferioribus subsessilibus, obovatis, obtusis, crenulatis, superioribus breviter lanceolatis linearibusve; floribus parvis, solitariis, longe pedunculatis; calycis glabri vel puberuli laciniis lanceolato-subulatis, uninerviis, inferne interdum 1—2 denticulatis, tubo obconico duplo longioribus; corolla lilacina, ad tertiam partem lobata; capsula erecta, turbinato-obconica, profunde sulcata, laevi, calycis laciniis ea vix longioribus superata. — Planta tenera, 5—15 cm. alta, floribus minutis, 6—7 mm. longis, hac nota et calycis laciniis abbreviatis, angustissimis, a sequentis formis parvis, capsula laevi interdum occurrentibus, discrepat; specimina tamen quaedam in herbario dubia.

In pascuis siccis Thessaliae rarissime: pr. Karditza (Heldr.), Sophates (Haussk.). — Maio. Jun. ⊙.

43. **C. ramosissima** S. et S. pr. I. p. 137, Fl. gr. III. p. 3 t. 204; Ch. et B. exp. p. 70, Fl. pelop. p. 14; Mazz. in ant. ion. V. p. 204; Marg. et R. fl. Zante p. 63; Bois. fl. or. III. p. 942; Spreitz. in z. b. G. 1877 p. 714 f. calyce glabro; Heldr. fl. cephal. p. 50; Hal. Beitr. fl. Epir. p. 31, Beitr. fl. Achaia p. 26, in ö. b. Z. 1896 p. 15, 1897 p. 286; Haussk. symb. p. 146, cum f. *glabrescens* (caulibus foliisque glabrescentibus, superne glaberrimis). — *C. Loreyi* Poll. elem. bot. II. p. 148, Fl. veron. I. p. 271 t. 2; Friedr. Reise p. 292; Gelmi in bull. soc. bot. it. 1889 p. 450. — Exsicc.: Orph. fl. gr. n. 654; Sint. it. thessal. n. 338, Baen. herb. europ. n. 9225. —

Pilosa vel glabrescens, a collo simplex vel multicaulis; caulibus erectis vel adscendentibus, ramosis vel simplicibus et unifloris; foliis crenatis, inferioribus obovato-spathulatis, in petiolum attenuatis, caulinis ovali-lanceolatis lanceolatisque, acutis vel obtusis, sessilibus; floribus saepius majusculis, solitariis, longe pedunculatis; calycis laciniis lanceo-

latis, plus minus manifeste trinerviis, inferne saepius denticulatis, tubo hemisphaerico, albo-strigoso, rarius glabro, multo longioribus, patentibus; corolla lilacina, ad $^2/_3$ lobata; capsula erecta, immatura sphaeroida, demum turbinato-obconica, profunde sulcata, papilloso-strigosa, rarius glabra, calycis laciniis accretis ea multo longioribus superata. — Species polymorpha, variat nempe indumento, caule solitario et simplici vel e collo pluribus, ramosis, foliorum forma, calyce nunc dense albo-strigoso, nunc glabro, laciniis denticulatis vel integris, floribus magnis, 3 cm. et parvis 1 cm. tantum longis.

In herbidis, saxosis regionis inferioris et montanae. Epirus: pr. Kalentini, Mazuki, Vulgarelion (Hal.); Thessalia: pr. Klinovo, Malakasi, ad monasterium Korona et Meteora (Haussk.), pr. Kalabaka (Sint.), in oropedio Neuropolis (Heldr.); Acarnania: pr. Agrinion (Nied.); Aetolia: pr. Aetolikon (Reis.), mt. Taphiassos (Hal.); mt. Parnassus (Sibth.); Achaia: pr. Patras (Hal.), mt. Olenos, Kyllene (Heldr.); Argolis: pr. Nauplia (Orph.), Port Tolon, peninsula Methana (Haussk.); Laconia: mt. Malevo (Leon.), Taygetos (Chaub.); Messenia: pr. Methone, Pylos, Phigalea (Chaub.), Diabolitzi (Heldr.), Pyrgos, Arcadia (Friedr.); Zante (Sibth.); Cephalonia: pr. Argostoli, in oropedio Omala (Heldr.); Ithaca (Spreitz.); Corcyra: pr. Kastrades (Maen.), Pelleca (Gelmi), Garizza, Anemomylo (Mazz.). — Apr. Jul. ⊙

Obs. Species dubiae: *C. pauciflora* Desf. choix pl. p. 36 t. 26 a Tournefort in Creta lecta, autoribus recentioribus incognita, sec. Bois. fl. or. III. p. 937 ad *C. Sprunerianum* referenda videtur. — *C. lilifolia* L. sp. p. 165, quae a S. et S. pr. I. p. 138 in Cretae montibus elatioribus nive obductis, sed probabiliter ex confusione quadam, indicatur. — *C. mollis* L. sp. ed. 2 p. 237, in DC. mon. camp. p. 238 e Creta (Tourn.), probabiliter etiam ex confusione. — *C. uniflora* L. sp. p. 163, species arctica, indicatur certissime erronee a Pieri corc. fl. p. 30 et Mazz. in ant. ion. V. p. 202 in Corcyra. — *C. bicaulis* Lap. fl. pyr. p. 13 = *C. speciosa* Pourr. act. tol. III. p. 309, a Mazz. in ant. ion. V. p. 202, ex confusione in Corcyra. — Dein species novae: *C. muscosa et pruinosa* Mazz. l. c. p. 204 e Corcyra, e descriptione incompleta non recognoscendae.

3. Specularia Heist. syst. gen. p. 8.

a. Corolla majuscula, calycis laciniis longior vel aequilonga.

1. **S. pentagonia** L. sp. p. 169; Mazz. in ant. ion. V. p. 206; (*Campanula*); DC. mon. camp. p. 44; Raul. cret. p. 804; Bois. fl. or. III. p. 959. — *Prismatocarpus pentagonius* L'Her. sert. angl. p. 2. — Icon: Desf. choix pl. t. 33. — Exsicc.: Reverch. pl. cret. a. 1884 n. 103, in Baen. herb. europ. n. 5691; Dörfl. fl. aeg. n. 2, pl. cret. n. 89.

Pubescenti-hirta vel glabriuscula, simplex vel a collo ramosa; caulibus foliatis, centrali erecto, lateralibus adscendentibus, simplicibus

vel apicem versus ramosis; foliis obsolete undulato-crenatis, inferioribus orbiculatis, obovatis vel oblongo-spathulatis, saepe petiolatis, ceteris sessilibus, oblongo-lanceolatis lanceolatisve; floribus sessilibus, solitariis raro binis, terminalibus et axillaribus; calycis laciniis lanceolatis, sub anthesi plerumque tubo 2—3plo brevioribus, demum patulis; corolla majuscula violacea, calycis lacinias aequante vel superante; capsula apice non constricta. — Corolla ampla, 15—18 mm. longa.

In cultis, inter segetes. Creta: pr. Kissamos (Rev.), Malaxa, Canea, Askyphos, Anopolis, Komitades (Raul.), Males distr. Hierapetra (Leon.); Cycladum insula Amorgos (Leon); indicatur quoque pr. Messongi et Lefchimo Corcyrae (Mazz.). — Mart. Maio. ⊙

2. **S. speculum** L. sp. p. 168; S. et S. pr. I. p. 142, Fl. gr. III. p. 12 t. 216; Ch. et B. exp. p. 71, Fl. pelop. p. 15; (*Campanula*); DC. mon. camp. p. 346; Marg. et R. fl. Zante p. 64; Bois. fl. or. III. p. 959; Spreitz. in z. b. G. 1877 p. 715; Heldr. fl. cephal. p. 50, Fl. Aegina p. 308; Haussk. symb. p. 146; Hal. in ö. b. Z. 1897 p. 286, in z. b. G. 1899 p. 190. — *Prismatocarpus speculum* L'Her. sert. angl. p. 2; Mazz. in ant. ion. V. p. 206; Friedr. Reise p. 274; Ung. Reise p. 125. — Exsicc.: Rev. pl. cret. a. 1883 n. 103; Sint. it. thessal. n. 337 et 906.

Scabrido-pubescens, simplex vel a collo ramosa; caulibus foliatis, centrali erecto, lateralibus adscendenti-diffusis, ramulosis; foliis undulato-crenatis, inferioribus obovatis, in petiolum attenuatis, ceteris oblongis, semiamplexicaulibus; floribus subsessilibus, ad apicem caulis fasciculato-cymosis, ramulorum saepius solitariis; calycis laciniis lineari-lanceolatis, sub anthesi tubo aequilongis vel parum brevioribus, demum patenti-reflexis; corolla mediocri, violacea, calycis lacinias aequante; capsula apice constricta. — Habitu magis diffuso, floribus fere duplo minoribus, calycis laciniis minoribus, demum patenti-reflexis et capsula apice constricta a praecedente distincta.

β. **calycina** DC. mon. camp. p. 347; Raul. cret. p. 804. — Calycis laciniae latiores, oblongae. — Exsicc.: N. v.

γ. **racemosa** Bois. fl. or. III. p. 959. — Inflorescentia ob ramos laterales brevissimos fasciculiformes, ut in *S. falcata*, racemosa. — Exsicc.: N. v.

In cultis, inter segetes regionis inferioris et montanae. Thessalia: pr. Malakasi, Meteora (Sint.); Attica: mt. Hymettus, pr. Stephani, insula Aegina (Heldr.); Achaia: pr. Patras (Chaub.); Argolis (Chaub.): insula Poros (Friedr.); Laconia: mt. Malevo (Leon.), in Maina pr. Androvista, usque Methone Messeniae (Chaub.); Elis: in silva Kapellis pr. Lala (Reiser); Cyclades (Sibt.); Creta: pr. Malaxa (Rev.); Zante: pr. Litakia (Marg.); Cephalonia (Ung.): pr. Argostoli (Spreitz.), Pessada, Aphrata (Heldr.); Corcyra: pr. Ascensione, Anemomylo (Mazz.); — β. Creta: pr. Khalepa, inter Askyphos et Anopolis (Raul.); — γ. in Peloponneso (Gittard.). — Apr. Jul. ⊙

b. Corolla parva, calycis laciniis duplo brevior.

3. **S. falcata** Ten. pr. fl. nap. p. 16, fl. nap. I. p. 77 t. 20; Friedr. Reise p. 274; (*Prismatocarpus*); DC. mon. camp. p. 345; Bois. fl. or. III. p. 960; Hal. in ö. b. Z. 1897 p. 96. — *Campanula falcata* Roem. et Schult. syst. V. p. 154. — Exsicc.: Heldr. et Hal. fl. sporad. a. 1896.

Glabra vel parce pubescens, simplex vel a collo ramosa; caulibus foliatis, centrali erecto, lateralibus adscendentibus, inferne saepe ramulosis, in spicam interruptam abeuntibus; foliis obsolete crenatis, inferioribus oblongis, in petiolum attenuatis; ceteris lanceolatis vel ovato-lanceolatis, semiamplexicaulibus; floribus sessilibus, ad axillas solitariis vel geminatis, inferioribus remotis, superioribus approximatis; calycis laciniis a basi lanceolato-lineari longe attenuato-acuminatis, arcuato-reflexis, tubo subbrevioribus; corolla coerulea; capsula apice non constricta.

In arvis, saxosis regionis inferioris et montanae, raro. Sporadum insula Peristeri et Skopelos (Leon.); mt. Parnes Atticae, Acrocorinthus (Heldr.); insula Poros (Friedr.); mt. Taygetos in Alagonia (Zahn). — Apr. Jul. ⊙

4. **S. hybrida** L. sp. p. 168; S. et S. pr. I. p. 143; Ch. et B. exp. p. 72, Fl. pelop. p. 15; (*Campanula*); DC. mon. camp. p. 348; Raul. cret. p. 804; Bois. fl. or. III. p. 960; Hal. Beitr. fl. Achaia p. 26; Haussk. symb. p. 146. — *Prismatocarpus hybridus* L'Hér. sert. angl. p. 2; Mazz. in ant. ion. V. p. 206. — Icon: Rchb. t. 255. — Exsicc.: Heldr. herb. norm. n. 89; Dörfl. pl. cret. n. 55.

Differt a praecedente foliis magis undulatis et crenatis, laciniis calycis acutis, basi angustatis, erectis, tubo 3—4 plo brevioribus, capsula apice constricta.

In arvis, saxosis regionis inferioris et montanae. Thessalia: mt. Ghavellu in Pindo (Haussk.); Attica: mt. Parnes, Lycabettus, Pateras (Heldr.), Kerata; ad Acrocorinthum (Haussk.); Achaia: mt. Panachaicon (Hal.); Argolis: pr. Nauplia (Sprun.), insula Hydra (Heldr.); Cycladum insula: Kythnos (Tunt.), Tenos (Leon.); Creta: inter Askyphos et Anopolis, mt. Ida (Raul.), pr. Males distr. Hierapetra (Leon.); Corcyra: pr. Ascensione, Anemomyla (Mazz.). — Apr. Jul. ⊙

4. Petromarula DC. mon. camp. p. 209.

1. **P. pinnata** L. sp. p. 171; S. et S. pr. I. p. 144, Fl. gr. III. p. 15 t. 220; Sieb. avis p. 3, rem. p. 3, in Flora I. p. 271; (*Phyteuma*); DC. l. c.; Weiss in z. b. G. 1869 p. 49; Raul. cret. p. 803; Bois. fl. or. III. p. 957; Bald. viagg. Creta p. 75. — Exsicc.: Rev. pl. cret. n. 101; Dörfl. pl. cret. n. 27 a et 132.

Rhizomate crasso; caulibus elatis, simplicibus vel ramosis, glabris superne puberulis, in racemum longum vel paniculam multifloram abeuntibus; foliis glabris, radicalibus numerosis caulinisque inferioribus longe petiolatis, pinnatisectis, segmentis utrinque 3—4 ovatis, grosse lobato-dentatis, terminali multo majore cordato, foliis superioribus diminutis, breviter petiolatis vel sessilibus; calycis glabri vel puberuli

laciniis lanceolatis, tubo hemisphaerico subduplo longioribus; corollae azureae laciniis linearibus, demum patenti-reflexis; capsula sphaerica. — Species pulcherrima, caulibus 40—120 cm. altis. Radix et folia edulia.

Ad rupes regionis maritimae et collinae Cretae (Belli in Clus. hist. p. 299): pr. Rumata (Bald.), Malaxa (Rev.), Canea (Weiss), Kissamos, Theriso, Almyros, Spili, Voriza, insula Dia, Kritsa, Males, Strovodoxare (Raul.). — Apr. Jun. ♃.

5. **Podanthum** Don gen. syst. III. p. 748.

a. Capsula nutans, poris basilaribus dehiscens.

1. **P. trichocalycinum** Ten. fl. nap. I. p. 67 t. 16; Sieb. avis p. 3, rem. p. 3, in Flora I. p. 275; (*Campanula*); Bois. fl. or. III. p. 955, suppl. p. 335; Heldr. chlor. Parn. p. 23; Haussk. symb. p. 146, cum var. *densiflora*. — *Campanula Pichleri* Vis. fl. dalm. suppl. p. 74 t. 6. — Exsicc.: Rev. pl. cret. n. 102, in Baen. herb. europ. n. 4857; Sint. it. thessal. n. 904.

Rhizomate fusiformi, carnoso; caulibus elatis, glabris vel inferne hirtis, in racemum brevem vel elongatum, simplicem vel inferne ramulis auctum abeuntibus; foliis tenuibus, glabris, sessilibus subsessilibusve, ovatis vel oblongis, grosse inaequaliter et acute dentato-serratis, superioribus acuminatis; floribus solitariis vel 2—3 nis, pedicellis tenuissimis, bractea setacea longioribus, suffultis; calycis glabri laciniis setaceis, patentibus, tubo ovato-hemisphaerico triplo longioribus; corolla glabra, coerulea, in lacinias lineares basin usque partita, calycis laciniis duplo longiore; capsula ovata, angulato-sulcata. — Caulis 25—80 cm. altus, folia usque ad 8 cm. longa 35 mm. longa, corolla angusta, circa 15 mm. longa.

In umbrosis, silvaticis regionis montanae et subalpinae, rare. Thessalia: mt. Zygos, Oxya (Haussk.); Phocis: mt. Parnassus pr. Carcaria (Heldr.); Creta: mt. Drakona et Volokia (Rev.) in mt. Sphacioticis (Sieb.). — Jun. Jul. ♃.

b. Capsula erecta, poris sub apicem sitis dehiscens.

α. Caulis tota longitudine foliosus.

2. **P. canescens** W. et K. pl. rar. hung. I. p. 12 t. 14 (*Phyteuma*); Bois. fl. or. III. p. 950; Hal. in ö. b. Z. 1897 p. 286. — Exsicc.: Heldr. pl. fl. hellen. a. 1896.

Canescenti-scabridum vel glabrescens; caulibus elatis, virgatis, in racemum brevem vel elongatum, simplicem vel inferne ramulis auctum abeuntibus; foliis crenulatis, subtus pallidioribus, inferioribus ovatis oblongisve, breviter petiolatis, ceteris sensim decrescentibus, oblongo-lanceolatis lanceolatisve, sessilibus, in bracteas lineares transeuntibus; floribus axillaribus, subsessilibus, solitariis vel 2—3 nis; calycis laciniis lanceolatis, tubo obconico vix longioribus; corolla pallide violacea, calycis laciniis triplo longiore; capsula ovoidea.

In saxosis regionis superioris mt. Korax Aetoliae (Tunt.). — Jul. Aug. ♃.

β. Caulis ima basi foliosus, ceterum subnudus.

3. **P. limonifolium** L. sp. ed. 2 p. 239 *(Campanula)*; Bois. fl. or. III. p. 951; Heldr. chlor. Parn. p. 23; Hal. Beitr. fl. Epir. p. 31; Beitr. fl. Thessal. p. 16, Beitr. fl. Achaia p. 26; Haussk. symb. p. 146, cum var. *ramosum* (f. caule ramoso); Bald. riv. coll. bot. Alb. 1895 p. 57, 1896 p. 77, in bull. herb. Bois. IV. p. 205; Form. in Ver. Brünn 1896 p. 36, 1897 p. 29, in D. bot. Monat. 1896 p. 77. — *Phyteuma limonifolium* S. et S. pr. I. p. 144, Fl. gr. III. p. 14 t. 218; Mazz. in ant. ion. V. p. 206; Clem. sert. p. 64; Hal. in z. b. G. 1888 p. 760. — *Ph. stricta* Sims. bot. mag. t. 2145. — *Campanula virgata* Lodd. bot. cat. t. 667, non Labill. ic. pl. dec. II. p. 11. — *Ph. virgata* Ch. et B. exp. p. 72, Fl. pelop. p. 15, non Willd. sp. I. p. 924. — Exsicc.: Heldr. herb. norm. n. 316 et 1563; Sint. it. thessal. n. 905; Bald. it. alb. epir. IV. n. 132; Dörfl. fl. gr. n. 218.

Pubescenti-scabridum vel glabrum; caulibus rigidis, virgatis, simplicibus vel ramosis, in spicas longas, tenues, laxas abeuntibus; foliis radicalibus numerosis, caulinisque imis lanceolatis vel lanceolato-linearibus repando-dentatis vel integris, acutis, in petiolum decurrentibus, ceteris paucis, linearibus, valde diminutis, bracteiformibus; floribus sessilibus, 2—3 fasciculatis vel solitariis; calycis laciniis lanceolatis, tubo obconico-oblongo aequilongis vel brevioribus; corolla violacea, calycis laciniis triplo longiore; capsula ellipsoidea, asperula, costis valde prominulis.

β. **repandum** S. et S. pr. I. p. 143; Hal. in z. b. G. 1888 p. 760; *(Phyteum)* pro sp. — *P. limonifolium β. alpinum* Bois. fl. or. III. p. 951; Haussk. symb. p. 146. — Caules humiles, nani, simplices; spica brevis. — Forma regionis alpinae, quae sine limites typo confluit. — Exsicc.: Hal. it. gr. II. a. 1893.

In dumosis, silvaticis, pascuis regionis montanae et alpinae. Epirus, mt. Tsumerka, pr. Kalarrytes, mt. Peristeri (Hal.), mt. Murga, Profeti: Elias, Nimercka (Bald.); Thessalia: mt. Tringia (Hartl.), pr. Tsungera Malakasi, Uranaeos, mt. Zygos, Ghavellu (Haussk.), pr. Miluna et Tafilvris in Olympo, mt. Othrys (Form.); mt. Kiona (Hal.), mt. Oeta, Parnassus, Pateras, Pentelicon, Hymettus (Heldr.); Achaia: mt. Omplo pr. Patras, Kyllene (Heldr.), Chelmos (Hal.); Arcadia: pr. Zatuna (Orph.); Laconia: mt. Taygetos (Chaub.); Leucas: pr. Eukluvi (Bald.); Corcyra: ad litora (Clem.), pr. Spartilla (Mazz.), Myrtiotissa (Bickn.). — Maio, Jul. ♃.

4. **P. Psaridis** Heldr. herb. gr. norm. n. 1564. —

Differt (an specifice) a praecedente caulibus gracilibus flexuosis; floribus semper solitariis; calycis laciniis sublongioribus; capsula ovoidea laeviuscula, costis parum prominulis.

In regione alpina mt. Taygetos l. d. Megala Zonaria (Zahn). — Jun. Jul. ♃.

Obs. *Phyteuma Micheli* All. fl. ped. I. p. 115, indicatur a Mazziari in ant. ion. V. p. 208 pr. Spartilla Corcyrae, sed probabiliter erronee

6. Diosphaera Buser in bull. herb. Bois. II. p. 519.

a. Caules breves vel elongati, pulvinaria non formantes; flores numerosi, in corymbum densum capitatum congesti.

1. **D. Jacquini** Sieb. avis p. 3, rem. p. 3, Reise I. p. 437, II. p. 316 t. 6, in Flora V. I. p. 247; Spreng. syst. I. p. 724; Raul. cret. p. 803; Bald. viagg. Creta p. 75 (*Phyteuma*); Buser in bull. herb. Bois. II. p. 520. — *Sedum aizoon* S. et S. pr. I. p. 309; Sieb. avis rem. p. 4, Flora I. p. 272; Spreng. syst. II. p. 433, quoad pl. cret.; Raul. cret. p. 757; non L. — *Campanula Jacquini* DC. mon. camp. p. 251. — *Trachelium Jacquini* Bois. fl. or. III. p. 961, snppl. p. 335. — Exsicc.: Rev. pl. cret. a. 1883; Bald. it cret. n. 187.

Rhizomate crasso, multicauli; caulibus erectis, ved adscendentibus, 5—15 cm. altis, subangulatis, glabris vel ad angulos vel undique breviter puberulis; foliis coriaceis, nitidis, glabris vel puberulis, ovatis vel ellipticis, remote crenulatis vel serratis, radicalibus in petiolum sensim attenuatis, caulinis sessilibus; floribus pedicellis calyce brevioribus suffultis; bracteolis minimis, linearibus; calycis pubescentis laciniis lanceolatis, tubo brevioribus vel aequilongis; corolla tubulosa, coerulea, rarius albido-carnea, fere ad medium in lacinias lineares partita, calyce 3—4 plo longiore; capsula 3—2 loculari.

In rupium fissuris regionis subalpinae et alpinae. Creta: mt. Drakona (Rev.) et supra Anoia (Raul.) in mt. Sphacioticis (Sibth.), mt. Ida Sieb.). — Jul. Nov. ♃.

2. **D. dubia** Friv. pl. turc. exp. 1836 n. 74 (*Phyteuma*); Buser in bull. herb. Bois. II. p. 522. — *Trachelium rumelianum* Hampe in Flora 1837 p. 234; Hal. in ö. b. Z. 1897 p. 96. — *Phyteuma rumelicum* Grieseb. spic. II. p. 291. — *Campanula rumeliana* Vatke in Linnaea 1874 p. 705. — *T. rumelicum* Bois. fl. or. III. p. 961. — Exsicc.: Heldr. herb. n. 2493; Orph. fl. gr. n. 662, Sint. et Bornm. it. turc. n. 1846.

Praecedenti maxime affinis et ab ea caulibus elongatis, 15—35 cm. altis, foliis minoribus, minus coriaceis, vix lucidis, argutius serratis, capitulis majoribus floribus numerosioribus compositis tantum discedit. — Variat uti illa, caulibus et foliis glabriusculis vel puberulis.

Ad parietes rupium regionis submontanae et subalpinae. Thessalia: pr. Litochoron (Orph.) et Hagios Dionysios (Heldr.) in Olympo; in Sporadum insula Jura — Gyaros veterum — (Leon.). — Jun. Sept. ♃.

b. Caules nani, numerosissimi, pulvinaria densa formantes; flores 1—5, in axillis foliorum summorum.

3. **D. asperuloides** Orph. in Fl. gr. exs. n. 248 (*Campanula*); Buser in bull. herb. Bois. II. p. 523. — *Trachelium asperuloides* Bois. et Orph. diagn. ser. 2. III. p. 117, Fl. or. III. p. 962.

Rhizomate crassiusculo, multicauli; caulibus angulatis, crebre foliatis; foliis minimis, nitidis, ciliatis, suborbiculatis vel ovato-spathulatis, sessilibus; pedicellis calyci subaequilongis, medium versus bibracteolatis; calycis glabri vel hispiduli laciniis subulatis, tubo subduplo longioribus; corolla tubulosa, carnea, ad tertiam partem in lacinias lineares, erecto-patulas fissa; capsula biloculari. — Species pulchella habitu *Asperulae* quaedam pulvinaris.

In rupibus regionis mediae mt. Chelmos Arcadiae pr. Stygem, rarissime (Orph.). — Jul. Aug. ♃.

2. Tribus. **WAHLENBERGIEAE** End. gen. pl. p. 514.

7. **Edrajanthus** DC. pr. VII. p. 448.

a. Folia radicalia elliptica vel oblonga, in petiolum longiusculum attenuata.

1. **E. parnassicus** Bois. et Spr. diagn. VII. p. 17, Fl. or. III. p. 916, suppl. p. 332; Hal. in ö. b. Z. 1888 p. 760, Heldr. chlor. Parn. p. 23; (*Campanula*); Hal. Beitr. fl. Epir. p. 31, in ö. b. Z. 1897 p. 286. — Exsicc.: Heldr. herb. a. 1857, it. gr. septentr. a. 1879, pl. hellen. a. 1896; Hal. it. gr. a. 1888; Reis. pl. gr. a. 1894.

Pubescens; rhizomate crassiusculo, pleiocephalo, rosulas steriles caulesque floriferos edente; caulibus adscendentibus vel decumbentibus, simplicibus, remote foliatis; foliis obsolete repando-crenatis, rosularum et caulinis inferioribus ellipticis vel oblongis, in petiolum saepe elongatum attenuatis, superioribus lanceolatis, sessilibus; floribus brevissime pedunculatis, 2—5 in fasciculos terminales congestis, rarius solitariis, bracteis a basi ovata caudatis, patulis, floribus aequilongis vel brevioribus cinctis; calycis hispiduli, exappendiculati laciniis lanceolatis, tubo duplo longioribus; corolla anguste campanulata, violacea, glabra vel parce hirtula, laciniis calycis duplo longiore, lobis erecto-patulis. — Auctores, qui probabiliter formas tantum putatas, facie alienas, viderunt, speciem hanc ob capsulae dehiscentiam in genere *Edrajantho* collocandam, *Campanulae radicosae* affinem esse dicunt, a qua tamen specimina saltem intacta non solum notis fere omnibus, sed etiam habitu toto distinctissima sunt.

In saxosis regionis alpinae, rare. Aetolia: mt. Tymphrestus (Samarit.), Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Sprun.), Oeta (Heldr.); Elis: mt. Olenos (Heldr.). — Jul. Aug. ♃.

β. Folia linearia.

2. **E. graminifolius** L. sp. p. 166; S. et S. pr. I. p. 158, Fl. gr. II. p. 5 t. 206; Marg. et R. fl. Zante p. 64 (*Campanula*); DC. pr. VII. p. 448; Bois. fl. or. III. p. 887; Hal in z. b. G. 1888 p. 760; Beitr. fl. Epir. p. 31, in ö. b, Z. 1897 p. 286; Heldr. chlor. Parn. p. 22; Bald. riv. coll. bot. Alb. 1895 p. 56, 1896 p. 76. — *Wahlenbergia graminifolia* DC. mon. camp. p. 130. — *E. tenui-*

folius Bois. fl. or. III. p. 887, non (W. et K.) DC., qui foliis angustioribus ad apicem usque ciliatis, floribus minoribus, bracteis e basi lata abrupte caudatis, denticulatis et calycis laciniis multo angustioribus, linearibus, denticulatis specifice discedit et in Graecia non crescit. — *Hedraeanthus graminifolius* Wettst. monogr. p. 16. — *H. Kitaibelii* Wettst. l. c. p. 12 quoad pl. e mt. Parnasso, non *Edrajanthus Kitaibelii* DC., qui foliis latioribus denticulatis, bracteis denticulatis, calycis laciniis longioribus, sinubus appendiculatis discedit et in Graecia non crescit. — Exsicc.: Orph. fl. gr. n. 661; Hal. it. gr. II. a. 1893; Sint. it. thessal. n. 669.

Rhizomate crassiusculo, pleiocephalo, rosulas steriles caulesque floriferos edente; caulibus pilosis, erectis vel adscendentibus, simplicibus, foliatis; foliis integris, basin versus patenter ciliatis, glabris vel hirtulis; floribus brevissime pedunculatis, 3—8 in fasciculos terminales congestis, bracteis ovatis, caudatis, patulis, parte inferiori glabris, superiori hirsutis, crispule albo-ciliatis, floribus aequilongis vel brevioribus cinctis; calycis parce crispule pilosi, exappendiculati, laciniis triangulari-lanceolatis, tubo sublongioribus; corolla tubuloso-campanulata, coeruleo-violacea, parce hirtula, laciniis calycis duplo longiore, lobis erecto-patulis. — Occurrit caulibus elatioribus 12—20 cm. et nanis 4—7 cm. longis (*f. elatus et pusillus* Wettst. mon. p. 17).

β. **australis** Wettst. monogr. p. 17. — In omnibus partibus vegetivis pilis albis retroflexis villosus. — Intermediis cum typo conjuncta est. — Exsicc.: Orph. fl. gr. n. 419; Heldr. herb. n. 2735, herb. gr. norm. n. 721.

In rupestribus regionis alpinae. Epirus: mt. Tsumerka, Gamila (Buld.), Peristeri (Hal.), mt. Plaka pr. Chaliki (Sint.); Thessalia: mt. Olympus (Orph.); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Orph.); Zante (Sibth.). — Jul. Aug. ♃.

8. Jasione L. gen. n. 1005.

1. **J. dentata** DC. pr. VII. p. 415; Bois. fl. or. III. p. 885; pro var. *J. montanae*. — *J. Heldreichii* Bois. et Orph. diagn. ser. 2 VI. p. 120; Haussk. symb. p. 147; Form. in Ver. Brünn 1897 p. 28. — *J. montana* Form. in Ver. Brünn 1896 p. 36. non L. — Exsicc.: Orph. fl. gr. n. 663 (mt. Corfiati Macedoniae).

Biennis, pilis albis plus minus hispida; radice simplici, caules adscendentes vel erectos, simplices vel parce ramosos, inferne foliosos edente; foliis sessilibus, lineari-lanceolatis, undulatis; pedunculis elongatis, nudis; floribus in capitulum terminalem congestis; bracteis lanceolatis, setaceo-acuminatis, utrinque setaceo-dentatis, capitulo brevioribus; pedicellis calyci subaequilongis; calycis laciniis subulatis, tubo triplo longioribus, corolla cyanea dimidio brevioribus. — *J. montanae* L., bracteis ovatis integris vel dentatis, dentibus acutis nec setaceo-aristatis et pedicellis brevioribus, ideo capitulis densioribus diversae, proxima.

In rupestribus regionis montanae. Thessalia: pr. Tsungeri, Uranaeos, Malakasi (Haussk.), Konisko, Kerasia Sina, mt. Cuka, Hagios Elias, Phlambures et Mitrica in mt. Chassia, pr. Hepdominta Aderfia et Phlamburo in mt. Oxya, pr. Patsios in Olympo (Form.). — Jun. Jul. ⊙ N. v.

2. **J. supina** Sieb. in Spreng. syst. I. p. 810; Bald. riv. coll. bot. Alb. 1896 p. 76. – *J. orbiculata* Griseb. Reise I. p. 304, quae ex auctore ipso in spic. II. p. 293 synonimum speciei Sieberiani. —

Perennis, glabriuscula, caespitosa; radice simplici, caudiculos foliosos caulesque diffusos vel adscendentes, humiles, foliosos, simplices edente; foliis integris subcrenatisve, radicalibus spathulato-lanceolatis, ciliatis, in petiolum brevem decurrentibus, caulinis oblongis lanceolatisve, sessilibus; pedunculis superne nudis; floribus in capitulum terminalem congestis; bracteis late ovatis, acute dentatis, capitulum subaequantibus; pedicellis calyce brevioribus; calycis laciniis lanceolato-linearibus, tubo sesquilongioribus, corolla cyanea dimidio brevioribus.

In pascuis alpinis mt. Smolika supra Kerasovo distr. Konitza in Epiro (Bald.). — Jul. Aug. ♃. N. v.

LXV. Ordo. **Lobeliaceae** Juss. ann. mus. 18 p. 1.

1. **Laurentia** Mich. nov. gen. p. 18. t. 14.

1. **L. tenella** Biv. cent. I. p. 53 t. 2 (*Lobelia*); DC. pr. VII. p. 410; Weiss in z. b. G. 1869 p. 49; Raul. cret. p. 802; Bois. fl. or. III. p. 884; Bald. viagg. Creta p. 73. — *Lobelia laurentia* Willd. sp. I. p. 948 p. p.; Sieb. avis p. 3, rem. p. 3, in Flora I. p. 276, non L. — *Lob. setacea* S. et S. pr. I. p. 145, Fl. gr. III. p. 16 t. 221; Mazz. in ant. ion. V. p. 208. — *Lob. Bivonae* Tin. cat. hort. Pan. 1827 p. 279. — *Lob. cretica* Juss. herb. ex DC. pr. VII. p. 410. — Exsicc.: Rev. pl. cret. n. 100; Bald. it. cret. n. 31.

Acaulis, nana, glabra; foliis radicalibus, oblongo-spathulatis, integris, in petiolum attenuatis; scapis unifloris, folia multo superantibus, bracteolis 1—2 minimis obsitis; calycis laciniis lanceolatis, tubo duplo longioribus; corolla pallide coerulea, calyce subduplo longiori.

In locis uliginosis, ad scaturigines. Creta: ad Kastelli et Lusakies distr. Kissamos (Bald.), pr. Platania (Rev.), Canea (Weiss), Ipos, Malaxa, Almyros, Rhodakino, Francocastello, Embaro, Hierapetra, Turtuli (Raul.), in fauce Serisso (Sieb.); indicatur quoque pr. Manduchio et San Pantaleone Corcyrae (Mazz.). — Apr. Jun. ♃.

LXVI. Ordo. **Ericaceae** DC. fl. fr. III. p. 675.

1. Tribus. *Ericeae* Don gen. syst. III. p. 843. — Corolla marcescens; fructus capsularis, loculicide vel rarius septicide dehiscens.

1. Erica L. gen. n. 192.

1. **E. arborea** L. sp. p. 353; S. et S. pr. I. p. 256, Fl. gr. t. 351; Sieb. avis p. 3, rem. p. 4, in Flora I. p. 271; Ch. et B. exp. p. 111, Fl. pelop. p. 25; Marg. et R. fl. Zante p. 64; Friedr. Reise p. 264, 268 et 277; Fraas fl. class. p. 194; Ung. Reise p. 129; Raul. cret. p. 805; Bois. fl. or. III. p. 970; Heldr. fl. cephal. p. 50, in Sitzungsb. acad. Wiss. Berlin 1883 p. 4 et 5; Spreitz. in z. b. G. 1877 p. 728, 1887 p. 664; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal. Beitr. fl. Epir. p. 31; Haussk. symb. p. 147; Form. in Ver. Brünn 1895 p. 37. — *E. scoparia* Pieri corc. fl. p. 49, non L. — Exsicc.: Heldr. herb. dimorph. n. 70; Baen. herb. europ. n. 9253.

Fruticosa, elata; ramulis hirtis; foliis 3—4 nis, glabris, dorso sulcatis; corolla breviter campanulata, alba, calyce duplo longiore; antheris inclusis, basi appendiculatis; stylo exserto, stigmate peltato.

In fruticetis regionis inferioris et montanae passim per totam Graeciam. — Mart. Jun. ♄.

2. **E. verticillata** Forsk. descr. pl. fl. aeg. arab. p. 210; Ch. et B. exp. p. 110, Fl. pelop. p. 25; Link in Linnaea 1834 p. 578; Marg. et R. fl. Zante p. 64; Fraas fl. class. p. 194; Clem. sert. p. 65; Ung. Reise p. 129; Weiss in z. b. G. 1869 p. 752; Raul. cret. p. 805; Bois. fl. or. III. p. 970; Spreitz. in z. b. G. 1877 p. 728; Heldr. fl. cephal. p. 51, in Sitzungsb. acad. Wiss. Berlin 1883 p. 5, Fl. Aegina p. 308, chlor. Thera p. 16; Bald. in nuovo giorn. bot. ital. 1894 p. 98, viagg. Creta p. 76; Haussk. symb. p. 147. — *E. manipuliflora* Salisb. in trans. linn. VI. p. 344; S. et S. pr. I. p. 257. — *E. cinerea* Pieri corc. fl. p. 58, non L. — *E. mediterranea* Sieb. avis p. 3, rem. p. 4, in Flora I. p. 270; Fraas fl. class. p. 194; non L. — *E. cretica* Tausch in Flora XXII. p. 637. — *E. vagans* Benth. in DC. pr. VII. p. 667 p. p.; Friedr. Reise p. 264, non L. — Huc probabiliter: *E. multiflora* S. et S. pr. I. p. 256; Ch. et B. exp. p. 111, Fl. pelop. p. 25; Fraas fl. class. p. 194; Bois. fl. or. III. p. 970, quoad pl. graecam; Spreitz. in z. b. G. 1887 p. 664; vix L. — Exsicc.: Orph. fl. gr. n. 215; Heldr. herb. norm. n. 950 et 955, in Magn. fl. sel. n. 3059, in Baen. herb. europ. n. 5520; Rev. pl. cret. n. 104; Bald. it. bot. penins. balkan. a. 1889, it. cret. n. 186.

Fruticosa, humilis vel elata; ramulis glabris; foliis ternis, glabris, dorso convexis; corolla campanulata, carnea, calyce plus duplo longiore; antheris exsertis, non appendiculatis; stylo exserto, stigmate crassiusculo, stylo vix latiore.

In fruticetis regionis inferioris et montanae passim per totam Graeciam. — Aug. Mart. ♄.

Obs. *E. herbacea* L. sp. p. 352; S. et S. pr. I. p. 257; Fraas fl. class. p. 195; = *E. carnea* L. sp. p. 355. — Sec. Sibthorp „in Graecia ubique vulgaris", certe ex confusione quadam indicatur.

2 Tribus. *Arbuteae* DC. pr. VII. p. 580. — Corolla decidua; fructus baccatus, indehiscens.

2. Arbutus L. gen. n. 552.

1. **A. unedo** L. sp. p. 395; S. et S. pr. I. p. 274, Fl. gr. t. 373; Pieri corc. fl. p. 53; Urv. enum. p. 44; Ch. et B. exp. p. 115, Fl. pelop. p. 26; Marg. et R. fl. Zante p. 64; Friedr. Reise p. 272; Fraas fl. class. p. 195; Clem. sert. p. 65; Ung. Reise p. 130; Heldr. Nutzpfl. p. 39, fl. cephal. p. 50, in Sitzungsb. acad. Wiss. Berlin 1883 p. 5; Raul. cret. p. 805; Bois. fl. or. III. p. 966; Spreitz. in z. b. G. 1877 p. 728; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 147; Bald. viagg. Creta p. 76; Form. in Ver. Brünn 1895 p. 37, 1897 p. 55. — Exsicc.: Heldr. herb. norm. n. 211 et 956; Orph. fl. gr. n. 489.

Fruticosa, cortice rugoso, fusco; foliis coriaceis, persistentibus, glabris, obovatis, oblongis vel lanceolatis, acutis vel obtusis, acute vel obtuse serratis, breviter petiolatis; racemis nutantibus, glabris, subpaniculatis; calycis lobis breviter triangularibus; corolla ovoideo-urceolata, cerino-albida, obtuse 5loba; baccis globosis, amplis, muricato-verrucosis, coccineis, carnosis.

β. **turbinata** Pers. syn. I. p. 482; Bois. fl. or. III. p. 966; Haussk. symb. p. 147. — Baccae turbinatae, majores. — Exsicc.: N. v.

In regione inferiori et montanae, fruticeta sempervirentia efformans, per totam Graeciam. — Sept. Maio. ♄

2. **A. andrachne** L. sp. ed. 2 p. 566; S. et S. pr. I. p. 274, Fl. gr. t. 374; Sieb. avis rem. p. 4; Urv. enum. p. 44; Ch. et B. exp. p. 115, Fl. pelop. p. 26; Friedr. Reise p. 264, 268 et 273; Fraas fl. class. p. 195; Clem. sert. p. 65; Ung. Reise p. 130; Heldr. Nutzpfl. p. 39, fl. Cephalon. p. 50; Bois. fl. or. III. p. 966; Spreitz. in z. b. G. 1877 p. 728; Chlor. Waldverh. p. 30; Hal. in ö. b. Z. 1890 p. 39, Beitr. fl. Epir. p. 31; Haussk. symb. p. 147; Bald. riv. ooll. bot. Alb. 1896 p. 77. — *A. integrifolia* Lam. dict. I. p. 227; Sieb. avis p. 3, in Flor. V, 1 p. 247. — *A. Sieberi* Klotzsch in Linnaea XXIV. p. 9; Raul. cret. p. 805. — Exsicc.: Orph. fl. gr. n. 78; Heldr. herb. norm. n. 957; Sint. et Bornm. it. or. n. 1356 et 1927; Bald. it. alb. epir. IV. n. 133.

Fruticosa vel arborea, cortice laevi, rubro; foliis coriaceis, persistentibus, glabris, ovatis, integris, longiuscule petiolatis; racemis erectis, parce glanduloso-pilosis, paniculatis; calycis lobis rotundatis; corolla ovoidea, cerina, obtuse 5 loba; baccis globosis, parvis, reticulato-rimosis, aurantiacis, duris.

In fruticetis sempervirentibus regionis inferiores et montanae per totam Graeciam passim; sed in insulis Jonicis in Cephalonia tantum indicatur. — Mart. Maio. ♃

1 × 2, **A. unedo × andrachne** Bois. fl. or. III. p. 966. — *A. hybrida* Ker. bot. reg. t. 619. — *A. andrachnoides* Link en. I. p. 395. — *A. intermedia* Heldr. in Flora 1844 p. 14. — *A. nothocomaros* Heldr. herb. norm. n. 959. — *A. andrachne × unedo* Haussk. symb. p. 147. — Hybrida, characteribus et habitu nunc exacte intermedia, nunc vel ad unum alterum parentum vergens. — Attica: mt. Pentelicon, pr. Paphi (Heldr.); Argolis: mt. Chelone in peninsula Methana (Haussk.). —

LXVII. Ordo. **Pyrolaceae** Lindl. syst. bot. p. 62.

1. **Pyrola** L. gen. n. 554.

1. **P. chlorantha** Sw. in vet. acad. handl. 1810 p. 190 t. 5; Bois. fl. or. III. p. 974; Hal. in z. b. G. 1899 p. 190. — Icon: Fl. dan. t. 1693. — Exsicc.: Reis. fl. gr. a. 1898.

Glabra, rhizomate tenui, repente; caulibus erectis, ad basin foliatis, ceterum squamis 1—2 minutis munita; foliis ovatis vel orbiculatis, obtusis, integris; racemo laxo, brevi, vago; calycis laciniis ovatis, corolla 4 plo brevioribus; petalis obovatis, albo-viridulis; stylo inferne declinato, apice arcuato-adscendente incrassato et annulato, stigmatibus erectis, concretis.

In silvaticis montanis rarissime. Thessalia: mt. Olympus (Heldr.); Laconia: mt. Xerovuni pr. Langhada (Reiser). — Maio, Jul. ♃

2. **P. secunda** L. sp. p. 396; Bois. fl. or. III. p. 973; Haussk. symb. p. 147. — Icon: Fl. dan. t. 402. — Exsicc.: Orph. fl. gr. n. 797.

Glabra; rhizomate tenui, repente; caulibus erectis vel adscendentibus, inferne foliatis, ceterum squamis 1—4 minutis munitis; foliis ovatis vel ellipticis, acutis, tenuiter serrulatis; racemo denso, elongato, secundo; calycis laciniis ovatis, corolla 4 plo brevioribus; petalis ovalibus, albidis; stylo recto, apice nec incrassato, nec annulato, stigmatibus depressis.

In silvaticis subalpinis, rarissime. Thessalia: mt. Ghavellu in Pindo (Haussk.), mt. Olympus supra Hagios Dionysios (Orph.). — Jun. Aug. ♃

LXVIII. Ordo. **Monotropaceae** Nutt. gen. north. amer. pl. I. p. 272.

1. **Monotropa** L. gen. n. 536.

1. **M. hypopitys** L. sp. p. 387; Bois. fl. or. III. p. 975, suppl. p. 337; Heldr. fl. Cephal. p. 51. — *Hypopitys multiflora* Scop. fl. carn. ed. 2 I. p. 285. —

Aphylla; radice grumulosa; caule erecto, simplici, carnoso, squamis ovatis pellucidis obsito; racemo sub anthesi nutante, densifloro, demum erecto, elongato, floribus breviter pedicellatis; sepalis spathulato-lanceolatis; petalis obovato-cuneatis, pallide fusco-luteis, basi saccatis. — Planta aphylla, sicca nigricans, facie *Orobanches*.

α. **glabra** Roth tent. I. p. 180; Haussk. symb. p. 147. — *M. hypophegea* Wallr. sched. p. 191. — Icon: Rchb. f. 675. — Glabra. — Exsicc.: Orph. fl. gr. n. 335.

β. **hirsuta** Roth l. c.; Form. in Ver. Brünn 1897 p. 55. — Bracteae, sepala, petala, stamina et stylus hirti. — Exsicc.: N. v.

In silvaticis montanis, rare. Thessalia: in oropedio Neuropolis (Haussk.), pr. Phlambures et Mitrica in mt. Chassia (Form.), mt. Olympus (Heldr.); Peloponnesus: mt. Chelmos (Orph.), in faucibus Langada in mt. Taygetos (Zahn); Cephalonia: mt. Aenos (Letourn.). — Jul. Aug. ♃

III. Subclassis **Corolliflorae** DC. pr.

LXIX. Ordo. **Styraceae** DC. pr. VIII. p. 244.

1. **Styrax** Tourn. inst. p. 598.

1. **S. officinalis** L. sp. p. 444; S. et S. pr. I. p. 275, t. 375; Sieb. avis. p. 3, rem. p. 4; Ch. et B. fl. pelop. p. 30; Fraas fl. class. p. 194; Clem. sert. p. 80; Heldr. Nutzpfl. p. 38; Raul. cret. p. 807; Bois. fl. or. IV. p. 35; Chloros Waldverh. p. 37; Bald. viagg. Creta p. 76. — Exsicc.: Orph. fl. gr. n. 5; Heldr. herb. norm. n. 497; Rev. pl. cret. n. 106, in Baen. herb. europ. n. 4895, in Magn. fl. sel. n. 1756; Dörfl. fl. aeg. n. 106.

Fruticosa vel arborescens; foliis petiolatis, ovatis vel ellipticis, integris, supra viridibus glabriusculis, subtus cano-tomentosis; cymis terminalibus, paucifloris; pedicellis calyce truncato longioribus, eoque canotomentosis; corollae albae, tomentosae laciniis lanceolatis, calyce 3—4plo longioribus; fructu globoso, calyce persistente basi adnato. — Flores suaveolentes.

Ad rivos, in fruticetis regionis inferioris et montanae. Boeotia: pr. Orchomenos (Fraas); Attica: mt. Parnes, ad Cephissum, mt. Pentelicon (Fraas); mt. Parnassus (Sprun.); Achaia: pr. Dervenion (Hal.); Messenia: pr. Andrusa (Chaub.); Creta (Sibth.): pr. Topolia (Bald.), Platania, Rumata, Askyphos, Retymo, Rustika, Dramia, Vrisinas, Perama, Damasta, insula Dia (Raul.); Cycladum insula Naxos (Leon.). — Apr. Maio. ♄

LXX. Ordo. **Ebenaceae** Vent. tabl. p. 445.

1. **Diospyros** L. gen. n. 1151.

1. **D. lotus** L. sp. p. 1057. — Icon: Pall. fl. ross. t. 58. — Exsicc.: Heldr. it. gr. septentr. a. 1879.

Arborea; foliis petiolatis, oblongis, acuminatis, integris, supra glabriusculis, subtus pubescentibus, discoloribus; floribus minutis, axillaribus, brevissime pedicellatis; calyce brevi, acute 4 lobo, ciliato et intus villoso; corolla mascula calyce triplo longiore, urceolato-campanulata, rubro-viridula, ad tertiam partem in lobos reflexos, rotundatos, ciliatos fissa; bacca fulva, calyce aucto suffulta.

Phthiotis: pr. Hypata ad radices mt. Aeta (Heldr.); vix spontanea. — Maio, Jun. ♄

LXXI. Ordo. **Jasminaceae** R. Br. pr. p. 520.

1. **Jasminum** Tourn. inst. p. 597.

1. **J. fruticans** L. sp. p. 7; Heldr. Nutzpfl. p. 29; Form. in D. bot. Monat. 1890 p. 24, in Ver. Brünn 1896 p. 59, 1897 p. 43; Bald. it. alb. epir. 1896 p. 77. — Icon: Rchb. t. 36. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Fruticosum, glabrum, ramis angulatis; foliis trifoliatis, rarius simplicibus, foliolis oblongis, obtusis, integris; floribus 2—4, terminalibus; calycis dentibus subulatis, tubo 2—3 plo longioribus; corollae flavae tubo calyce duplo longiore, laciniis obtusis.

In dumetis, rupestribus, silvaticis regionis inferioris et montanae. Epirus: pr. Mesojefira et Vromonero distr. Ljaskovik (Bald.); Thessalia: ad monasterium Korona (Heldr.), pr. Chaliki, Koturi, Velitsena, Kataphygi, Vlazda in Pindo, pr. Palaeokastro, Pharsalus, Suilismen, Diodendron, Nezeros, Rapsani in Olympo, in valle Tempe, mt. Pelion (Form.); Messenia (Heldr.). — Apr. Jun. ♄

2. **J. humile** L. sp. p. 7; Ch. et B. exp. p. 14, Fl. pelop. p. 1; Bois. fl. or. IV. p. 42. — Icon: Bot. reg. t. 350. —

Fruticosum, glabrum, ramis teretiusculis; foliis trifoliolatis, foliolis ovatis, integris, terminali acutato; floribus 1—3, subterminalibus; calyce brevissime dentato; corollae flavae tubo calyce multoties longiore, laciniis obtusis.

In collibus siccis pr. Methone Messeniae (Chaub.). — Apr. Jun. ♄ N. v.

Obs. In hortis coluntur: *J. sambac* L. sp. p. 6; Heldr. Nutzpfl. p. 29. — *J. odoratissimum* L. sp. p. 7; Dallap. prosp. p. 1; Heldr. l. c. — *J. grandiflorum* L. sp. ed. 2 p. 9; Dallap. l. c.; Heldr. l. c. — *J. officinale* L. sp. p. 7; Heldr. l. c. —

LXXII. Ordo. **Oleaceae** Lindl. syst. bot. p. 67.

1. Tribus. *Oleineae* Don in Loud. arb. p. 283. — Fructus carnosus.

1. Olea L. gen. n. 20. — Corolla induplicato-valvata, subrotata; stamina exserta; drupa putamine osseo. Inflorescentia axillaris.

1. **O. europaea** L. sp. p. 8; S. et S. pr. I. p. 4, Fl. gr. I. p. 3 t. 3; Sieb. avis p. 2, rem. p. 2; Dallap. prosp. p. 2; Urv. enum. p. 1; Ch. et B. exp. p. 14, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 432—440, ubi 35 s. d. species enumerantur; Marg. et R. fl. Zante p. 65; Friedr. Reise p. 263 et 269; Fraas fl. class. p. 154; Ung. Reise p. 126; Heldr. Nutzpfl. p. 30, in Sitzungs. acad. Wiss. Berlin 1883 p. 5 et 7, Fl. cephal. p. 51, Fl. Aegina p. 379, chlor. Thera p. 27; Raul. cret. p. 807; Bois. fl. or. IV. p. 36; Chlor. Waldverh. p. 33; Hal. in z. b. G. 1888 p. 760, Beitr. fl. Epir. p. 31, Beitr. fl. Thessal. p. 16, in ö. b. Z. 1896 p. 15, 1897 p. 97; Form. in D. bot. Mon. 1890 p. 24, in Ver. Brünn 1895 p. 31, 1896 p. 59, 1897 p. 43; Haussk. symb. p. 148; Bald. viagg. Creta p. 77. — Exsicc.: Heldr. herb. norm. n. 858—861; Baen. herb. europ. n. 9294.

Arborea vel fruticosa, sempervirens, foliis coriaceis, glabris, oblongis vel lanceolatis, integris, mucronulatis, supra cinereo-virentibus, subtus lepidoto-canis; floribus in racemis axillaribus; corolla alba; drupis globosis vel ellipsoideis.

α. **typica.** — *O. sativa* Hoffm. et Lk. fl. port. I. p. 387. — Arborea, ramis inermibus, foliis lanceolatis, fructu majore; magis carnoso. — Valde variat quoad fructus magnitudinem et formam, nec non putaminis superficiem.

β. **oleaster** Hoffm. et Lk. fl. port. I. p. 387. — Arborea vel fruticosa, ramis plus minus indurato-spinescentibus, foliis oblongis praecipue in formis fruticosis saepe diminutis, fructu minore, minus carnoso.

Stirps silvestris in collibus apricis regionis calidae totius Graeciae; — *α.* colitur abunde in formis innumeris. — Maio, Jun. ♄

2. Phillyrea Tourn. inst. t. 367. — Corolla imbricata, subrotata; stamina exserta; drupa putamine chartaceo, fragili. Inflorescentia axillaris.

1. **P. media** L. sp. ed. 2 p. 10; S. et S. pr. I. p. 3; Pieri corc. fl. p. 2; Sieb. in Flora I. p. 271, avis rem. p. 2; Ch. et B. exp. p. 14, Fl. pelop. p. 1; Marg. et R. fl. Zante p. 65; Friedr. Reise p. 271; Fraas fl. class. p. 153; Ung. Reise p. 126; Heldr. Nutzpfl. p. 29, in Sitzungs. acad. Wiss. Berl. 1883 p. 7, Fl. cephal. p. 51; Raul. cret. p. 807; Bois. fl. or. IV. p. 37; Chloros Waldv. p. 34; Gelmi in bull. soc. bot. it. 1889 p. 450; Form. in D. bot. Monat. p. 24, in Ver. Brünn 1896 p. 59, 1897 p. 43; Hal. Beitr. fl. Epir.

p. 31; Haussk. symb. p. 148. — *P. latifolia* S. et S. pr. I. p. 4, Fl. gr. I. p. 2 t. 2; Pieri corc. fl. p. 2; Sieb. avis p. 2, rem. p. 2; Mazz. in ant. ion. II. p. 440; Fraas fl. class. p. 153; Heldr. Nutzpfl. p. 29; Raul. cret. p. 807; Chlor. Waldverh. p. 34; non L.; quae folia cordato-ovata habet. — *P. pedunculata* Ch. et B. exp. p. 14, Fl. pelop. p. 1 (f. pedunculis elongatis, fructu duplo longioribus). — *P. spinosa* Tausch in Flora XIV. p. 642 (f. foliis ovatis vel lanceolatis, spinuloso-serratis, serraturis patulo-recurvatis). — *P. angustifolia* Mazz. in ant. ion. II. p. 440; Ch. et B. fl. pelop. p. 1; non L., quae folia lineari-lanceolata integerrima habet. — *P. obliqua* Willd. sp. pl. I. p. 43; Mazz. in ant. ion. II. p. 440 (f. foliis lanceolato-oblongis, serratis, oblique flexis). — Exsicc.: Heldr. reliqu. Orph. a. 1886; Bald. it. bot. penins. balk. a. 1889; Sint. et Bornm. it. turc. n. 1364.

Fruticosa vel arborea; foliis coriaceis, glabris, ovatis, oblongis vel ovato-lanceolatis, obtusis acutisve, serratis integrisve, breviter petiolatis vel subsessilibus; floribus ad axillas fasciculatis, pedicellis brevissimis vel eis sublongioribus suffultis; calycis lobis obtusis; corolla viridi-alba, lobis ovatis; drupa globosa, mutica vel apiculata. — *P. latifolia et angustifolia* L. a. variis auctoribus in Graecia indicata, ibi non occurrunt et sec. Bois. fl. or. IV. p. 37 in Oriente toto deesse videntur.

In fruticetis et silvaticis regionis inferioris et montanae per totam Graeciam. — Maio, Jun. ♄.

3. Ligustrum L. gen. n. 22. — Corolla subinduplicato-valvata, infundibuliformis; stamina inclusa; bacca, loculis membrana tenuissima vestitis. Inflorescentia terminalis.

1. **L. vulgare** L. sp. p. 7; Pieri corc. fl. p. 1; Ch. et B. exp. p. 14, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 432; Fraas fl. class. p. 153; Bois. fl. or. IV. p. 37; Hal. Beitr. fl. Achaia p. 26; Haussk. symb. p. 148. — *L. sempervirens* Mazz. in ant ion. II. p. 432 (f. foliis persistentibus). — Icon: Rchb. t. 33. — Exsicc.: Sint. et Bornm. it. turc. n. 1365.

Fruticosa; foliis subcoriaceis, glabris, oblongis ellipticisve, acutis obtusisve, integris, breviter petiolatis; floribus in paniculam thyrsoideam dispositis, breviter pedicellatis; calyce breviter et obtuse dentato; corolla alba, tubo calyce longiore; bacca pisiformi, nigra.

Ad sepes, in dumosis regionis montanae, rare. Thessalia: in oropedio Neuropolis (Haussk.), pr. Litochori in Olympo (Sint.); Euboea: mt. Telethrion (Heldr.); Boeotia: ad Cephissum, pr. Lebadea (Fraas); Achaia: mt. Chelmos (Orph.), pr. Kalavryta (Heldr.), Sudena, Planideri (Hal.); Arcadia (Chaub.); Corcyra (Pieri): pr. Synarades, Calefactiones, Fano, in scopulis Diapulo et Diascopo (Mazz). — Apr. Jun. ♄.

2. Tribus. *Fraxineae* Bartl. ord. nat. p. 218. — Fructus siccus, samaroideus.

4. Fraxinus L. gen. n. 1166.

1. Sectio. *Ornus* Pers. syn. I. p. 9 pro gen.; Pers. syn. II. p. 605. — Flores corollati.

1. **F. ornus** L. sp. p. 1057; S. et S. pr. I. p. 5, Fl. gr. I. p. 4 t. 4; Ch. et B. exp. p. 14, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 440; Friedr. Reise p. 281; Fraas fl. class. p. 153; Landerer in Flora XV. p. 394; Heldr. Nutzpfl. p. 31, in Sitzungsb. acad. wiss. Berlin 1883 p. 4 et 7; Bois. fl. or. IV. p. 39; Haussk. symb. p. 148; Hal. in ö. b. Z. 1896 p. 15; Bald. riv. coll. bot. Alb. 1895 p. 58; Form. in Ver. Brünn 1896 p. 59, 1897 p. 43. — *F. rotundifolia* Lam. dict. II. p. 546; Fraas fl. class. p. 153; (f. foliolis rotundatis). — *Ornus europaea* Pers. syn. I. p. 9. — Exsicc.: Orph. fl. gr. n. 705 et 706; Heldr. it. thessal. n. 105; Sint. it. or. a. 1889 n. 1930, it. thessal. n. 415; Sint. et Bornm. it. turc. 1366; Bald. it. alb. epir. III. n. 190.

Arborea; follis synanthiis, imparipinnatis, foliolis 3—4 jugis, petiolulatis, ovali-lanceolatis, acuminatis, crenatis, glabris vel subtus ad nervum barbatis; floribus in cymas racemoso-paniculatas, terminales et axillares dispositis; petalis albis, linearibus, calyce multo longioribus; samaris lineari-lanceolatis, emarginatis vel truncatis.

In dumetis nemoribusque regionis montanae. Epirus: pr. Philippiada (Bald.); Thessalia: pr. Chaliki, Klinovo, in oropedio Neuropolis, pr. Meteora (Haussk.), Kalabaka (Sint.), mt. Mitrica in mt. Chassia, pr. Patsios (Form.) et Hagios Dionysios (Orph.) in Olympo, in valle Tempe, pr. Laspochory, mt. Ossa (Heldr.); Euboea: pr. Mantudi (Fraas); Aetolia: pr. Aetolikon (Reis.); mt. Parnassus (Sibth.); Attica: mt. Parnes pr. Panagia Kliston (Heldr.); Arcadia: pr. Trikala (Fraas), mt. Diaforti pr. Andrizena (Friedr.), pr. Phigalea (Chaub.); Laconia: mt. Malevo (Fraas), Taygetos (Orph.); Corcyra: mt. San Stefano pr. Benizze (Mazz.). — Apr. Maio. ♄.

2. Sectio. *Fraxinaster* DC. pr. VIII. p. 276. — Flores apetali.

2. **F. excelsior** L. sp. p. 1057; Fraas fl. class. p. 154; Bois. fl. or. IV. p. 39, suppl. p. 343; Heldr. in Sitzungsb. acad. wiss. Berl. 1883 p. 5; Chloros Waldverh. p. 32; Hal. Beitr. fl. epir. p. 31; Haussk. symb. p. 148. — Icon: Rchb. t. 31. —

Arborea; foliis hysteranthiis, imparipinnatis, foliolis 4—6 jugis, subsessilibus, oblongo-lanceolatis, acuminatis, serratis, glabris vel subtus ad nervum puberulis; floribus nigricantibus, in cymas ad ramorum apicem erectas, dein pendulas dispositis; samaris ellipticis, obtusis vel retusis. —

In silvis montanis, rare. Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.); Thessalia: mt. Pindus (Fraas) in oropedio Neuropolis, pr. Karditza (Haussk.), mt. Oeta, Ossa (Heldr.); Aetolia (Chloros). — Mart. Apr. ♄.

LXXIII. Ordo. **Asclepiadaceae** R. Br. in Wern. soc. I. p. 12.

Dispositio generum:

a. Corolla 5 partita, fauce corona, e squamis in aristam uncinatam elongatis constante, munita; filamenta omnino vel superne saltem libera; antherae apice cohaerentes, dorso barbatae.

1. Periploca L.

b. Corolla 5 loba vel 5 partita, fauce corona 5—10 loba vel 5 phylla munita; filamenta connata; antherae appendice membranacea terminatae.

α. Folliculi laeves.

× Corona scutelliformis, carnosa, lobis 5—10, rotundatis vel obscure apiculatis.

2. Vincetoxicum Moench.

×× Corona tubuloso-campanulata, laciniis 10, ligulatis.

3. Cynanchum L.

××× Corona phyllis 5, liberis, oblongo-linearibus constans.

4. Cionura Griseb.

β. Folliculi echinati; corona phyllis 5, gynostegio adnatis, cucullaribus vel navicularibus constans.

5. Gomphocarpus R. Br.

1. Periploca L. gen. n. 300.

1. **P. graeca** L. sp. p. 211; Desf. arb. I. p. 199; Raul. cret. p. 808; Bois. fl. or. IV. p. 49; Haussk. symb. p. 148. — Icon: Fl. gr. t. 249. — Exsicc.: Heldr. herb. a. 1878.

Fruticosa, volubilis; foliis ovatis vel ovato-lanceolatis, obtusis vel acutis, integris, glabris, breviter petiolatis; cymis ramulos terminantibus, laxifloris; corollae fuscae, viridi marginatae, laciniis oblongis, patulis, superne barbatis, subtus glabris; parapetalis basi biauriculatis, in aristam, petalorum dimidium aequantem abeuntibus; folliculis cylindraceis, 9—10 cm. longis.

Ad sepes, rivulos regionis inferioris, rarissime. Acarnania (Heldr.); Aetolia: pr. Aetolikon (Heldr.); Argolis: peninsula Methana (Haussk.); Creta (Desf.). — Maio, Jun. ♃.

2. Vincetoxicum Moench meth. p. 317.

a. Corolla intus hirsuta.

α. Corolla flava.

1. **V. canescens** Willd. nov. act. nat. cur. III. p. 418 (*Asclepias*); Decaisne in DC. pr. VIII. p. 523; Raul. cret. p. 808; Bois. fl. or. IV. p. 52. — *V. luteum* Sieb. avis rem. p. 3. — *Cynanchum luteum* Spreng. syst. I. p. 852. — Exsicc.: Orph. herb. n. 343 (Chios).

Tomentello-canescens; caulibus erectis, flexuosis, simplicibus vel inferne ramosis, foliis ovatis, acutis, breviter petiolatis, umbellis sub-

sessilibus, axillaribus et terminalibus; floribus brevissime pedicellatis; corollae laciniis oblongo triangularibus; coronae lobis 5, obtusis, concaviusculis; folliculis inferne inflatis.

In petrosis regionis inferioris montium. Creta mt. Lassiti (Sieb.). — Maio, Jun. ♃. N. v.

β. Corolla atropurpurea.

2. **V. speciosum** Bois. et Sprun. diagn. IV. p. 39, Fl. or. IV. p. 53; Haussk. symb. p. 53. — *V. triste* Griseb. spic. II. p. 68. — *Asclepias Dioscoridis* Fraas fl. class. p. 159. — Exsicc.: Heldr. it. thessal. IV. a. 1885, reliqu. Orph. a. 1886; Sint. it. thessal. n. 1182.

Caulibus elatis, simplicibus, crispule velutinis; foliis pubescentibus, brevissime petiolatis et sessilibus, cordato-ovatis, obtusis vel acutis, superioribus oblongis, acuminatis; umbellis axillaribus, subsessilibus; pedicellis crispo-lanatis, flori aequilongis; calycis lanati laciniis lanceolatis; corollae majusculae laciniis lanceolatis; coronae 5 fidae lobis latis, obtusis, apice incrassatis; folliculis tomentellis, apice longe attenuatis. — Caulis bipedalis et longior, folia saepe 10 cm. longa, 5—6 cm. lata, corolla diametro 1 cm. lata.

In regione montana et alpina. Thessalia: in oropedio Neuropolis, mt. Ghavellu (Haussk.), supra Prioni pr. Sermeniko (Sint.); Aetolia: mt. Tymphrestus (Sprun.); Euboea: mt. Dirphys (Fraas). — Jun. Jul. ♃.

3. **V. nigrum** L. sp. p. 216 (*Asclepias*); Moench meth. p. 317. — *Cynanchum nigrum* R. Br. mem. wern. I. p. 147. — Exsicc.: Sint. it. thessal. n. 690; Dörfl. fl. gr. n. 287.

Caulibus erectis, saepe volubilibus, puberulis; foliis breviter petiolatis vel sessilibus, ovatis vel ovato-lanceolatis, acutis vel acuminatis, ad nervos puberulis, inferioribus basi interdum cordatis; umbellis axillaribus, pedunculatis vel sessilibus; pedicellis puberulis, flore 2—3 plo longioribus; calycis puberuli laciniis lanceolatis; corollae parvae laciniis oblongis, obtusiusculis; coronae 10 lobae lobis obtusis, alternatim brevioribus; folliculis glabrescentibus, apice longe attenuatis.

In subalpinis mt. Plaka supra Chaliki Thessaliae (Sint.) et mt. Korax Aetoliae (Leon.). — Jun. Jul. ♃.

b. Corolla glabra.

4. **V. nivale** Bois. et Heldr. diagn. VII. p. 20, Fl. or. IV. p. 55; Heldr. chlor. Parn. p. 23. — Huc forsan e loco: *Asclepias vincetoxicum* S. et S. pr. I. p. 166, vix L.; nec non *V. fuscatum* Hal. in z. b. G. 1888 p. 760 (specimina deflorata). — Exsicc.: Pichl. pl. gr. a. 1876; Dörfl. fl. gr. n. 287.

Caulibus erectis, simplicibus, pubescentibus; foliis subsessilibus sessilibusve, ad nervos puberulis, inferioribus a basi truncata vel subcordata ovatis, superioribus oblongis, acuminatis; umbellis axillaribus, pedunculatis; pedicellis puberulis, flore aequilongis; calycis glabri laciniis lanceolatis; corollae luteae laciniis oblongis; corona breviter in lobos 5 crassiusculos, obtusos fissa; folliculis ignotis. — Caules 10—35 cm. alti.

β. **undulatum** Heldr. in Bois. diagn. ser. 2 VI. p. 121. — Folia petiolata, margine subundulata, corolla intus pilis nonnullis obsoletis obsita. — Exsicc.: N. v.

In saxosis regionis superioris. Thessalia: mt. Olympus (Heldr.); Aetolia: mt. Korax (Leon.); mt. Parnassus (Sprun.); Peloponnesus: mt. Chelmos ad fontem Stygis (Orph.), mt. Taygetos ad nives cacuminis Hagios Paraskevi (Heldr.); — β. Eurytania: mt. Tymphrestus loco Cancellia dicto (Samar.). — Jun. Jul. ♃.

Obs. Ulterius observandum anne *V. fuscatum* Bois. fl. or. IV. p. 55 e mt. Parnasso ad speciem praecedentem pertineat. — Quid sit *V. laxum* Sieb. avis p. 3 e Creta et *Asclepias vincetoxicum* Dallap. prosp. p. 35, nescio.

3. Cynanchum L. gen. n. 301.

1. **C. acutum** L. sp. p. 212; S. et S. pr. I. p. 165, Fl. gr. III. p. 45 t. 250; Urv. enum. p. 27; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 60; Heldr. fl. cephal. p. 52, in Sitzungsb. acad. Wiss. Berl. 1883 p. 9; Form. in D. bot. Monat. 1891 p. 24, 1898 p. 78, in Ver. Brünn 1895 p. 32, 1896 p. 59, 1897 p. 43; Bald. in nuov. giorn. bot. ital. 1894 p. 98; Haussk. symb. p. 148. — *C. monspeliacum* L. sp. p. 212; S. et S. pr. I. p. 165, Fl. gr. III. p. 46 t. 251; Ch. et B. exp. p. 79, Fl. pelop. p. 17; Marg. et R. fl. Zante p. 66; Fraas fl. class. p. 160; Ung. Reise p. 126. — Exsicc.: Heldr. herb. norm. n. 119; Orph. fl. gr. n. 37; Sint. et Bornm. it. turc. n. 1367.

Glaucum, caulibus elongatis, volubilibus, glabris vel parce puberulis; foliis a basi cordata ovatis vel oblongis, acutis, glabris; petiolis puberulis, limbo brevioribus; floribus in cymas axillares pedunculatas dispositis; calycis pubescentis laciniis oblongis; corollae albae vel roseae laciniis oblongis, obtusis; folliculis acuminatis, glabris.

In dumetis, ad sepes, in ruderatis, olivetis, maritimis. Epirus: pr. Luros, Ranca (Form.), pr. Prevesa (Bald.); Thessalia: pr. Kalabaka, Trikala, Velestino, Larissa, Makrichori, Baba, Kapurna, Koryza, Volo, Lechonia, Agria, Stylida, Lamia, Miluna in Olympo (Form.), Litochoron (Sint.); Boeotia: pr. Lebadea (Fraas); Attica: frequens pr. Athenas, ad Phaleron, Piraeum (Heldr.), pr. Eleusis (Ung.); Peloponnesus: ad fl. Eurotas, in planitie Helos (Chaub.); Cycladum insula Syra (Weiss), Melos (Urv.); Zante (Marg.); Cephalonia: pr. Lixuri (Heldr.). — Jun. Aug. ♃.

4. Cionura Griseb. spic. II. p. 69.

1. **C. erecta** L. sp. p. 213; S. et S. pr. p. 166; Sieb. avis p. 3, rem. p. 3; Urv. enum. p. 27; Ch. et B. exp. p. 78, Fl. pelop. p. 16; Fraas fl. class. p. 160; (*Cynanchum*); Griseb. l. c.; Bois. fl. or. IV. p. 62; Hal. in z. b. G. 1888 p. 760, Beitr. fl. Thessal. p. 16. —

Marsdenia erecta R. Br. mem. Wern. I. p. 29; Heldr. Nutzpfl. p. 31, in ö. b. Z. 1898 p. 184; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 808; Bald. viagg. Creta p. 77; Haussk. symb. p. 148; Hal. in z. b. G. 1899 p. 190. — Huc sec. Raul. l. c.: *Cynanchum acutum* Sieb. avis rem. p. 3, in Flora I. p. 272, non L. — Icon: Jacq. misc. t. 2. — Exsicc.: Orph. fl. gr. n. 88; Heldr. herb. norm. n. 258, in Magn. fl. sel. n. 2526; Rev. pl. cret. n. 108.

Glabra, basi suffrutescens; caulibus superne interdum subvolubilibus; foliis petiolatis, cordato-ovatis, breviter acuminatis; cymis pedunculatis, axillaribus et terminalibus; calycis laciniis ovatis; corollae albae laciniis linearibus; folliculis oblongis, apice attenuatis.

In collibus siccis, olivetis, arenosis maritimis. Thessalia: pr. Kalabaka (Hal.); Phocis: pr. Amphissa (Hal.); Boeotia (Fraas); Attica: pr. Athenas, ad Cephissum, pr. Oropos, Phaleron (Heldr.); Euboea: pr. Chalkis (Pichl.); Achaia: pr. Patras (Heldr.); Laconia: pr. Sotirianika (Zahn), Kalamata, Armyros, Zarnate, Megalopolis, Gortys, Sparta, ad fl. Eurotas (Chaub.); Cycladum insula: Tenos (Weiss), Syra (Urv.), Naxos (Heldr.); Creta: pr. Kissamos (Rev.), Canea (Weiss), Kladiso, Platania, Hagios Paulos, Hagios Nicolaos supra Samaria, Theriso (Raul.). — Maio, Jul. ♄.

5. Gomphocarpus R. Br. mem. Wern. I. p. 38.

1. **G. fruticosus** L. sp. p. 216; Ch. et B. exp. p. 78, Fl. pelop. p. 16; (*Asclepias*); R. Br. l. c.; Marg. et R. fl. Zante p. 66; Heldr. Nutzpfl. p. 31, in ö. b. Z. 1898 p. 184, chlor. Thera p. 30; Bois. fl. or. IV. p. 61; Spreitz. in z. b. G. 1890 p. 297; Hal. in ö. b. Z. 1896 p. 15, in z. b. G. 1899 p. 190. — Icon: Jacq. misc. t. 2. — Exsicc.: Dörfl. fl. gr. n. 175.

Glabrescens vel minute pubescens, basi suffrutescens; ramis virgatis, stricte foliosis; foliis lineari-lanceolatis, subpetiolatis; floribus umbellatis; umbellis pedunculatis, axillaribus; calycis laciniis lanceolatis; corollae albae laciniis ovatis, ciliatis; folliculis ovato-acuminatis, brevissime puberulis, muricibus linearibus elongatis obsitis.

Ad vias, rivulos regionis inferioris, olim cultus, nunc subspontaneus. Zacynthus (Marg.); Peloponnesus (Guerin): pr. Gaitzies (Zahn); Cyclades: Naxos (Chaub.), Andros, Thera et fere in omnibus ceteris insulis (Heldr.); Creta: pr. Hagia Rumeli (Spreitz.). — Jun. Aug. ♄.

LXXIV. Ordo. Apocynaceae R. Br. mem. wern. I. p. 59.

1. Nerium L. gen. n. 181.

1. **N. oleander** L. sp. p. 209; S. et S. pr. I. p. 164, Fl. gr. III. p. 42 t. 248; Sieb. avis p. 3, in Flora I. p. 273; Dallap. prosp. p. 34; Urv. enum. p. 27; Ch. et B. exp. p. 78, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 66; Friedr. Reise p. 264; Fraas fl. class. p. 159;

Heldr. Nutzpfl. p. 31, Fl. cephal. p. 52, chlor. Thera p. 16, chlor. Mykon. p. 247; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 808; Bois. fl. or. IV. p. 47; Spreitz. in z. b. G. 1887 p. 664; Hal. in z. b. G. 1888 p. 760; Bald. viagg. Creta p. 76; Haussk. symb. p. 148; Form. in Ver. Brünn 1895 p. 32, 1897 p. 43. — Exsicc.: Heldr. herb. norm. n. 960; Rev. pl. cret. n. 107.

In torrentium alveis et ad ripas rivulorum regionis inferioris frequens; deesse videtur in Epiro. — Maio, Aug. ♄.

2. Vinca L. gen. n. 295.

a. Folia ovata, basi rotundata vel subcordata, apice attenuata.

1. **V. major** L. sp. p. 209; Raul. cret. p. 808; Bois. fl. or. IV. p. 45. — Icon: Rchb. t. 22. — Exsicc.: Pichl. pl. gr. a. 1876; Sagb. pl. corc. n. 42.

Caulibus sterilibus prostratis, non radicantibus, floriferis erectis; foliis breviter petiolatis, margine ciliatis, ceterum glabris; pedunculis ex axillis superioribus solitariis, folio brevioribus; calycis laciniis linearibus, acuminatis, ciliatis, tubum corollae saepe subaequantibus; corolla violacea. — Sequentibus omnibus partibus multo major.

In umbrosis regionis inferioris, rarissime. Corcyra: pr. urbem (Sagb.), an sponte; Attica: pr. Marathon (Pichler), sed dubito an specimina re vera in Attica lecta sint; Creta: ad radices mt. Karadagh pr. Candia (Heldr.). — Febr. Apr. ♃.

a. Folia elliptica vel ovato-lanceolata, utrinque attenuata.

2. **V. minor** L. sp. p. 209. — Icon: Rchb. t. 21. — Exsicc.: Heldr. fl. thessal. a. 1883.

Caulibus sterilibus prostratis, radicantibus, floriferis erectis; foliis brevissime petiolatis, omnino glabris; pedunculis ex axillis superioribus solitariis, folio aequilongis vel longioribus; calycis laciniis lanceolatis, acutis, glabris, tubo corollae multo brevioribus; corolla violacea.

In nemorosis Thessaliae pr. Velestinos (Heldr.); indicatur quoque a S. et S. pr. I. p. 164, in Arcadia, agro Eliensi et Argolica, nec non a Fraas fl. class. p. 159 in mt. Kyllene, sed loca probabiliter ad sequentem speciem spectant. — Mart. Maio. ♃.

3. **V. herbacea** W. et K. pl. rar. Hung. I. p. 8 t. 9; Ch. et B. exp. p. 78, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 66; Friedr. Reise p. 279; Ung. Reise p. 126; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 45; Heldr. fl. cephal. p. 52, chlor. Parn. p. 23; Haussk. symb. p. 148; Hal. in ö. b. Z. 1897 p. 286. — Exsicc.: Heldr. herb. norm. n. 698 et 1059; Orph. fl. gr. n. 397.

Caulibus omnibus procumbentibus vel declinatis, non radicantibus; foliis brevissime petiolatis, margine brevissime scabridis; pedunculis ex axillis superioribus solitariis, folio aequilongis vel longioribus; calycis laciniis lineari-lanceolatis, acutis, margine breviter ciliato-scabridis, tubo

corollae duplo brevioribus. — Planta graeca ab austro-hungarica saepius foliis latioribus, minus manifeste ciliato-scabris et calycis laciniis glabris discedit et his notis *V. mixtae* Vel. fl. bulg. p. 380 et 646 accedit.

In fruticetis, umbrosis siccis, circa vineta regionis inferioris et montanae. Thessalia: in oropedio Neuropolis (Haussk.); Aetolia: pr. Mesolongion (Reis.); mt. Parnassus, Cithaeron, Parnes Atticae, pr. Assos Corinthiae (Heldr.); Achaia: mt. Kyllene pr. Flamburitza (Orph.); Arcadia: mt. Parthenion pr. Tripoliza (Friedr.); Messenia: pr. Navarin, Methone, mt. Kupe, insula Sapienza (Chaub.); Laconia: mt. Selitza pr. Kalamata (Zahn), mt. Malevo (Orph.); Zante (Marg.); Cephalonia: pr. Lixuri (Ung.). — Febr. Maio. ♃

LXXV. Ordo. Gentianaceae Juss. gen. p. 141.

Dispositio generum:

1. Tribus. *Gentianeae* Griseb. in DC. pr. IX. p. 38. — Corollae aestivatio contorta; testa membranacea; folia saepius opposita.

a. Stylus 1 vel subnullus, stigmata sessilia, in capsula persistentia.

1. Gentiana L.

b. Stylus conspicuus, deciduus.

α. Calyx 6—8 partitus; corolla hypocrateriformis, limbo 6—8 partito; stamina 6—8.

2. Chlora L.

β. Calyx 5 fidus; coralla infundibuliformis, limbo 5 fido; stamina 5.

3. Erythraea Ren.

γ. Calyx 4 dentatus vel 4 partitus; corolla infundibuliformis, limbo 4 partito; stamina 4.

4. Cicendia Adans.

2. Tribus. *Menyantheae* Endl. gen. p. 605. — Corollae aestivatio induplicativa; testa lignosa; folia alterna.

5. Limnanthemum Gmel.

1. Gentiana L. gen. n. 322.

a. Corolla lutea.

1. **G. symphyandra** Murb. Beitr. fl. Südbosn. p. 89. — Exsicc.: Orph. fl. gr. n. 1001 (Macedonia); Heldr. it. gr. septentr. a. 1879.

Rhizomate longo, cylindrico, ramoso, crasso, multicipite; caule erecto, elato, crasso, simplici; foliis glaucescentibus, infimis ellipticis, in petiolum attenuatis, caulinis mediis et superioribus ovatis, acutis; floribus pedicellatis, fasciculatis, in pseudoverticillos, bracteis cucullatis involucratos dispositis; calyce membranaceo, spathaceo-dimidiato, inaequaliter-dentato; corolla calycem duplo excedente, in lobos 5—7, lanceolato-lineares, acutos profunde partita, fauce plicis destituta; antheris in tubum connatis; stigmatibus erecto-patentibus. — Snbspecies orientalis *G. luteae* L., quae floribus subminoribus longius pedicellatis, antheris liberis et stigmatibus spiraliter revolutis discedit.

In pascuis subalpinis, rarissime, hucusque solum in mt. Oeta Phthiotidis loco Terratza dicto (Heldr.). — Jul. Aug. ♃

b. Corolla azurea.

2. **G. asclepiadea** L. sp. p. 227; Bois. fl. or. IV. p. 76. — Icon: Jacq. fl. austr. IV. t. 328. — Exsicc.: Heldr. it. gr. septentr. a. 1879.

Rhizomate indurato, fibroso, multicipite; caulibus erectis, elatis, simplicibus, foliosis; foliis viridibus, decussatis, a basi ovata vel subcordata lanceolato-acuminatis, radicalibus nullis; floribus 1—3 in axillis foliorum superiorum, subsessilibus, spicam foliosam formantibus; calycis tubulosi, corolla 4 plo brevioris, apice truncati, dentibus linearibus, brevibus; corolla tubuloso-campanulata, lobis ovatis, acutis, plicis brevibus auctis; antheris coalitis; stigmatibus distinctis.

In subalpinis, rarissime. Thessalia: mt. Olympus et in faucibus Ruphias mt. Oeta Phthiotidis (Heldr.). — Jul. Sept. ♃

3. **G. verna** L. sp. p. 228; Bois. fl. or. IV. p. 73; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 9. — *Hippion aestivum* Schm. fl. boem. cent. II. p. 20. — *G. aestiva* R. et Sch. sept. VI. p. 156. — *G. verna v. aestiva* Bald. riv. coll. bot. Alb. 1896 p. 77. — Exsicc.: Orph. herb. n. 3699; Bald. it. alb. epir. IV. n. 185. — Caespitosa; rhizomate tenui, multicipite, ramos breves floriferos et steriles rosuliferos edente; caulibus pumilis, unifloris; foliis sessilibus, radicalibus rosulatis, breviter oblongo-lanceolatis, acutis, caulinis 2—4, multo minoribus; calycis tubulosi, anguste-alati dentibus lanceolatis, acutis, corollae tubo brevioribus; corolla hypocraterimorpha, lobis ovatis, obtusiusculis, plicis bifidis auctis; antheris liberis; stigmatibus contiguis. — Specimina a me visa propter calycem anguste alatam ad speciem Linnaeanam, nec ad *G. tergestinam* Beck spectant.

In herbidis alpinis, rarissime. Epirus: mt. Smolika, Papignon et Gamila (Bald.); Thessalia: mt. Olympus (Orph.). — Jul. Aug. ♃

2. Chlora L. gen. n. 1258.

C. perfoliata L. sp. p. 232 (*Gentiana*); L. syst. ed. 12 II. p. 257; S. et S. pr. I. p. 255; Pieri corc. fl. p. 49; Sieb. avis p. 3, rem. p. 4; Ch. et B. exp. p. 110. Fl. pelop. p. 24; Marg. et R. fl. Zante p. 66; Friedr. Reise p. 254; Clem. sert. p. 66; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 809; Bois. fl. or. IV. p. 66; Spreitz. in z. b. G. 1877 p. 716, 1887 p. 664; Heldr. fl. Cephal. p. 52; Haussk. symb. p. 148; Form. in Ver. Brünn 1895 p. 32; Hal. in ö. b. Z. 1897 p. 97. 1899 p. 25. — *C. intermedia* Ten. syll. p. 565. Haussk. symb. p. 148. — Icon: Rchb. t. 19 f. 1. — Exsicc.: Orph. herb. n. 653.

Glauca; caule erecto, in cymam corymbosam plus minus multifloram abeunti, rarius unifloro; foliis radicalibus rosulatis, subpetiolatis, obovatis, obtusis, caulinis triangulari-ovatis, tota sua latitudine connatis;

calycis, ad basin usque partiti, laciniis subulatis, uninerviis, corolla brevioribus; corollae flavae laciniis oblongis, obtusis.

In humidis, pinguibus regionis inferioris et montanae. Attica: mt. Parnes pr. Tatoi (Heldr.), mt. Pentelicon, pr. Sunium (Haussk.); Achaia: pr. Patras (Hal.); Corinthus (Haussk.); Argolis (Sibth.): pr. Nauplia (Friedr.); Sporadum insula Jura (Leon.); Euboea: pr. Steni (Orph.); Cycladum insula Naxos (Reis.); Creta: pr. Canea (Weiss), Suda, Ipos, Acroteri, promontorium Meleka, inter Francocastello et Rhodakino (Raul.); in Strophadum insula majore (Reis.); Zante (Marg.); Cephalonia: pr. Pessada, Lordata, Piscardo (Heldr.); Corcyra (Pieri): pr. Alipu, Kanali, Kontokali (Form.), Benizze, Palaeokastrizza (Spreitz.). — Maio, Jul. ⊙

2. **C. serotina** Koch in Rchb. icon. III. p. 6 f. 351; Marg. et R. fl. Zante p. 66; Bois. fl. or. IV. p. 66; Haussk. symb. p. 148. — Huc forsan: *C. imperfoliata* Ch. et B. fl. pelop. p. 25; an L. fil. suppl. p. 218, quae calycis laciniis ad quartam partem connatis, foliis inferioribus non connatis discedit et in Graecia vix crescit. — Exsicc.: Heldr. pl. fl. hellen. a. 1897; Baen. herb. europ. n. 9241.

Differt a praecedente foliis ovatis vel ovato-lanceolatis, basi subangustiore ovata connatis, calycis profunde partiti laciniis lanceolato-subulatis, subtrinerviis, corollam subaequantibus, corollae laciniis acutiusculis.

In maritimis, pinetis regionis inferioris, pr. Pharsalum Thessaliae; Nauplia Argolidis (Haussk.); Messene (Chaub.); pr. Kunupeli Elidis (Heldr.); Zante (Marg.); Corcyra: pr. Kanone (Baen.); et forsan loca nonnulla ad praecedentem enumerata, quacum facile commutatur. — Maio, Jul. ⊙

3. Erythraea Ren.

a. Corolla rosea (rarissime alba).

α. Flores in paniculam plus minus fastigiatam dispositi; stylus indivisus, stigmate bilobo.

× Folia ima rosulata; corolla majuscula.

1. **E. centaurium** L. sp. p. 229 excl. var.; Pieri corc. fl. p. 35; Dallap. prosp. p. 36; (*Gentiana*); Pers. syn. I. p. 283; Ch. et B. exp. p. 75, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 66; Fraas fl. class. p. 160; Heldr. Nutzpfl. p. 32, Fl. cephal. p. 52, chlor. Parn. p. 23; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 809; Bois. fl. or. IV. p. 68; Spreitz. in z. b. G. 1887 p. 664; Form. in D. bot. Monat. 1891 p. 24, in Ver. Brünn 1895 p. 32, 1896 p. 59, 1897 p. 43; Hal. Beitr. fl. Epir. p. 31, Beitr. fl. Aetol. p. 8, in ö. b. Z. 1896 p. 15, 1897 p. 97, 1899 p. 25; Haussk. symb. p. 148; Bald. riv. coll. bot. Alb. 1896 p. 77. — *Chironia centaurium* Curt. fl. lond. IV. t. 22; S. et S. pr. I. p. 156; Sieb. avis rem. p. 3. — Exsicc.: Heldr. it. thessal. n. 1084 b.

Caule erecto, quadrangulari, saepe inde a basi opposite ramoso; foliis integris, radicalibus rosulatis, obovato vel spathulato-oblongis, ob-

tusis, in petiolum brevem attenuatis, caulinis sessilibus, ovato-oblongis vel lanceolatis, acutis; floribus sessilibus, ad apicem ramorum fasciculatis, corymbum saepius compactum formantibus; corollae tubo sub anthesi calyce subduplo longiore lobis ellipticis, obtusis vel acutiusculis.

β. **grandiflora** Pers. syn. I. p. 283. — *E. grandiflora* Biv. ap. Roem. et Sch. syst. IV. p. 167; Raul. cret. p. 809. — Major, limbo majusculo, lobis fere duplo majoribus, obtusis, 6—7 mm. longis. — Intermediis ad typum transit. — Exsicc.: Rev. pl. cret. a. 1883.

In collibus dumosis, silvaticis, pascuis praesertim montanis passim per totam Graeciam, in insulis Jonicis, in Creta et Sporadibus; non indicatur in Cycladibus, sed in iis probabiliter etiam provenit; — *β.* Creta: pr. Canea (Rev.), Enneachoria, inter Kastelliana et Pyrgos in Messara, pr. Potamies (Raul.). — Maio, Jul. ⊙.

×× Folia ima non rosulata; corolla parva.

2. **E. tenuiflora** Hoffm. et Link fl. port. I. p. 354 t. 67; Ung. Reise p. 126; Heldr. Nutzpfl. p. 32. — *E. latifolia* Bois. fl. or. IV. p. 67; Heldr. fl. cephal. p. 53, Fl. Aegina p. 379; Haussk. symb. p. 148; Form. in Ver. Brünn 1895 p. 32; Hal. in ö. b. Z. 1897 p. 97; vix Sm. engl. bot. I. p. 321. — *E. latifolia v. tenuiflora* Griseb. spic. II. p. 62; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 809. — Exsicc.: Orph. fl. gr. n. 198.

Caule erecto, quadrangulari, superne stricte ramoso, cymis densis; foliis oblongis, obtusis, sessilibus, saepius approximatis et internodium subaequantibus; floribus subsessilibus, ad apicem ramorum fasciculatis, corymbum saepe compactum formantibus; corollae tubo tenuissimo, sub anthesi calycem parum superante, lobis lanceolatis, acutiusculis; capsula calyce aequilongo vel parum longiore. — Specimina typica cymis densis et foliis approximatis a sequente facile diagnoscuntur, attamen occurrunt formae, cymis laxioribus, quae saltem in herbariis dubiae manent.

In pascuis regionis inferioris et montanae, in maritimis ut videtur per totam Graeciam. — Apr. Jul. ⊙.

3. **E. pulchella** Sw. in vet. acad. handl. 1783 p. 85 t. 3 (*Gentiana*); Fr. nov. fl. suec. p. 30; Ch. et B. fl pelop. p. 16; Weiss in z. b. G. 1869 p. 744; Hal. in ö. b. Z. 1897 p. 97; Form. in Ver. Brünn 1897 p. 43; Heldr. fl. Aegina p. 379, in ö. b. Z. 1898 p. 184. — *E. ramosissima* Pers. syn. I. p. 283; Heldr. Nutzpfl. p. 32; Raul. cret. p. 808; Bois. fl. or. IV. p. 67; Spreitz. in z. b. G. 1887 p. 664; Haussk. symb. p. 148; Bald. riv. coll. bot. Alb. 1896 p. 77. — Exsicc.: Sint. it. thessal. n. 985; Dörfl. pl. cret. n. 91.

Caule erecto, quadrangulari, a basi vel superne ramosissimo, laxe cymoso-fastigiato; foliis oblongis vel oblongo-linearibus, acutiusculis, sessilibus, saepius remotis et internodio multo brevioribus; floribus sessilibus et pedicellatis, solitariis, cymam laxam subfastigiatam formantibus; corollae tubo tenui, sub anthesi calyce sublongiore, lobis lanceolatis, acutiusculis; capsula calyce longiore.

In locis similibus, quibus praecedens, per totam ut videtur Graeciam. — Apr. Jul. ⊙.

β. Flores in spicas longas, laxiusculas, simplices vel bifidas dispositi; stylus indivisus, stigmate infundibuliformi.

4. **E. spicata** L. sp. p. 230 (*Gentiana*); Pers. syn. I. p. 283; Ch. et B. exp. p. 76, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 67; Friedr. Reise p. 273; Heldr. Nutzpfl. p. 32, Fl. cephal. p. 52, in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 247; Raul. cret. p. 809; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 69; Hal. in ö. b. Z. 1890 p. 39, 1896 p. 15; Haussk. symb. p. 149; Form. in Ver. Brünn 1895 p. 32. — *Chironia spicata* Willd. sp. I. p. 1069; S. et S. pr. I. p. 156, Fl. gr. III. p. 32 t. 238; Sieb. avis rem. p. 3. — Exsicc.: Orph. fl. gr. n. 192; Sint. it. or. a. 1889 n. 1923; Dörfl. fl. aeg. n. 248.

Caule erecto, quadrangulari, superne stricte ramoso; foliis oblongis lanceolatisve, sessilibus, superioribus acutis; floribus subsessilibus; corollae tubo, calyci aequilongo, lobis lanceolatis, obtusiusculis.

In uliginosis, salsuginosis, graminosis regionis inferioris et submontanae. Epirus: pr. Prevesa, Agiotoma (Form.); Thessalia: pr. Trikala, Larissa (Haussk.), Caterina (Sint.); Aetolia (Nied.); Doris: ad radices mt. Kiona (Reis.); Attica: pr. Athenas (Sibth.), Phaleron, Podoniphti (Heldr.), Sunium (Chaub.); Argolis: pr. Argos (Chaub.), Poros (Friedr.); Laconia: ad fl. Eurotas (Chaub.); Cycladum insula: Syra (Weiss), Naxos (Heldr.), Rhenea (Leon.); Creta (Sieb.): pr. Suda (Rev.); Zante (Sibth.); Cephalonia: pr. Kutavo, Drapano, Same (Heldr.); Corcyra: pr. Kanali (Form.). — Maio, Sept. ⊙.

b. Corolla flava; cyma dichotoma; stylus bicruris.

5. **E. maritima** L. mant. I. p. 55 (*Gentiana*); Pers. syn. I. p. 383; Sieb. avis p. 3; Ch. et B. exp. p. 76, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 67; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 809; Bois. fl. or. IV. p. 68; Hal. Beitr. fl. Aetol. p. 8. — *Chironia maritima* Willd. sp. I. p. 1069; S. et S. pr. I. p. 156, Fl. gr. III. p. 31 t. 237. — Exsicc.: Dörfl. pl. cret. n. 54.

Caule erecto, quadrangulari, ramoso vel simplici et unifloro; foliis sessilibus, inferioribus ellipticis ovatisve, superioribus ovato-lanceolatis lanceolatisve; cyma pauciflora, laxa; floribus pedunculatis; corollae tubo calyci subbreviore, lobis ovatis, acutiusculis; capsula calyce longiore.

In arenosis maritimis, herbidis regionis inferioris et montanae, rarior. Aetolia (Nied.): pr. Antirrhion (Hal.); Attica: mt. Pentelicon (Heldr.); Elis: pr. Kunupeli, Lintzi (Heldr.); Messenia (Sibth.): inter Arcadia et Pylos, pr. Methone (Chaub.); Zante (Sibth.); Creta (Sieb.): pr. Canea (Weiss), Nerokuru, Suda, Francocastello, Melidoni, inter Rukaka et Istakria (Raul.), pr. Males distr. Hierapetra (Leon.) — Apr. Jun. ⊙.

Obs. Species dubia: *E. virescens* Willd. in Roem. et Schult. syst. veg. IV. p. 787, = *Centaurium minus* Sieb. Reise p. 297. — Creta (Sieb.). —

4. Cicendia Adans. fam. II. p. 503.

1. **C. filiformis** L. sp. p. 231 (*Gentiana*); Delarb. fl. arv. I. p. 20. — *Exacum filiforme* S. et S. pr. I. p. 98; Ch. et B. fl. pelop. p. 10; Mazz. in ant. ion. IV. p. 946. — Icon: Fl. dan. t. 324.

Pusilla; caule erecto, simplici et unifloro vel a basi dichotome plurifloro; foliis imis 4—6 approximatis, oblongo-lanceolatis, caulinis paucis, linearibus; calycis dentibus triangulari-lanceolatis; corolla flava, calyce duplo longiore; capsula uniloculari.

In pratis humidis. Elis (Sibth.); Corcyra: ad lacum Stavropotamo (Mazz.); a recentioribus ut videtur non lecta. — Apr. Jul. ⊙.

5. Limnanthemum Gmel. act. petrop. 1769 XV. p. 567.

1. **L. nymphoides** L. sp. p. 145; S. et S. pr. I. p. 128; (*Menyanthes*); Hoffm. et Link fl. port. I. p. 344; Bois. fl. or. IV. p. 65; Form. in Ver. Brünn 1896 p. 60. — *Villarsia nymphoides* Vent. choix pl. p. 9. — Icon: Fl. dan. t. 339. —

Caule elongato, submerso, radicante; foliis suboppositis, petiolatis, natantibus, cordato-suborbiculatis; floribus in axillis foliorum cymoso fasciculatis, pedunculatis, natantibus; calycis laciniis lanceolatis; corollae flavae laciniis obovatis, breviter fimbriolatis; capsula ovata, acuminata.

In aquis Graeciae (Sibth.), in lacu Nezeros ad Olympum (Form.). — Jul. Aug. ♃. N. v.

Obs. Ex ordine *Sesamaceae* DC. colitur et hinc inde subspontaneum: *S. orientale* L. sp. p. 634, Dall. prosp. p. 87; Fraas fl. class. p. 187; Clem. sert. p. 67; Heldr. Nutzpfl. p. 38; = *S. indicum v. subindivisum* DC. pr. IX. p. 250; Raul. cret. p. 809. — Dein: *S. indicum* L. l. c.; Sieb. avis p. 4; Urv. enum. p. 76; Ch. et B. exp. p. 178, Fl. pelop. p. 40; Bois. fl. or. IV. p. 81; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 7, chlor. Thera p. 29; Bald. viagg. Creta p. 77, —

LXXVI Ordo Cyrtandraceae Jack. in trans. Linn. soc. XIV. p. 23.

1. Ramonda Pers. syn. I. p. 216.

1. **R. Nathaliae** Panc. et Petr. in fl. agri Nyss. p. 574, in Glasnik 1885 p. 121 t. 2. — Exsicc.: Bald. it. alb. epir.

Rhizomate fibroso, collo rufo-lanato; foliis rosulatis, obovatis vel obovato-orbiculatis, grosse crenatis, in petiolum attenuatis, rugosis, supra virentibus, griseo-hirsutis, subtus dense ferrugineo-lanatis; scapis glanduloso-hirtis, 1—3 floris, ebracteatis, pedicellis calyce longioribus; calycis glandulosi laciniis ovatis, obtusis, corollae tubo longioribus; corolla violacea, rotata, fauce aurantiaco-villosa, fere ad basin in lobos 4, rarissime 5 subaequales, obovato-orbiculatos partita; staminibus 4, filamentis

rectis, anthera obtusa brevioribus; capsula glanduloso-pubescente, calyce duplo longiore. — *R. pyrenaicae* Pers. floribus pentameris et antheris acutis praesertim diversae habitu similis. Species tertia generis *R. serbica* Panc., floribus minoribus, pentameris corolla subcampanulata et filamentis longioribus a nostra discedit.

In fissuris rupium umbrosarum. Epirus: mt. Micikeli distr. Zagorion et mt. Konitza supra Gorica (Bald.). — Maio, Jul. ♃.

2. **R. Heldreichii** Bois. diagn. ser. 2 III. p. 141 (*Haberlea*); Benth. et Hook. gen. II. p. 1024; Fritsch in Engl. nat. Pflanzenfam. IV. 3b p. 144. — *Jankaea Heldreichii* Bois. dec. prima p. 4, Fl. or. IV. p. 82. — Icon: Clarke cyrtand. mon. V. t. 18. — Exsicc.: Orph. fl. gr. n. 625; Sint. it. or. a. 1889 n. 1855.

Rhizomate fibroso, collo rufo-lanato; foliis rosulatis, ovatis ellipticisve, integris, basi breviter attenuatis, supra adpresse albo-sericeis, subtus ferrugineo-lanatis; scapis glanduloso-hirtis, 1—3 floris, ebracteatis; pedicellis calyce longioribus; calycis glandulosi laciniis ovatis, obtusis, corollae tubo brevioribus; corolla violacea, campanulata, fauce non aurantiaco-villosa, ad medium in lobos 4, subaequales, obovatos partita; staminibus 4, filamentis rectis, anthera obtusa aequilongis; capsula glanduloso-pubescente, calyce duplo longiore. — Species pulcherrima, foliis integris albo-sericeis, corolla campanulata a praecedenti diversissima.

In rupibus regionis mediae mt. Olympus supra Hagios Dionysios, loco unico (Heldr.). — Jun. Aug. ♃.

LXXVII. Ordo. Polemoniaceae Vent. tabl. II. p. 398.

1. Polemonium L. gen. n. 217.

1. **P. coeruleum** L. sp. p. 162; Form. in Ver. Brünn 1896 p. 61. — Icon: Fl. dan. t. 255. —

Glabrum vel superne glanduloso-puberulum; caule elato; foliis pinnatis, segmentis ovato-lanceolatis lanceolatisve, integris; panicula elongata; calycis laciniis ovatis, acutis; corolla coerulea, raro alba, calyce triplo longiore; antheris ovatis, aurantiacis.

In montosis, rarissime. Thessalia: pr. Patsios in mt. Olympus (Form.). — Jun. Jul. ♃. N. v.

LXXVIII. Ordo. Convolvulaceae Vent. tabl. II. p. 394.

1. Ipomaea L. gen. n. 216. —

Corolla praefloratione contorta; stamina inclusa; stylus unicus, stigmate capitato; capsula 2—5 valvis.

1. **I. sagittata** Poir. voy. barb. II. p. 122. — Icon: Cav. ic. pl. II. t. 107. —

Glabra; caulibus scandentibus; foliis petiolatis, integris, inferioribus cordato-suborbiculatis, ceteris sagittatis, sensim angustioribus, lanceolatis

et lineari-lanceolatis, acutis, auriculis saepe longissimis; pedunculis axillaribus, folio brevioribus, 1—2 floris; sepalis ellipticis, obtusis; corolla purpurea; capsula globosa.

In paludosis ad Actium Acarnaniae (Bald.). — Maio, Aug. ♃. N. v.

Obs. *Batatas litoralis* L. sp. ed. 2 p. 227; Pieri corc. fl. p. 29 (*Convolvulus*); Chois. conv. rar. p. 124. — In Corcyra ex confusione indicatur. — *B. edulis* Chois. conv. or. p. 53; Heldr. Nutzpfl. p. 35. — Colitur ob tubera edulia in insulis Jonicis, pr. Pyrgos, Athenas et in insula Naxos (Heldr.). — Exsicc.: Heldr. a. 1855.

2. Convolvulus L. gen. n. 215. —

Corolla praefloratione contorta; stigmata inclusa; stylus unicus, stigmatibus 2, distinctis; capsula 2—4 valvis.

1. Sectio. *Calystegia* R. Br. pr. fl. nov. holl. p. 483. — Calyx bracteis 2—4, foliaceis involucratus; stigmata ovata, planiuscula; capsula subunilocularis.

a. Caulis scandens; corolla alba.

1. **C. sepium** L. sp. p. 153; S. et S. pr. I. p. 123; Pieri corc. fl. p. 28; Dall. prosp p. 22; Mazz. in ant. ion. V. p. 200; Fraas fl. class. p. 170. — *Calystegia sepium* R. Br. p. 483; Marg. et R. fl. Zante p. 67; Raul. cret. p. 811; Bois. fl. or. IV. p. 111; Heldr. fl. cephal. p. 52; Haussk. symb. p. 149; Form. in Ver. Brünn 1897 p. 44. — Icon: Fl. dan. t. 458. — Exsicc.: Heldr. pl. fl. hellen. a. 1879, Fl. thessala a. 1883.

Glaber; caulibus proceris, volubilibus; foliis petiolatis, ovatis vel oblongis, acutis vel acuminatis, basi cordato-sagittatis, auriculis integris vel angulatis; pedunculis longis, unifloris; bracteis ovatis ovalibusque, basi subcordatis, planiusculis, apice attenuatis.

Ad sepes, fossas. Thessalia: pr. Trikala (Form.), Ormanmagula (Haussk.), Paxedes pr. Volo (Heldr.); Boeotia (Fraas); Attica: ad Cephissum, Phaleron (Fraas); Creta: pr. Candia (Heldr.), Platania (Raul.); Zante (Marg.); Cephalonia (Dallap.); Corcyra (Pieri): pr. Garizza (Mazz.); et certe alibi. — Maio, Sept. ♃.

2. **C. silvaticus** W. et K. pl. rar. hung. III. p. 390 t. 261; Ung. Reise p. 128; Hal. Beitr. fl. Epir. p. 32. — *C. silvestris* Willd. en. I. p. 202. — *Calystegia silvestris* R. et Schult. syst. IV, p. 183; Form. in Ver. Brünn 1895 p. 33, 1896 p. 61, 1897 p. 44. — *Calystegia silvatica* Chois. in DC. pr. IX. p. 433; Raul. cret. p. 811; Bois. fl. or. IV. p. 111; Gelmi in bull. soc. bot. it. 1889 p. 450; Form. in D. bot. Monat. 1891 p. 25; Haussk. symb. p. 149. — Exsicc.: Heldr. it. Thessal. IV. a. 1885.

Differt a praecedente bracteis ovatis, basi inflato-ventricosis, apice rotundatis.

Ad sepes, in silvaticis. Epirus: pr. Kalentini ad septentrionem urbis Arta (Hal.), pr. Philippiades, ad lacum Janina, pr. Chartmut,

Han Levka (Form.); Thessalia: ad monasterium Korona (Heldr.), pr. Trikala, Velestinos, Lamia (Form.); Laconia: pr. Kastanitza (Heldr.); Creta: pr. Haghia, Platania, Kaenuriokorio (Raul.); Corcyra (Ung.). — Maio, Sept. ♃

b. Caulis prostratus; corolla purpurea.

3. C. **soldanella** L. sp. p. 159; S. et S. pr. I. p. 136; Ch. et B. exp. p. 69, Fl. pelop. p. 14; Mazz. in ant. ion. V. p. 200; Fraas fl. class. p. 171. — *Calystegia soldanella* R. Br. pr. p. 433; Marg. et R. fl. Zante p. 67; Bois. fl. or. IV. p. 112; Heldr. in ö. b. Z. 1898 p. 184. — Icon: Rchb. t. 140. — Exsicc.: Sint. et Bornm. it. turc. n. 1373.

Glaber vel interdum caulibus pedunculisque pubescens; rhizomate tenui, longissimo; caulibus procumbentibus, 5—25 cm. longis; foliis carnosulis, longe petiolatis, reniformibus, integris vel repandis; pedunculis unifloris; bracteis ovatis, concavis.

In arenosis maritimis. Thessalia: pr. Litochoron (Sint.); Attica: ad Phaleron (Fraas), Pikrodaphni (Heldr.); Messenia: pr. Methone et in Laconia (Chaub.); Cycladum insula Naxos (Chaub.); Zante (Sibth.): ad Tsilivi (Marg.); Corcyra: pr. Sidari, Messongi (Mazz.). — Maio, Jul. ♃

2. Sectio. *Euconvolvulus* Neilr. fl. Niederöst. p. 530. — Calyx nudus; stigmata lineari-cylindrica, saepe revoluta, capsula bilocularis.

a. Caulis non volubilis; folia nec hastata, nec palmatipartita.

α. Perennes.

× Rami vetusti spinescentes; ovarium glabrum.

4. C. **lanatus** Vahl. symb. I. p. 16; S. et S. pr. I. p. 136, Fl. gr. III. p. 1 t. 202; Bois. fl. or. IV. p. 89. — *C. sericeus* Chois. in DC. pr. IX. p. 400; Raul. cret. p. 809; non L. —

Fruticosus, humilis, ramosus; caule erecto, ramis rigidis, erecto-patulis, junioribus tomentosis, vetustis denudatis, spinescentibus; foliis cinerascenti-tomentosis, inferioribus oblongo-vel lineari-spathulatis, in petiolum attenuatis, superioribus sessilibus, breviter lanceolatis; floribus fasciculatis, fasciculis capituliformibus, 5—10floris, sessilibus, ad axillas foliorum superiorum spicas interruptas formantibus; bracteis ovatis; sepalis lanceolatis, hirsutis; corolla albo-rosea, sericea, calyce $2^1/_2$ plo longiore. — Folia inferiora 20—25 mm. longa, 6—10 mm. lata, corolla 15—17 mm. longa.

In collibus siccis Cretae (Sibth.), sine loci specialis indicatione; a recentioribus non lectum. — Maio. ♄ N. v.

5. C. **dorycnium** L. sp. ed. 2 p. 224; S. et S. pr. I. p. 136, Fl. gr. III. p. 1 t. 201; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 70, Fl. pelop. p. 14; Fraas fl. class. p. 171; Clem. sert. p. 67; Raul. cret. p. 810; Bois. fl. or. IV. p. 91; Spreitz. in z. b. G. 1890 p. 297; Haussk. symb. p. 149; Bald. viagg. Creta p. 77; Heldr. fl.

Aegina p. 380, chlor. Thera p. 17; Form. in D. bot. Monat. 1898 p. 78. — Exsicc.: Orph. fl. gr. n. 205; Heldr. herb. norm. n. 438 et 1264, herb. dimorph. n. 36; Rev. pl. cret. n. 109; Dörfl. fl. aeg. n. 44.

Erinaceo-dumosus; caule erecto, ramis rigidis, patule villosis, divaricatim ramosissimis, amplissime paniculatis, ramulis demum spinescentibus; foliis sessilibus, villosis, inferioribus oblongo-spathulatis vel ellipticis, superioribus linearibus; floribus breviter pedicellatis, in cymis terminalibus, trifloris vel abortu saepe unifloris dispositis; bracteis minutis; sepalis obovato-rotundis, sericeis; corolla rosea, extus ad angulos hirsuta, calyce multo longiore. — Caules 40—60 cm. alti, dumos latos, subglobosos, intricatos efformans, corolla 15 mm. longa.

In aridis regionis inferioris. Attica: pr. Athenas (Orph.), ad Phalerum, Piraeum, insula Aegina (Heldr.); pr. Megara (Chaub.); Corinthus (Sibth.); Creta: pr. Kissamos, Platania (Rev.), Perivolia (Bald.), Canea, Sina, Hagia Rumeli, Alikampos, Askyphos, Rethymo, Hierapetra (Raul.); Cycladum insula: Thera (Heldr.), Anaphi (Leon.). — Maio, Jul. ♃

×× Rami inermes; ovarium hirsutum.

○ Folia lanceolata, flores 1—3, cymosi.

. Fruticosus, ramis adultis lignosis, hornotinis herbaceis.

6. **C. oleaefolius** Desf. enc. III. p. 552; Urv. enum. p. 24; Raul. cret. p. 810; Bois. fl. or. IV. p. 93; Haussk. symb. p. 149; Hal. in ö. b. Z. 1896 p. 15; Heldr. fl. Aegina p. 380, chlor. Thera p. 17. — *C. Tournefortii* Sieb. avis. p. 3, rem. p. 3. — *C. lineatus* S. et S. pr. I. p. 135, Fl. gr. II. p. 81 t. 199, non L. — *C. cneorum* L. sp. p. 157 pro p. quoad pl. creticam; Ch. et B. exp. p. 70, Fl. pelop. p. 14. — Exsicc.: Orph. fl. gr. n. 481; Heldr. herb. norm, n. 439 et 1158, herb. dimorph. n. 37; Dörfl. fl. aeg. n. 144.

Adpresse argenteo-sericeus, basi fruticosus; ramis herbaceis, erectis, elongatis, simplicibus vel stricte ramosis; foliis lineari-lanceolatis, angustis, obtusiusculis; floribus terminalibus, cymoso-subcapitatis; pedicellis calyce brevioribus; bracteis subulatis; sepalis lanceolatis, villosis; corolla rosea, ad angulos sericea, calyce 4 plo longiore.

In collibus saxosis, maritimis. Attica: pr. Phaleron, Piraeum, in scopulo Hagios Georgios pr. Salamis, in insula Arpedoni, Aegina et adjacente scopulo Metopi (Heldr.), pr. Megara (Orph.), isthmus Corinthiacus (Chaub.); Euboea (Wild); Laconia (Chaub.); Cycladum insula Makaria pr. Naxos (Leon.), Caloyero (Sibth.), in scopulo St. Istad pr. Cimolos (Urv.), Thera (Heldr.); Creta: ad promontorium Meleka (Raul.). — Maio. Jun. ♄

. . Rami e rhizomate suffrutescente herbacei.

7. **C. lineatus** L. sp. ed. 2 p. 224; Raul. cret. p. 810; Hal. in ö. b. Z. 1899 p. 25. — Icon: Rchb. t. 134. — Exsicc.: Reiser fl. gr. a. 1898.

Caulibus herbaceis, adpresse sericeis, e rhizomate pluricipiti adscendentibus, simplicibus vel parum ramosis, 5—20 cm. altis, interdum brevissimis; foliis lanceolatis vel lineari-lanceolatis, utrinque adpresse sericeis, infimis in petiolum attenuatis, superioribus subsessilibus; cymis 1—4 floris; breviter pedunculatis; floribus breviter pedicellatis; bracteis linearibus; sepalis oblongo-lanceolatis, sericeis; corolla rosea, ad angulos sericea, calyce 4 plo longiore.

In collibus siccis regionis inferioris. In Strophadum insula majore (Reis.); indicatur quoque in Creta, sed probabiliter ex confusione. — Maio, Jul. ♃

8. **C. cantabricus** L. sp. p. 158; S. et S. pr. I. p. 135; Pieri corc. fl. p. 29; Ch. et B. exp. p. 70, Fl. pelop. p. 14; Friedr. Reise p. 284; Raul. cret. p. 810; Bois. fl. or. IV. p. 95; Heldr. fl. cephal. p. 52; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 33, 1896 p. 61, 1897 p. 45; Hal. Beitr. fl. Epir. p. 32, Beitr. fl. Thessal. p. 16; Haussk. symb. p. 149. — Icon: Jacq. fl. austr. III. t. 296. — Exsicc.: Orph. herb. n. 2878; Sint. it. thessal. n. 359.

Caulibus herbaceis, patule rarius adpresse hirsutis, e rhizomate indurato adscendentibus, 15—40 cm. altis; foliis lanceolatis vel lineari-lanceolatis, utrinque hirsutis, inferioribus in petiolum attenuatis, superioribus sessilibus; cymis 1—4 floris, longe pedunculatis, paniculam laxam formantibus; floribus breviter pedicellatis; bracteis lineari-subulatis; sepalis lanceolatis, hirsutis; corolla rosea, extus ad angulos hirsuta, calyce 3 plo longiore. — A praecedenti indumento patulo, caulibus saepius elatis, cymis longe pedunculatis, paniculam effusam formantibus diversissimus.

In collibus siccis, saxosis regionis inferioris et montanae. Epirus: pr. Sadovica, ad monasterium Hagios Paraskevi (Form.), mt. Tsumerka (Hal.); Thessalia: pr. Kastania (Hal.), mt. Baba (Form.), in Pindo dolopico (Haussk.), pr. Kalabaka (Sint.), Pharsalus, Aivali (Heldr.), mt. Chassia, pr. Selicany in mt. Ossa, pr. Plessidi, Portaria, Volo, mt. Othrys (Form.); Attica: mt. Cithaeron (Heldr.), Lycabettus (Orph.), pr. Athenas, Laurion (Haussk.); Achaia: mt. Kyllene (Heldr.); Argolis: pr. Nauplia (Friedr.), Argos (Chaub.); Laconia et Messenia (Chaub.); Creta (Raul.); Cephalonia: pr. Drapano, Rasata (Heldr.); Corcyra (Pieri). — Maio, Jul. ♃

9. **C. radicosus** Heldr. et Sart. in Bois. diagn. ser. 2 III. p. 124, Fl. or. IV. p. 99. — Exsicc.: Heldr. herb. norm. n. 961.

Caulibus herbaceis, adpressiuscule hirsutis, e rhizomate sublignoso pumilis, dense caespitosis, 1—3 cm. longis, saepius brevissimis; foliis linearibus vel lanceolato-spathulatis, subsessilibus, supra glabris, subtus adpresse hirsutis; cymis 1—3 floris, breviter pedunculatis; floribus breviter pedicellatis; bracteis folio conformibus; sepalis ovato-oblongis, hirsutis; corolla rosea, an angulos hirsuta. — Praecedenti valde affinis, cujus formam abbreviatam alpinam forsan sistit.

Achaia: in oropedio Livadi regionis superioris mt. Kyllene, in terra argillosa compacta (Heldr.). — Jun. Jul. ♃.

○○ Folia ovata, minima; flores solitarii.

10. **C. cochlearis** Griseb. spic. II. p. 76; Bois. fl. or. IV. p. 98, suppl. p. 348; Heldr. chlor. Parn. p. 23. — *C. parnassicus* Bois. et Orph. diagn. ser. 2 III. p. 125. — Exsicc.: Orph. fl. gr. n. 480; Heldr. pl. fl. hellen. a. 1876 et 1878.

Dense pulvinatus, adpresse argyreo-sericeus; rhizomate crasso, lignoso, multicipiti; foliis minimis, ovatis, cuneatis, petiolo eis subaequilongo suffultis; floribus solitariis, in caespite subsessilibus; sepalis ovatis, acuminatis; corolla purpurea, ad angulos sericea. — Foliorum limbus 4—5 mm. longus.

In saxosis regionis abietinae, rare. Attica: mt. Parnes (Heldr.), Pateras (Bartholomatos); mt. Parnassus ad Hagios Nicolaos et Carcaria (Orph.). — Jun. Jul. ♃.

b. Annui.

α. Capsula pubescens.

11. **C. tricolor** L. sp. p. 158; Marg. et R. fl. Zante p. 67. — Icon: Rchb. t. 137.

Pubescens; caulibus adscendentibus vel decumbentibus, subsimplicibus; foliis obovatis, oblongis vel oblongo-lanceolatis, inferioribus in petiolum brevem attenuatis, ceteris sessilibus; pedunculis axillaribus, solitariis, unifloris, folium aequantibus vel superantibus, minute bibracteolatis, fructiferis recurvis; sepalis ovatis, hirsutis; corolla calyce 3—4 plo longiore, limbo coeruleo, tubo flavescente, ad angulos pubescente.

In sepibus insulae Zacynthus (Marg.). — Mart. Maio. ⊙ N. v.

β. Capsula glabra.

12. **C. pentapetaloides** L. syst. nat. ed. 12 III. p. 229; S. et S. pr. I. p. 134, Fl. gr. II. p. 80 t. 197; Ch. et B. exp. p. 69, Fl. pelop. p. 14; Bois. fl. or. IV. p. 110; Heldr. fl. Aegina p. 380. —

Pubescens; caulibus adscendentibus vel decumbentibus, subsimplicibus; foliis radicalibus oblongo-spathulatis, in petiolum longe attenuatis, superioribus oblongo-lanceolatis, sessilibus; pedunculis axillaribus, solitariis, unifloris, folio brevioribus, minute bibracteolatis, fructiferis recurvis; sepalis ovatis, glabris; corolla calyce duplo longiore, limbo coeruleo, tubo flavescente, pubescente. — Differt a praecedente, cui habitu similis, praesertim sepalis et capsulis glabris et corolla multo minori.

In rupestribus siccis et graminosis. Messenia usque ad mt. Ithome (Chaub.); Argolis (Sprun.); insula Aegina: pr. Janitzio (Heldr.); Archipelagus (Sibth.). — Apr. Maio. ⊙. N. v.

13. **C. siculus** L. sp. p. 156; S. et S. pr. I. p. 134, Fl. gr. II. p. 79 t. 196; Ch. et B. exp. p. 69, Fl. pelop. p. 14; Friedr. Reise p. 271; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 810; Bois. fl. or. IV. p. 109; Haussk. symb. p. 149; Hal. in ö. b. Z. 1897

p. 97; Heldr. chlor. Mykon. p. 247. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889.

Pubescens; caulibus adscendentibus vel decumbentibus, subsimplicibus; foliis petiolatis, ovatis, basi cordatis vel truncatis; pedunculis axillaribus, solitariis, unifloris, folio brevioribus, bracteis binis calycem aequantibus vel superantibus obsitis, fructiferis recurvis; sepalis ellipticis, hirsutis; corolla calyce vix duplo longiore, limbo coeruleo, tubo flavescente, pubescente. — Foliis subcordato vel truncato-ovatis; omnibus petiolatis et calyce hirsuto a praecedente statim diagnoscitur.

In rupestribus, arenosis siccis regionis inferioris. Attica (Sprun.); Argolis: pr. Nauplia, Tolon (Haussk.), Poros (Friedr.); Messenia: pr. Pylos (Fauché), Methone, Corone (Chaub.); Creta (Sibth.); Cycladum insula Mykonos (Sart.), Cimolos (Heldr.), Syra (Weiss); Sporadum insula Peristeri (Leon.). — Mart. Maio. ⊙.

b. Caulis volubilis; folia hastata vel (superiora palmatipartita).

α. Folia hastata, nunquam palmatipartita.

14. **C. arvensis** L. sp. p. 253; S. et S. pr. I. p. 123; Pieri corc. fl. p. 29; Ch. et B. exp. p. 68, Fl. pelop. p. 14; Marg. et R. fl. Zante p. 67; Mazz. in ant. ion. V. p. 196; Friedr. Reise p. 275; Fraas fl. class. p. 170; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 810; Bois. fl. or. IV. p. 108; Heldr. fl. cephal. p. 52, Fl. Aegina p. 380, chlor. Thera p. 17, chlor. Mykon. p. 247; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 32, 1896 p. 61; Haussk. symb. p. 149. — Icon: Rchb. t. 136.

Glaber vel puberulus; caulibus prostratis vel volubilibus, saepe valde elongatis, ad 1 m. longis; foliis petiolatis, triangulari-ovatis vel oblongis, acutis, obtuse vel acute sagittato-hastatis; pedunculis axillaribus, solitariis, subunifloris, folium superantibus, bibracteolatis; sepalis ovato-oblongis, obtusis; corolla alba vel rosea, glabra, calyce multo longiore; capsula glabra.

β. **Cherleri** Ag. in Roem. et Schult. syst. IV. p. 261 pro sp.; Heldr. fl. Aegina p. 380. — *C. auriculatus* Mazz. in ant. ion. V. p. 196, an Desr. . — *C. arvensis β. linearifolius* Chois. in DC. pr. IX. p. 407. — Foliorum limbus elongatus, late linearis vel linearilanceolatus, auriculis longioribus, divaricatis. — Exsicc.: Heldr. pl. fl. hellen. a. 1897.

In arvis, ad vias, sepes regionis inferioris et montanae per omnem Graeciam. — Maio, Aug. ♃.

15. **C. hirsutus** Stev. in M. a. B. fl. taur. cauc. I. p. 422; Sieb. avis rem. p. 3; Ch. et B. exp. p. 68, Fl. pelop. p. 14; Bois. fl. or. IV. p. 105; Haussk. symb. p. 149; Form. in Ver. Brünn 1897 p. 45. — Icon: M. a. B. cent. rar. I. t. 6. — Exsicc.: Heldr. pl. fl. hellen. a. 1890.

Hirsutus; caulibus decumbentibus vel scandentibus; foliis petiolatis, ovatis ovalibusque, cordato-hastatis, sinuato-subdentatis, acutis vel ob-

tusiusculis, auriculis rotundatis; pedunculis axillaribus, solitariis, 1—3 floris, folio multo longioribus; pedicellis bibracteolatis; sepalis ovato-oblongis, acuminatis; corolla albo-rosea vel flavida, ad angulos pilosa, calyce 4plo longiore; capsula hirsuta. — Species polymorpha, praecedenti omnibus partibus major, a quo insuper indumento, sepalis acuminatis et capsula hirsutissima discedit.

β. **Sibthorpii** Roem. et Schult. syst. IV. p. 285; Mazz. in ant. ion. V. p. 196; pro sp. — *C. sagittifolius* S. et S, pr. I. p. 133, Fl. gr. II. p. 77 t. 193; Haussk. symb. p. 149; non Mich. — Minus hirsutus, foliis a basi sagittato-hastato triangulari-lanceolatis, in apicem angustissimum longe productis, auriculis divaricatis, obtuse bilobulatis, sepalis angustioribus longioribusque. — Intermediis ad typum transit. — Exsicc.: Heldr. it. IV. Thessal. a. 1883.

In agris, collibus herbosis, ad sepes regionis inferioris. Thessalia: ad monasterium Meteora, pr. Pharsalus, Aivali, Persufli (Haussk.), Trikala, in Phthiotide (Heldr.); Boeotia (Sprun.); Euboea: pr. Xerochori (Heldr.); Peloponnesus (Chaub.): pr. Nauplia (Haussk.); Archipelagus (Sibth.); Creta (Sieb.); Corcyra: pr. Ermones, Castrades (Mazz.). — Maio, Aug. ♃

β. Folia inferiora cordato-ovata, superiora palmatipartita.

16. **C. althaeoides** L. sp. p. 156, excl. var. β.; S. et S. pr. I. p. 133, Fl. gr. II. p. 78 t. 194; Urv. enum. p. 23; Mazz. in ant. ion. V. p. 196; Marg. et R. fl. Zante p. 67; Friedr. Reise p. 202; Fraas. fl. class. p. 171; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 742; Bois. fl or. IV. p. 106; Heldr. fl. cephal. p. 52, Fl. Aegina p. 380; Spreitz. in z. b. G. 1887 p. 664; Haussk. symb. p. 149. — *C. alceaefolius* Ch. et B. exp. p. 68, Fl. pelop. p. 14, non Lam. — Exsicc.: Heldr. herb. norm. n. 962; Dörfl. fl. aeg. n. 70.

Villoso-hirsutus, opacus; caulibus diffusis, apice volubilibus; foliis petiolatis, inferioribus cordato-ovatis, obtuse crenatis vel lobatis, superioribus subpedatim palmatifidis partitisve, laciniis lanceolatis vel linearibus, integris vel dentatis, terminali longiore; pedunculis axillaribus, solitariis, uni rarius bifloris, folio multo longioribus; pedicellis bibracteolatis; sepalis ovato-oblongis, obtusis, margine scariosis; corolla rosea, ad angulos parce pilosa, calyce 4—5plo longiore; capsula ellipsoidea, glabra.

β. **italicus** Roem. et Sch. syst. IV. p. 266; Sieb. avis. p. 3; Raul. cret. p. 810; pro sp. — Caules et folia parce et adpresse pubescentia. — Exsicc.: Rev. pl. cret. n. 110.

In herbidis, maritimis, dumosis regionis inferioris. Attica: mt. Hymettus (Fraas), pr. Laurion (Haussk.), Porto Rhaphti, insula Aegina (Heldr.); Petalium insula Megalonisi (Holzm.); Corinthus (Heldr.), Acrocorinthus, Nauplia (Haussk.); Elis: pr. Pyrgos, Arcadia (Friedr.); Messenia: pr. Methone, Corone (Chaub.); Cycladum insula: Syra, Tenos (Weiss), Naxos (Leon.), Melos (Urv.); Creta: pr. Canea (Weiss); Zante (Marg.); Cephalonia (Ung.); Leucas: ad monasterium Hagios Georgios

(Spreitz.); Corcyra: pr. Spartilla (Mazz.); — β. Cycladum insula Keos (Heldr.); Creta: pr. Akroteri (Rev.), Khalepa, Lutro (Raul.). — Apr. Jun. ♃

17. **C. tenuissimus** S. et S. pr. I p. 134, Fl. gr. II. p. 7 t. 195; Sieb. avis p. 3, rem. p. 3; Mazz. in ant. ion. V. p. 196; Marg. et R. fl. Zante p. 67; Friedr. Reise p. 271; Clem. sert. p. 67; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 811; Bois. fl. or. IV. p. 107; Heldr. fl. Cephal. p. 52, Fl. Aegina p. 380, chlor. Thera p. 17; Spreitz in z. b. G. 1877 p. 715; Hal. in z. b. G. 1888 p. 760; Gelmi in bull. soc. bot. ital. 1889 p. 450; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 32, 1896 p. 61, 1897 p. 45; Haussk. symb. p. 149. — *C. althaeoides* β. L. sp. p. 156; Ch. et B. exp. p. 69, Fl. pelop. p. 14. — *C. argyreus* DC. fl. fr. V. p. 423. — *C. althaeoides v. argyreus et pedatus* Chois. in DC. pr. IX. p. 409. — Exsicc.: Orph. fl. gr. n. 482; Sint. it. thessal. n. 131; Dörfl. pl. cret. n. 92.

Adpresse argyreo-sericeus, nitens; caulibus diffusis, apice volubilibus; foliis petiolatis, inferioribus cordato-oblongis, obtuse crenatis vel lobatis, superioribus palmatim vel pedatim partitis, laciniis-linearibus, plerisque integris, terminali longiore; pedunculis axillaribus, solitariis, unifloris, folio multo longioribus; pedicellis bibracteolatis; sepalis ovato-oblongis, obtusis, margine scariosis; corolla rosea, ad angulos parce pilosa, calyce 4—5 plo longiore; capsula subglobosa, glabra. — Indumento nitide sericeo, foliis caulinis angustius et profundius divisis et corolla subminore a praecedente specifice differt.

β. **virescens.** — Foliis viridibus, breviter adpresse pilosis. — A var. β. speciei praecedentis, foliis caulinis tenuiter partitis, corolla minori et capsula ellipsoidea discedit. — Exsicc.: Heldr. pl. fl. hellen. a. 1900.

In collibus siccis, dumosis regionis inferioris et montanae per totam Graeciam; — β. mt. Pentelicon Atticae (Heldr.). — Apr. Jul. ♃

Obs. *C. suffruticosus* Desf. atl. I. p. 175 et *C. farinosus* L. mant. II. p. 203, a S. et S. pr. I. p. 133 et 135 in Graecia probabiliter ex confusione quadam indicantur.

3. Cressa L. gen. n. 179. — Corolla praefloratione imbricata; stigmata exserta; styli 2, stigmatibus capitatis; capsula 2 valvis.

1. **C. cretica** L. sp. p. 223; S. et S. pr. I. p. 171, Fl. gr. III. p. 51 t. 256; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 81, Fl. pelop. p. 17; Fraas fl. class. p. 171; Raul. cret. p. 811; Bois. fl. or. IV. p. 114; Spreitz. in z. b. G. 1887 p. 664; Haussk. symb. p. 149; Form. in Ver. Brünn 1895 p. 32, 1896 p. 61; Heldr. in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 247. — Exsicc.: Orph. fl. gr. n. 479; Dörfl. fl. aeg. n. 257.

Villosula, cinerea; caulibus erectiusculis vel diffusis, a basi ramosissimis; foliis minutis, ovato-lanceolatis, sessilibus, floribus sessilibus, ad ramorum apicem spicato-capitatis; sepalis ovato-lanceolatis, corolla parva, profunde 5 fida, flava, parum brevioribus; capsula ovata, monosperma.

In argillosis salsis, maritimis. Epirus: pr. Prevesa (Form.); Thessalia: ad Karacair pr. Larissa (Form.); Attica: ad Phaleron (Fraas), Pikrodaphni (Heldr.); Peloponnesus: Argolis, in planitie Helos (Chaub.), pr. Kunupeli Elidis (Heldr.); Cycladum insula: Kythnos, Rhenea (Tunt.), Delos, Antiparos, Naxos, Melos (Chaub.); Creta (Sieb.): pr. Hierapetra (Raul.); Corcyra: pr. Potamo (Spreitz.), Alipu, Kanali, Kipuria, Kanoni (Form.). — Jul. Aug. ♃

LXXIX. Ordo. **Cuscutaceae** Endl. gen. pl. p. 655.

1. **Cuscuta** L. gen. n. 170.

1. Sectio. *Monogynella* Engelm. gen. cusc. sp. p. 74. — Stylus unicus; stigmata capitata vel ovata; capsula regulariter circumscissa.

1. **C. monogyna** Vahl symb. II. p. 32; Fraas fl. class. p. 770; Bois. fl. or. IV. p. 121; Haussk. symb. p. 149; Form. in Ver. Brünn 1897 p. 45. — Icon: Fl. gr. t. 257. — Exsicc.: Orph. fl. gr. n. 281; Sint. it. thessal. n. 1328.

Floribus racemoso-spicatis; calycis laciniis ovatis, obtusis; corollae tubo calyce vix longiore, laciniis tubo dimidio brevioribus, ovatis, obtusis, erectis; squamulis hippocrepidiformibus; denticulatis, medio tubo adnatis; antheris cordato-ovatis, paulo sub fauce subsessilibus; stylo stigmati aequilongo, ovario multo breviore. — Floribus demum manifeste pedicellatis et capsula magna (fere magnitudine pisi) insignis.

Ad frutices parasitica. Thessalia: mt. Pelion (Haussk.), Othrys (Form.); Attica: ad Phaleron (Heldr.); Corinthia: pr. Lutraki (Heldr.); Laconia: mt. Malevo (Orph.). — Jun. Aug. ⊙

2. Sectio. *Eucuscuta* Engelm. l. c. p. 10. — Styli duo, stigmata filiformia; capsula regulariter circumscissa.

a. Styli ovario breviores.

2. **C. europaea** L. sp. p. 124; Friedr. Reise p. 275; Bois. fl. or. IV. p. 117; Haussk. symb. p. 149. — Icon: Fl. dan. t. 199. — Exsicc.: Sint. it. thessal. n. 1428.

Floribus sessilibus, globoso-capitatis, calycis laciniis ovatis, obtusis; corollae tubo calyce sublongiore, laciniis tubo brevioribus, ovatis, obtusis, erecto-patulis, squamulis parvis, tenuissime bifidis vel truncatis, tubo adpressis; stylis stigmatibus et ovario multo brevioribus.

Ad frutices et herbas parasitica. Thessalia: pr. Kriovrisi in valle Penei (Sint.), ad monasterium Korona, pr. Palaeokastron, Karditza (Haussk.),

Pharsalus (Heldr.); Peloponnesus (Bois.): insula Poros (Friedr.) — Jul. Aug. ⊙

b. Styli ovario longiores.

α. Flores pentameri.

3. **C. epithymum** L. sp. p. 124; S. et S. pr. I. p. 173; Sieb. avis p. 3, rem. p. 3; Dallap. prosp. p. 19; Fraas fl. class. p. 170; Bois. fl. or. IV. p. 115; Heldr. fl. cephal. p. 52; Hal. Beitr. fl. Epir. p. 32. — *C. minor* Ch. et B. exp. p. 58, Fl. pelop. p. 11; Marg. et R. fl. Zante p. 67; Ung. Reise p. 128; Raul. cret. p. 811. — Exsicc.: N. v.

Floribus sessilibus, globoso-capitatis; calycis laciniis, ovatis, acutis, patulis; corollae tubo calycem superante, limbi tubo brevioris laciniis ovatis, acutis, patulis, demum reflexis; squamulis majusculis, conniventibus, fimbriatis.

β. **macranthera** Heldr. et Sart. in Bois. diag. ser. 2 III. p. 126; Haussk. symb. p. 149; pro sp; Engelm. l. c. p. 11; Bois. fl. or. IV. p. 115. — *C. Calliopes* (sphalmate *Cassiopes*) Heldr. et Sart. in Bois. diagn. ser. 2 III. p. 128; Heldr. chlor. Parn. p. 123. — *C. epithymum γ. Calliopes* Bois. fl. or. IV. p. 115. — Flores brevissime pedicellati; calyx basi longius obconicus, laciniis acutis vel obtusiusculis; antherae submajores, magis exsertae. — Exsicc.: Orph. herb. n. 743.

γ. **alba** Presl del. Prag. p. 87; Weiss in z. b. G. 1869 p. 742; Haussk. symb. p. 149; pro sp.; Engelm. l. c. p. 13. — *C. epithymum δ. micrantha* Bois. fl. or. IV. p. 116. — Flores minores; calyx corollae tubum subaequans, ejus et corollae lobi acutiores. — Exsicc.: Heldr. herb. norm. n. 237.

Ad herbas diversas parasitica. Indicatur in Graecia (Bois.) et ejus insulis (Sibth.); Attica: mt. Hymettus (Ung.); Laconia pr. Skardamyle, in Maina (Chaub.); Creta (Sieb.): ad Akroteri (Raul.); Zante (Marg.); Cephalonia (Dallap.): mt. Aenos, pr. Lixuri (Heldr.); — β. Epirus: pr. Kalarrytes ad radices mt. Peristeri (Hal.); Thessalia: ad monasterium Korona (Haussk.); mt. Parnassus et Taygetos (Heldr.); — γ. Thessalia: ad monasterium Korona (Haussk.), mt. Oeta Phthiotidis (Heldr.); Attica: mt. Lycabettus (Haussk.), Hymettus (Heldr.); Elis: in colle castri Chlemutzi (Heldr.); Argolis: peninsula Methana (Haussk.); Cycladam insula Syra et Tenos (Weiss). — Maio. Jul. ⊙.

4. **C. planiflora** Ten. syll. p. 128, Fl. neap. t. 220; Weiss. in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 116. —

Floribus sessilibus, globoso-capitatis; calycis laciniis ovatis, obtusiusculis, patulis; corollae tubo calycem superante, limbi tubo longioris laciniis ovatis, obtusis, breviter acuminatis, patulis, nunquam reflexis; squamulis majusculis, conniventibus, fimbriatis. — Praecedenti proxima et ab ea saepe aegre discernenda; differt corollae subrotatae limbo tubo proportione longiore, valde expanso, lobis obtusioribus, staminibus stylisque magis exsertis.

α. **Tenorii** Engelm. l. c. p. 17; Bois. fl. or. IV. p. 116. — Calycis et corollae lobi turgidi, apice saepe cucullati; flores minores; styli ovario parum longiores, capsulae aequilongi vel breviores. — Exsicc.: Orph. fl. gr. n. 517.

β. **approximata** Babingt. in ann. nat. hist. XIII. p. 253, pro sp.; Engelm. l. c. p. 15; Bois. fl. or. IV. p. 116; Haussk. symb. p. 149; Form. in Ver. Brünn 1897 p. 45. — *C. leucosphaera* Bois. et Heldr. in Heldr. herb. norm. n. 33, Heldr. chlor. Parn. p. 23, in Sitzungsb. acad. Wiss. Berlin 1883 p. 4. — *C. epispartos* Bois. et Orph. in Orph. fl. gr. n. 518. — Calycis laciniae membranaceae, vix carnosae; corollae lobi apice tantum turgidi; flores et capitula majora; styli ovario subduplo longiores. — Intermediis ad typum transit. — Exsicc.: Orph. l. c.; Heldr. l. c., fl. Thessal. n. 4; Sint. et Bornm. it. turc. n. 1374; Sint. it. thessal. n. 1045.

Ad herbas varias parasitica. Thessalia: ad monasterium Korona (Heldr.); Aetolia: mt. Korax (Heldr.); Peloponnesus: mt. Kyllene (Heldr.), Malevo (Orph.); Cycladum insula Syra (Weiss); — β. Thessalia: ad monasterium Korona et Meteora, pr. Kalabaka, Orman Magula, Aivali, Pharsalus (Haussk.), mt. Chassia (Form.), Olympus (Sint.), Pelion (Heldr.); Laconia: mt. Malevo (Orph.). Jun. — Aug. ⊙.

β. Flores plerumque tetrameri, saepe tantum trimeri (flos centralis tantum saepe 5merus).

5. **C. globularis** Bert. fl. ital. VII. p. 625; Haussk. symb. p. 150; Hal. Beitr. fl. Aetol. p. 8; Heldr. fl. Aegina p. 380. — *C. micrantha* Tin. in Guss. syn. II. p. 887, non Chois. — *C. palaestina* Bois. diagn. XI. p. 86, Fl. or. VI. p. 116; Bald. viagg. Creta p. 77. — Exsicc.: Heldr. pl. fl. hellen. a. 1877 et 1878.

Floribus sessilibus, in capitula minima glomeratis; calycis carnosuli laciniis latis, brevibus, carinato-costatis, obtusis; corollae calycem vix excedentis laciniis breviter ovatis, obtusis; squamulis majusculis, late spathulatis, denticulatis.

Ad herbas varias parasitica. Aetolia: mt. Taphiassos (Hal.); Attica: in colle Turcovuni (Haussk.), Lycabettus, in Pharmacusarum insula Megali Kyra, in insula Aegina (Heldr.); Achaia: pr. Patras (Hal.), Trika (Heldr.), mt. Malevo (Orph.); Creta (Sieb.): mt. Hagion Pneuma, Ida (Bald.); Archipelagus (Engelm.). — Apr. Jun. ⊙.

3. Sectio. *Clistogrammica* Engelm. l. c. p. 47. — Styli duo, stigmata capitata; capsula nunquam basi dehiscens, baccata.

6. **C. breviflora** Vis. fl. dalm. II. p. 231. — *C. obtusiflora* Humb. Bonpl. et Kunth nov. gen. III. p. 122 var. *breviflora* Engelm. l. c. p. 47; Bois. fl. or. IV. p. 121. — Exsicc.: Orph. fl. gr. n. 615 (*C. atheniensis*).

Floribus pedicellatis; laxiuscule capitato-glomeratis, pedicellis calyci aequilongis vel longioribus; calycis laciniis ovatis, obtusis; corollae calyce sublongioris, tubo brevi scutellato, lobis ovatis, obtusis, mox reflexis

squamulis minimis, bifidis; stylis crassis, ovario brevioribus; capsula irregulariter rumpente. — Caules intense aurantiaci, flores saepe tetrameri, suaveolentes.

In *Ocimo* et *Solano melongena* parasitans. Graecia (Zucc.): pr. Athenas (Orph.). — Jul. Sept. ⊙.

LXXX. Ordo. **Borraginaceae** Juss. gen. p. 128.

Dispositio tribuum generumque.

1. Tribus. *Heliotropeae* Endl. gen. p. 646. — Stylus terminalis.

1. Heliotropium L.

2. Tribus. *Borragineae* Endl. gen. p. 647. — Stylus a basi centro gynobaseos ortus.

1. Subtribus. *Cerintheae* DC. pr. X. 2. — Nuculae 2, biloculares, areola plana toro insidentes, a gynobasi vel stylo liberae.

2. Cerinthe L.

2. Subtribus. *Anchuseae* Koch. syn. p. 497. — Nuculae 4, (raro abortu 3—2) uniloculares, basi excavata toro insidentes et ibidem saepius annulo tumido instructae, a gynobasi vel stylo liberae.

a. Corolla rotata; filamenta sub apice appendice obtusa extus aucta.

3. Borrago L.

b. Corolla infundibuliformis vel cylindrica; filamenta exappendiculata.

α. Corolla breviter cylindrica, limbo in lacinias lineares, tubo multo longiores partita; filamenta longe exserta.

4. Trachystemon Don.

β. Corolla infundibuliformis, laciniis latis, brevibus; filamenta non exserta.

× Nuculae basi annulo tumido cinctae.

○ Corolla fauce aperta, squamulis minutis, pilosis vel laciniatis obsita.

5. Nonnea Med.

○○ Corolla fauce fornicibus, obtusis, papillosis clausa.

6. Anchusa L.

○○○ Corolla fauce fornicibus 5, subulatis, in conum collocatis clausa.

7. Symphytum L.

×× Nuculae basi planae; corolla fornicibus destituta.

8. Pulmonaria L.

3. Subtribus. *Lithospermeae* Koch syn. p. 500. — Nuculae 4, rarius 2, uniloculares, areola plana toro insidentes, a gynobasi vel stylo liberae.

a. Antherae basi sagittato-bicalcaratae.

9. Onosma L.

b. Antherae ovales vel oblongae.

α. Corollae limbus irregularis, subbilabiatus.

10. Echium L.

β. Corollae limbus regularis.

α. Stamina exserta.

11. Moltkia Lehm.

β. Stamina inclusa.

× Corolla fauce aperta.

3. **H. Bocconi** Guss. ind. sem. hort. Bocc. 1825, pr. fl. sic. I. p. 204; Link in Linnaea IX. p. 577; Bois. fl. or. IV. p. 131; Heldr. fl. cephal. p. 53; Hal. Beitr. fl. Aetol. p. 8. — Icon: Bocc. sic. t. 49. — Exsicc.: Hal. it. gr. sec. a. 1893.

Pilis brevibus adpressis obsitum, viride; caule erecto, saepe a basi ramoso; foliis petiolatis, ovatis, obtusis; spicis solitariis et conjugatis, tandem elongatis; calycis hirsuti, corollae tubo subbrevioris, laciniis linearibus, fructiferis subpatulis; corolla alba, extus hirta, limbo tubo aequilongo; stigmate e stylo brevissimo, glabro, a basi conica longe subulato-attenuato; nuculis glabris, verrucosis. — A praecedente, cui habitu et corollis magnis valde affine, stigmate longe subulato-attenuato, nec breviter conico; a sequentibus corolla magna diversum. — Flores odori.

In ruderatis, cultis, collibus siccis. rare. Acarnania: pr. Kravassaras (Hal.); Aetolia: pr. Mesolongion (Nied.); Peloponnesus (Link); Cephalonia: pr. Hagia Euthymia, Piscardo, in oropedio Omala (Heldr.). — Jun. Aug. ⊙.

β. Corolla parva, limbo 2—3 mm. lato; stigma puberulum.

4. **H. europaeum** L. sp. p. 130; S. et S. pr. I. p. 111; Pieri corc. fl. p. 22; Sieb. avis p. 3; Ch. et B. exp. p. 60, Fl. pelop. p. 12; Mazz. in ant. ion. V. p. 180; Link in Linnaea IX. p. 577; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 263 et 275; Clem. sert. p. 67; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 130; Heldr. fl. cephal. p. 53, Fl. Aegina p. 380, chlor. Thera p. 17; Form. in D. bot. Monat. 1891 p. 24, 1898 p. 78, in Ver. Brünn 1895 p. 32; Haussk. symb. p. 150. — *H. europaeum β. tenuiflorum* Guss. fl. sic. syn. I. p. 212; Bois. fl. or. IV. p. 130; Heldr. fl. cephal. p. 53; Spreitz. in z. b. G. 1890 p. 297; Form. in Ver. Brünn 1896 p. 60, 1897 p. 44; (f. indumento magis canescenti, corolla minore lobis erectiusculis, stigmate subbreviore, quae notae autem mihi valde inconstantes videntur et saltem in herbario saepe vix percipiendae sunt). — Exsicc.: Heldr. herb. norm. n. 863a et b, it. Thessal. n. 89; Sint. et Bornm. it. turc. n. 1400, 1402 et 1904; Sint. it. thessal. n. 545.

Pilis brevibus adpressis obsitum, viride vel subcanescens; caule erecto, saepe a basi ramoso; foliis petiolatis, ovatis vel ellipticis, obtusis, basi saepius attenuatis; spicis solitariis et conjugatis, tandem elongatis; calycis hirsuti, corollae tubo aequilongi, laciniis linearibus, fructiferis subpatulis; corolla alba, extus hirta, limbo tubo subbreviore; stigmate e stylo brevi elongato-conico, basi tenuissime puberulo ceterum glabro; nuculis glabris vel parce hispidulis, scrobiculato-rugosis. Flores inodori.

In ruderatis, cultis, arvis regionis inferioris et montanae. Epirus: pr. Prevesa, Ephemiades, Han Mescho, Janina, Hag. Paraskevi, mt. Micikeli (Form.); Thessalia: mt. Baba, pr. Makrichori, Kalabaka, Trikala, Karditza Velestinos, Larissa, pr. Kalyvia, Volo (Form.), Aivali, Pharsalus (Haussk.), mt. Ossa (Heldr.), pr. Litochori, Caterine (Sint.); Euboea: pr. Chalkis (Heldr.); Attica: ubique in agro Atheniensi, mt.

Pentelicon, insula Aegina (Heldr.); Achaia: pr. Patras (Link); pr. Carithena Arcadiae, pr. Chimova in Maina (Chaub.); Argolis: pr. Poros (Friedr.); Cyclades (Sibth.): insula Syra (Weiss), Jos, Thera (Heldr.); Creta (Sieb.): pr. Platania (Rev.), Hagia Rumeli (Spreitz.); Zante (Marg.); Cephalonia: pr. Drapano, Same, Pyrgi, Piscardo, Hagia Euthymia (Heldr.); Corcyra (Pieri): pr. urbem (Friedr.). — Jun. Sept. ⊙.

5. **H. dolosum** Notar. rep. fl. lig. p. 319; Haussk. symb. p. 150; Heldr. in ö. b. Z. 1898 p. 184, chlor. Thera p. 17. — *H. macrocarpum* Guss. en. pl. inar. p. 214, t. 7. — *H. Eichwaldi* Bois. fl. or. IV. p. 131, suppl. p. 350; Spreitz. in z. b. G. 1887 p. 664; non Steud. nom. I. p. 744, quod synon. *H. elliptici* Led. — *H. ellipticum* Nym. consp. p. 508, quoad pl. graecam, non Led. — Exsicc.: Heldr. herb. norm. n. 537 et 864; Sint. et Bornm. it. thessal. n. 1401.

Adpresse canescens; caule erecto, saepe a basi ramoso; foliis petiolatis, ovatis, obtusis, basi rotundatis; spicis solitariis et conjugatis; calycis hirsuti, corollae tubo aequilongi, laciniis oblongis, fructiferis fructu applicatis; corolla alba, extus hirta, limbo tubo subbreviore; stigmate e stylo subnullo elongato-conico, hispidulo; nuculis glabris, minutissime verruculosis. — Flores odori, folia saepe minora quam in specie praecedente.

In arenosis maritimis. Thessalia: pr. Litochori (Sint.); Phthiotis: pr. Lamia (Heldr.); Attica: pr. Athenas (Haussk.), ad Phalerum (Heldr.); Doris: pr. Amphissa (Heldr.); Cycladum insula: Syra, Siphnos, Naxos, Thera (Heldr.); Cythaera: pr. Hagios Nicolaos (Spreitz.). — Jul. Sept. ⊙.

2. Sectio. *Piptoclaina* Don gen. syst. gard. IV. p. 364. — Calyx cum fructu deciduus; nuculae in caryopsem unicam concretae.

6. **H. supinum** L. sp. p. 130; S. et S. pr. I. p. 111, Fl. gr. II; p. 48 t. 157; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 60, Fl. pelop. p. 12; Mazz. in ant. ion. V. p. 180; Fraas fl. class. p. 161. Raul. cret. p. 811; Bois. fl. or. IV. p. 127; Heldr. fl. cephal. p. 53. in Sitzungsb. acad. Wiss. Berl. 1883 p. 8, in ö. b. Z. 1898 p. 184, chlor. Thera p. 17; Form. in D. bot. Monat. 1891 p. 24, in Ver. Brünn 1896 p. 60; Bald. in nuovo giorn. bot. ital. 1894 p. 99; Haussk. symb. p. 150. — Exsicc.: Orph. fl. gr. n. 235; Rev. pl. cret. n. 111; Sint. it. or. a. 1889 n. 1895; Dörfl. fl. aeg. n. 256 et 443.

Cano-tomentellum; caule erecto, a basi ramoso, lateralibus saepe prostratis; foliis petiolatis, ovatis, obtusis; spicis lateralibus et terminalibus, subsolitariis, tandem elongatis; calyce 5 dentato, corollam subaequante, fructifero fructum cingente; corolla minuta, alba, extus hirta, lobis ovatis; stigmate elongato-conico, stylo aequilongo; nucula unica, verrucosa, glabra.

In maritimis, arenosis, ruderatis regionis inferioris. Epirus: pr. Prevesa (Bald.); Thessalia: pr. Karditza, Sophates (Haussk.), ad ripas Penei pr. Larissa (Heldr.), ad Karacair, Velestinos, Volo (Form.), Katerina (Sint.); Attica: ad Phalerum (Orph.); Argolis (Fraas); Laconia: ad Vasilipotamos, Trinissa (Chaub.); Cycladum insula: Tenos, Naxos

(Chaub.), Amorgos (Sibth.), Thera (Heldr.); Creta: pr. Kissamos, Canea (Rev.), Aradhena, Hierapetra (Raul.); Cephalonia: pr. Drapano (Heldr.); Corcyra (Mazz.). — Jun. Nov. ⊙.

2. Tribus. *Borragineae* Endl. gen. p. 647.

1. Subtribus. *Cerintheae* DC. pr. X. p. 2. —

2. Cerinthe L. gen. n. 186.

a. Corolla 5 dentata, dentibus breviter triangularibus, demum reflexis.

1. **C. major** L. sp. p. 136; Pieri corc. fl. p. 25; Mazz. in ant. ion. V. p. 186; Bois. fl. or. IV. p. 149; Heldr. fl. cephal. p. 53. — *C. aspera* Roth cat. I. p. 33; S. et S. pr. I. p. 119, Fl. gr. II. p. 59 t. 170; Sieb. avis p. 3, rem. p. 3; Tausch in Flora XII. p. 641; Ch. et B. exp. p. 66, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 188; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 266; Fraas fl. class. p. 164; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 743; Spreitz. in z. b. G. 1877 p. 715; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 152. — *C. strigosa* Rchb. pl. crit. VIII. f. 984, Fl. exc. p. 340. — Exsicc.: Heldr. herb. norm. n. 431; Baen. herb. europ. n. 9239.

Glauca; caulibus glabris, erectis vet adscendentibus, simplicibus vel ramosis; foliis ellipticis vel obovato-oblongis, tuberculis albis interdum setiferis plus minus adspersis et praesertim apicem versus saepius ciliatis, inferioribus in petiolum attenuatis, superioribus basi cordatis, rotundato-auriculatis; racemis scorpioideis, densis; bracteis cordatis, viridibus, calyces aequantibus vel superantibus; calycis laciniis oblongis; corolla lata, recta, flava, ad staminum insertionem zona purpurea picta, calyce plus duplo longiore; antheris inclusis, filamento subaequilongis; carpellis laevibus. — Corolla 2 cm. longa, 1 cm. lata.

β. **pallida** Guss. pl. rar. p. 85; Raul. cret. p. 812, pro var. *C. asperae*. — Corollis flavis, concoloribus. — Exsicc.: Rev. pl. cret. n. 191.

In cultis, vinetis, inter segetes. Attica: pr. Athenas (Fraas), ad Illyssum (Friedr.); Boeotia (Orph.); Elis: pr. Lechaena (Heldr.); Messenia: pr. Pylos usque Monsusta et mt. Kupe (Chaub.); Cycladum insula: Syra (Weiss), Melos (Leon.); Creta: pr. Kissamos (Rev.), Canea (Weiss), Perivolia (Raul.); Zante (Sibth.); Cephalonia: pr. Palike, Lixuri (Heldr.); Corcyra (Pieri). — Febr. Jun. ⊙.

2. **C. retorta** S. et S. pr. I. p. 120, Fl. gr. II. p. 60 t. 171; Ch. et B. exp. p. 66, Fl. pelop. p. 13; Friedr. Reise p. 283; Bois. fl. or. IV. p. 149; Hal. in ö. b. Z. 1896 p. 15; 1897 p. 286; Haussk. symb. p. 152. — *C. purpurea* Vis. in bot. Zeit. Ergbl. 1829 I. p. 8, Fl. dalm. t. 21; Clem. sert. p. 68. — Exsicc.: Orph. fl. gr. n. 484;

Heldr. herb. norm. n. 327 et 1357, herb. fl. hellen. n. 59; Rev. pl. cret. n. 192; Sint. it. thessal. n. 507; Dörfl. fl. gr. n. 307.

Glauca; caulibus erectis, ramosis, glabris; foliis ovatis, ellipticis vel oblongo-spathulatis, tuberculis albis plus minus obsitis, inferioribus in petiolum attenuatis, superioribus basi cordatis, late rotundato-auriculatis; racemis scorpioideis, densis; bracteis cordatis, violaceis, calyces aequantibus vel superantibus; calycis laciniis oblongis; corolla angusta, ochroleuca, apice subincurva, violacea, calyce duplo longiore, fauce constricta; antheris inclusis, filamento sublongioribus; carpellis laevibus. — Floribus in racemos violaceos, exsiccatione atropurpureos vel cyaneos digestis, insignis. — Corolla 15 mm. longa, 3 mm. lata.

In rupestribus regionis inferioris et montanae, rare. Thessalia: pr. Guvelzi (Sint.); Acarnia: pr. Astacos (Nied.), insula Oxya Echinadum (Reis.); Aetolia: pr. Aetolikon (Reis.); Attica: in faucibus Hymetti (Heldr.); Acrocorinthus (Orph.); Elis: mt. Sekhi pr. Alithura (Friedr.); Messenia: pr. Messene (Chaub.), mt. Ithome (Heldr.); Creta: pr. Malaxa (Rev.); Corcyra: mt. Pantocrator (Clem.). — Febr. Jun. ⊙.

b. Corolla ultra tertiam partem in lacinias lanceolatas erecto-conniventes fissa.

3. **C. minor** L. sp. p. 137; S. et S. pr. I. p. 120; Mazz. in ant. ion. V. p. 186; Bois. fl. or. IV. p. 148; Bald. riv. coll. bot. Alb. 1896 p. 78; Form. in Ver. Brünn 1897 p. 44. — *C. cleiostoma* Bois. et Spr. diagn. IV. p. 44, quae a *C. minori* pedicellis fructiferis erectis, sepalis ovatis auriculato-cordatis et corollae laciniis subulatis, inter se arcte conniventibus (notis meo sensu valde vagis), differre dicitur, sed postea ab auctore ipso Boissier in Fl. or. IV. p. 148 inter synonima *C. minoris* enumeratur. — *C. maculata* Raul. cret. p. 812, vix L. sp. p. 137, quae species dubia et ab auctoribus nunc ad *C. majorem*, nunc ad *C. alpinam* ducitur vel ab aliis pro mera *C. minoris* forma declaratur. — *C. macrophylla* Bois. et Heldr. in Heldr. pl. exs. n. 384; Heldr. herb. norm n. 1457 (f. caule foliisque majoribus DC. pr. X. p. 3). — Icon: Jacq. fl. austr. t. 224. — Exsicc.: Heldr. pl. exs. n. 384, herb. norm. n. 432 et 1457; Orph. fl. gr. n. 191, 1006 et 1157; Sint. it. thessal. n. 634.

Glauca; caulibus glabris, erectis, ramosis; foliis ellipticis vel obovato-oblongis, tuberculis albis plus minus adspersis, inferioribus in petiolum attenuatis, superioribus basi cordatis, rotundato-auriculatis; racemis scorpioideis, fructiferis valde elongatis; bracteis cordatis, viridibus, calyces superantibus; calycis laciniis inaequalibus; corolla angusta, recta, flava, sub limbo interdum maculis purpureis picta (f. *maculata*) calyce sesquilongiore; antheris filamento brevissimo suffultis, corolla subbrevioribus; carpellis laevibus.

In faucibus, rupestribus umbrosis regionis montanae, rare. Epirus: pr. Kastrica distr. Janina (Bald.); Thessalia: in valle Negerli pr. Chaliki (Sint.), mt. Ghavellu (Form.), Olympus (Heldr.); Euboea: mt. Telethrion (Heldr.); Attica: mt. Hymettus (Sprun.); Peloponnesus: mt. Kyllene pr. Hagios

Vlasios, mt. Malevo pr. Platanos (Orph.), mt. Taygetos (Heldr.); Creta: pr. Sphinkia (Heldr.); Corcyra (Mazz.). — Apr. Jul. ⊙ vel ♃.

2. Subtribus. *Anchuseae* Koch. syn. p. 497.

3. Borrago L. gen. n. 182.

1. **B. officinalis** L. sp. p. 137; S. et S. pr. I. p. 122; Pieri corc. fl. p. 26; Dallap. prosp. p. 21; Ch. et B. exp. p. 63, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 188; Marg. et R. fl. Zante p. 69; Ung. Reise p. 128; Heldr. Nutzpfl. p. 34, Fl. cephal. p. 53; Raul. cret. p. 813; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 150; Spreitz. in z. b. G. 1877 p. 715. — Icon: Rchb. t. 101. — Exsicc.: Heldr. pl. fl. hellen. a. 1883.

Patule setosa; caule crasso, succulento, erecto, superne paniculato-corymboso, foliis inferioribus ovatis, petiolatis, floralibus oblongis, sessilibus; calycis laciniis linearibus; corolla azurea, fornicibus albis; antheris exsertis, violaceis; nuculis rugosis.

In ruderatis, cultis regionis inferioris, colitur ad usum culinarem et saepius ut videtur quasi sponte tantum proveniens. Indicatur: Attica: pr. Athenas (Heldr.); Peloponnesus (Sibth.): pr. Astros (Chaub.); Cycladum insula: Syra (Weiss), Tenos, Naxos (Chaub.); Creta: pr. Canea (Raul.); Zante (Marg.); Cephalonia (Dallap.); Ithaca: pr. Vathy (Spreitz.); Corcyra (Pieri). — Mart. Jun. ⊙.

4. Trachystemon Don in Edinb. new. phil. journ. XIII. p. 259.

1. **T. creticum** Willd. sp. I. p. 778; S. et S. pr. I. p. 123, Fl. gr. II. p. 64 t. 176; Sieb. avis p. 3; Ch. et B. exp. p. 63, Fl. pelop. p. 13; Marg. et R. fl. Zante p. 69; (*Borrago*); Don. l. c. — *Psilostemon creticum* DC. pr. X. p. 36; Raul. cret. p. 813; Bois. fl. or. IV. p. 177. — Exsicc.: Heldr. pl. cret. n. 1339.

Puberula breviterque setuloso-hispida; caule erecto vel adscendente, fere a basi ramoso; foliis ovatis vel oblongis, petiolatis, summis diminutis sessilibus; racemis scorpioideis, demum elongatis; calycis laciniis lanceolatis; corollae albidae tubo ima basi intus glanduloso, limbo tubo subtriplo longiore, in lacinias lineares partito, fornicibus linearibus, petaloideis, papillosis, longe exsertis, sed laciniis corollae brevioribus; filamentis squamula ciliata ima basi obsitis, ceterum glabris, rubris, fornices superantibus; nuculis areolato-rugosis.

In rupestribus umbrosis. Euboea (Auch.); Zacynthus (Sibth.); Messenia: inter Arcadia et Gargaliano, insula Sapienza (Chaub.); Creta (Sibth.): ad Acrotiri (Heldr.), Theriso, Almyros, in fauce Katholiko, inter Selia et Hagios Joannes Kaimenos, insula Dia (Raul.). — Mart. Apr. ♃.

5. **Nonnea** Med. phil. bot. I. p. 31.

1. **N. obtusifolia** Willd. sp. I. p. 780 (*Lycopsis*); Roem. et Sch. syst. IV. p. 71; Bois. fl. or. IV. p. 164; Haussk. symb. p. 152. — *Anchusa leiosperma* Fauch. et Ch. in Fl. pelop. p. 12 t. 7. — *N. lamprocarpa* Griseb. spic. II. p. 93, ubi erronee mt. Parnassus pro mt. Parnes ut locus natalis indicatur. — Huc forsan: *Lycopsis vesicaria* Friedr. Reise p. 277, non L. — Exsicc.: Orph. fl. gr. n. 166.

Setulis brevibus, basi tuberculatis aspera, multicaulis; caulibus erectiusculis vel decumbentibus, simplicibus vel subsimplicibus, foliis obovato-oblongis oblongisve, radicalibus basi attenuatis, caulinis sessilibus, breviter decurrentibus; floribus axillaribus, foliis superatis, racemos laxos formantibus; calycis 5 fidi, fructiferi cernui, inflati, laciniis lanceolatis, acutis; corolla coerulescente, calycem vix superante; nuculis oblongis, erectis, basi affixis, acutiusculis, atris, laevigatis, nitidis, parce puberulis.

In rupestribus umbrosis regionis inferioris et submontanae, rare. Attica: ad Cephissum pr. Chelidonu, pr. Haschia (Heldr.) in mt. Parnes (Sprun.), pr. Asteri in mt. Hymettus (Orph.), pr. Kamariza Laurii (Haussk.); Elis: pr. Olympia (Daenzer); in insula Poros, si *L. vesicaria* Friedr. huc pertineat. — Apr. Maio. ⊙.

2. **N. ventricosa** S. et S. pr. I. p. 117, Fl. gr. II. p. 58 t. 168; Ch. et B. fl. pelop. p. 12; Mazz. in ant. ion. V. p 182; (*Anchusa*); Griseb. spic. II. p. 93; Raul. cret. p. 813; Bois. fl. or. IV. p. 169; Haussk. symb. p. 152. — *Lycopsis Sibthorpiana* Roem. et Sch. syst. IV. p. 770; Lehm. asper. p. 258. — *N. Sibthorpiana* Don gen. syst. IV. p. 336. — Exsicc.: Heldr. herb norm. n. 557 et 1062.

Setulis brevibus, basi tuberculatis asperula, multicaulis; caulibus adscendentibus vel decumbentibus, simplicibus vel subsimplicibus; foliis oblongis oblongo-lanceolatisve, radicalibus basi attenuatis, caulinis sessilibus, breviter decurrentibus; racemo terminali, simplici vel bifido, tandem laxiusculo; calycis 5 fidi, fructiferi cernui, subvesiculosi, laciniis lanceolatis, acutis; corolla albida, calycem parum superante; nuculis reniformibus, depressis, ad medium sinus renalis affixis, obtusis, brunneis, areolato-rugosis, opacis, glabris. — Nucularum forma a praecedente statim distinguenda.

In arenosis, vinetis, incultis regionis inferioris, rare. Attica: pr. Haschia in mt. Parnes, ad Phalerum (Heldr.); Corinthia: pr. Asso (Heldr.), Kalamaki (Haussk.); Messenia (Chaub.); Creta: mt. Sphakia (Lehm.); Corcyra: pr. Synarades (Mazz.). — Mart. Maio. ⊙.

Obs. *N. pulla* L. syst. ed. 10 p. 916 (*Lycopsis*); DC. fl. fr. III. p. 626; Raul. cret. p. 813; in Creta ex confusione indicatur.

6. **Anchusa** L. gen. n. 182.

Dispositio specierum:

1. Sectio. *Caryolopha* Fisch et. Mey. ind. tert. hort. petrop. p. 31. — Corollae tubus rectus, limbus regularis; nuculae basi in stipitem areoligeram contractae.

1. A. sempervirens L.

2. Sectio. *Euanchusa* Gr. et Godr. fl. fr. II. p. 512. — Corollae tubus rectus, limbus regularis; nuculae basi in stipitem non contractae.

a. Calyx 5 fidus.

α. Folia integra vel subrepanda, margine non undulato-crispa.

× Calycis laciniae acutae.

2. A. officinalis L. **3. A. parnassica** Bois. et Orph.

×× Calycis laciniae obtusae.

4. A. ochroleuca M. a. B. **5. A. Gmelini** Led.

β. Folia sinuato-dentata vel undulato-crispa.

6. A. hybrida Ten. **7. A. undulata** L. **8. A. aspera** Bois.

b. Calyx 5 partitus.

α. Nuculae rectae.

× Corolla coerulea.

9. A. italica Retz. **10. A. thessala** Bois. et Spr.

×× Corolla flava.

11. A. aegyptiaca (L.).

β. Nuculae in rostrum obliquum abeuntes.

12. A. stylosa M. a. B. **13. A. caespitosa** Lam.

γ. Nuculae horizontaliter curvatae.

14. A. aggregata Lehm.

3. Sectio. *Lycopsis* Griseb. spic. II. p. 95. — Corollae tubus ad medium inflexo-curvatus, limbus subirregularis, obliquus; nuculae basi in stipitem non contractae.

15. A. arvensis L. **16. A. variegata** L.

1. Sectio. *Caryolopha* Fisch. et Mey. ind. tert. hort. petrop. p. 31. — Corollae tubus rectus, limbus regularis; nuculae basi in stipitem areoligeram contractae.

1. **A. sempervirens** L. sp. p. 134. — *Caryolopha sempervirens* Fisch. et Mey. l. c.; DC. pr. X. p. 41. — *A. Adami* Mazz. in ant. ion. V. p. 182, sec. DC. l. c. — Icon: Rchb. t. 105.

Scabrido-pubescens, breviterque setoso-hispidula; caulibus erectis, superne ramosis; foliis ovali-lanceolatis, radicalibus in petiolum longum contractis, persistentibus, superioribus sessilibus, summis diminutis, cordato-ovatis; racemis brevibus, laxifloris; bracteis foliaceis, lanceolatis; pedicellis calyce brevioribus; calycis laciniis lanceolatis; corolla calyce paulo longiore, coerulea; nuculis ovatis, rectis, rostratis, reticulato-rugosis.

In umbrosis humidis. Corcyra: pr. Arcudilla (Mazz.); species pro Graecia dubia, a recentioribus non lecta. — Maio, Jun. ♃. N. v.

2. Sectio. *Euanchusa* Gr. et Godr. fl. fr. II. p. 512. — Corollae tubus rectus, limbus regularis; nuculae basi in stipitem non contractae.

a. Calyx 5 fidus.

α. Folia integra vel subrepanda, margine non undulato-crispa.

× Calycis laciniae acutae.

2. **A. officinalis** L. sp. p. 133; Pieri corc. fl. p. 22; Dallap. prosp. p. 19; Mazz. in ant. ion. V. p. 180; Friedr. Reise p. 279; Bois. fl. or. IV. p. 151 p. p.; Heldr. fl. cephal. p. 53; Form. in D. bot. Mon. 1891 p. 24, in Ver. Brünn 1895 p. 32; Bald. riv. coll. bot. Alb. 1896 p. 78. — *A. angustifolia* L. sp. p. 133; ?Mazz. in ant. ion. V. p. 182. — Icon: Fl. dan. t. 572. — Exsicc: Sint. et Bornm. it. turc. n. 1397; Bald. it. alb. epir. IV. n. 249.

Pilis patulis brevibus e tuberculis ortis hispida; caulibus erectis, saepe elatis, superne paniculatis; foliis oblongo-lanceolatis lanceolatisve, integris vel subrepandis, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; racemis tandem laxifloris; bracteis triangulari-lanceolatis; pedicellis calyce brevioribus; calycis hispidi, fructiferi accreti, laciniis lanceolatis; corollae violaceo-coeruleae tubo calycem subsuperante; fornicibus ovatis, velutinis; nuculis oblique ovatis et rostratis, latere altero tuberculato-rugosis.

β. **macrocalyx** Haussk. in Mitth. thür. bot. Ver. V. 3 p. 71, symb. p. 150; Form. in Ver. Brünn 1897 p. 44; pro sp. — Robusta, tuberculis majoribus obsita; calyce majore, laciniis latioribus; corollae laete cyaneae lobis longioribus; nuculis majoribus, brevissime rostratis, grosse plicato-rugosis. — Exsicc.: Sint. it. thessal. n. 531 et 544.

In herbidis regionis inferioris et montanae. Epirus: pr. Ruvilastro, Han Mescho, Janina, mt. Micikeli, Prosgoli, in valle Dipotami (Form.), mt. Smolika (Bald.); Thessalia: pr. Chaliki et Koturi in Pindo, mt. Ossa (Form.), mt. Olympus (Heldr.); Aetolia: mt. Tymphrestus (Samar.); Euboea: mt. Dirphys (Orph.); Peloponnesus: pr. Lintzi Elidis (Heldr.), mt. Kyllene (Orph.), pr. Carithena Arcadiae (Friedr.), mt. Taygetos ad Hagios Dimitri (Heldr.); Cephalonia (Dallap.); Corcyra (Pieri): pr. Platiterra, Cunicades (Mazz.); sed loca nonnulla, praesertim alpina probabiliter ad speciem sequentem spectant; — *β.* mt. Gionskala et Ghavellu in Pindo (Haussk.), mt. Mavrika et Katasara in mt. Othrys (Form.). — Maio, Aug. ⊙ et ♃.

3. **A. parnassica** Bois. et Orph. diagn. ser. 2 III. p, 134. — *A. officinails* Bois. fl. or. IV. p. 151 p. p., non L.. — Icon: Hal. it. gr. II. a. 1893; Tunt. et Leon. pl. gr. a. 1896.

Pilis albis, patulis, elongatis, plurimis basi non tuberculatis hispida; caulibus erectis, subsimplicibus; foliis lanceolatis lineari-lanceolatisve, integris vel subrepandis, inferibribus in petiolum attenuatis, superioribus semiamplexicaulibus; racemis densis; bracteis lanceolatis; pedicellis calyce brevioribus; calycis dense longeque albo-hispidi, fructiferi accreti, laciniis anguste lineari-lanceolatis; corollae azureae tubo calycem subsuperante; fornicibus oblongis, velutinis; nuculis oblique ovatis et rostratis, latere altero tuberculato-rugosis — Praecedenti affine, sed ab ea habitu minus elato (caulibus interdum 10 cm. tantum altis), bracteis lanceolatis, calycis longioris densissime albo-setosi, profundius fissi, laciniis angustioribus, corolla pulcherrime azurea et nuculis minoribus, specifice meo sensu distincta.

In herbidis lapidosis regionis mediae et superioris. Aetolia: mt. Korax (Tunt.); mt. Parnassus (Orph.); Achaia: mt. Panachaicon (Hal.); et probabiliter loca nonnulla ad praecedentem enumerata. — Jun. Aug. ♃

×× Calycis laciniae obtusae.

4. **A. ochroleuca** M. B. fl. taur. cauc. I. p. 125; Bois. fl. or. IV. p. 153. — *A. angustifolia* S. et S. pr. I. p. 115, Fl. gr. II. p. 55 t. 164; Urv. enum. p. 19; Tausch in Flora 1824 p. 226; Ch. et B. fl. pelop. p. 12; Raul. cret. p. 814. —

Pilis rigidis, tuberculo demum aucto insidentibus, undique hispida; caulibus erectis, ramosis, paniculatis; foliis integerrimis, infimis lineari-oblongis, basi attenuatis, ceteris lanceolatis linearibusve, semiamplexicaulibus; racemis demum laxis; bracteis parvis, oblongis vel ovato-oblongis, calyce brevioribus; pedicellis brevissimis; calycis hispidi, fructiferi inflati, campanulati, laciniis oblongis, obtusis; corollae ochroleucae, albae, roseae vel coerulescentis tubo calyce subduplo longiore; fornicibus ovatis, valde exsertis, pilis crassis dense obtectis; nuculis oblique ovatis et rostratis, latere altero areolato-rugosis, crebre et minute tuberculatis.

In neglectis, collibus argillosis, ad ripas. Archipelagus (Sibth.): insula Melos (Urv.); Creta (Sieb.); sed specimina non vidi, ideoque nescio, an planta graeca re vera ad *A. ochroleucum* M. B. pertineat. — Maio, Jul. ♃ N. v.

5. **A. Gmelini** Led. in Panders beitr. naturk. I. p. 62, Fl. ross. III. p. 118; Haussk. symb. p. 151. — *A. linearifolia* Urv. enum. p. 20. — Huc spectare videtur ex descriptione et icone: *A. obliqua* Vis. in mem. ist. venet. I. p. 13 t. 5; Bois. fl. or. IV. p. 153. —

Caulibus erectis, parce strigoso-hispidis, simplicibus vel ramosis; foliis anguste lanceolatis linearibusve, integerrimis, supra setis crebris, tuberculo insidentibus strigoso-hispidis, subtus setis rarioribus adspersis vel praeter carinam glabris, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; racemis densis; bracteis parvis, ovato-lanceolatis lanceolatisve, calyce brevioribus; pedicellis brevissimis; calycis setosi, fructiferi parum accreti, laciniis lineari-oblongis, obtusis; corollae azureae tubo calyce subduplo longiore; fornicibus ovatis, exsertis, margine dense villosis; nuculis oblique ovatis et rostratis, latere altero areolato-rugosis, tuberculatis. — Foliis angustis praecedenti similis, ab ea foliis subtus parce hispidis, racemis elongatis, calycibus parvis, fructiferis vix accretis et florum colore specifice discedit.

In cretaceis isthmi Corinthiaci pr. Kalamaki (Haussk.); Cycladum insula Tenos (Sart.), si *A. obliqua* Bois. re vera huc pertineat. — Maio, Jul. ♃. N. v.

β. Folia sinuato dentata vel undulato-crispa.

6. **A. hybrida** Ten. fl. nap. I. p. 45 t. 11; Ch. et B. exp. p. 62, Fl. pelop. p. 12; Friedr. Reise p. 267; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 152; Heldr. fl. Aegina p. 380, chlor. Thera p. 17; Form. in D. bot. Monat. 1898 p. 78. — Exsicc.: Heldr.

herb. norm. n. 307 et 1061; Rev. pl. cret. n. 114; Sint. it. thessal. n. 50; Baen. herb. europ. n. 9202.

Plus minus hispida, virescens; caulibus erectis vel adscendentibus, superne paniculatis; foliis oblongis oblongo-lanceolatisve, obtusis, margine sinuato-dentatis, subundulatis, praecipue supra plus minusve tuberculatis, inferioribus in petiolum attenuatis, superioribus sessilibus; racemis tandem laxiusculis; bracteis ovato-lanceolatis lanceolatisve, calyce brevioribus; pedicellis calyce brevioribus; calyce fructifero inflato, saepius ad medium 5 fido, laciniis triangulari-lanceolatis, obtusis; corollae coeruleae tubo calyce subduplo longiore; fornicibus ovatis, velutinis; nuculis oblique ovatis et subrostratis, latere altero crebrius tuberculato-rugosis. — Inter *A. officinalem* et *A. undulatam* quasi intermedia; ab illa foliis undulato-dentatis et calycis laciniis obtusis praesertim distincta.

β. **setigera.** — Tota setis patulis, elongatis, albis obsita. — Intermediis ad typum transit. — Exsicc.: Dörfl. fl. aeg. n. 78, pl. cret. n. 145.

In cultis, vinetis, arvis herbidis regionis inferioris. Corcyra: mt. Pantocrator (Baen.); Thessalia: ad Agria pr. Volo (Sint.); Attica: pr. Athenas (Heldr.), ad Phalerum (Friedr.); Achaia: pr. Patras (Hal.); Messenia: pr. Diabolitzi (Heldr.), Methone (Chaub.); Laconia pr. Marathonisi, Damala (Chaub.); insula Poros (Wied.), Aegina, Tenos, Melos, Thera (Heldr.); Creta: pr. Kissamos (Rev.), Canea (Weiss). — *β.* Cycladum insula Paros et pr. Anatoli Cretae (Leon.). — Mart. Jul. ⊙

7. **A. undulata** L. sp. p. 133; S. et S. pr. I. p. 116, Fl. gr. II p. 56 t. 165; Urv. enum. p. 19; Friedr. Reise p. 267; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 814; Bois. fl. or. IV. p. 152; Spreitz. in z. b. G. 1877 p. 715; Gelmi in bull. soc. bot. ital. 1889 p. 450; Haussk. symb. p. 151; Form. in Ver. Brünn 1896 p. 61; Bald. riv. coll. bot. Alb. 1896 p. 78; Heldr. fl. Aegina p. 380, chlor. Thera p. 17. — Exsicc.: Orph. fl. gr. n. 62; Heldr. herb. norm. n. 308.

Cinerascens, adpressiuscule tomentello-aspera; caulibus erectis vel adscendentibus, superne paniculatis; foliis lineari-oblongis et anguste linearibus, obtusis, margine undulato-crispis, praecipue supra plus minusve tuberculatis, inferioribus in petiolum attenuatis, superioribus saepe breviter decurrentibus; racemis tandem laxiusculis; bracteis ovato-lanceolatis lanceolatisve, calyce brevioribus; pedicellis calyce brevioribus; calyce fructifero inflato, saepius ad tertiam partem 5 fido, laciniis triangulari-lanceolatis, acutis vel acutiusculis; corollae coeruleae tubo calyce subduplo longiore; fornicibus ovatis, velutinis; nuculis oblique ovatis et subrostratis, latere altero crebrius tuberculato-rugosis. — Praecedenti maxime affinis et formis transitoriis, interdum difficillime distinguendis cum ea ut videtur conjuncta; formae typicae attamen foliis linearibus, angustis, manifeste undulato-crispis et calycibus plerumque ad tertiam partem in lacinias acutiores 5 fidis facile diagnoscuntur. — Cf. cetero-

qui Guss. pl. rar. p. 82 et Haussk. symb. p. 151. — Variat uti praecedens:

β. **maritima** Heldr. herb. norm. n. 1265. — Tota setis patulis, elongatis, albis obsita. — Intermediis ad typum transit. — Exsicc.: Heldr. l. c.; Dörfl. fl. aeg. n. 171.

In collibus aridis, arvis, arenosis regionis inferioris et montanae. Corcyra: mt. Pantocrator (Spreitz.); Epirus: mt. Micikeli (Bald.); Thessalia: pr. Klinovo, Kalabaka (Form.), mt. Pelion (Heldr.); Boeotia (Sprun.); Attica: pr. Athenas (Sprun.), Daphni (Pichl.), Phaleron (Friedr.); insula Aegina: pr. Mesagros (Heldr.); Tenos (Weiss), Melos (Sibth.), Thera (Heldr.); — β. in arenosis praesertim maritimis ad Phaleron, Raphina (Heldr.); Cycladum insula Naxos (Reis.). — Loca nonnulla ad speciem praecedentem spectare videntur. — Mart. Oct. ⊙

8. **A. aspera** Bois. in ann. sc. nat. IV. 2 p. 243; Haussk. symb. p. 151. — *A. ochroleuca* β. *canescens* Bois. fl. or. IV. p. 154 p. p., sec. Haussk. l. c. — *A. Sartorii* Heldr. herb. norm. n. 1060. —

Tota setis patulis, albis, elongatis hispidissima, canescens; caulibus e collo plurimis, decumbentibus vel adscendentibus, saepius mox supra basin in racemos elongatos abeuntibus; foliis lanceolatis, obtusis, margine undulato-crispis, utrinque crebre tuberculatis, inferioribus in petiolum attenuatis, superioribus sessilibus; racemis multifloris, densiusculis; bracteis lanceolatis, calyce brevioribus; pedicellis brevissimis; calyce fructifero inflato, ad medium 5 fido, laciniis lanceolatis, acutis; corollae coeruleae tubo calyce duplo longiore; fornicibus ovatis, velutinis; nuculis oblique ovatis et subrostratis, tuberculato-rugosis. — Varietati β. praecedentis affinis, ab ea caulibus prostratis, in racemos densos multifloros mox abeuntibus, calycis laciniis angustioribus acutis, corolla minore, e calyce longius exserta et indumento hispidissimo, diversa. — Variat calycibus parvis — fructiferis 7 mm. longis — et fere duplo majoribus.

In agris argillosis. Attica: ad radices mt. Parnes pr. Varipopi (Heldr.), pr. Athenas (Hal.), ad Phalerum, pr. Kamariza Laurii; ad radices Acrocorinthi (Haussk.); indicatur quoque a Form. in Ver. Brünn 1897 p. 44 pr. Konisko, Jerakari et in valle fl. Longi in Thessalia, sed determinationibus saepe falsis hujus auctoris loca thessala ad interim dubia manent. — Maio, Jul. ⊙

b. Calyx 5partitus.

α. Nuculae rectae.

× Corolla coerulea.

9. **A. italica** Retz obs. I. p. 12; Mazz. in ant. ion. V. p. 182; Fraas fl. class. p. 162; Raul. cret. p. 814; Bois. fl. or. IV. p. 154; Heldr. fl. cephal. p. 53, Fl. Aegina p. 380; Spreitz. in z. b. G. 1887 p. 664; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1896 p. 61. — *A. paniculata* Ait. hort. Kew. I. p. 177; S. et S. pr. I. p. 115, Fl. gr. II. p. 53 t. 163; Ch. et B. exp. p. 62, Fl. pelop. p.

121; Marg. et R. fl. Zante p. 69; Friedr. Reise p. 277. — Exsicc.: Heldr. pl. fl. hellen. a. 1878 et 1896.

Perennis, setis tuberculatis hispida; caule crasso, elato, ramoso, paniculato; foliis oblongis vel lanceolatis, integris vel subrepando-denticulatis, inferioribus in petiolum attenuatis, superioribus sessilibus; racemis saepius conjugatis, demum laxis; bracteis lanceolato-linearibus, pedicello brevioribus; calycis laciniis lineari-acuminatis, demum accretis; corollae azureae tubo calyce non longiore; fornicibus dense penicillatis, exsertis; nuculis angulato-trigonis, elevato-rugosis, inter rugas tenuissime pappulosis.

In arvis, ad vias regionis inferioris et montanae. Thessalia: pr. Trikala, Ambelakia, Baba, Velestino (Form.); Aetolia: mt. Korax (Tunt.); Attica: mt. Parnes, pr. Megara (Fraas), insula Metopi pr. Aegina (Heldr.); Corinthus (Fraas), insula Poros (Friedr.); Creta: pr. Canea, Perama (Raul.); Zante (Marg.); Cephalonia: pr. Drapano, Spilaea, Argostoli (Heldr.); Corcyra: pr. Manducchio (Mazz.). — Maio. Jul. ♃.

10. **A. thessala** Bois. et Sprun. diagn. XI. p. 99, Fl. or. IV. p. 160. — Icon: Rouy ill. pl. eur. rar. t. 267. — Exsicc.: Aznavour fl. constatip. a. 1897 (Makriköy).

Annua, setis tuberculatis hispida; caule gracili, erecto, parce ramoso; foliis integris, vix repandis, inferioribus oblongis, in petiolum attenuatis, superioribus lineari-lanceolatis, sessilibus; racemis saepius conjugatis, demum laxis; bracteis lanceolatis, calyce aequilongis; calycis laciniis linearibus, demum accretis; corollae pallide coeruleae tubo calyce sesquilongiore; fornicibus papillosis, subexsertis; nuculis semioblongis, parce rugosis et tuberculatis. — Habitu et florum magnitudine *A. stylosae* simillima, sed distinctissima stylo calyce fructifero breviore, nec longiore, calycis laciniis fructu duplo longioribus et praesertim nuculis rectis.

Inter segetes in Thessalia pr. Zeitun (Sprun.). — Apr. Jun, ⊙ N. v.

×× Corolla flava.

11. **A. aegyptiaca** L. sp. p. 138 (*Lycopsis*); DC. pr. X. p. 48; Raul. cret. p. 814; Heldr. in ö. b. Z. 1877 p. 157; Bois. fl. or. IV. p. 159, suppl. p. 351, — *Asperugo aegyptiaca* L. sp. ed. 2 1. 198. — Icon: Jacq. hort. vind. III. t. 21. — Exsicc.: Heldr. pl. fl. hellen. a. 1876.

Annua, setis tuberculatis hispida; caulibus flaccidis, diffusis, vel adscendentibus simplicibus vel ramosis; foliis oblongo-lanceolatis, utrinque attenuatis, denticulatis, inferioribus petiolatis, superioribus sessilibus; racemis laxissimis, foliatis; pedicellis calyci aequilongis; calycis laciniis lanceolatis, demum accretis; corollae tubo calyce breviore; fornicibus velutinis, exsertis; nuculis semiovatis, elevatim reticulatis. — Caules interdum pedem et amplius longi, in speciminibus ex insula Salamis nani, 5—8 cm. tantum longi.

In saxosis maritimis, rarissime. Attica: insula Salamis (Heldr.); Creta: ad Hierapetra (Raul.). — Mart. Maio. ⊙.

β. Nuculae in rostrum obliquum abeuntes.

12. **A. stylosa** M. a. B. fl. taur. cauc. I. p. 123. — Icon: Rchb. pl. cret. VIII. f. 966. — Exsicc.: Heldr. fl. thessal. a. 1885.

Annua, setis tuberculatis parce obsita; caule gracili, erecto, superne ramoso-paniculato; foliis subnudulato-repandis, inferioribus oblongis, in petiolum attenuatis, superioribus semiamplexicaulibus; racemis saepius conjugatis, tandem laxis; bracteis pedicello brevioribus; pedicellis calyci aequilongis vel longioribus; calycis laciniis linearibus, obtusiusculis; corollae coeruleae tubo duplo vel subtriplo longiore; fornicibus papillosis, subexsertis; stylo calyce duplo subtriplo longiore; nuculis oblique ovatis, in rostrum breve subcurvatis, areolato-rugosis, minute tuberculatis.

β. **major** DC. pr. X. p. 48; Bois. fl. or. IV. p. 160. — *A. Spruneri* Bois. diagn. XI. p. 98; Haussk. symb. p. 152; Heldr. fl. Aegina p. 380. — Corollae tubus calyce sublongior; stylus calyce parum longior. — Exsicc.: Heldr. herb. norm. n. 208.

In campis, vineis, arvis regionis inferioris. Thessalia: pr. Velestinos (Heldr.); — β. Attica: pr. Eleusis (Haussk.), Athenas, insula Aegina (Heldr.); Acrocorinthus (Haussk.). — Mart. Maio. ⊙.

13. **A. caespitosa** Lam. enc. I. p. 498; S. et S. pr. I. p. 117, Fl. gr. II. p. 59 t. 169; Sieb. avis p. 3, in Flora I. p. 275; Raul. cret. p. 814; Bois. fl. or. IV. p. 156; Spreitz. in z. b. G. 1890 p. 297; Bald. viagg. Creta p. 78. — Exsicc.: Spreitz. in cret. a. 1882; Bald. it. cret. n. 81.

Perennis, e rhizomate sublignoso, pluricipite caespitosa, nana, setis tuberculatis hispida; caulibus brevissimis, racemo 3—5 floro, interdum in rosula subsessili, terminatis; foliis rosulatis, linearibus, undulato-dentatis vel integris, obtusis, caules superantibus; pedicellis calyce subbrevioribus; calycis laciniis linearibus, obtusis; corollae cyaneae tubo calyce subduplo longiore; fornicibus penicillatis, exsertis; stylo calyce longiore; nuculis oblique ovatis, in rostrum elongatum abeuntibus, areolato-rugosis. — Species insignis, foliis solo adpressis.

In saxosis regionis alpinae Cretae: mt. Volakia, Stravopodia, Hagios Theodoros, Hagion Pneuma (Raul.) et Mavrus Lakus (Spreitz.) in mt. Sphacioticis (Sibth.). — Jun. Aug. ♃

γ. Nuculae horizontaliter curvatae.

14. **A. aggregata** Lehm. asp. p. 219 t. 47; Mazz. in ant. ion. V. p. 182; Bois. fl. or. IV. p. 157. — *A. parviflora* S. et S. pr. I. p. 117, Fl. gr. II. p. 57 t. 167, non Willd. — *A. micrantha* R. et Sch. syst. IV. p. 93 et 771. — *Lycopsis glomerata* Urv. en. p. 22.

Annua, setis tuberculatis hispidissima; caulibus erectis vel diffusis, ramosis; foliis lineari-oblongis linearibusve, repandis, inferioribus in petiolum attenuatis, superioribus sessilibus; racemis densissimis, corymboso-capituliformibus; floribus subsessilibus, bracteis linearibus, calyce longioribus; calycis laciniis demum accretis, triangulari-lanceolatis; corollae cyaneae parvae tubo e calyce breviter exserto; fornicibus penicillatis,

exsertis; nuculis lateraliter compressis, breviter rostratis, reticulato-rugosis.

In arenosis maritimis. Attica: pr. Athenas (Sibth.); Corcyra: pr. Carumades (Mazz.); species pro Graecia dubia, a recentioribus non lecta. — Apr. Maio. ⊙ N. v.

3. Sectio. *Lycopsis* Griseb. spic. II. p. 95. — Corollae tubus ad medium inflexo-curvatus, limbus subirregularis, obliquus; nuculae basi in stipitem non contractae.

15. **A. arvensis** L. sp. p. 139; Pieri corc. fl. p. 26; Ch. et B. fl. pelop. p. 13; (*Lycopsis*); M. a. B. fl. taur. cauc. I. p. 123; Mazz. in ant. ion. V. p. 188; Bois. fl. or. IV. p. 160. — Icon: Fl. dan. t. 485.

Setis tuberculatis hispida; caulibus erectis, superne ramosis, corymboso-paniculatis; foliis lanceolatis, repando-dentatis, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; racemis saepius bifidis, tandem laxis, bracteis calyces aequantibus vel superantibus foliatis; pedicellis calyce brevioribus; calycis 5 partiti laciniis lanceolatis; corollae coeruleae tubo calyce sublongiore, medium versus curvato; fornicibus papillosis; nuculis magnis, transverse ovatis, curvato-rostratis, elevatim reticulato-areolatis, tuberculatis.

In arvis, incultis. Corcyra (Pieri); Peloponnesus: austro-occidentalis (Chaub.), pr. Nisi et Kalamata (Fauché); a recentioribus non lecta. — Maio, Jul. ⊙ N. v.

16. **A. variegata** L. sp. p. 138; S. et S. pr. I. p. 123, Fl. gr. II. p. 66 t. 178; Ch. et B. exp. p. 63, Fl. pelop. p. 13; Marg. et B. fl. Zante p. 69; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 814; Spreitz. in z. b. G. 1877 p. 715; Haussk. symb. p. 152; Heldr. fl. Aegina p. 380; (*Lycopsis*); Lehm. asp. p. 223; Mazz. in ant. ion. V. p. 180; Bois. fl. or. IV. p. 161, suppl. p. 351; Heldr. fl. cephal. p. 53; Fors. in bull. herb. Bois. III. p. 88; Form. in D. bot. Mon. 1898 p. 78. — *L. bullata* Cyr. pl. rar. I. p. 34 t. 11; Mazz. l. c. p. 188. — Huc probabiliter: *A. ovata* Mazz. l. c. p. 182; Friedr. Reise p. 265. — Exsicc.: Sint. it. thessal. n. 46; Baen. herb. europ. n. 9285.

Hispidula setisque sparsis tuberculatis obsita; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis oblongis, repando-dentatis, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; racemis solitariis, tandem laxiusculis, saepe elongatis; bracteis lanceolatis, calyce brevioribus; pedicellis calyce brevioribus; calycis 5 partiti laciniis lanceolato-linearibus; corollae cyaneae, saepe albo-variegatae, rarissime albae tubo calyci aequilongo, antrorsum curvato; fornicibus papillosis; nuculis parvis, transverse ovatis, breviter acutiuscule rostratis, tenuiter rugulosis.

In apricis, herbidis, ad vias regionis inferioris et montanae. Thessalia: pr. Kalabaka (Sint.), Pharsalus (Haussk.); Aetolia: mt. Mesolongion,

Krioneri (Reis.); Attica: pr. Athenas (Friedr.), mt. Kerata (Haussk.), Hymettus, Pharmacusarum insula Megali Kyra, insula Aegina (Heldr.); Peloponnesus (Sibth.); Acrocorinthus (Haussk.); pr. Manesi, Hagios Vlasios ad radices mt. Olenos Achaiae (Hal.), pr. Methone Messeniae (Chaub.); Creta (Sibth.): pr. Malaxa (Rev.), Canea (Weiss); Archipelagus (Sibth.): in insulae Syra (Weiss) mt. Pyrgos (Fors.); Zante: ad colles Acroteri, pr. Litakia (Marg.); Cephalonia: pr. Argostoli, mt. Aenos (Heldr.); Ithaca (Spreitz.); Corcyra: pr. Socracchi, Sidari, Magulades (Mazz.), Kanone (Baen.). — Febr. Jun. ⊙

7. Symphytum L. gen. n. 185.

a. Fornices inclusae.

1. **S. officinale** L. sp. p. 136; S. et S. pr. I. p. 119; Bois. fl. or. IV. p. 171; Form. in D. bot. Mon. 1891 p. 25. — Icon: Fl. dan. t. 664.

Praeter pubem minutam, setis albis, saepius tuberculatis asperum; radice fusiformi, ramosa; caulibus erectis, elatis, alatis, superne ramosis; foliis ovato-lanceolatis lanceolatisve, acuminatis, integris, inferioribus in petiolum decurrentibus, superioribus sessilibus, longe lateque decurrentibus; calycis hispiduli laciniis lanceolatis, tubo subduplo longioribus; corollae violaceae, rarius albidae, calyce triplo longioris, obconico-tubulosae, dentibus breviter triangularibus, obtusis, recurvis; fornicibus inclusis, antheras filamentis 2—3 longiores subsuperantibus; nuculis obliquis, ovato-trigonis, laevibus, lucidis.

In Graeciae provinciis et insulis (Sibth.); Thessalia: pr. Chaliki in Pindo, pr. Selicany in mt. Ossa (Form.); species pro ditione nostra dubia. — Maio, Jul. ♃.

2. **S. anatolicum** Bois. diagn. IV. p. 43, Fl. or. IV. p. 173. — Exsicc.: Dörfl. fl. aeg. n. 27.

Praeter pubem minutam, setis albis, saepius tuberculatis hirsutum; radice fusiformi, ramosa; caulibus flaccidulis, exalatis, fere a basi ramosis; foliis oblongis, acutis, subrepandis, inferioribus in petiolum attenuatis, superioribus sessilibus, non vel brevissime decurrentibus; calycis dense albo-strigosi laciniis lanceolato-linearibus, tubo 3—4plo longioribus; corollae albae vel violaceae, calyce duplo longioris, tubuloso-infundibuliformis, dentibus brevibus ovatis, obtusis, erectis; fornicibus inclusis, antheras filamentis aequilongas vix superantibus; nuculis rectiusculis, oblongis, areolato-nervosis, minute tuberculatis, subopacis.

In humidis umbrosis, in fauce Chalandra insulae Naxos (Leonis). — Apr. Maio. ♃.

b. Fornices longe exsertae.

3. **S. bulbosum** Schimp. in Flora VIII (1825) p. 17; Marg. et R. fl. Zante p. 68; Ung. Reise p. 127; Spreitz. in z. b. G. 1877 p. 715, 1887 p. 664; Bois. fl. or. IV. p. 176; Heldr. fl. cephal. p. 53, chlor. Parn. p. 24; Gelmi in bull. soc. bot. ital. 1889 p. 450; Hal.

in ö. b. Z. 1896 p. 15, 1897 p. 97; Haussk. symb. p. 150; Form. in Ver. Brünn 1896 p. 60. — S. *tuberosum* S. et S. pr. I. p. 119; Mazz. in ant. ion. V. p. 186,; non L. — *S. officinale* Pieri corc. fl. p. 24; Dallap. prosp. p. 20; non L. — *S. macrolepis* Gay. in Rchb. fl. germ. exc. p. 347; Friedr. Reise p. 280. — *S. brochum* Ch. et B. exp. p. 65 t. 7, Fl. pelop. p. 13 t. 9; Fraas fl. class. p. 163. — Probabiliter huc ex DC. pr. X. p. 52: *Anchusa bulbosa* Mazz. in ant. ion. V. p. 182. — Exsicc.: Orph. fl. gr. n. 863; Sint. it. thessal. n. 49 et 637.

Praeter pubem minutam, setis albis, saepius tuberculatis hirsutum; rhizomate gracili, elongato, tubera subglobosa gerente; caulibus erectis, exalatis, simplicibus vel parce ramosis; foliis ovato-oblongis, acutis, integris, inferioribus in petiolum attenuatis, superioribus sessilibus, interdum breviter decurrentibus; calycis viscidi laciniis lanceolatis, tubo subduplo longioribus; corolla ochroleuca, calyce duplo longiore, dentibus ovatis, obtusis, erectis; fornicibus denticulatis, dimidia corolla longioribus; antheris filamento aequilongis; nuculis obliquis tuberculatis, annulo basilari basi dentato.

In umbrosis regionis montanae et subalpinae. Thessalia: pr. Chaliki (Form.), mt. Ghavellu, in oropedio Neuropolis (Haussk.), pr. Kalabaka, Lechonia ad Volo (Sint.), mt. Pelion (Heldr.); Sporadum insula Skiathos (Leon.); mt. Parnassus, in Boeotia (Heldr.); Attica: ad Cephissum (Fraas); Achaia: pr. Patras (Reis.), mt. Panachaicon (Hal.); Messenia: pr. Methone, mt. Kupe, pr. Messene (Chaub.); Arcadia: mt. Diaforti pr. Andrizena (Friedr.); Laconia: mt. Malevo pr. Hagios Petros (Orph.); Zante (Sibth.); Cephalonia: in oropedio Omala (Heldr.); Leucas: pr. monasterium Hagios Georgios (Spreitz.); Corcyra: pr. San Deca (Spreitz.), Socracchi, Benizze, Casopitza, Figareto (Mazz.). — Apr. Jun. ♃.

4. **S. ottomanum** Friv. in Flora 1836 p. 439; Ung. Reise p. 127; Bald. riv. coll. bot. alb. 1896 p. 78. — Icon: Rchb. t. 104. — Exsicc.: Bald. it. alb. ep. IV. n. 189; Sint. it. thessal. n. 636.

Differt a praecedente radice fusiformi; caulibus valde ramosis; corolla brevi, dentibus parum e calyce emersis; fornicibus valde exsertis corolla aequilongis; antheris filamento longioribus.

In silvaticis subalpinis, rarissime. Epirus: ad Bocikopulon distr. Ljaskovik (Bald.); Thessalia: in valle Negerli pr. Chaliki (Sint.); Euboea: pr. Aqueductum (Ung.). — Apr. Jul. ♃.

8. **Pulmonaria** L. gen. n. 184.

1. **P. auriculata** Bois. diagn. XI. p. 95 (*Nonnea*). — *P. officinalis* S. et S. pr. I. p. 118; Ch. et B. exp. p. 63, Fl. pelop. p. 13. — *P. mollis* Bois. fl. or. IV. p. 170, vix Wolff. — *Anchusa Cesatiana* Fenzl et Friedr. in Friedr. Reise p. 282; sec. Bois. fl. or. IV. p. 162 et 1199. — Exsicc.: Heldr. pl. fl. hellen. a. 1899.

Rhizomate crasso, cylindrico, obliquo; caule erecto, simplici, breviter glanduloso-pubescenti, setulis albis intermixtis; foliis setulis brevibus,

adpressis scabridis, supra parce glandulosis, radicalibus oblongo-ovatis, acutis, in petiolum longum, alatum decurrentibus, caulinis paucis, oblongis, sessilibus, paulo supra basin utrinque rotundato-auriculatam, subangustatis, floralibus breviter ovatis, acutioribus, basi rotundatis; floribus in paniculam terminalem contractum dispositis; calyce glanduloso, ad tertiam partem acute 5 lobo, post anthesin inflato; corollae violaceae, glabrae, tubo calyci aequilongo, lobis rotundatis. — Species mihi ex unico specimine deflorato tantum nota, *P. molli* Wolff certe maxime affinis, ab ea tamen foliis caulinis paucis, oblongis, supra basin late rotundato-auriculatam subangustatis diversa videtur.

In silvaticis montanis, rarissime. Arcadia: in quericetis pr. Bassae (Friedr.); Messenia: in humidis vallis fl. Neda pr. Phigalea (Chaub.); Laconia: in regione abietina inter Langada et Neraidovuno (Zahn). — Mart. Maio. ♃.

3. Subtribus. *Lithospermeae* Koch. syn. p. 500.

9. Onosma L. gen. n. 159.

1. Sectio. *Asterotricha* Bois. fl. or. IV. p. 180. — Foliorum setae tuberculo stellatim piloso insidentes.

a. Caules herbacei; indumentum caulis adpressissimum.

1. **O. Spruneri** Bois. diagn. XI. p. 109, Fl. or. IV. p. 202. — Exsicc.: Orph. fl. gr. n. 744.

Totum setis brevibus, adpressissimis, e tuberculo minute stellatim piloso ortis cinereum; caulibus e radice verticali, simplici, 1—3, erectis, elatis, simplicibus; foliis longe lanceolatis, acutis, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; floribus breviter pedicellatis; calyce adpressissime setoso, laciniis lanceolatis; corolla albida, pubescenti, calyce subtriplo longiore, dentibus triangulari-ovatis, deflexis; antheris inclusis, filamentorum parte libera aequilongis; nuculis magnis oblongis, obtusiusculis, obscure granulato-tuberculatis, nitidis. — Caules 30—35 cm. alti, folia ad 10 cm. longa, 1 cm. lata, corolla 25 mm. longa, ore fere 1 cm. lata. — Caulibus herbaceis et indumento adpressissimo insigne.

Ad radices mt. Cithaeron Boeotiae pr. Kriekuki et in regione inferiori mt. Kyllene Achaiae, rarissime (Orph.). — Maio. Jun. ⊙.

b. Caules basi suffrutescentes; indumentum caulis patulum.

α. Antherae inclusae.

2. **O. tauricum** Pall. in nov. act. Petrop. 1792 p. 306. — *O. laconicum* Bois. et Orph. in Orph. fl. gr. exs. n. 860. — *O. stellulatum v. angustifolium* Bois. fl. or. IV. p. 201 p. p.; non *O. angustifolium* Lehm. pl. asperif. p. 361, quod foliis flavido-cinerascentibus, superioribus basi dilatatis discedit. — Exsicc.: Orph. l. c.; Heldr. herb. norm. n. 1458.

R lticaule; caulibus humilibus, erectiusculis vel espitosis, praeter pubem minutam, setis

sparsis patulis obsitis; foliis linearibus, oblongo-linearibusve, obtusiusculis, planiusculis, latitudine multoties longioribus, setis longis, adpressis, e tuberculo stellatim piloso ortis, obsitis, cinereis, radicalibus confertissimis, in petiolum attenuatis, caulinis sessilibus; floribus subsessilibus; calycis laciniis anguste lanceolatis, obtusiusculis, margine setis albis, longis, tuberculo non insidentibus, dense ciliatis, dorso breviter et adpressissime hirtis; corolla glabra, citrina, calyce duplo subtriplo longiore, dentibus triangulari-ovatis, deflexis; antheris filamentorum parte libera longioribus; nuculis ovato-trigonis, subinflexo-rostratis, obscure granulato-tuberculatis, nitidis. — Differt ab affine *O. stellulato* W. et K. foliis angustioribus, longioribus et floribus subsessilibus, nec pedicello post anthesin calycem aequante suffultis. — Specimina graeca cum tauricis a Callier it. taur. n. 81 distributis optime quadrant. Caules 10—25 cm. alti.

In rupestribus regionis montanae. Laconia: mt. Malevo pr. Hagios Joannes (Orph.), in valle fl. Nedon et in mt. Xirovuni Alagoniae (Zahn.). — Apr. Maio. ♃.

Obs. Nescio an huc spectent: *O. angustifolium* Form. in Ver. Brünn 1895 p. 32, ex Epiro pr. Ephemiades; et *O. stellulatum v. angustifolium* Bois. fl. or. IV. p. 201, e mt. Cithaeron, Kyllene et e Megaspilaeon, nec non Form. in Ver. Brünn 1896 p. 60, e Patsios in Olympo.

3. **O. erectum** S. et S. pr. I. p. 121, Fl. gr. II. p. 62 t. 173; Sieb. avis rem. p. 3; ?Urv. enum. p. 20; Raul. cret. p. 815. — *O. stellulatum v. erectum* Bois. fl. or. IV. p. 202 p. p.; Spreitz. in z. b. G. 1890 p. 297. — Exsicc.: Rev. pl. cret. p. 115; Dörfl. pl. cret. n. 64.

Basi suffrutescens, multicaule; caulibus humilibus, erectiusculis vel adscendentibus, simplicibus, caespitosis, praeter pubem minutam, setis patulis obsitis; foliis anguste linearibus oblongo-linearibusve, obtusiusculis, margine revolutis, latitudine multoties longioribus, setis longis adpressis et subpatulis, e tuberculo stellatim piloso ortis, obsitis, cinereis, radicalibus confertissimis, in petiolum attenuatis, caulinis sessilibus; floribus subsessilibus; calycis laciniis anguste lanceolatis, obtusiusculis, undique setis albis, longis, plerisque tuberculo non insidentibus, adpressis et subpatulis obsitis; corolla glabra, flava, calyce duplo longiore, dentibus triangulari-ovatis, deflexis; antheris filamentorum parte libera longioribus; nuculis ovato-trigonis, subinflexo-rostratis, laevibus, nitidis. — Differt a praecedente setis tuberculo majore longius stellatim piloso insidentibus obsitis, calycis laciniis undique fere aequaliter setosis, setis minus adpressis et colore corollae saturatiore.

β. **pubiflorum.** — *O. erectum* Clem. sert. p. 68. — Corolla pubescens. — Exsicc.: Heldr. herb. norm. n. 558 et 1358; Orph. fl. gr. n. 60.

In rupestribus montanis. Creta: mt. Lakus, in oropedio Omalo (Rev.), pr. Askyphos (Spreitz.), ad Hagia Rumeli, promontorium Meleka,

pr. Malaxa, mt. Ida (Raul.); ? Cycladum insula Melos (Urv.); — β. Euboea: pr. Steni (Pichl.); Attica: mt. Hymettus, Pateras (Heldr.); Achaia: mt. Panachaicon (Hal.). — Apr. Jun. ♃.

4. **O. echioides** L. sp. p. 137 (*Cerinthe*); L. sp. ed. 2 p. 196; Pieri. fl. corc. p. 25; Mazz. in ant. ion. V. p. 188; Friedr. Reise p. 271 (v. *sparsiflora*); Fraas fl. class. p. 164; Fors. in bull. herb. Bois. III. p. 88; Raul. cret. p. 815; non Jacq. fl. austr. III. t. 295, nec ant. mult. — *O. stellulatum* Ch. et B. exp. p. 64, Fl. pelop. p. 13, Raul. cret. p. 815; Form. in D. bot. Monat. 1891 p. 24; Bald. riv. coll. bot. Alb. 1895 p. 58. — *O. pallidum* Bois. diagn. XI. p. 112; Haussk. symb. p. 153. — *O. Orphanidis* Bois. in Heldr. herb. norm. n. 729; Ung. Reise p. 128. — *O. stellulatum v. genuinum* Bois. fl. or. IV. p. 201. — *O. stellulatum v. pallidum* Bois. l. c.; Heldr. chlor. Parn. p. 24; Form. in Ver. Brünn 1896 p. 61. — Exsicc.: Heldr. herb. norm. n. 729; Orph. fl. gr. n. 862 et 1005. —

Basi suffrutescens, multicaule; caulibus erectis, saepius elatis, praeter pubem minutam, setis patulis obsitis, foliis oblongo-linearibus oblongisve, obtusis, planiusculis, setis longis adpressis, e tuberculo stellatim piloso ortis obsitis, virescentibus, radicalibus sub anthesi saepe emarcidis, in petiolum attenuatis, caulinis sessilibus; floribus subsessilibus; calycis laciniis anguste lanceolatis, undique setis albis, longis, plerisque tuberculo non insidentibus, erecto-patulis obsitis; corolla pubescente, pallide flava, calyce 2—3 plo longiore, dentibus triangulari-ovatis, deflexis; antheris filamentorum parte libera longioribus; nuculis ovato-trigonis, subinflexo-rostratis, laevibus, nitidis. — Species ab auctoribus multis cum *O. stellulato* W. et K. confusa, quod autem egregie differt pedicellis post anthesin calycem aequantibus, nec brevissimis.

α. **typicum.** — Folia oblongo-linearia, latitudine multo longiora, virescentia, radicalia sub anthesi emarcida, corolla calyce duplo longior. — Exsicc.: Orph. fl. gr. n. 862 et 1005; Heldr. herb. norm. n, 719.

β. **brachyphyllum.** — Folia oblonga latitudine triplo longiora, radicalia sub anthesi emarcida; corolla elongata calyce triplo longior. — Exsicc.: Dörfl. fl. gr. n. 285.

γ. **brachycalyx.** — Folia anguste oblongo-linearia vel linearia, latitudine multoties longiora, radicalia caespitosa; corolla elongata, calyce triplo longior. — An forte species propria. — Exsicc.: Hal. it. gr. II. a. 1893.

In saxosis regionis montanae. Epirus: mt. Micikeli (Bald.); Thessalia: ad monasterium Meteora, pr. Malakasi, Uranaeos, Tsungeri in Pindo, mt. Pelion (Haussk.), pr. Kapurna, pr. Miluna in Olympo (Form.); Euboea: mt. Dirphys (Heldr.); Aetolia: mt. Arapocephala supra Agrinion (Heldr.); Phocis (Sprun.): mt. Parnassus pr. Rachova (Orph.); Boeotia: pr. Lebadea (Orph.); Attica: mt. Cithaeron (Heldr.), Parnes, Hymettus (Fraas); Achaia: mt. Kyllene (Fraas); Argolis: pr. Poros (Friedr.);

Arcadia: pr. Carithena, Androvista (Chaub,); Cycladum insula Syra: mt. Pyrgos (Fors.); Creta: ad promontorium Meleka (Raul.); Corcyra: pr. Spagus, Ducade, Afiona (Mazz.); sed ob confusionem permultam hujus speciei loca indicata nonnulla vix huc pertinent; — β. Aetolia: mt. Korax pr. Musinitza (Leon.); — γ. Achaia: mt. Kastro pr. Kalavryta (Hal.). — Maio, Jun. ♃.

β. Antherae exsertae.

5. **O. leptanthum** Heldr. herb. norm. 1565.

Basi suffrutescens, multicaule; caulibus humilibus, erectiusculis vel adscendentibus, simplicibus, caespitosis, patule setosis; foliis oblongo-linearibus, obtusis, planis, latitudine multoties longioribus, setis patulis, e tuberculo stellatim piloso ortis obsitis, cinerascentibus, radicalibus confertis, in petiolum attenuatis, caulinis sessilibus; floribus pedicellatis, pedicellis post anthesin calyce dimidio brevioribus; calycis laciniis anguste lanceolatis, acutiusculis, setis patulis longis undique obsitis; corolla flava, glabra, calyce parum longiore, dentibus triangularibus, deflexis; antheris filamentorum parte libera sublongioribus; nuculis ovato-trigonis, subinflexo-rostratis, obscure granulato-tuberculatis, nitidis. — Species distinctissima, habitu *O. frutescentis*, sed asterotricha.

In rupestribus regionis mediae et superioris mt. Taygetos, l. d. Megalo Stephani, Chalasmeno (Zahn) et Hagia Barbara (Pichl.). — Jun. Jul. ♃.

2. Sectio. *Heterotricha* Bois. fl. or. IV. p. 180. — Foliorum setae tuberculo ad folia inferiora saepius glabro, ad superiora praesertim in pagina inferiori stellatim piloso insidentes.

6. **O. helveticum** DC. pr. X. p. 62 pro var. *O. echioides*; Bois. diagn. XI. p. 111; Hal. Beitr. fl. Epir. p. 32. — Exsicc.: Hal. it. gr. II. a. 1893.

Totum setis longis, patulis, e tuberculo glabro vel sparse et brevissime stellatim piloso ortis obsitum, virescens; caulibus e radice verticali simplici, herbaceis, erectiusculis, mediocribus, simplicibus vel parce ramosis; foliis lanceolato-linearibus, acutiusculis, inferioribus in petiolum attenuatis, superioribus semiamplexicaulibus; floribus breviter pedicellatis; calycis subpatule et longe setosi laciniis lanceolatis; corolla pallide flava, puberula, calyce subduplo longiore, dentibus triangularibus, brevibus, deflexis; antheris inclusis, filamentorum parte libera aequilongis, nuculis ovato-trigonis, subinflexo-rostratis, sublaevibus, nitidis. — Indumento mixto inter sectionem praecedentem et sequentem media, habitu *O. arenaria* W. et K. simile.

In rupestribus regionis superioris mt. Strungula supra Melisurgi in Epiro (Hal.); Creta: ad promontorium Meleka (Sieb.). — Jun. Jul. ⊙.

3. Sectio. *Haplotricha* Bois. fl. or. IV. p. 179. — Foliorum setae tuberculo glabro insidentes.

a. Herbacea, biennia.

7. **O. graecum** Bois. diagn. XI. p. 106, Fl. or. IV. p. 182; Weiss in z. b. G. 1869 p. 742. — *O. echioides* Sieb. avis p. 3. — *O. tinctorium* Ch. et B. exp. p. 65, Fl. pelop. p. 13; non M. a. B. Huc forsan: *O. arenarium* Ung. Reise p. 128 (e Boeotia ad lacum Hylike), non W. et K. —

Totum setis adpressis, tuberculo minuto glabro insidentibus pubeque brevi, velutina cinerascens, vix strigosum; caule erecto, crassiusculo, mediocri, superne ramoso-paniculato; foliis radicalibus lanceolato-linearibus, obtusis, basi attenuatis, superioribus sessilibus, a basi rotundata oblongis; floribus subsessilbus; calycis fructiferi valde accreti laciniis linearibus; corolla anguste tubulosa, pubescenti, lutea, demum violascenti, calyce subduplo longiore, dentibus brevissimis; antheris summo apice vix exsertis, filamento brevissimo; nuculis trigonis, profunde scrobiculatis et tuberculatis. — Rarius pede dimidio elatius, corolla anguste tubulosa et nuculis profunde scrobiculato-tuberculatis insigne.

In siccis regionis submontanae. Attica: mt. Hymettus (Bois.); Messenia: pr. Methone (Chaub.), Navarin (Fauché); Creta: pr. Canea (Weiss), Malaxa (Raul.); Cycladum insula Syra (Bois). — Maio. ⊖. N. v.

8. **O. taygeteum** Bois. et Heldr. diagn. XI. p. 101, Fl. or. IV. p. 182. — Icon: Rouy. ill. t. 18. — Exsicc.: Heldr. pl. fl. hellen. n. 421.

Totum setis longis patentibus, tuberculo majusculo glabro insidentibus strigosissimum, virescens; caule erecto, crasso, a basi ramosissimo; foliis radicalibus linearibus, longis, obtusis, caulinis abbreviatis, lanceolatis ovato-lanceolatisve, sessilibus; floribus pedicellatis, pedicellis calyce 3—4plo brevioribus; calycis fructiferi valde accreti laciniis lanceolatis, binis vel ternis saepius coalitis; corolla clavata, pubescenti, pallida, calyce vix longiore, dentibus brevissimis; antheris apice breviter exsertis, filamento brevissimo; nuculis trigonis, magnis, rugoso-tuberculatis, nitidis, in rostrum anceps, acutum, rectum abrupte angustatis. — *O. setoso* Led. (*O. echioides* Jacq. non L. = *O. Visianii* Clem. = *O. calycinum* Stev.) affine, quod foliis summus basi valde dilatatis semiamplexicaulibus et antheris inclusis differe videtur — *O. rhodopeum* Vel. in Sitzungsb. böhm. Ges. Wiss. 1894 p. 22 vix a nostro discedit.

Inter lapides in alveo exsiccato torrentis inter Varvara et Hagios Paraskevi in regione silvatica mt. Taygetos (Heldr.). — Jul. ⊖.

b. Suffruticosum.

9. **O. frutescens** Lam. ill. gen. n. 1837; Raul. cret. p. 815; Bois. fl. or. IV. p. 190; Hal. Beitr. fl. Aetol. p. 8, Beitr. fl. Achaia p. 26; Haussk. symb. p. 153. — *O. echioides* S. et S. pr. I. p. 121, Fl. gr. t. 172; Ch. et B. exp. p. 65, Fl. pelop. p. 13, non L. — *O. orientale* Lehm. asp. p. 376; non Pall. quod = *Arnebia cornuta* Fisch. et Mey.; nec *Cerinthe orientalis* L. am. ac. IV. p. 267, quae species dubia. — Exsicc.: Orph. fl. gr. n. 61; Heldr. herb. norm. n. 367 et 1359.

Basi suffruticosum; multicaule, totum setis, tuberculo glabro insidentibus hispidum, cinerascens; caulibus hornotinis herbaceis, humilibus, erectiusculis vel adscendentibus, simplicibus; foliis oblongis vel oblongo-lanceolatis, basi attenuatis; floribus pedicellatis, pedicellis calyce brevioribus; calycis fructiferi valde accreti laciniis lanceolatis; corolla glabra, flava, apice violacea, ampliata, calyce tertia parte longiore, dentibus breviter triangularibus, deflexis; antheris exsertis, filamentorum parte libera longioribus, nuculis subglobosis, breviter rostratis, laevibus, nitidis, marmoratis.

In rupium fissuris regionis inferioris et montanae. Acarnania (Nied.); Aetolia: mt. Chalkis et Taphiassos ad sinum Patranum (Hal.); Boeotia: pr. Drachmano (Heldr.); Attica: mt. Cithaeron, Parnes, Lycabettus, Hymettus (Heldr.), Kerata (Haussk.), in Pharmacusarum insula Lero (Heldr.); Acrocorinthus (Sprun.); Achaia: pr. Psatopyrgos (Heldr.), Kalavryta, in valle Voreikos pr. Megaspilaeon (Hal.), mt. Kyllene (Orph.); Argolis: in peninsula Vromolimni, mt. Palamidi pr. Nauplia (Haussk.); Arcadia, Messenia (Chaub); Laconia: pr. Sparta, Leondari (Bois.); Creta: pr. Acroteri, Malaxa (Raul.). — Apr. Jun.

Obs. *O. simplicissimum* L. sp. ed. 2 p. 196; Raul. cret. p. 815. — In Creta certissime erronee indicatur.

10. Echium L. gen. n. 191.

a. Stamina exserta.

α. Corolla albida.

1. **E. italicum** L. sp. p. 139; S. et S. pr. I. p. 124; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 64, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 190; Marg. et R. fl. Zante p. 68; Fraas fl. class. p. 164; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 205; Heldr. fl. cephal. p. 53, Fl. Aegina p. 380; Spreitz. in z. b. G. 1887 p. 665; Hal. in z. b. G. 1888 p. 760, Beitr. fl. Epir. p. 32, Beitr. fl. Aetol. p. 8, Beitr. fl. Thessal. p. 16; Form. in D. bot. Monat. 1891 p. 24, in Ver. Brünn 1895 p. 32, 1896 p. 60, 1897 p. 44; Haussk. symb. p. 152. — *E. pyramidatum* DC. pr. X. p. 23; Raul. cret. p. 813. — *E. altissimum* Raul. l. c, vix Jacq. fl. austr. V. p. 35, quod panicula ob ramos florales abbreviatos in racemum elongatum spiciformem abeunti, spiculis brevibus, folio florali brevioribus, calyce minori et corolla calyce duplo longiore discedit et in Graecia vix crescit. — Exsicc.: Dörfl. fl. aeg. n. 72.

Caule erecto, setoso-hispidissimo, in paniculam amplam pyramidatam abeunti; foliis lineari-lanceolatis, setis basi tuberculatis, adpressiusculis hispidis; spiculis elongatis, inferioribus saepius bifidis trifidisve, folia floralia superantibus; calycis laciniis lanceolatis; corolla hirta, sub anthesi calycem parum excedente; nuculis trigonis, acuminatis, muricato-tuberculatis.

In aridis, cultis derelictis regionis inferioris et submontanae per totam Graeciam. — Apr. Jul. ♃

β. Corolla coerulea, violacea vel punicea.

× Folia linearia vel lanceolata, basi non cordata.

○ Biennia; corolla coerulea; nuculae breviter tuberculatae.

2. **E. vulgare** L. sp. p. 139; S. et S. pr. I. p. 125; Pieri corc. fl. p. 26; Ch. et B. exp. p. 64. Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 190; Fraas fl. class. p. 163; Hal. in z. b. G. 1888 p. 763. — Icon.: Fl. dan. t. 445. — Exsicc.: Hal. it. gr. a. 1888.

Viride, setis homomorphis, basi tuberculatis obsitum; caule unico vel pluribus, erectis, elatis, in paniculam angustam vel pyramidatam abeuntibus; foliis lanceolatis vel lineari-lanceolatis; spiculis brevibus, inferioribus interdum bifidis, folia floralia superantibus; calycis adpresse setoso-strigosi laciniis lineari-lanceolatis, erectis; corolla hirta, calyce duplo subtriplo longiore; nuculis subtrigonis, acutis, tenuiter et inaequaliter tuberculatis.

In arvis arenosis, rupestribus regionis inferioris, rerissime. Thessalia (Fraas); Attica: pr. Tatoi (Hal.); Laconia (Sibth.); Corcyra (Pieri): pr. Stavro (Mazz.). — Maio, Jul. ⊙

3. **E. pustulatum** S. et S. pr. I. p. 125, Fl. gr. II. p. 68 t. 180; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 271; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 206.

Differt a praecedente, cui valde affine, indumento setis dimorphis, aliis minutis adpressiusculis, aliis longis e tuberculo valido ortis, strigoso-hispido, cinerascenti, panicula pyramidata laxiuscula, corolla majori et nuculis magis tuberculatis.

In arvis, herbidis regionis inferioris, rarissime. Zacynthus (Marg.); insula Poros (Friedr.), Syra et pr. Canea Cretae (Weiss). — Apr. Jul. ⊙ N. v.

○○ Perennia; corolla punicea (sicca violacea); nuculae acute tuberculatae.

4. **E. elegans** Lehm. asper. p. 459; Weiss in z. b. G. 1869 p. 742. — *E. hispidum* S. et S. pr. I. p. 125, Fl. gr. II. p. 68 t. 181; Link. in Linnaea IX. p. 578; Haussk. symb. p. 152; Heldr. in ö. b. Z. 1898 p. 104, chlor. Thera p. 17; non Burm. prodr. p. 5, quod *E. capitatum* L. mant. p. 42 = *Lobostemon capitatus* Buek in Linnaea 1837 p. 143; nec Thunb. in Schrad. journ. 1806 p. 40 = *Lobostemon hispidus* DC. pr. X. p. 10; ambae species Capenses. — *E. Sibthorpii* Roem. et Sch. syst. IV. p. 26. — *E. diffusum* Ch. et B. exp. p. 339, non S. et S. — *E. setosum* Ch. et B. fl. pelop. p. 13; non Vahl symb. II. p. 35, quod species aegyptiaca. — *E. rubrum* Fraas fl. class. p. 163, non Jacq. — *E. sericeum v. hispidum* Bois. fl. or. IV. p. 207; Form. in D. bot. Monat. 1898 p. 78. — Exsicc.:

Heldr. herb. norm. n. 240 et 1160; Orph. fl. gr. n. 864; Rev. pl. cret. n. 112; Dörfl. fl. aeg. n. 125, pl. cret. n. 146.

Pluricaule, totum setis patulis, basi tuberculatis hispidissimum, canum; caulibus erectis, elatis, basi interdum lignosis, in paniculam angustam, laxam abeuntibus; foliis linearibus vel lineari-lanceolatis; spiculis simplicibus, brevibus, demum elongatis, folia floralia superantibus; calycis laciniis linearibus, erectis; corolla pubescente, praeterea setis longioribus obsita, calyce subtriplo longiore.

In arenosis praesertim maritimis. Attica: pr. Athenas (Orph.), in colle Turcovuni (Haussk.), ad Phalerum (Heldr.); Achaia: pr. Exocampos (Heldr.); Sycionia, Corinthus (Chaub.); Argolis: pr. Nauplia (Sibth.), insula Hydra (Pichl.); Laconia: pr. Monembasia, Vatica (Fraas); Cycladum insula Syra (Weiss), Naxos, Thera (Heldr.); Creta: pr. Kissamos (Rev.), Canea (Weiss), Anatoli distr. Hierapetra (Leon.). — Apr. Jun. ♃

5. **E. sericeum** Vahl symb. II. p. 35; Bois. fl. or. IV. p. 207. — *E. elegans v. incanum* Bois. et Heldr. diagn. XI. p. 93; Raul. cret. p. 812. — Exsicc.: Heldr. pl. cret. a. 1846.

Differt a praecedente indumento e pube duplici, altera brevissima densa, altera e setis elongatis constanti, adpressissime cinereo; foliis margine subundulatis, floralibus abbreviatis; calycis laciniis obtusioribus.

In arenosis maritimis Cretae: pr. Canea (Heldr.), Khalepa, Retymo, Candia (Raul.). — Apr. Jun. ♃

×× Folia radicalia ovata vel oblonga, superiora e basi cordata semiamplexicauli attenuata.

6. **E. plantagineum** L. mant. alt. p. 202; S. et S. pr. I. p. 124, Fl. gr. II. p. 67 t. 179; Sieb. avis p. 3; Ch. et B. exp. p. 64, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 188; Friedr. Reise p. 271; Raul. cret. p. 813; Bois. fl. or. IV. p. 208; Heldr. fl. cephal. p. 53, Fl. Aegina p. 380, chlor. Mykon. p. 247, prosth. chlor. Thera p. 3; Gelmi in bull. soc. bot. it. 1889 p. 450; Form. in D. bot. Monat. 1891 p. 24, in Ver. Brünn 1895 p. 32, 1896 p. 60, 1897 p. 44; Haussk. symb. p. 152; Hal. Beitr. fl. Epir. p. 32, in ö. b. Z. 1897 p. 97. — *E. violaceum* Urv. enum. p. 22; Tausch in Flora XII. p. 641; Mazz. in ant. ion. V. p. 190; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 277; Ung. Reise p. 127; Weiss in z. b. G. 1869 p. 742; non L. mant. p. 42, quod sec. Gr. et Godr. fl. fr. II. p. 524 probabiliter ad *E. rubrum* Jacq., sec. alios ad *E. pustulatum* S. et S. spectat. — Huc forsan: *E. maritimum* Sieb. avis rem. p. 3, vix Willd. sp. I. p. 788. — Exsicc.: Sint. it. thessal. n. 1492 et 299 b. (*v. versicolor* Haussk. symb. p. 152, virescens sparse setosum, caules elongati laxe paniculati, flores minores); Dörfl. fl. aeg. n. 136.

Pilis subadpressis, flexibilibus, basi tuberculatis, molliter hispidum; caulibus solitariis vel pluribus, erectis vel adscendentibus, ramosis vel simplicibus, laxiuscule paniculatis; foliis radicalibus ovatis oblongisve, in petiolum attenuatis, praeter nervum medium nevis lateralibus

conpicuis munitis, caulinis saepius valde decrescentibus, semiamplexicaulibus; racemis dissitifloris, demum valde elongatis: calycis laciniis lineari-lanceolatis; corolla violacea, calyce 3—4 plo longiore, tubo in limbum amplum, superne paucisetosum abrupte ampliato: nuculis grosse tuberculato-asperis, obtusiusculis. — Stirps sat polymorpha, a speciebus aliis nostris tamen foliis radicalibus ovatis, nervosis et floribus magnis facile distinguenda. — Variat interdum floribus albis: *E. niveum* Mazz. in ant. ion. V. p. 190. — *E. plantagineum v. albiflorum* Raul. cret. p. 813.

In herbidis, pratis, collibus aridis regionis inferioris per totam Graeciam. — Mart. Jun. ♃

β. Stamina inclusa.

7. **E. diffusum** S. et S. pr. I p. 125, Fl. gr. II. p. 69 t. 182; Sieb. avis p. 2, rem. p. 3, in Flora V. 2 p. 639; Guss. cat. pl. Bocc. a. 1821 p. 23. — *E. arenarium* Guss. pl. rar. p. 88 t. 17, pr. fl. sic. I. p. 227, fl. sic. syn. I. p. 233; Clem. sert. p. 68; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 812; Bois. fl. or. IV. p. 210; Spreitz. in z. b. G. 1887 p. 665; Haussk. symb. p. 152; Heldr. fl. Aegina p. 380; chlor. Mykon. p. 247. — ? *E. setosum β.* Ch. et B. fl. pelop. p. 13. — *E. sericeum v. diffusum* Bois. fl. or. IV. p. 207. — Exsicc.: Heldr. herb. norm. n. 339; Rev. pl. cret. n. 113; Dörfl. fl. aeg. n. 143.

Setis basi tuberculatis hispidum; caulibus adscendentibus vel prostratis, simplicibus vel ramosis; foliis radicalibus oblongo-spathulatis, obtusis, in petiolum attenuatis, caulinis sessilibus, a basi subdilatata, saepe semiamplexicauli lanceolatis; spicis demum elongatis; calycis patule strigoso-hispidi, fructiferi feri immutati, laciniis lanceolatis; corolla parva, hirta, calyce subduplo longiore; nuculis acuminatis, acute tuberculatis. — *E. diffusum* S. et S., ex descriptione et icone meo sensu ab *E. arenario* Guss. non differt, ideoque nec ad formas *E. sericei*, nec ad eas *E. hispidi* pertinet.

In arenosis maritimis. Attica: ad Piraeum, Phalerum, pr. Raphina (Heldr.), Laurion (Haussk.); Argolis (Sprun.); insula Aegina, Tenos, Syra (Weiss), Cythnos (Heldr.), Rhenea (Tunt.), Naxos (Sart.), Makaria (Leon.), Thera (Heldr.); Creta (Sibth.): pr. Kissamos, Platania (Rev.), Francocastello, insula Gaidaronisi (Raul.); Cythaera (Spreitz.). — Apr. Jul ⊙ et ⊙.

8. **E. parviflorum** Moench meth. p. 423. — *E. calycinum* Viv. ann. bot. I. 2 p. 164; Urv. enum p. 22; Mazz. in ant. ion. V. p. 190; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 268 et 277; Ung. Reise p. 128; Raul. cret. p. 812; Bois. fl. or. IV. p. 210; Haussk. symb. p. 152; Heldr. fl. Aegina p. 380. — *E. creticum* S. et S. pr. I. p. 69 t. 183; Pieri corc. fl. p. 27; Ch. et B. 13; an L. sp. p. 139. — *E. prostratum* 60 t. 12; Mazz. in ant. ion. V. p. 192; erb. norm. n. 523.

Setis basi tuberculatis piloso-hispidum; caulibus adscendentibus vel prostratis, simplicibus, rarius parce ramosis; foliis radicalibus oblongis oblongo-spathulatisve, obtusis, in petiolum attenuatis, caulinis sessilibus, interdum semiamplexicaulibus; spicis demum elongatis; calycis adpresse setosi, fructiferi valde aucti, laciniis foliaceis, oblongis; corolla parva, hirta, calyce subduplo longiore; nuculis acuminatis; acute tuberculatis. A praecedente indumento molliori, foliis latioribus, calyce adpresse setoso, laciniis foliaceis distinguitur.

β. **erectum** DC. pr. X. p. 21. — Caulibus erectis. — Exsicc.: Heldr. pl. fl. hellen. a. 1878.

In collibus siccis, arenosis maritimis, aridis regionis inferioris. — Attica: ad Rheitro pr. Eleusis (Ung.), mt. Kerata (Haussk.), in colle Lycabetto, ad Piraeum, Phalerum, in Pharmacusarum insula Lero (Heldr.), pr. Laurion (Haussk.); insula Aegina, Poros (Friedr.); Messenia: pr. Navarin (Chaub.); Cycladum insula: Melos (Urv.), Naxos (Chaub.); Creta: pr. Lutro (Raul.); Zante: mt. Scopo (Marg.); Corcyra (Pieri); pr. Stavropotamo, Chiefaloipso (Mazz.). — Mart. Maio. ⊙

11. Moltkia Lehm. in act. soc. nat. scrut. halens. III. 2. p. 3.

1. **M. petraea** Tratt. thesaur. bot. p. 8 t. 34 (*Echium*); Griseb. spic. II. p. 515; Bois. fl. or. IV. p. 221; Bald. riv. coll. bot. alb. 1895 p. 58, 1896 p. 79. — *Lithospermum petraeum* DC. pr. X. p. 82. — Exsicc.: Bald. it. alb. epir. III. a. 1875.

Suffruticosum, ramosissimum; foliis oblongo-linearibus linearibusve, obtusis, supra strigulosis, subtus incanis, margine subrevolutis; racemis cymosis, confertis; calycis laciniis linearibus, cano-sericeis; corolla azurea, rarissime alba, tubuloso-infundibuliformi, calyce subtriplo longiore; nuculis ovatis, laevissimis, oblique attenuato-rostratis.

In rupium fissuris regionis montanae, rarissime. Epirus: pr. Vonicko, ad Klissura distr. Philippiada (Bald.); mt. Oeta pr. Thermopylas (Sprun.). — Maio, Jul. ♃

12. Macrotomia DC. in Meisn. comm. p. 190.

1. **M. cephalotes** DC. pr. X. p. 96 (*Arnebia*); Bois. fl. or. IV. p. 213. — *Munbya cephalotes* Bois. diagn. XI. p. 116. — Exsicc.: *Munbya jasminiflora* Orph. fl. gr. n. 112.

Collo crasso, valde squamoso; caule erecto, simplici; foliis adpressissime setosis, canis, radicalibus crebris, longe lineari-lanceolatis, caulinis brevioribus, acuminatis; racemis terminalibus, in capitulum amplum congestis; calycis albo-villosi laciniis lineari-acuminatis; corollae aureae, pubescentis, calyce parum vel sesqui-longiore, limbo amplo, lobis subrotundis; nuculis scrobiculato-punctatis, obsolete tuberculatis, triquetro-ovatis, apice attenuatis, acutis, ancipitibus. — Planta pulcherrima, ad 30 cm. alta, foliis radicalibus ad 10 cm. longis, 1 cm. latis, corymbo multifloro, corolla circa 3 cm. longa.

In rupestribus regionis mediae mt. Chelmos pr. Stygem, rare (Orph.). — Jun. Jul. ♃

13. Alkanna Tausch in Flora 1824 p. 234.

a. Nuculae valde curvatae, vermiformes, sinu basilari angusto, rostro fere deorsum flexo.

α. Folia undulato-crispa.

1. **A. orientalis** L. sp. p. 133 (*Anchusa*); Bois. diagn. IV. p. 46, Fl. or. p. 228. — *Lithospermum orientale* L. syst. veg. p. 156; S. et S. pr. I. p. 114, Fl. gr. p. 51 t. 160; Ch. et B. exp. p. 61, Fl. pelop. p. 12.

Tota indumento brevi glanduloso-viscida, insuper setis brevibus tuberculatis interdum sparsissimis obsita; caulibus erectiusculis, superne ramosis; foliis acutis, rosularum elongato-lanceolatis, inferne longe attenuatis, caulinis semiamplexicaulibus, inferioribus oblongo-lanceolatis lanceolatisve, superioribus ovato-lanceolatis, in bracteas sursum diminutas abeuntibus; racemis fructiferis valde elongatis; calycis fructiferi accreti, breviter pedicellati, deflexi, laciniis lanceolatis; corolla extus glabra, saturate lutea, calyce duplo longiore, limbo majusculo; nuculis scrobiculato-reticulatis, areola basilari breviter stipitata.

α. **typica.** — Bracteae ovatae, obtusae; racemi fructiferi densiusculi. — Exsicc.: N. v.

β. **hellenica** Bois. fl. or. IV. p. 228; Hal. Beitr. fl. Aetol. p. 9; Haussk. symb. p. 155. — Bracteae breviter lanceolatae, acutae; racemi fructiferi laxiusculi. — Exsicc.: Heldr. herb. norm. n. 204; Orph. fl. gr. n. 297; Dörfl. fl. aeg. n. 379.

In rupestribus regionis inferioris et montanae. Archipelagus (Sibth.); — β. Aetolia: mt. Arakynthos (Heldr.), pr. Mesolongion (Nieder), mt. Chalkis (Hal.); Phocis: pr. Delphi (Leon.); Peloponnesus: Acrocorinthus (Bois.), mt. Malevo (Sprun.), Argolis, in Maina, pr. Messene, Corone, Methone, mt. Kupe (Chaub.). — Febr. Maio. ♃

β. Folia non undulato-crispa.

× Calyx glanduloso-hispidus.

○ Corolla extus pubescens.

2. **A. calliensis** Heldr. in Bois. fl. or. suppl. p. 353. — Exsicc.: Heldr. it. gr. sept. a. 1879; Dörfl. fl. gr. n. 337.

Tota indumento brevi glanduloso-viscida, insuper setis brevibus, tuberculatis, interdum sparsissimis obsita; caulibus erectiusculis, superne ramosis; foliis inferioribus oblongis, obtusis, basi attenuatis, superioribus basi subdilatata semiamplexicaulibus, oblongis oblongo-lanceolatisve, acutis, in bracteas sursum diminutas abeuntibus; racemis fructiferis valde [illegible]ifloris; calycis fructiferi subaccreti, brevissime pedicellati, [illegible] lanceolatis; corolla pubescente, lutea, tubo calycem [illegible] limbo parvo; nuculis scrobiculato-reticulatis, areola

basilari stipitata. — A praecedente foliis non undulatis, racemis densifloris, calyce et corolla fere duplo minore discedit.

In regione abietina mt. Korax pr. Musinitza Aetoliae (Heldr.). — Maio. Jul. ♃

3. **A. pindicola** Haussk. in Mitth. thür. bot. Ver. 1888 p. 32, symb. p. 153. — *A. pulmonaria* Form. in D. bot. Mon. 1891 p. 24; Hal. Beitr. fl. Thessal. p. 16; vix Griseb. spic. II. p. 90 (cf. Griseb. l. c. p. 515 et Haussk. l. c. p. 154). — Huc probabiliter: *A. scardica* Form. in Ver. Brünn 1896 p. 60, vix Griseb. — Exsicc.: Sint. it. thessal. n. 1494.

Tota indumento brevi glanduloso-viscida, insuper setis brevibus, tuberculatis, interdum sparsissimis obsita; caulibus erectiusculis, superne ramosis; foliis acutis, inferioribus oblongo-lanceolatis, basi attenuatis, superioribus basi dilatata semiamplexicaulibus, oblongis vel ovato-lanceolatis in bracteas sursum diminutas abeuntibus; racemis fructiferis laxiusculis; calycis fructiferi subaccreti, brevissime pedicellati, deflexi, laciniis lanceolatis; corolla pubescente, sordide flavida, striis violascentibus longitudinaliter percursa, tubo longe exserto, calyce duplo longiore, limbo parvo; nuculis scrobiculato-reticulatis, areola basilari breviter stipitata. — A praecedente foliis tenuioribus, majoribus, omnibus acutis, corolla longe exserta, pallida, violaceo-striata discedit.

β. **conferta** Haussk. symb. p. 154. — Caules breviores, dodrantales, tenuiores, densius foliosi; folia minora; racemi fructiferi conferti; nuculae apice breviter carinatae. — Exsicc.: N. v.

In saxosis regionis inferioris et montanae Thessaliae: pr. Malakasi, Said Pascha, Korona, Vlazda, Hagios Paraskevi, Kritsotades, Konisko, Kastri, Kalabaka (Form.), mt. Ghavellu (Haussk.); — β. ad Gionskala supra Sermeniko (Haussk.). — Maio, Jun. ♃

4. **A. methanaea** Haussk. in Mitth. thür. bot. Ver. 1888 p. 32, symb. p. 153. — *A. pulmonaria* Hal. Beitr. fl. Achaia p. 26, vix Griseb. — *A. graeca* Hal. in z. b. G. 1899 p. 190, non Bois. et Spr. — Exsicc.: Haussk. it. gr. a. 1885; Heldr. pl. fl. hell. a. 1887; Hal. it. gr. sec. a. 1893.

Caulibus erectiusculis; mox supra basin vel a medio ramosis, glanduloso-viscidis, pilisque longioribus saepe tuberculatis obsitis; foliis acutis, pilis setaceis, minute tuberculatis strigulosis, radicalibus elongato-lanceolatis, basi attenuatis, caulinis semiamplexicaulibus, inferioribus oblongis oblongo-lanceolatisve, superioribus saepe basi dilatatis, in bracteas anguste lanceolatas sursum abeuntibus; racemis fructiferis laxifloris, elongatis: calycis fructiferi accreti, pedicellati, deflexi, laciniis lanceolatis, molliter glanduloso-villosis; corolla pubescente, flavida, tubo fuscescente, calyce longiore, limbo mediocri; nuculis scrobiculato-reticulatis, areola basilari stipitata. — Habitu et characteribus magis ad sequentem, nec ad praecedentes spectat, sed corolla extus pubescens, ab ea insuper bracteis angustioribus, saepe anguste lanceolatis specifice differre videtur.

In saxosis montanis. In peninsula Methana pr. Vromolimni, ad Acrocorinthum (Haussk.); Achaia: in faucibus Phlamburitza mt. Kyllene (Heldr.), pr. Megaspilaeon (Hal.); Elis: in silva Kapellis pr. Lala (Weiss). — Maio, Jul. ♃

○○ Corolla extus glabra.

5. **A. graeca** Bois. et Spr. diagn. IV. p. 47, Fl. or. IV. p. 230; Spreitz. in z. b. G. 1877 p. 715, 1887 p. 665; Heldr. fl. cephal. p. 53; Bald. riv. coll. bot. alb. 1896 p. 79. — *Nonnea lutea* Ch. et B. fl. pelop. p. 12, non (Lam.) Rchb.. — *Anchusa lutea* Friedr. Reise p. 277, non M. a. B. — Exsicc.: Orph. fl. gr. n. 298.

Caulibus erectiusculis, simplicibus vel ramosis, glanduloso-viscidis, pilisque longioribus saepe tuberculatis obsitis; foliis acutis, pilis setaceis, minute tuberculatis, adpressis strigulosis, radicalibus elongato-lanceolatis, basi attenuatis, caulinis semiamplexicaulibus, inferioribus oblongis oblongo-lanceolatisve, superioribus basi dilatatis, in bracteas ovato-lanceolatas, sursum decrescentes abeuntibus; racemis fructiferis laxifloris, elongatis; calycis fructiferi accreti, pedicellati, deflexi, laciniis lanceolatis, molliter glanduloso-villosis; corolla extus glabra, lutea, tubo fuscescente, calyce longiore, limbo majusculo; nuculis scrobiculato-reticulatis, areola basilari stipitata.

In saxosis montanis. Phthiotis: in silva Muntzuraki ad mt. Kukkos (Heldr.); Achaia: mt. Kyllene (Orph.); Argolis (Bois.): pr. Potamia (Friedr.); Laconia: mt. Malevo (Orph.); Messenia: pr. Andrusa (Chaub.); Cephalonia: mt. Aenos (Spreitz.); Leucas: ad monasterium Hagios Georgios (Spreitz.). — Maio, Jul. ♃

×× Calyx dense albo-villosus.

6. **A. boeotica** DC. pr. X. p. 98; Hal. in ö. b. Z. 1897 p. 286. — *A. graeca* Ung. Reise p. 128, non Bois. et Spr.. — *A. graeca v. hispidior* Bois. fl. or. IV. p. 230; Heldr. chlor. Parn. p. 24. — Exsicc.: Heldr. herb. norm. n. 1 et 1459; Orph. fl. gr. n. 859, herb. n. 490.

Caulibus erectiusculis, simplicibus vel superne ramosis, patule pilosis, eglandulosis; foliis acutis, pilis setaceis, minute tuberculatis, adpressis strigulosis, radicalibus elongato-lanceolatis, basi attenuatis, caulinis semiamplexicaulibus, inferioribus oblongis oblongo-lanceolatisve, superioribus basi dilatatis, in bracteas sursum abeuntibus; racemis fructiferis laxifloris, elongatis; calycis fructiferi accreti, pedicellati, deflexi, laciniis lanceolatis, albo-villosis, eglandulosis; corolla extus glabra, lutea vel coeruleo-lutescenti, tubo calyce longiore, limbo mediocri; nuculis scrobiculato-reticulatis, areola basilari stipitata. — *A. methaneae et A. graecae* valde affinis, ab ambabus indumento praesertim inflorescentia dense albo-villosa, eglandulosa, a priori insuper corolla glabra discrepat.

β. **versicolor** Bois. fl. or. IV. p. 231; Bald. riv. coll. bot. alb. 1896 p. 79; pro var. *A. graecae*. — Inflorescentia praeter indumentum albo-villosum, parce glandulifera. — Exsicc.: Orph. fl. gr. n. 858.

In saxosis montanis. Euboea (Ung.): mt. Telethrion (Heldr.), Dirphys (Orph.); Aetolia: ad lacum Trichonis (Reis.); mt. Parnassus (Heldr.); Boeotia (Auch.); Attica: mt. Cithaeron (Heldr.), Parnes (Orph.); mt. Xerovuni Taygeti Alagoniae (Zahn.); — β. Epirus: mt. Maria distr. Ljaskovik (Bald.); Argolis: mt. Artemision supra Argos (Heldr.); mt. Malevo Laconiae (Orph.). — Apr. Jul. ♃

7. **A. Stribrnyi** Vel. fl. bulg. p. 647. — *A. graeca* Baen. prosp. herb. europ. 1897 p. 6, non Bois. et Sprun. — Exsicc.: Baen. herb. europ. n. 9199.

Caulibus erectiusculis, simplicibus vel superne ramosis, foliisque patule hirtis, pilis glanduliferis intermixtis, praetera setis firmis tuberculatis aspera; foliis radicalibus oblongo-lingulatis, apice rotundatis, caulinis sessilibus, obtusis, inferioribus oblongis vel oblongo-linearibus, superioribus basi saepe dilatatis, sursum in bracteas abeuntibus; racemis fructiferis densiusculis, elongatis; calycis fructiferi accreti, breviter pedicellati, deflexi, laciniis lanceolatis, patule albo-setosis, glandulosis; corolla extus glabra vel tenuiter puberula, coeruleo-lutescenti, tubo calyce sublongiore, limbo mediocri; nuculis scrobiculato-reticulatis, areola basilari stipitata. — A praecedentibus indumento aspero-setoso, foliis omnibus obtusis egregie discedit, cum speciminibus bulgaricis ab auctore ipso et a Stribrny in Baen. herb. europ. n. 7270 distributis bene congruit.

β. **pelia.** — Vix glandulosa, setis mollioribus obsita, ideo minus aspera. — Exsicc.: Heldr. it. thessal. a. 1883.

In saxosis montanis. Corcyra: inter Scripero et jugum Panteleimon (Baen.); — β. Thessalia: in regione media mt. Pelion (Heldr.). — Apr. Maio. ♃

b. Nuculae parum curvatae, capiti arietis similes, sinu basilari lato, rostro horizontali.

8. **A. Sartoriana** Bois. et Heldr. diagn. ser. 2 III. p. 134, Fl. or. IV. p. 229; Haussk. symb. p. 155. — Exsicc.: Heldr. pl. fl. hellen. n. 1721.

Praeter pubem brevissimam, setis patulis basi saepissime tuberculatis hispida, glanduloso-viscidula; caulibus adscendentibus vel prostratis, simplicibus vel parce ramosis; foliis obtusis, inferioribus subrepandis, oblongis, in petiolum attenuatis, superioribus oblongo-lanceolatis, basi rotundato-cuneata sessilibus, in bracteas sursum abeuntibus; racemis densifloris; calycis fructiferi accreti, brevissime pedicellati, patuli, laciniis lanceolatis; corolla extus glabra, albida, siccatione pallide flavescenti, tubo calycem non superante, limbo mediocri; nuculis rugoso-tuberculatis, tuberculis in rugas subquinas concentricas confluentibus, areola basilari substipitata. — A sequentibus visciditate, foliis repandis, floribus pallidis et nuculis non irregulariter tuberculatis differt.

In arenosis regionis inferioris. Argolis: inter Nauplia et Tolon (Heldr.). — Apr. Maio. ♃

9. **A. Sieberi** DC. pr. X. p. 99; Raul. cret. p. 815; Bois. fl. or. IV. p. 228. — Exsicc.: Heldr. pl. cret. n. 1341; Dörfl. pl. cret. n. 69.

Praeter pubem brevissimam, setis patulis, basi saepissime tuberculatis, aspera, cinerea, eglandulosa; caulibus adscendentibus vel prostratis, subsimplicibus; foliis oblongis oblongo-lanceolatisve, obtusis, radicalibus basi attenuatis, superioribus basi subdilatata semiamplexicaulibus, sursum in bracteas abeuntibus; racemis fructiferis laxifloris; calycis fructiferi accreti, breviter pedicellati, deflexi, laciniis lanceolatis; corolla extus glabra, coerulescente, tubo fusco, calyce duplo longiore, limbo luteolo, parvo; nuculis irregulariter rugoso-tuberculatis, areola basilari substipitata. — Sequenti affinis, ab ea indumento rariore, magis patulo, aspero, e setis validioribus constante, foliis brevioribus sublatioribusque, corolla versicolori, tubo longiore discedens.

In arenosis praesertim maritimis Creta: ad promontorium Maleka (Sieb.), Acroteri, pr. Hagia Triada, Francocastello, Males, mt. Lassiti (Raul.), pr. Anatoli distr. Hierapetra (Leon.). — Mart. Majo ♃.

10. **A. tinctoria** L. sp. p. 132; Ch. et B. exp. p. 62, Fl. pelop. p. 12; Link in Linnaea IX. p. 578; Friedr. Reise p. 266 et 273; (*Lithospermum*); Tausch in Flora 1824 p. 234; Clem. sert. p. 68; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 742 cum var *incana;* Raul. cret. p. 816; Bois. fl. or. IV. p. 227; Spreitz. in z. b. G. 1887 p. 665; Gelmi in bull. soc. bot. ital. 1889 p. 451; Haussk. symb. p. 155; Fors. in bull. herb. Bois. III. p. 88; Form. in D. bot. Monat. 1898 p. 78; Heldr. fl. Aegina p. 381, chlor. Thera p. 17. — *Anchusa tinctoria* ? L. sp. ed. 2 p. 192; S. et S. pr. I. p. 116, Fl. gr. II. p. 36 t. 166; Pieri corc. fl. p. 23; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 62, Fl. pelop. p. 12; Mazz. in ant. ion. V. p. 182: Fraas fl. class. p. 162. — *Alkanna Matthioli* Tausch in Flora 1824 p. 235, („simillima *A. tinctoriae*, quacum saepe confusa, sed differt magnitudine minori, caulibus procumbentibus, foliis angustioribus, tubo corollae duplo longiore, duplove angustiore, denique hirsutie minore, non incanescente"). — Exsicc.: Heldr. herb. norm. n. 704.

Praeter pubem velutinam, setis adpressis vel patulis, basi saepe tuberculatis plus minus strigosa, canescens, eglandulosa; caulibus adscendentibus vel prostratis, subsimplicibus; foliis lanceolatis, obtusis, radicalibus basi attenuatis, superioribus basi subdilatata semiamplexicaulibus, sursum in bracteas abeuntibus; racemis fructiferis laxiusculis; calycis fructiferi accreti, breviter pedicellati, subdeflexi, laciniis lanceolatis; corolla azurea, extus glabra, tubo calycem aequante vel paulo longiore, limbo parvo; nuculis irregulariter rugoso-tuberculatis, areola basilari substipitata.

β. **Lehmani** Tin in Guss. fl. sic. syn. II. p. 791. — Magis strigosa, canescens; corollae tubus calycem excedens. — Exsicc.: Heldr. reliqu. Orph. a. 1887.

In arenosis, rupestribus aridis regionis inferioris et montanae. Attica: frequens pr. Athenas, in colle Lycabettus, Turcovuni, mt. Kerata, Pentelicon, Hymettus, ad Piraeum, Phalerum; in insula Aegina, Thera (Heldr.), Syra, Tenos (Weiss); Peloponnesus (Sibth.): in isthmo Dara Argolidis (Friedr.), mt. Taygetos (Chaub.); Creta: ad promontorium Meleka, pr. Francocastello, Candia (Raul.); Leucas: pr. Amaxichi (Spreitz.); Corcyra (Pieri), mt. Deca (Ung.), ad Scripero (Gelmi), cap Bianco (Spreitz.), pr. Menades, Sidari, Tiflo (Mazz.); — β. ad saxa Scyronica pr. Megara (Orph.). — Mart. Maio. ♃

14. Lithospermum L. gen. n. 181.

1. Sectio. *Lithodora* Griseb. spic. II. p. 85. — Corollae faux plicis destituta; nuculae ovato-triquetrae, laeves vel minntissime tuberculatae. Frutices vel suffrutices.

1. **L. Zahnii** Heldr. herb. gr. norm. n. 1360; Hal. in z. b. G. 1899 p. 190. — *L. fruticosum* β. L. sp. p. 133, nam planta ex insula Samos haud dubie huc spectat; S. et S. pr. I. p. 114, Fl. gr. II. p. 52 t. 161. — *L. rosmarinifolium* Bois. fl. or. IV. p. 220, non Ten. fl. nap. III. p. 178, quod foliis latioribus, minus revolutis, subtus canis, supra adpresse pubescentibus et corolla extus pubescente specifice differt. — Exsicc.: Heldr. l. c.

Fruticosum, ramosissimum; ramis patulis, vetustis denudatis, novellis crebre foliosis, patule hirsutis; foliis lineari-lanceolatis, obtusis, supra viridibus, nitidis, setis basi tuberculatis, praesertim ad marginem valde revolutam obsitis, subtus pallidioribus, adpresse pilosis; floribus 5—10 inter folia superiora sessilibus, congestis; calycis laciniis lineari-lanceolatis, adpresse strigulosis; corolla azurea, glabra, calyce duplo longiore, lobis majusculis; nuculis laevibus. — Frutex sempervirens, foliis 3—4 cm. longis, eis *Roris marini* similibus.

Ad rupes excelsas in mt. Selitza inter Mantinia et Kampos et pr. Sotirianika Laconiae boreo-occidentalis (Zahn.); a Sibthorpio in montosis Graeciae et Archipelagi insulis indicatur. — Mart. Apr. ♃

2. **L. hispidulum** S. et S. pr. I. p. 114, Fl. gr. II. p. 53 t. 162; Raul. cret. p. 815; Bois. fl. or. IV. p. 219. — Exsicc.: Sint. et Rigo it cypr. n. 102 (Cyprus).

Suffruticosum, intricatim ramosissimum; ramis vetustis denudatis, novellis crebre foliosis, adpressissime cano-sericeis; foliis elliptico-oblongis, obtusis, obscure viridibus, opacis, setulis adpressissimis, ad marginem subrevolutum et nervum validioribus, basi tuberculatis obsitis; floribus 1—4 inter folia superiora sessilibus, congestis; calycis laciniis lineari-lanceolatis, adpresse strigulosis; corolla purpureo-coerulescente, calyce duplo longiore, lobis mediocribus; nuculis minutissime tuberculatis. — Dumi semipedales, folia parva 5—12 mm. longa, flores eis praecedentis minores.

In saxosis maritimis Cretae, in littore meridionali inter Hagios Paulos et Hagia Rumeli (Heldr.); occurrit quoque in insula Karpathos (Pichl.). — Mart. Apr. ♃ N. v.

2. Sectio. *Eulithospermum* Griseb. l. c. — Corollae faux sub lobis gibbis vel plicis velutinis instructa; nuculae ovatae, laevissimae. Herbae perennes.

3. **L. purpureocoeruleum** L. sp. p. 132; S. et S. pr. I. p. 114; Ch. et B. fl. pelop. p. 12; Mazz. in ant. ion. V. 180; Friedr. Reise p. 281; Ung. Reise p. 128; Spreitz. in z. b. G. 1877 p. 715; Bois. fl. or. IV. p. 218, suppl. p. 352; Haussk. symb. p. 155; Hal. in ö. b. Z. 1896 p. 16. — Icon: Jacq. fl. austr. t. 14. — Exsicc.: Orph. fl. gr. n. 1158.

Caulibus patule hispidis, florigeris erectis, apice 2—3 fidis, sterilibus diffusis; foliis lanceolatis, acutiusculis, adpresse hirtis; floribus laxiuscule racemosis, breviter pedicellatis; calycis laciniis anguste linearibus; corolla violacea tandem azurea, extus hirta, calyce duplo longiore, limbo amplo; nuculis ovato-globosis, albis.

In nemorosis, silvaticis, praesertim montanis, rare. Thessalia: pr. Malakasi, Kalabaka, Neuropolis, Orman Magula (Haussk.), mt. Pelion pr. Makrinitza et Portaria (Heldr.); Euboea: mt. Telethrion (Heldr.), pr. Achmet Aga (Orph.); Achaia; pr. Patras (Reiser); Arcadia: pr. Andrizena, mt. Diaforti (Friedr.); Messenia: pr. Andrusa (Chaub.); Corcyra: pr. Castagna, Calefactiones (Mazz.), Bragagniotica (Ung.), mt. Deca (Spreitz.). — Apr. Maio. ♃

4. **L. officinale** L. sp. p. 132; Bois. fl. or. IV. p. 218; Form. in Ver. Brünn 1896 p. 61. — Icon: Fl. dan. t. 1084.

Caule firmo, erecto, scabro, superne ramosissimo; foliis lanceolatis, acutis, scaberrimis; floribus numerosissimis, subsessilibus, racemos longos formantibus; calycis laciniis lanceolatis; corolla albido-flavida, extus hirta, calycem parum superante, limbo parvo; nuculis ovato-globosis, albis.

Indicatur in valle Tempe Thessaliae (Form.) et in Peloponneso (Bois.). — Maio, Jul. ♃ N. v.

3. Sectio. *Rhytispermum* Link handb. II. p. 579. — Corollae faux plicis longitudinalibus, velutinis obsita; nuculae ovato-triquetrae, tuberculatae, scrobiculatae vel echinatae. Herbae annuae.

a. Flores albi vel coerulei.

α. Nuculae ovoideo-trigonae, acutae, scrobiculatae.

5. **L. arvense** L. sp. p. 132; S. et S. pr. I. p. 113; Ch. et B. exp. p. 61, Fl. pelop. p. 12; Mazz. in ant. ion. V. p. 180; Marg. et R. fl. Zante p. 68; Friedr. Reise p. 267; Weiss in z. b. G. 1869 p. 742; Bois. fl. or. IV. p. 216; Heldr. fl. cephal. p. 53, fl. Aegina p. 381; Haussk. symb. p. 155; Form. in Ver. Brünn 1896 p. 61; Bald. riv. coll. bot. Alb. 1896 p. 79. — *L. Leithneri* Heldr. et Sart. in

Heldr. herb. norm. n. 146. — Icon: Fl. dan. t. 456. — Exsicc.: Orph. fl. gr. n. 16; Heldr. herb. norm. n. 146 et 146 b, in Baen. herb. europ. n. 4186; Sint. it. thessal. n. 39; Dörfl. fl. gr. n. 458.

Adpresse strigulosum; caulibus erectis vel adscendentibus, saepe a basi ramosis; foliis inferioribus obovato-oblongis, in petiolum attenuatis, superioribus lanceolatis; racemis fructiferis laxis; pedicellis fructiferis brevissimis, vix incrassatis; bracteis flores superantibus; corollae albae, rarius coerulescentis, tubo hirto, calyce non longiore; nuculis scrobiculatis et acute tuberculatis.

β. **Sibthorpianum** Griseb. spic. II. p. 86; Clem. sert. p. 68; Raul. cret. p. 815; Bois. fl. or. IV. p. 216; Haussk. symb. p. 155; pro sp. — *L. tenuiflorum* S. et S. fl. gr. II. p. 50 t. 159, non L. — Canescens, a collo ramosum, ramis brevibus, prostratis; nuculis scrobiculatis, subtuberculatis, minoribus. — Exsicc.: Orph. fl. gr. n. 1116.

In arvis, inter segetes regionis inferioris et montanae. Epirus: mt. Micikeli (Bald.); Thessalia: mt. Said Pascha (Form.), pr. Volo (Sint.); Attica: ad Phalerum (Friedr.), pr. Athenas, mt. Pentelicon, Hymettus, insula Aegina (Heldr.); insula Hydra (Pichl.), Syra (Weiss), Melos (Leon.); Messenia: pr. Methone, Stenicleros (Chaub.); Zante (Marg.); Cephalonia: pr. Francata (Heldr.); Corcyra (Mazz.); — β. Attica: mt. Parnes, Pentelicon, Hymettus (Heldr.), ad Phalerum (Clem.); Creta: pr. Canea (Raul.). — Febr. Maio. ⊙

6. **L. incrassatum** Guss. pr. fl. sic. I. p. 211; Friedr. Reise p. 279; Raul. cret. p. 815; Bois. fl. or. IV. p. 217; Heldr. fl. cephal. p. 53; Hal. Beitr. fl. Epir. p. 32; Haussk. symb. p. 155. — Exsicc.: Dörfl. fl. gr. n. 344.

Adpresse strigulosum; caulibus erectis vel decumbentibus, saepe a basi ramosis; foliis inferioribus obovatis, in petiolum attenuatis, superioribus lanceolatis linearibusve; racemis fructiferis laxis; pedicellis fructiferis valde incrassatis, obconicis, apice calyci aequilatis; corollae coerulescentis vel albae, tubo hirto, calyce non longiore; nuculis scrobiculatis, subtuberculatis.

In herbidis montanis in regionem alpinam adscendens. Epirus: mt. Peristeri (Hal.); Thessalia: pr. Pharsalus (Haussk.); Aetolia: mt. Korax (Leon.); Attica: mt. Parnes (Sprun.); Achaia: mt. Panachaicon (Heldr.); Arcadia: pr. Tripoliza (Friedr.); Laconia: mt. Taygetos (Bois.); Creta: pr. Askyphos, Theodori, Mavri in mt. Sphacioticis, mt. Ida, Lazaro in Lassiti (Raul.); Cephalonia: mt. Aenos (Heldr.). — Apr. Jul. ⊙

β. Nuculae ovato-trigonae, lateraliter saepe bigibbae, in rostrum breve contractae.

7. **L. tenuiflorum** L. fil. suppl. p. 130; Fraas fl. class. p. 162; Clem. sert. p. 69; Bois. fl. or. IV. p. 217; Haussk. symb. p. 155. — Icon: Jacq. ic. rar. II. t. 313. — Exsicc.: Orph. fl. gr. n. 15; Heldr. herb. norm. n. 472, in Baen. herb. europ. n. 2401.

Adpresse strigulosum; caule erecto, superne ramoso; foliis inferioribus oblongo-spathulatis, breviter petiolatis, superioribus oblongo-linearibus; racemis fructiferis laxiusculis; pedicellis fructiferis brevissimis, non incrassatis; bracteis flores parum superantibus; corollae coeruleae vel albae, tubo hirto, calyce sublongiore; nuculis tuberculatis.

In collibus et cultis, rare. Euboea: pr. Chalkis (Sprun.); Attica: pr. Eleusis (Haussk.), Athenas (Orph.), mt. Hymettus (Heldr.); Argolis (Sprun.). — Mart. Maio ⊙.

b. Flores lutei.

8. **L. apulum** L. sp. p. 131; Vahl. symb. II. p. 32; S. et S. pr. I. p. 118, Fl. gr. II. p. 49 t. 158; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 61, Fl. pelop. p. 12; Friedr. Reise p. 270; Fraas fl. class. p. 162; Weiss in z. b. G. 1869 p. 742; Raul. cret. p. 815; Bois. fl. or. IV. p. 218; Haussk. symb. p. 155; Heldr. fl. Aegina p. 381. — Exsicc.: Dörfl. pr. cret. n. 52.

Patule hispidum; caule erecto, apice ramoso; foliis oblongo-linearibus, inferioribus in petiolum attenuatis; racemis subcorymbosis, densis; pedicellis fructiferis brevissimis; bracteis flores superantibus; corollae tubo pubescente, calycem superante; nuculis ovato-acuminatis, gibbosis, tuberculatis, dorso laevibus.

In siccis apricis, saxosis. Attica: pr. Athenas (Sprun.); Eleusis (Haussk.); Corinthia: pr. Kalamaki, Acrocorinthus (Haussk.); Achaia: pr. Megaspilaeon (Hal.); Messenia: pr. Arcadia, mt. Kupe, pr. Messene (Chaub.); insula Aegina: ad monasterium (Heldr.), promontorium Perdicca (Friedr.); Cyclades: Syra (Weiss), Melos, Kimolos (Leon.); Creta: cap Meleka (Sieb.), ad Canea, Akroteri, Malaxa, Francocastello (Raul.), Males distr. Hierapetra (Leon.). — Mart. Jun. ⊙

15. Myosotis L. gen. n. 180.

Dispositio specierum:

a. Calycis pili adpressissimi, nunquam uncinati.
  α. Perennes vel biennes; corollae limbus planus.
    1. **M. palustris** L. 2. **M. lingulata** Lehm.
  β. Annuae; corollae limbus concavus.
    × Racemi ebracteati.
      ○ Planta parce et adpressissime hirta.
        3. **M. sicula** Guss.
      ○○ Planta inferne subpatentim, superne adpressiuscule hirta.
        . Racemi fructiferi elongati, laxiusculi.
          4. **M. idaea** Bois. et Heldr.
        .. Racemi fructiferi breves, imbricatim densiflori.
          5. **M. littoralis** Stev.
    ×× Racemi bracteati
        6. **M. pusilla** Lois.
b. Calycis pili patuli, baseos uncinati.
  α. Pedicelli fructiferi erecti vel patuli.

× Racemi aphylli.
○ Corollae limbus planus.
7. M. olympica Bois. **8. M. silvatica** Hoffm.
○○ Corollae limbus concavus.
. Corollae tubus calyce aequilongus.
9. M. arvensis L. **10. M. collina** Hoffm.
.. Corollae tubus calyce duplo longior.
11. M. versicolor Pers.
×× Racemi inferne foliati.
12. M. stricta Link.
β. Pedicelli fructiferi refracti.
13. M. refracta Bois.

a. Calycis pili adpressissimi, nunquam uncinati.

α. Perennes vel biennes; corollae limbus planus.

1. **M. palustris** L. sp. p. 131 pro var. *M. scorpioides*; With. arr. brit. pl. II. p. 225; Mazz. in ant. ion. IV. p. 960; Bald. riv. coll. bot. alb. 1896 p. 79. — Icon: Rchb. t. 119.

Caule erecto vel adscendente, angulato, ramoso, glabriusculo vel adpresse, inferne saepius patule hirto, foliis oblongo-lanceolatis lanceolatisve, acutiusculis, adpresse hirtis; racemis nudis, laxifloris; pedicellis post anthesin horizontalibus, fructiferis calyce aperto longioribus; corollae coeruleae, majusculae limbo 6—8 mm. diametro lato, plano; stylo calyci aequilongo; nuculis ovatis. — Occurit interdum floribus albis (*M. alba* Mazz. in ant. ion. IV. p. 960).

α. **typica.** — Caule crassiusculo, robustiore, adscendente, patentipiloso. — Exsicc.: N. v.

β. **strigulosa** Rchb. in Sturm. deutschl. fl. h. 42 pro sp.; Gr. et Godr. fl. fr. II. p. 529. — Gracilior, erecta, glabriuscula vel adpresse pilosa. — Exsicc.: Heldr. it. gr. septentr. a. 1879.

In humidis, ad fossas regionis inferioris et montanae, rarissime. Indicatur in Corcyra: ad lacum Stavropotamo et S. Onufrio (Mazz.); dein in Epiro: ad lacum Janina (Bald.); — β. in subalpinis mt. Oeta Phthiotidis l. c. Livadies (Heldr.). — Maio, Jul. ♃

2. **M. lingulata** Lehm. asper. p. 110; Hal. Beitr. fl. Aetol. p. 9. — *M. caespitosa* Schultz. fl. starg. suppl. p. 11. — Huc probabiliter: *M. commutata* Roem. et Sch. syst. IV. p. 102; Mazz. in ant. ion. IV. p. 960. — Icon: Rchb. t. 120. — Exsicc.: Hal. it. gr. sec. a 1893.

Adpresse hirta vel glabriuscula; caule flaccido, tereti, longe ramoso; foliis oblongo-linearibus vel linguaeformibus, obtusiusculis; racemis laxifloris, basi plerumque paucifoliatis; pedicellis post anthesin horizontalibus, florum infimorum demum calyce aperto 2—3 plo longioribus; corollae pallide coeruleae, parvae, limbo 2—4 mm. diametro lato, plano; stylo brevissimo; nnculis late ovatis, basi truncatis.

In paludibus Aetoliae: ad lacum Lysimachiae pr. Sikya (Heldr.), in palude pr. Krioneri ad radices mt. Chalkis ad sinum Patranum (Hal.); Corcyra: mt. Abramo (Mazz.). — Maio, Jun. ⊙

β. Annuae; corollae limbus concavus.

× Racemi ebracteati.

○ Planta parce et adpressissime hirta.

3. **M. sicula** Guss. syn. I. p. 214; Ch et B. exp. p. 61, Fl. pelop. p. 12; Bois. fl. or. IV. p. 235; Haussk. symb. p. 155; Bald. riv. coll. bot. alb. 1896 p. 79. — *M. micrantha* Guss. pr. I. p. 207, non Pall. — Icon: Rchb. t. 120. — Exsicc.: Bald. it. alb. epir. IV. p. 86.

Caule erectiusculo, valde ramoso, inferne saepe radicante; foliis oblongo-lanceolatis, obtusis; racemis nudis, demum laxis; pedicellis patentibus deflexisque, fructiferis inferioribus calyce parum longioribus; calycis fructiferi dentibus obtusis, subconniventibus; corollae parvae, coeruleae limbo concavo. — Praecedenti affinis, sed annua, racemi basi non foliati, pedicelli breviores, corolla minor limbo concavo.

In pratis humidis. Epirus: pr. Janina (Bald.); Thessalia: pr. Sermeniko, Neuropolis (Haussk.); Argolis: pr. Tyrinthum (Orph.); Messenia: inter Methone et Corone, mt. Taygetos (Chaub.); Cycladum insula Naxos (Chaub.). — Maio. Jun. ⊙

○○ Planta inferne subpatentim, superne adpressiuscule hirta.

. Racemi fructiferi elongati, laxiusculi.

4. **M. idaea** Bois. et Heldr. diagn. XI. p. 21, Fl. or. IV. p. 236; Raul. cret. p. 816; Haussk. symb. p. 155; Fors. in bull. herb. Bois. III. p. 88. — *M. cadmea* Hal. Beitr. fl. Achaia p. 26, non Bois. — Huc forsan: *M. cretica* Ung. Reise p. 127, vix Bois. et Heldr.; — nec non *M. stricta* Weiss in z. b. G. 1869 p. 742, vix Lk. —

Caulibus solitariis vel plus minusve numerosis, erectis vel diffusis, pumilis vel elongatis, simplicibus vel jam a basi ramosis; foliis radicalibus obovato-spathulatis, caulinis oblongo-linearibus; pedicellis fructiferis erecto-patentibus vel patenti-subdeflexis, saepe incrassatis, calyci aequilongis vel longioribus; calycis fructiferi dentibus acutiusculis, erectis, non conniventibus; corollae parvae, coeruleae vel albae limbo concavo.

α. **typica.** — Corolla exigua, limbo tubum non aequante; pedicelli fructiferi plerumque fere horizontales, calyci aequilongi vel sesquilongiores, parum incrassati. — Exsicc.: Orph. fl. gr. n. 218.

β. **boeotica** Reut. in Orph. fl. gr. n. 1002. — *M. idaea v. grandiflora* Bois. fl. or. IV. p. 236, p. p. — *M. idaea v. cadmea* Bois. l. c., quoad pl. boeoticam; Heldr. fl. Aegina p. 381; non *M. cadmea* Bois. diagn. XI. p. 122, quae ex diagnosi autoris et speciminibus byzantinis (Aznav. pl. de Turqu. a. 1896) corollae limbo amplo explanato, eo *M. palustris* aequilato, speciem propriam distinctissimam

sistit. — Corolla parva, limbo tubum aequante vel superante; pedicelli fructiferi horizontales vel subdeflexi, calyce sesquilongiores, vix incrassati. — Exsicc.: Orph. l. c.; Heldr. herb. norm. n. 158 et 865.

γ. **Kiesenwetteri** Heldr. in sched. — Huc forsan: *M. stricta* Friedr. Reise p. 268, vix Link. — Corolla parva, limbo tubum subaequante; pedicelli fructiferi erecti vel erecto-patuli, calyce aequilongi, manifeste incrassati. — Exsicc.: Orph. fl. gr. n. 219; Heldr. in Baen. herb. europ. n. 4192.

In collibus saxosis regionis inferioris et montanae. Attica: pr. Athenas (Orph.), mt. Kerata, Pentelicon (Haussk.), Hymettus, in colle Ardetto, Lycabetto, mt. Cithaeron, insula Aegina (Heldr.); Acrocorinthus (Haussk.); Achaia: mt. Panachaicon (Hal); Laconia: mt. Malevo (Orph.); Cycladum insula: Cythnos (Tunt.), Syra (Fors.); Creta: pr. Askyphos, Anopolis, mt. Ida, Lassiti (Raul.). — β. Thessalia: pr. Velestinos, mt. Pelion (Heldr.); Attica: mt. Parnes, Pentelicon, Hymettus (Heldr.); Boeotia: mt. Helicon (Orph.); mt. Parnassus (Guicc.); Euboea: mt. Dirphys (Orph.); Laconia: mt. Malevo (Orph.); — γ. Attica: mt. Pentelicon (Heldr.); Cycladum insula: Cythnos (Tunt.), Tenos (Leon.). — Mart. Jun. ⊙

.. Racemi fructiferi breves, imbricatim densiflori.

5. **M. littoralis** Stev. cat. hort. gorenk. a. 1812 p. 27; Clem. sert. p. 69; Bois. fl. or. IV. p. 236; Heldr. fl. Aegina p. 381. — *M. stricta* Friedr. Reise p. 270, sec. Heldr. fl. Aegina p. 381, non Link. — Huc probabiliter: *M. pusilla* Clem. sert. p. 69, ex loco. — Exsicc.: Orph. fl. gr. n. 220; Heldr. herb. norm. n. 554.

Nana, ad folia patule, secus inflorescentiam adpresse hispida; caulibus simplicibus vel a collo ramosis; foliis radicalibus aggregatis, obovatis obtusis, caulinis oblongis; pedicellis brevissimis, strictis, demum incrassatis, calyce aequilongis; calycis fructiferi dentibus acutiusculis, erectis, non conniventibus; corollae exiguae, coeruleae vel albae, limbo concavo, tubum non aequante. — Plantula 1—2 pollicaris, a collo saepe ramosissima.

In arenosis praesertim maritimis, rarissime. Attica: pr. Stadion (Clem.), ad Phalerum, insula Aegina (Heldr.). — Febr. Maio. ⊙

×× Racemi bracteati.

6. **M. pusilla** Lois. in Desv. journ. II. p. 260, t. 8; Bois. fl. or. IV. p. 236; Heldr. fl. cephal. p. 53. — *M. cretica* Bois. et Heldr. diagn. XI. p. 121; Raul. cret. p. 816. — *M. littoralis* Ung. Reise p. 127, non Stev. — *M. littoralis v. aenesia* Heldr. apud Spreitz. in z. b. G. 1877 p. 715. — *M. aenesia* Heldr. fl. cephal. n. 3563. — Icon: Mor. fl. sard. t. 96. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889; Dörfl. pl. cret. n. 27.

Pumila, ad folia patule, secus inflorescentiam adpresse hispida; caulibus simplicibus vel a collo ramosis; foliis radicalibus obovatis, obtusis, caulinis oblongis; racemis fructiferis laxiusculis; pedicellis fructiferis erecto-patulis, vix incrassatis, calyci aequilongis vel sublongi-

oribus; calycis fructiferi dentibus acutiusculis, erectis, non conniventibus; corollae exiguae, coeruleae vel albae, limbo concavo, tubum non aequante. — Plantula nana, praecedenti affinis, ab ea racemis fructiferis laxiusculis, bracteatis, pedicellis longioribus, erecto-patulis diversa.

In apricis montosis, in regionem abietinam adscendens. Cephalonia: mt. Aenos (Ung.); Creta: mt. Volakia (Raul.), Ida, Lassiti (Heldr.); insula Melos (Leon.). — Mart. Maio. ⊙

b. Calycis pili patuli, baseos uncinati.

a. Pedicelli fructiferi erecti vel patuli.

× Racemi aphylli.

○ Corollae limbus planus.

7. **M. olympica** Bois. diagn. IV. p. 50; Haussk. symb. p. 156. — *M. alpestris* Hal. Beitr. fl. Epir. p. 32; Bald. riv. coll. bot. Alb. 1895 p. 58; Form. in Ver. Brünn 1896 p. 60, non Schmidt fl. bohem. III. p. 26, quae calyce pilis adpressiusculis, paucis uncinatis et nuculis rotundato-ovatis parum discedit. — Exsicc.: Orph. herb. n. 713; Sint. it. thessal. n. 641.

Perennis; adpressiuscule hispida; caulibus pumilis, parce ramosis; foliis crassiusculis, radicalibus oblongo-spathulatis, in petiolum attenuatis, caulinis oblongis, sessilibus; racemis fructiferis laxis; pedicellis fructiferis erecto-patulis, calyce subbrevioribus; calyce pilis patulis uncinulatis albo-strigoso, laciniis fructiferis erectis; corollae majusculae azureae limbo plano, tubo longiore; nuculis oblongis, compressis, utraque facie subcarinatis. — Planta nostra ab ea ex Olympo bithynico pedicellis fructiferis minus strictis parum discedit, ab affini *M. pyrenaica* Pourr. indumento breviore et rariore tantum, a *M. alpestri* calycis pilis patulis plerisque uncinatis, nec adpressiusculis paucis uncinatis differt.

In regione subalpina et alpina. Epirus: mt. Peristeri, Tsumerka (Hal.), Kakardista (Bald.); Thessalia: mt. Karava (Haussk.); Doris: mt. Kiona (Hal.); Peloponnesus: pr. Zatuna (Orph.). — Jun. Jul. ♃

8. **M. silvatica** Hoffm. deutschl. fl. I. p. 85; Bois. fl. or. IV. p. 237; Hal. in z. b. G. 1888 p. 760, Beitr. fl. Epir. p. 32, Beitr. fl. Thessal. p. 16; Heldr. chlor. Parn. p. 24; Haussk. symb. p. 155; Form. in Ver. Brünn 1896 p. 60. — Icon: Rchb. t. 121. — *M. cyanea* Bois. et Heldr. fl. gr. exs. a. 1855, forma montium elatiorum, racemis abbreviatis, pedicellis brevioribus et floribus minoribus, quae sine limites in typum transit. — *M. alpestris* Ung. Reise p. 127, non Schmidt. — Exsicc.: Heldr. herb. norm. n. 157 et 157b; Orph. fl. gr. n. 483 et 1003; Sint. it. thessal. n. 638; Dörfl. fl. gr. n. 227.

Biennis; adpressiuscule hispida; caulibus saepius elatis, parce ramosis; foliis tenuibus, radicalibus oblongo-obovatis, in petiolum attenuatis, caulinis oblongis, sessilibus; racemis fructiferis laxis; pedicellis fructiferis patentibus, calycem aequantibus vel eo duplo longioribus; calyce pilis patulis uncinulatis albostrigoso, laciniis fructiferis erectis; corollae majusculae vel mediocris, azureae, rarius albae limbo plano, tubo longiore; nuculis ovatis, submarginatis, altera facie obscure marginatis. — Prae-

cedenti saepius elatior, folia tenuiora, indumentum mollius, habitus laxior, racemi elongati, pedicelli longiores magis patuli, calyces minores.

In silvaticis et umbrosis humidis regionis montanae et subalpinae. Epirus: pr. Mazuki (Hal.); Thessalia: pr. Chaliki, mt. Oxya (Hal.), Zygos (Form.), pr. Witomo (Sint.), mt. Ghavellu, ad Gionscala, Sermeniko, Neuropolis, Korona (Haussk.), mt. Ossa, Pelion (Heldr.); Euboea: mt. Dirphys (Orph.); Aetolia: mt. Korax (Heldr.); mt. Kiona (Hal.), Parnassus, Helicon (Orph.); Attica: mt. Cithaeron, Parnes (Heldr.); Achaia: mt. Panachaicon (Heldr.), Kyllene (Orph.), Taygetos (Zahn). — Maio, Aug. ⊙

Obs. *M. Mathildae* Haussk. symb. p. 158. — *M. silvatica v. Mathildae* Form. in Ver. Brünn 1897 p. 44. — „Ex affinitate *M. silvaticae*, sed habitu expanso, caulibus a basi ramosis duris non herbaceis, teretibus non angulosis, floribus minoribus, nuculis majoribus, ovato-oblongis, nec ovatis pallidioribus discedens," mihi ignota est, sed ex diagnosi ad ejus formas pertinere videtur. — Indicatur in mt. Ghavellu Pindi dolopici (Haussk.).

○○ Corollae limbus concavus.

. Corollae tubus calyce aequilongus.

9. **M. arvensis** L. sp. p. 131 pro var. *M. scorpioides;* Roth bot. Abhandl. p. 20; Urv. enum. p. 19; Ch. et B. exp. p. 60, Fl. pelop. p. 12; Mazz. in ant. ion. IV. p. 960; Friedr. Reise p. 269; Hal. Beitr. fl. Epir. p. 32, Beitr. fl. Achaia p. 26; Haussk. symb. p. 158; Bald. riv. coll. bot. alb. 1896 p. 79. — *M. scorpioides* L. sp. p. 131 p. p.; S. et S. pr. I. p. 112 p. p.; ? Mazz. l. c. p. 958. — *M. intermedia* Link en. hort. berol. I. p. 164; ? Mazz. l. c.; Bois. fl. or. IV. p. 239. — Icon: Rchb. t. 122. — Exsicc.: Bald. it. alb. epir. IV. n. 40; Dörfl. fl. aeg. n. 116.

Biennis, pilis patulis hispida; caulibus erectis, elatis, parce ramosis; foliis inferioribus spathulato-oblongis, in petiolum attenuatis, superioribus oblongo-lanceolatis, sessilibus; racemis laxifloris; pedicellis fructiferis patentibus, calyce duplo longioribus; calyce pilis uncinatis obsito, fructifero subclauso; corolla parva, coerulea.

In cultis et arenosis, praesertim montanis. Epirus: ad Kalentini pr. Arta (Hal.), mt. Micikeli pr. Janina (Bald.); Thessalia: in oropedio Neuropolis pr. Pezula (Heldr.), pr. Korona (Haussk.); Attica: mt. Parnes (Heldr.); Peloponnesus (Chaub.): mt. Panachaicon (Hal.), Malevo (Orph.); insula Aegina (Friedr.), Naxos in mt. Koroni (Leon.), Melos (Urv.); Corcyra: pr. Carocollo, Missostrati, Stratie, mt. Abramo (Mazz.). — Maio, Jul. ⊙

10. **M. collina** Hoffm. deutschl. fl. p. 61; Mazz. in ant. ion. IV. p. 960; Hal. in ö. b. Z. 1897 p. 97; Heldr. fl. Aegina p. 381. — *M. hispida* Schlecht. mag. nat. Berlin VIII. p. 230; Spreitz. in z. b. G. 1877 p. 715; Bois. fl. or. IV. p. 239, suppl. p. 353; Heldr. fl. cephal. p. 54. — Icon: Fl. dan. t. 2284. — Exsicc.: Sint. it. thessal. n. 40 b.

Annua, pilis patulis hispida; caulibus erectiusculis, gracilibus, saepius a medio ramosis; foliis inferioribus spathulato-oblongis, in petiolum attenuatis, superioribus oblongis, sessilibus; racemis laxifloris; pedicellis fructiferis patentibus, calyce aequilongis vel brevioribus; calyce pilis uncinatis obsito, fructifero aperto; corolla exigua, coerulea.

β. **grandiflora** Bois. et Heldr. diagn. XI. p. 123, Fl. or. IV. p. 239; Raul. cret. p. 816. — Corollae limbus major calycem manifeste superans. — Exsicc.: N. v.

γ. **gracillima** Losc. ser. inconf. p. 72; Willk. ill. hisp. II. t. 162; pro sp.; Haussk. symb. p. 158. — Racemi valde elongati, laxissimi; calyces minores. — Exsicc.: Heldr. et Hal. fl. Sporad. a. 1896. —

In collibus herbidis regionis inferioris et montanae. Thessalia: pr. Kalabaka (Sint.), Velestinos (Heldr.); Euboea (Orph.); Attica: mt. Parnes, pr. Kephissia, Buliasmene, mt. Hymettus, insula Salamis, Aegina (Heldr.); Cycladum insula Cythnos (Tunt.); Creta: mt. Lakus (Rev.); Cephalonia: mt. Aenos (Heldr.); Corcyra: mt. Deca (Spreitz), pr. Rodostamo (Mazz.); — β. Creta: pr. Platania, Canea, Murnies, Nerokuru, Tuzla (Raul.); — γ. Thessalia: mt. Zygos (Haussk.); Sporadum insula Scopelos (Leon.); mt. Hymettus et Kerata Atticae (Haussk.); Boeotia: mt. Helicon (Heldr.). — Mart. Maio. ⊙

.. Corollae tubus demum calyce duplo longior.

11. **M. versicolor** Pers. syn. I. p. 156 pro var. M. *arvensis*; Schlecht. mag. nat. Berlin VIII. p. 230; Mazz. in ant. ion. IV. p. 960. — Icon: Rchb. t. 124.

Annua, pilis patulis hispida; caulibus erectis, ramosis; foliis inferioribus spathulato-oblongis, in petiolum attenuatis, superioribus lineari-oblongis, sessilibus; racemis laxifloris; pedicellis fructiferis erecto-patulis, calyce brevioribus; calyce pilis uncinatis obsito, fructifero clauso; corolla parva, primo lutea, deinde coerulea, demum violacea.

Corcyra: pr. urbem (Mazz.); a recentioribus non lecta. — Mart. Maio ⊙ N. v.

×× Racemi inferne foliati.

12. **M. stricta** Link. en. hort. berol. I. p. 164; Hal. Beitr. fl. Epir. p. 32; Bald. riv. coll. bot. Alb. 1896 p. 80. — Icon: Fl. dan. t. 2285. Exsicc.: Hal. it. gr. II. a. 1893.

Annua, pilis patulis hispida; caulibus erectis, parce ramosis; foliis inferioribus spathulato-oblongis, in petiolum attenuatis, superioribus oblongis, sessilibus; racemis laxifloris; pedicellis fructiferis erecto-patulis, calyce brevioribus; calyce fructifero clauso, ad medium usque pilis uncinatis obsito; corolla exigua, coerulea, limbo concavo.

In herbidis regionis montanae et subalpinae, rarissime. Epirus: mt. Smolika (Bald.), mt. Tsumerka supra Theodoriana (Hal.). — Jun. Jul. ⊙

β. **Pedicelli fructiferi refracti.**

13. **M. refracta** Bois. voy. esp. p. 433 t. 125, Fl. or. IV. p. 240; Hal. Beitr. fl. Achaia p. 27; Haussk. symb. p. 158; Bald. riv. coll. bot. alb. 1895 p. 58. — Huc spectare videtur: *M. hispida v. pygmaea* Hal. Beitr. fl. Achaia p. 27 (specimina juvenilia, minutissima). — Exsicc.: Heldr. pl. fl. hellen. a. 1878 et 1895, it. thessal. IV. a. 1885. —

Annua, pilis patulis hispida; caulibus erectiusculis, saepe a basi ramosis; foliis oblongis; racemis laxiusculis; pedicellis brevissimis; calyce pilis uncinatis obsito, fructifero elongato; corolla minuta, coerulea, limbo concavo.

In lapidosis regionis subalpinae et alpinae, rarissime. Epirus: mt. Olycika (Bald.); Thessalia: mt. Zygos (Heldr.), Karava (Haussk.); Attica: mt. Parnes (Heldr.); Achaia: mt. Chelmos (Hal.); Creta: mt. Ida (Heldr.). — Maio, Jul. ⊙

Obs. Quid sit *M. nana* S. et S. pr. I. p. 112; Ch. et B. exp. p. 61, Fl. pelop. p. 12; in Argolide et Arcadia indicata, nescio; certe non Vill. fl. delph. II. p. 459, quae in Graecia non crescit.

4. Subtribus. *Cynoglosseae* DC. pr. X. p. 117.

16. Omphalodes Tourn. inst. t. 58.

1. **O. verna** Moench meth. p. 420. — *Cynoglossum omphalodes* L. sp. p. 135; Mazz. in ant. ion. V. p. 186. — Icon: Rchb. t. 125.

Pilis adpressis parce hirta, viridis; caulibus adscendentibus vel erectis; foliis inferioribus longe petiolatis, ovatis, acuminatis, basi cordatis, superioribus breviter petiolatis, ovato-lanceolatis; floribus in axillis foliorum superiorum breviter racemosis; pedicellis fructiferis elongatis, calyce multo longioribus, cernuis vel recurvis; calycis adpresse strigosi, fructiferi aucti, laciniis ovato-lanceolatis, acuminatis; corollae coeruleae limbo tubo multo longiore; nuculis laevibus, margine introflexo, integro.

In herbosis pr. Signes Corcyrae (Mazz.). — Mart. Maio ♃ N. v.

2. **O. Luciliae** Bois. diagn. IV. p. 41, Fl. or. IV. p. 267; Heldr. chlor. Parn. p. 24; Hal. in ö. b. Z. 1896 p. 16. — Icon: Jaub. et Sp. ill. t. 366. — Exsicc.: Heldr. fl. gr. n. 2977; Reiser fl. gr. a. 1894.

Glabra, glaucovirens; caulibus adscendentibus; foliis inferioribus longe petiolatis, ovatis vel oblongis, obtusiusculis, basi rotundatis vel cuneatis, superioribus breviter petiolatis, oblongo-ellipticis; floribus in axillis foliorum superiorum breviter racemosis; pedicellis fructiferis elongatis, calyce multo longioribus, cernuis vel recurvis; calycis glabri, fructiferi aucti, laciniis ovato-oblongis, obtusiusculis; corollae coeruleae limbo tubo multo longiore; nuculis ventre scrobiculato-rugosis, margine non introflexo, integro. — Glaucedine et glabritie, foliis basi non cordatis, calycis glabri laciniis ovatis et corolla majori a praecedente specifice diversa.

In fissuris rupium regionis abietinae et superioris rarissime, hucusque tantum in mt. Kiona (Reiser) et loco dicto Gurna in mt. Parnasso pr. Rachova (Heldr.). — Jun. Jul. ♃

Obs. *O. linifolia* L. sp. p. 134; Mazz. in ant. ion. V. p. 186; *(Cynoglossum)*; Moench. meth. p. 419. — Colitur in hortis Corcyrae (Mazz.).

17. **Paracaryum** Bois. diagn. XI. p. 128.

1. **P. myosotoides** Lab. pl. syr. dec. II. p. 6 t. 2; S. et S. pr. I. p. 118; Sieb. in Flora I. p. 276; (*Cynoglossum*); Bois. l. c. p. 130; Raul. cret. p. 817; Bald. viagg. Creta p. 78. -- *Omphalodes myosotoides* Schrank. in denkschr. acad. wiss. Münch. 1811 p. 222. — *P. Sibthorpianum* Bois. diagn. XI. p. 130.

Pilis adpressis, basi tuberculatis cinerascens; caulibus adscendentibus pumilis, simplicibus vel fere a basi racemiferis; foliis radicalibus oblongis petiolatis, caulinis angustis, sessilibus; racemis fructiferis laxis; corollae coerulea-violaceae, parvae, limbo tubo subbreviore; pedicellis fructiferis patenti-recurvis, calyce brevioribus; nuculis disco echinatis, rarius laevibus, membrana subconcava, nitida, margine denticulis horizontaliter curvatis obsita, rarius integra instructis; stylo nucula 5 plo breviore.

In regione alpina mt. Mavri, Theodori, et Hagion Pneuma (Raul.) in mt. Sphacioticis (Sibth.) Cretae. — Jul. ♃ N. v.

18. **Mattia** Schult. obs. p. 30.

1. **M. graeca** Bois. et Heldr. diagn. VII. p. 30 *(Rindera)*; DC. pr. X. p. 168; Bois. fl. or. IV. p. 273; Heldr. chlor. Parn. p. 24; Bald. riv. coll. bot. Alb. 1896 p. 80; Hal. in z. b. G. 1899 p. 190. — *M. Schmidtii* Heldr. ind. sem. Athen. a. 1860; Ung. Reise p. 128. — Exsicc.: Heldr. herb. norm. n. 575 et 1567; Orph. fl. gr. n. 168; Bald. it. alb. epir. IV. n. 186.

Adpresse argento-sericea; rhizomate caespitoso; caulibus erectis, simplicibus, corymboso-umbellatis; foliis radicalibus linearibus lineari-lanceolatisque, basi longe attenuatis, caulinis praeter infima paucis squamaeformibus; pedicellis floriferis brevibus, fructiferis fructu multo longioribus; calycis lanati laciniis linearibus; corolla purpurea, calyce sesquilongiore, ultra tertiam partem in lacinias oblongo-spathulatas fissa; antheris et praecipue stylo ultra faucem exsertis; nuculis laevibus, ala eis sublatiore, margine repando cinctis. — Stirps pulchra, caulibus floriferis 5—10 cm., fructiferis elongatis ad 20 cm. altis, nuculae 15 mm. latae saepe pulchre coerulescentes.

In rupestribus regionis abietinae et superioris. Epirus: mt. Gamila distr. Zagorion (Bald.); mt. Parnassus (Guicc.), Cithaeron (Heldr.); Euboea: mt. Dirphys (Schmidt); Peloponnesus: mt. Chelmos (Orph.), Kyllene, Malevo (Heldr.), Taygetos (Reis.); insula Poros (Wied. ex Bois. fl. or.). — Maio, Jul. ♃

19. Cynoglossum L. gen. n. 183.

a. Nuculae immarginatae.

α. Perenne, rhizomate squamoso, ramoso.

1. **C. sphacioticum** Bois. et Heldr. diagn. XI. p. 125, Fl. or. IV. p. 263; Raul. cret. p. 816, t. 16. — Exsicc.: Heldr. pl. cret. n. 1633.

Molliter hirsutum, cinerascenti-virens; caulibus nanis, erectis, in racemum simplicem vel bifidum, pauciflorum, laxum, brevem abeuntibus; foliis inferioribus oblongis, in petiolum attenuatis, ceteris a basi semiamplexicauli oblong-linearibus; corolla atroviolacea, calyce sesquilongiore; pedicellis erectis, fructu subaequilongis; nuculis plano-convexis, undique dense glochidiatis. — Species pulchella *Paracaryo myosotoidi* habitu similis, caulibus gracilibus, 8—12 cm. altis, foliis radicalibus 15—25 mm. longis, 3—6 mm. latis, floribus parvis.

In saxosis regionis superioris mt. Stravopodia, Mavri et Theodori in mt. Sphacioticis Cretae, rarissime (Heldr.). — Jul. Aug. ♃

β. Biennia, radice fusiformi.

× Folia caulina basi cordato-semiamplexicaulia.

2. **C. nebrodense** Guss pr. I. p. 216; Bois. fl. or. IV. p. 265, suppl. p. 354; Haussk. symb. p. 158. — Icon: Rchb. t. 131. — Orph. fl. gr. n. 1117.

Molliter hirsutum, cinerascenti-virens; caule erecto, paniculato; foliis inferioribus oblongis, in petiolum attenuatis, ceteris a basi semiamplexicauli lanceolatis; racemis ebracteatis; fructiferis elongatis, laxiusculis; corolla rubra, calyce parum longiore; pedicellis fructiferis patenti-recurvis, fructu brevioribus; nuculis depressis, planiusculis, undique dense aculeato-glochidiatis.

In rupestribus subalpinis alpinisque. Thessalia: mt. Ghavellu, in oropedio Neuropolis (Haussk.), mt. Pelion (Heldr.); Euboea: mt. Dirphys (Heldr.); Achaia: mt. Chelmos (Orph.). — Apr. Jun. ⊙

3. **C. pictum** Ait. hort. Kew. I. p. 179; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 63, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 186; Marg. et R. fl. Zante p. 69; Friedr. Reise p. 273; Fraas fl. class. p. 162; Ung. Reise p. 128; Weiss in z b. G. 1869 p. 743; Raul. cret. p. 816; Spreitz. in z. b. G. 1877 p. 715, 1887 p. 665; Bois. fl. or. IV. p. 265; Heldr. fl. cephal. p. 54, Fl. Aegina p. 381; Haussk. symb. p. 158. — *C. creticum* Vill. fl. dauph. II. p. 157. — Huc probabiliter: *C. apenninum* S. et S. pr. I. p. 118; Pieri fl. corc. p. 24; Marg. et R. l. c.; vix L.. — Icon: Rchb. t. 130. — Exsicc.: Sint. it. thessal. n. 44; Baen. herb. europ. n. 9248.

Tomento tenui, molli canescens; caule elato, paniculato; foliis inferioribus oblongis, in petiolum longum attenuatis, ceteris a basi cordato-semiamplexicauli lanceolatis; racemis ebracteatis, fructiferis elongatis, densiusculis; corolla pallide coerulea, reticulatim picta, calyce sesqui-

longiore; pedicellis fructiferis arcuato-recurvatis, fructu subbreviores; nuculis plano-convexis, undique dense aculeato-glochidiatis. — Corolla pallide coerulea, venis saturate coeruleis picta insigne.

Ad vias, saepe, in arvis, ruderatis regionis inferioris et submontanae, passim per totam Graeciam; non indicatur in insulis maris Aegaei, sed sine dubio ibi etiam occurrit. — Apr. Jul. ♃

// Folia caulina basi rotundato-angustata.

4. **C. albanicum** Deg. et Bald. in Bald. riv. coll. hex. Alb. 1896 p. 80. — Exsicc.: Bald. it. alb. IV. n. 188.

Tomento tenui, molli canescens; caule elato, crasso, stricte et anguste paniculato; foliis inferioribus ignotis, superioribus ovato-lanceolatis vel ovato-rhombeis, basi rotundato-angustata sessilibus; racemis ebracteatis, fructiferis parum elongatis, densis; corolla ignota; pedicellis fructiferis erecto-patulis, brevissimis; nuculis paulo depressis, undique aculeis glochidiatis crassis obsitis. — Praeter foliorum formam a praecedentibus nuculis magnis, crasse aculeatis, a sequente et a *C. officinale* nuculis immarginatis diversum. — Ulterius observandum.

In aridis alpestribus ad fontem Bocikopulon distr. Pogoni in Epiro boreali (Bald.). — Maio, Jun. ♃

b. Nuculae margine elevato, denticulato-glochidiato cinctae.

5. **C. Columnae** Ten. fl. nap. pr. p. 14, Fl. nap. t. 116; Marg. et R. fl. Zante p. 69; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 816; Bois. fl. or. IV. p. 265; Heldr. fl. cephal. p. 54; Hal. Beitr. fl. Epir. p. 32; Haussk. symb. p. 159; Form. in Ver. Brünn 1897 p. 44. — Huc spectant saltem pro maxima parte: *C. officinale* S. et S. pr. I. p. 117; Pieri corc. fl. p. 23; Dallap. prosp. p. 20; Mazz. in ant. ion. V. p. 184; Friedr. Reise p. 279; Bois. fl. or. IV. p. 264, quoad pl. graecam; Form. in D. bot. Monat. 1891 p. 24, in Ver. Brünn 1895 p. 92; non L. sp. p. 134, quod corolla sordide rubra, pedicellis infimis bracteatis, fructiferis longioribus, nuculis undique aequaliter glochidiatis, margine crassiusculo prominulo cinctis, differt. — *C. montanum* Ch. et B. exp. p. 63, Fl. pelop. p. 13; non Lam. quod in Graecia certe non provenit. — *C. silvaticum* Mazz. l. c., non Haenke. — *C. cheirifolium* Mazz. l. c. p. 186; Raul. cret. p. 816; non L. sp. p 134, quod praeter alias notas, racemis bracteatis discedit. — Exsicc.: Orph. fl. gr. n. 865; Heldr. herb. dimorph. n. 38; Sint. it. thessal. n. 633.

Tomento molli canescens vel canum; caule erecto, paniculato; foliis inferioribus oblongis, in petiolum attenuatis, ceteris a basi cordato-semiamplexicauli breviter lanceolatis; racemis ebracteatis, fructiferis elongatis; corolla rubra vel rarius albida; pedicellis fructiferis erecto-patulis, fructu brevioribus; nuculis plano-compressis, aculeato-glochidiatis, margine elevato cin-

ak. symb. p. 159. — Obscure virens, breviter
tiores, valde obtusatae; corolla violacea. —
V. a. 1883.

Ad sepes, vias, in herbidis regionis inferioris et montanae passim per totam Graeciam; — *β*. Epirus: in regione abietina mt. Tsumerka pr. Vulgarelion (Hal.); Thessalia: mt. Karava (Heldr.), ad monasterium Korona in Pindo (Haussk.). — Mart. Jul. ⊙

20. Lappula Gilib. fl. lith. I. p. 25.

1. **L. echinata** Gilib. l. c.; Form. in Ver. Brünn 1896 p. 60. — *Myosotis lappula* L. sp. p. 131; S. et S. pr. I. p. 112; Ch. et B. exp. p. 61, Fl. pelop. p. 12. — *L. myosotis* Moench meth. p. 417. — *Echinospermum lappula* Lehm. asp. p. 121; Bois. fl. or. IV. p. 249; Haussk. symb. p. 159; Form. in Ver. Brünn 1895 p. 32. — *M. echinosperma* Mazz. in ant. ion. IV. p. 960. — *L. vulgaris* Mazz. l. c. V. p. 180. — Icon: Fl. dan. t. 692. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Pilis adpressiusculis, basi tuberculatis hirsuta; caule erecto, rigido, ramoso, ramis in racemos bracteatos, demum elongatos abeuntibus; foliis lanceolatis, inferioribus in petiolum attenuatis, ceteris sessilibus; pedicellis fructiferis erectis, calyce brevioribus; corolla parva, coerulea; nuculis facie tuberculatis, margine 2—3 seriatim glochidiatis.

Ad vias, in ruderatis, saxosis regionis inferioris, in subalpinam adscendens, rare. Epirus: mt. Prosgoli (Bald.); Thessalia: pr. Chaliki, mt. Zygos (Haussk.), Dokimi (Form.), pr. Malakasi (Heldr.); Argolis (Sibth.); Laconia, pr. Andrusa Messeniae (Chaub.); Corcyra: pr. Castrades, Mattia (Mazz.). — Maio, Aug. ⊙

21. Asperugo L. gen. n. 189.

1. **A. procumbens** L. sp. p. 138; S. et S. pr. I. p. 123, Fl. gr. II. p. 65 t. 177; Ch. et B. exp. p. 63, Fl. pelop. p. 13; Mazz. in ant. ion. V. p. 192; Friedr. Reise p. 266; Fraas fl. class. p. 161; Clem. sert. p. 69; Bois. fl. or. IV. p. 275; Haussk. symb. p. 159. — *A. alba* Mazz. l. c.; f. floribus albis. — Icon: Fl. dan. t. 552. — Exsicc.: Heldr. pl. fl. hellen. a. 1892.

Setis rigidis aspera; caule decumbente, a basi ramosa; foliis elliptico-oblongis, inferioribus in petiolum attenuatis, ceteris sessilibus; floribus axillaribus, brevissime pedicellatis, 3—4 fasciculatis, foliis oppositis; pedicellis fructiferis cernuis; corolla minuta, coerulea, rarissime alba; nuculis verruculosis.

In ruderatis, ad vias regionis inferioris. Boeotia: pr. Lebadea (Fraas); Attica: pr. Athenas (Friedr.), mt. Hymettus (Orph.); Corinthus (Fraas); Argolis (Sibth.): pr. Argos (Fraas); Messenia: pr. Pylos (Chaub.). — Mart. Maio ⊙

22. Rochelia Rchb. in Flora 1824 p. 243.

1. **R. disperma** L. sp. ed. 2 p. 191; S. et S. pr. I. p. 115; (*Lithospermum*); Stapf in bot. Ergebn. Polak exp. Pers. I. p. 31. — *R. stellulata* Rchb. pl. cret. II. t. 123.

Setulis tuberculatis adpressis canescens; caule erecto, apice vel a basi ramoso; foliis inferioribus oblongo-spathulatis, in petiolum attenuatis, ceteris linearibus, sessilibus; racemis cymosis vel paniculatis, densifloris, bracteatis; pedicellis fructiferis recurvatis, calyce aequilongis; calycis laciniis linearibus, hamato-incurvis; corolla minuta, coerulea; nuculis glochidibus stellatis, sessilibus tuberculatis.

Thessalia: mt. Ossa (Sibth.), sed a recentioribus non lecta. — Mart. Maio ⊙

Obs. *Cordia myxa* L. sp. p. 190 ex ordine *Cordiacearum* R. Br., colitur in hortis pr. Athenas et in insula Syra (Heldr. Nutzpfl. p. 35).

LXXXI. Ordo. Solanaceae Juss. gen. p. 124.

Dispositio tribuum generumque:

1. Tribus. *Datureae* Endl. gen. p. 663. — Ovarium incomplete quadriloculare; fructus capsularis, apice valvis dehiscens.

1. Datura L.

2. Tribus. *Solaneae* Don syst. IV. p. 400. — Ovarium biloculare.

a. Fructus capsularis, operculo circumscisse dehiscens.

2. Hyosciamus L.

b. Fructus baccatus indehiscens.

α. Calyx fructifer valde auctus, baccam plus minus includens.

× Calyx fructifer baccam includens; corolla anguste campanulata. Frutex.

3. Withania Pauq.

×× Calyx fructifer baccam omnino tegens; corolla subrotata. Herba.

4. Physalis L.

β. Calyx fructifer parum auctus, baccam basi cingens.

× Antherae rima longitudinali dehiscentes.

○ Herbae acaules, floribus radicalibus, longe pedunculatis.

5. Mandragora Tourn.

○○ Herbae cauligerae.

6. Atropa L.

○○○ Frutices.

7. Lycium L.

×× Antherae poris 2 dehiscentes.

8. Solanum L.

1. Tribus. DATUREAE Endl. gen. p. 663.

1. Datura L.

1. **D. stramonium** L. sp. p. 179; S. et S. pr. I. p. 152; Pieri corc. fl. p. 32; Ch. et B. exp. p. 7. Fl. pelop. p. 15; Mazz. in ant. ion. V. p. 214; Fraas fl. class. p. 166; Clem. sert. p. 70; Heldr. Nutzpfl. p. 37; Bois. fl. or. IV. p. 292; Heldr. fl. cephal. p. 54, in Sitzungsb.

acad. wiss. Berl. 1883 p. 163; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 62, 1896 p. 33, 1897 p. 45; Haussk. symb. p. 159. — Icon: Fl. dan. t. 436. — Exsicc.: Heldr. it. gr. septentr. n. 1879.

Sparsim papillosa; caule erecto, ramoso; foliis petiolatis, ovatis oblongisve, inaequaliter sinuato-dentatis; floribus solitariis, axillaribus, erectis; corolla infundibuliformi, alba, calyce pentagono subduplo longiore, lobis tenuiter mucronatis; capsula oblonga, erecta, aculeata. — Planta foetida, floribus magnis, 5—10 cm. longis, capsula 3—5 cm. longa.

In locis cultis, ruderatis regionis inferioris et submontanae. Frequens in Epiro et Thessalia; Aetolia: pr. Musinitza (Heldr.); Attica: pr. Eleusis, Athenas (Fraas), ad Cephissum (Clem.); Arcadia (Sibth.): pr. Tripoliza (Fraas); Messenia (Sibth.): pr. Methone (Chaub.); Laconia: ad fl. Eurotas, in planitie Helos (Chaub.); Cephalonia: mt. Phalaris, pr. Lordata, Lixuri (Heldr.); Corcyra (Pieri): pr. Garizza, San Rocco (Mazz.). — Jun. Sept. ⊙

Obs. *D. metel* L. sp. p. 179; Ch. et B. fl. pelop. p. 15. — Dense pubescens, cinerascens, foliis repandis integrisve, corollae limbo purpurascente, capsula pendula a praecedente discedens, occurrit introducta ad ostia fl. Eurotas pr. Limonia et in planitie Helos (Chaub.).

2. Tribus. **SOLANEAE** Don syst. IV. p. 400.

2. **Hyosciamus** L. gen. n. 247.

a. Folia caulina sessilia.

1. **H. niger** L. sp. p. 179; S. et S. pr. I. p. 152; Dallap. prosp. p. 24; Mazz. in ant. ion. V. p. 214; Fraas fl. class. p. 166; Heldr. Nutzpfl. p. 37; Raul. cret. p. 817; Bois. fl. or. IV. p. 294; Hal. Beitr. fl. Epir. p. 32; Form. in D. bot. Monat. 1891 p. 25, in Ver. Brünn 1895 p. 33. — Icon: Fl. dan. t. 1452. — Exsicc.: Sint. it. thessal. n. 434.

Villosus, viscidulus; caule erecto, simplici vel ramoso; foliis ovatis vel oblongis, sinuatis vel sinuato-pinnatifidis, inferioribus in petiolum attenuatis, ceteris semiamplexicaulibus; floribus sessilibus, in spicam foliosam, unilateralem, scorpioideam, tandem elongatam, rectam dispositis; foliis floralibus sessilibus, oblongo-lanceolatis, calyces superantibus, superioribus integris; calyce fructifero stricto, supra medium constricto, dentibus erecto-patulis, triangularibus, mucronatis; corolla subaequali, pallida, violaceo-reticulata, fundo violaceo; staminibus faucem parum excedentibus.

β. **pallidus** Kit. in Willd. en. hort. berol. I. p. 227, pro sp.; Koch syn. p. 509; Haussk. symb. p. 159; Form. in Ver. Brünn 1897 p. 45. — Floribus unicoloribus, pallidis. — Exsicc.: N. v.

In ruderatis, arvis regionis inferioris et montanae. Epirus: pr. Sadovica, Hagios Paraskevi (Form.), Kalarrytes (Hal.); Thessalia: pr. Said Pascha, Malakasi, Velestinos (Form.), Kalabaka (Sint.); Euboea: pr. Kastaniotissa (Heldr.), Steni (Fraas); Laconia: pr. Kastanitza in mt. Malevo (Fraas); Creta: pr. Neochori distr. Apokorona (Raul.); Cephalonia (Dallap.); Corcyra: mt. Abramo (Mazz.); — *β*. Thessalia: mt. Zygos, monasterium Korona, pr. Karditza, Pharsalus, Volo (Haussk.), Mavreli (Form.). — Apr. Jul. ⊙ et ⊙

2. **H. reticulatus** L. sp. ed. 2 p. 257; Raul. cret. p. 817. — Icon: Jaub. et Sp. ill. V. t. 416. — Exsicc.: Heldr. herb. n. 695 (Pisidia).

Adpresse puberulus; caule erecto, simplici vel ramoso; foliis oblongis vel lanceolatis; sinuato-dentatis vel pinnatifidis, inferioribus in petiolum attenuatis, ceteris sessilibus, vix amplexicaulibus; floribus sessilibus, in spicam foliosam, unilateralem, scorpioideam, tandem elongatam, rectam dispositis; foliis floralibus sessilibus, oblongis vel lanceolatis, integris, flores superantibus; calyce fructifero stricto, non constricto, dentibus patulis, spinosis; corolla aequali, purpureo-violacea, reticulata; staminibus fauce sublongioribus. — Corolla saturate purpureo-violacea insignis.

Indicatur a Linnaeo in Creta, sed a recentioribus non lecta. — Apr. Jul. ♃ et ⊙ N. v.

b. Folia omnia petiolata.

3. **H. albus** L. sp. p. 180; S. et S. pr. I. p. 153, Fl. gr. III. p. 24 t. 230; Pieri corc. fl. p. 33; Dallap. prosp. p. 23; Ch. et B. exp. p. 73. Fl. pelop. p. 15; Mazz in ant. ion. V. p. 214; Marg. et R. fl. Zante p. 70; Friedr. Reise p 271; Fraas fl. class. p. 166; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 295; Heldr. fl. cephal. p. 54; Fl. Aegina p. 381 *v. major*. in ö. b. Z. 1898 p. 184, chlor. Thera p. 17, chlor. Mykon. p. 248; Bald. viagg. Creta p. 79; Haussk. symb. p. 159 *v. minor*; Form. in Ver. Brünn 1895 p. 33, 1896 p. 62. — *H. major* Mill. dict. n. 2; Heldr. Nutzpfl. p. 37; Raul. cret. p. 818; f. corollae fundo violaceo. — *H. minor* Mill. dict. n. 4; Weiss in z. b. G. 1869 p. 744; f. corolla fundo virescente. — *H. varians* Vis. fl. dalm. t. 24. — *H. graecus* Heldr. chlor. Mykon. p. 248, nomen solum. — Exsicc.: Sint. et Bornm. it. turc. n. 1377; Sint. it. thessal. n. 548; Bald. it. cret. alt. n. 351; Dörfl. fl. aeg. n. 38.

Villoso-viscosus; caulibus erectis vel decumbentibus; foliis petiolatis, a basi subcordata, truncata vel breviter cuneata ovato-orbiculatis, grosse triangulari-lobatis vel obtuse repando-lobatis; floribus praeter infima, saepe breviter pedunculata, sessilibus, in spicam foliosam, unilateralem, scorpioideam, tandem elongatam, rectam dispositis; foliis floralibus petiolatis, flores superantibus, cordatis, ovatis, dentatis vel saepius oblongis, integris, basi attenuatis; calyce fructifero stricto, non constricto, dentibus erectis, breviter triangularibus, acutis, pungentibus; corolla pallida, non reticulata, fundo virente vel violaceo, lobis subinaequalibus; staminibus

fauce sublongioribus. — Variat caule plus minus robusto, indumento plus minus copioso, foliorum forma et florum magnitudine et colore, sed folia diversa et corollae basi virescentes vel violaceae saepe in uno eodemque specimine occurrunt, quamobrem hae formae, ab autoribus nonnullis ut species propositae, nequidem ut varietates distinguendae sunt. Cf. Vis. fl. dalm. II. p. 234, Bois. fl. or. IV. p. 296 et Parl. fl. ital. VI. p. 672.

In ruderatis, ad vias, muros regionis inferioris. Thessalia: pr. Kalabaka, Litochori (Sint.), Trikala (Haussk.), Volo (Haussk.), mt. Pelion (Heldr.); Attica: in Acropoli Athenarum; insula Aegina (Heldr.), Poros (Heldr.); Cycladum insula: Tenos, Syra (Weiss), Keos, Cythnos (Tunt.), Mykonos (Sart.), Naxos, Kameni pr. Thera (Heldr.), Amorgos (Leon.); pr. Sparta et in Messenia littorali (Chaub.); Creta: pr. Canea, Kissamos (Raul.), pr. Purphuras ad radices Idae, pr. Myron distr. Malevisi (Bald.); Zante (Marg.); Cephalonia (Dall.): pr. Argostoli, Lordata, Same, Piscardo, Lixuri (Heldr.); Corcyra (Pieri): pr. urbem, Evropuli, Afra (Form.). — Febr. Oct. ⊙ et ♃

4. **H. aureus** L. sp. p. 180; S. et S. pr. I. p. 153, Fl. gr. III. p. 25 t. 231; Sieb. avis p. 3, rem. p. 3; Fraas fl. class. p. 166; Raul. cret. p. 818; Bois. fl. or. IV. p. 296. — Exsicc.: Sint. et Rigo it. cypr. n. 9.

Villoso-viscosus; caulibus decumbentibus vel pendulis, ramosis; foliis petiolatis, cordatis, ovatis vel orbiculatis, triangulari-lobatis, lobis acute dentatis; floribus omnibus pedunculis calyce brevioribus suffultis, in racemum foliosum, unilateralem, scorpioideum, tandem elongatum, rectum dispositis; foliis floralibus petiolatis, flores superantibus, triangulari-ovatis vel oblongis, acute dentatis; calyce fructifero erecto-patulo vel nutante, non constricto, membranaceo, dentibus erectis, triangulari-lanceolatis, acutis non pungentibus; corolla aurea, fundo violaceo, lobis valde inaequalibus; staminibus exsertis.

Ad moenia et muros regionis inferioris Cretae: pr. Candia (Heldr.); erronee copiose dicitur a Sibthorpio in muris et ruderatis Graeciae. — Mart. Jun. ♃

3. Withania Pauquy diss. bellad. in Feruss. bull. sc. nat. V. p. 254.

1. **W. somnifera** L. sp. p. 182; S. et S. pr. I. p. 154, Fl. gr. III. p. 27 t. 233; Sieb. avis p. 3, rem. p. 3; Ch. et B. exp. p. 74, Fl. pelop. p. 15; Fraas fl. class. p. 165; Raul. cret. p. 818; *(Physalis)*; Mor. fl. sard. III. p. 156; Bois. fl. or. IV. p. 287. — Exsicc.: Orph. fl. gr. n. 723; Rev. pl. cret. p. 116.

Fruticosa, parce tomentosa; ramis flexuosis; foliis petiolatis, ovatis, integris; floribus brevissime pedunculatis, axillaribus, aggregatis, rarius solitariis; calycis albo-lanati laciniis linearibus, brevibus, muticis; corolla parva, luteo-virenti; bacca globosa, rubra, calyce vesiculoso, virente, marcescente inclusa.

In collibus, ad sepes, vias regionis inferioris, rare. Euboea (Sibth.): pr. Chalkis (Fraas); Peloponnesus: pr. Epidaurus, Kalamata (Chaub.), Monembasia (Fraas); Creta: pr. Kissamos, Malaxa (Rev.), Khalepa, Candia (Heldr.), Perama (Bald.). — Apr. Oct. ♄

4. Physalis L. gen. n. 250.

1. **P. alkekengi** L. sp. p. 183; S. et S. pr. 1. p. 154, Fl. gr. III. p. 28 t. 234; Dallap. prosp. p. 27; Bois. fl. or. IV. p. 287; Heldr. in Sitzungsb. acad. wiss. Berlin 1883 p. 160; Hal. Beitr. fl. Epir. p. 32; Haussk. symb. p. 159. — Exsicc.: Heldr. it. thessal. n. 83; herb. norm. n. 1567.

Viridis, parce hirtula; rhizomate repente; caulibus erectis vel adscendentibus; foliis petiolatis, ovatis, repandis; floribus pedunculatis, axillaribus, solitariis; pedunculis fructiferis nutantibus; calycis villosuli laciniis breviter triangularibus, acuminatis; corolla albida; bacca globosa, rubra, calyce ample vesiculoso, ovato, igneo inclusa.

In umbrosis regionis montanae, rare. Epirus: pr. Mazuki (Hal.); Thessalia: ad monasterium Korona in Pindo (Haussk.), mt. Ossa pr. coenobium Hagios Demetrius (Heldr.), mt. Pelion pr. Zagora (Aphentulis); mt. Parnassus (Sibth.); Laconia: mt. Malevo (Orph.), pr. Megali Anastasova (Zahn); Cephalonia (Dall.). — Maio, Jul. ♃

Obs. *C. annuum* L. sp. p. 188; Dallap. prosp. p. 27; Ch. et B. exp. p. 74, Fl. pelop. p. 15; Marg. et R. fl. Zante p. 85; Heldr. Nutzpfl. p. 36, Fl. cephal. p. 54; Raul. cret. p. 818. — *C. grossum* Willd. en. hort. berol. I. p. 21; Dallap. p. 27. — *C. longum* DC. cat. hort. monsp. 1813 p. 86; Fraas fl. class. p. 169. — Coluntur ad usum culinarem.

5. Mandragora Tourn. inst. p. 76.

a. Corolla albo-virens; bacca globosa.

1. **M. officinarum** L. sp. p. 181; Sieb. avis p. 3; Dallap. prosp. p. 24; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 291; Boissieu in bull. soc. bot. fr. 1896 p. 287. — *Atropa mandragora* L. sp. ed. 2 p. 259; Sieb. avis p. 3, in Flora I. p. 271; Ch. et B. exp. p. 74, Fl. pelop. p. 15. — *M. vernalis* Bert. comm. mandrag. p. 9, t. 1; Raul. cret. p. 819; Heldr. Nutzpfl. p. 36, in Mitth. geogr. Ges. Thüring. V. p. 77, chlor. Mykon p. 248; Haussk. symb. p. 159. — Exsicc.: Heldr. pl. fl. hellen. a. 1899.

Radice crassa, carnosa, fusiformi, simplici vel ramosa; foliis rosulatis, breviter petiolatis, ovato-oblongis oblongisve, corrugato-crispis, saepe sinuato-dentatis, pilosulis vel glabris, inferioribus obtusis, ceteris acutis vel acuminatis; pedunculis e collo fasciculatis, unifloris, calyce longioribus, hirtulis; calycis laciniis lanceolatis, acutis, corolla duplo brevioribus, bacca subbrevioribus; bacca lutea. — Radix ad 60 cm. longa, extus albida; folia viridia, carnosula, ad 40 cm. longa, 20 cm. lata,

foetida; pedunculi 2—5 cm. longi; corolla 3 cm. longa; bacca 2—3 cm. diametro.

In apricis, incultis regionis inferioris. Thessalia: pr. Zeitun, Lamia (Sprun.); Corinthus (Heldr.); Messenia: pr. Pylos, Methone, insula Sapienza, in Maina (Chaub.); Cycladum insula Keos (Boissieu), Syra, Mykonos, Rhenea (Heldr.); Creta: pr. Suda (Weiss), in fauce Perivoglia (Sieb.), pr. Canea, Hierapetra, insula Dia (Heldr.); Cephalonia (Dallap.); sed loca nonnula forsan ad sequentem spectant. — Dec. Mart. ♃

1 × 2. **M. officinarum** × **Haussknechtii.** — *M. hybrida (Haussknechtii* × *vernalis)* Haussk. et Heldr. in Mitth. geogr. Ges. Thür. V. p. 77; Haussk. symb. p. 159. — Folia ovato-oblonga, rugulosa, eis *M. officinarum* longius petiolata; flores longe pedunculati, steriles; calycis laciniae acuminatae, sed eis *M. Haussknechtii* breviores; corollae pallide violaceae, magnitudine eis parentium intermedia. Specimen valde mancum tantum vidi. — Exsicc.: Heldr. pl. fl. hellen. a. 1885.

In arenosis maritimis pr. Corinthum, inter parentes (Haussk.).

b. Corolla violacea; bacca ellipsoidea.

2. **M. Hausknechtii** Heldr. in Mitth. geogr. Ges. Thür. V. p. 77, chlor. Mykon. p. 248; Haussk. sym. p. 159. — Exsicc.: Rev. pl. cret. n. 117, in Baen. herb. europ. n. 5050; Heldr. pl. fl. hellen. a. 1885.

Radice crassa, carnosa, fusiformi, simplici vel ramosa; foliis rosulatis, longiuscule petiolatis, oblongo-lanceolatis, acuminatis, corrugato-crispis, saepe sinuato-dentatis, pilosulis vel glabris; pedunculis e collo fasciculatis, unifloris, calyce multoties longioribus, hirtulis; calycis laciniis lanceolatis, acuminatis, corollae fere aequilongis, bacca longioribus; bacca lutea. — Folia ad 30—40 cm. longa, 5—9 cm. lata; pedunculi ad 15 cm. longi; corollae eis speciei praecedentis submajores; bacca 3 cm. longa, 2 cm. lata. — Differt a *M. officinarum* foliis acuminatis angustioribus, calycis laciniis longe acuminatis, corolla violacea et bacca ellipsoidea.

In arenosis maritimis, locis incultis. Corinthus (Haussk.); Cycladum insula Rhenea (Heldr.); Creta: pr. Canea (Rev.). — Mart. Apr. ♃

3. **M. autumnalis** Spreng. syst. I. p. 699; Bois. fl. or. IV. p. 291; Heldr. in Mitth. geogr. Ges. Thür. V. p. 77. — *Atropa mandragora* S. et S. pr. I. p. 153, Fl. gr. III. p. 26 t. 232; Fraas fl. class. p. 166; non L.. — *M. officinarum* Bert. comm. mandrag. p. 10 t. 2; Dun. in DC. pr. XIII. p. 466; Heldr. Nutzpfl. p. 36; non L.. — *M. microcarpa* Bert. l. c. p. 12 t. 3; Heldr. Nutzpfl. p. 36; Weiss in z. b. G. 1869 p. 744; Heldr. in Mitth. geogr. Ges. Thür. V. p. 77; Haussk. symb. p. 159. — Exsicc.: Orph. fl. gr. n. 75; Heldr. herb. norm. n. 257.

Radice crassa, carnosa, fusiformi, simplici vel ramosa; foliis rosulatis, longiuscule petiolatis, oblongis oblongo-lanceolatisve, subrepandis,

supra glabriusculis, subtus puberulis, infimis obtusis, ceteris acutis vel acuminatis; pedunculis e collo fasciculatis, unifloris, calyce longioribus, glabriusculis; calycis laciniis lanceolatis, corolla subtriplo brevioribus, baccam subaequantibus; bacca luteo-purpurascente. — **Radix** extus nigricans, minor; folia minora coerulescenti-viridia, subtus pallidiora, petioli subpurpurascentes; bacca parva, 2 cm. longa.

In locis incultis. Phthiotis: in valle fl. Sperchius (Fraas); Attica: pr. Athenas (Sibth.), Eleusis, Brahami (Heldr.); Elis (Sibth.); Archipelagus (Sibth.): insula Syra (Weiss); huc quoque forsan loca nonnulla ad *M. officinarum* indicata. — Sept. Nov. ♃

6. Atropa L. gen. n. 249.

1. **A. belladonna** L. sp. p. 181; Fraas fl. class. p. 166; Heldr. Nutzpfl. p. 36; Bois. fl. or. IV. p. 291; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1897 p. 45. — Icon: Fl. dan. t. 758.

Caule crasso, elato, herbaceo, viscidulo-pubescente; foliis breviter petiolatis, ovatis vel oblongis, integris, glabris vel puberulis; floribus axillaribus, solitariis, pedunculatis, nutantibus; corolla squalide violacea; bacca globosa, nigra.

In umbrosis, silvaticis montanis. Thessalia: mt. Strugo pr. Kastania in Pindo, pr. Hepdominta Aderfia in mt. Oxya (Form.), mt. Olympus (Heldr.), mt. Oeta supra Patradjik (Fraas); mt. Parnassus (Heldr.). — Jun. Jul. ♃ N. v.

7. Lycium L. gen. n. 262.

1. **L. europaeum** L. sp. p. 192; S. et S. pr. I. p. 155, Fl. gr. III. p. 30 t. 236; Ch. et B. exp. p. 75, Fl. pelop. p. 16; Mazz. in ant. ion. V. p. 214; Marg. et R. fl. Zante p. 69; Fraas fl. class. p. 166; Clem. sert. p. 70; Bois. fl. or. IV. p. 288; Heldr. fl. cephal. p. 54, Fl. Aegina p. 381, chlor. Thera p. 18; Spreitz in z. b. G. 1887 p. 665; Terrac. in Malpigh. 1890 p. 60; Bald. in Malpigh. 1894 p. 60; Haussk. symb. p. 159. — *L. mediterraneum* Dun. in DC. pr. XIII. p. 523. — Exsicc.: Heldr. herb. norm. n. 1266.

Fruticosum; ramis erectis, spinosis, glabris vel pruinosis; foliis fasciculatis, petiolatis, obovato-lanceolatis, integris, acutis vel obtusis, glabris; floribus e fasciculis solitariis vel geminis, pedunculatis; calyce subaequaliter 5 dentato, subbilabiato; corolla violacea, tubo infundibuliformi, lobis tubo triplo brevioribus; bacca globosa, rubra.

Ad sepes, vias. Epirus: pr. Prevesa (Bald.); Thessalia: pr. Velestinos, Aivali, Órman Magula (Haussk.); pr. Athenas, Corinthum (Heldr.); Laconia: pr. Astros (Fraas), ad fl. Eurotas (Chaub.); Cythaera: pr. Kapsali (Spreitz.); Cephalonia: pr. Lixuri, in oropedo Omalo (Heldr.); Corcyra: pr. Anemomylo, Garizza, Manducchio (Mazz.). — Febr. Jun. ♄

2. **L. barbarum** L. sp. p. 192 p. p.; S. et S. pr. I. p. 155; Fraas fl. class. p. 166. — *L. vulgare* Dun. in DC. pr. XIII. p. 509.

Differt a praecedente ramis evolutis virgatis, arcuato-nutantibus pendulisve, corollae lobis tubo dimidio longioribus et baccis ellipsoideis.

Indicatur a Fraas pr. Athenas et in insula Tenos, a Sibthorp in insula Naxos. — Febr. Jun. ♄ N. v.

Obs. *Cestrum Parqui* L'Her. stirp. I. p. 73. Colitur et hinc inde e. g. pr. Kastrades Corcyrae (Baen. herb. europ. n. 9240) subspontaneum.

8. Solanum L. gen. n. 251.

a. Rami volubiles; corolla violacea; bacca ellipsoidea, rubra.

1. **S. dulcamara** L. sp. p. 185; S. et S. pr. I. p. 154; Ch. et B. exp. p. 74, Fl. pelop. p. 15; Mazz. in ant. ion. V. p. 215; Fraas fl. class. p. 165; Land. in Flora 1856 p. 307; Heldr. Nutzpfl. p. 36, in Sitzungsb. acad. Wiss. Berlin 1883 p. 160; Bois. fl. or. IV. p. 285; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 33, 1896 p. 61, 1897 p. 45; Haussk. symb. p. 159. — Icon: Fl. dan. t. 607. — Exsicc.: Heldr. it. gr. septentr. a. 1879, it. thessal. n. 91.

Caule fruticoso, ramis herbaceis; foliis petiolatis, minute pubescentibus vel tomentellis, ovatis vel ovato-oblongis, saepe cordatis, integris vel superioribus hastatis vel utrinque lobulo auctis; cymis extraaxillaribus, longe pedunculatis, laxis; calycis lobis late triangularibus; corollae laciniis lanceolatis, saepe reflexis.

In humidis, ad ripas, salicetis regionis inferioris et montanae. Epirus: pr. Prevesa, Philippiades, Janina, Kacka (Form.); Thessalia: pr. Malakasi, Kastania, Neochorion (Form.), mt. Karava, pr. Trikala (Haussk.), Karditza, Larissa, Velestinos, Megalivris (Form.), mt. Ossa pr. Retziani (Heldr.); Euboea (Fraas); Aetolia: pr. Musinitza in mt. Korax (Heldr.); Boeotia: ad Cephissum, pr. Lebadea (Fraas); Argolis: in palude Lerna (Sart.); Achaia: pr. Kalavryta (Hal.); Laconia: ad Magula pr. Sparta (Chaub.); Cycladum insula Andros (Fraas). — Maio, Aug. ♄

Obs. *S. sodomaeum* L. sp. p. 187; Bald. in nuov. giorn. bot. it. I. p. 99; et *S. citrullifolium* A. Br. ind. sem. h. Frib. a. 1849; ramis foliisque aculeis validis rectis luteis pungentibus obsitis, foliis sinuato-pinnatilobis, coluntur ornamenti gratia et hinc inde, e. g. pr. Prevesa (Bald.), Athenas (Heldr.), subsponte ocurrunt.

b. Rami non volubiles; corolla alba; bacca globosa.

2. **S. suffruticosum** Schousb. in Willd. en. hort. berol. I. p. 236; Haussk. symb. p. 159.

Perenne; caule elato, lignescente, glabro, ramis herbaceis; foliis petiolatis, glabris, ovatis vel ovato-lanceolatis, repandis vel sinuato-dentatis; cymis pedunculatis, umbelliformibus; calycis lobis ovatis, corolla duplo brevioribus; bacca nigra.

In saxosis peninsulae Methana pr. Vromolimni Argolidis (Haussk.). — Maio, Sept. ♃ N. v.

3. **S. nigrum** L. sp. p. 186; S. et S. pr. I. p. 155; Pieri corc. fl. p. 34; Dallap. prosp. p. 25; Ch. et B. exp. p. 75, Fl. pelop. p. 16; Marg. et R. fl. Zante p. 71; Friedr. Reise p. 264 et 271; Clem. sert. p. 70; Fraas fl. class. p. 166; Raul. cret. p. 818; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 284; Form. in D. bot. Mon. 1891 p. 25, in Ver. Brünn 1895 p. 33, 1896 p. 61, 1897 p. 45; Haussk. symb. p. 159; Heldr. fl. Aegina p. 381, chlor. Thera p. 18, chlor. Mykon. p. 248. — Icon: Fl. dan. t. 460. — Exsicc.: Heldr. herb. norm. n. 586; Sint. et Bornm. it. turc. n. 1375; Dörfl. pl. cret. n. 137.

Annuum; caule erecto, herbaceo, ramoso, foliisque glabriusculo; foliis petiolatis, ovatis vel deltoideo-rhombeis, repandis vel sinuato-dentatis; cymis pedunculatis, umbelliformibus; calycis lobis ovatis, corolla 2—3 plo brevioribus; bacca nigra. — Species indumento et praesertim baccarum colore, in herbario vix distinguendo, variabilis.

β. **humile** Bernh. in Willd. en. hort. berol. I. p. 236; Clem. sert. p. 70; Raul. cret. p. 819; pro sp.; Bois. fl. or. IV. p. 284. — Bacca viridis vel viridi-lutescens.

γ. **flavum** Kit. in Schult. oestr. fl. ed. 2 I. p. 294, Fraas fl. class. p. 166; pro sp. — *S. chlorocarpum* Spenn. fl. frib. p. 1074. — *S. nigrum v. chlorocarpum* Bois. fl. or. IV. p. 284; Heldr. fl. cephal. p. 54. — Bacca ceraceo-flava.

δ. **miniatum** Bernh. in Willd. en. hort. berol. I. p. 236; Mazz. in ant. ion. V. p. 214; Marg. et R. p. 70; Bois. fl. or. IV. p. 284; pro sp. . — Bacca miniata.

ε. **villosum** L. sp. p. 186. — *S. villosum* Lam. ill. II. p. 18; Ch. et B. fl. pelop. p. 16; Raul. cret. p. 218; Bois. fl. or. IV. p. 285. — Undique patule tomentoso-villosum; bacca miniata vel crocea.

In ruderatis, cultis et incultis per totam Graeciam. — Jan. Jul ⊙

Obs. Late coluntur ad usum domesticum: *S. melongena* L. sp. p. 186. — *S. tuberosum* L. sp. p. 185. — *S. pseudocapsicum* L. sp. p. 184. — *Lycopersicum esculentum* Mill. gard. dict. n. 2. — Dein: *Nicotiana tabacum* L. sp. p. 180. *N. rustica* L. sp. p. 180 et ornamenti gratia *N. glauca* Grah in Edinb. phil. journ. 1828 p. 175.

LXXXII. Ordo. **Scrofulariaceae** Lindl. Key. syst. bot. p. 66.

Dispositio tribuum generumque:

A Corollae labium inferius praefloratione a labio superiore inclusum.

1. Tribus. Verbasceae Benth. in DC pr. X. p. 224. — Corolla rotata, subirregularis, tubo brevissimo; stamina 5 vel 4; capsula bivalvis.

a. Stamina 5.

Verbascum L.

b. Stamina 4.

2. Celsia L.

2. Tribus. *Cheloneae* Benth. in DC. pr. X. p. 298. — Corolla campanulata vel tubulosa, irregularis, non saccata nec calcarata; stamina 5 vel 4; capsula bivalvis.

a. Corolla oblique campanulata, tubo subgloboso; stamina fertilia 4, didynama, sterile 1, squamaeforme, ad tubi apicem inter lobos binos superiores insertum, interdum nullum; antherae 1 loculares, transverse dehiscentes.

3. Scrofularia L.

b. Corolla tubulosa, tubo cylindrico-elongato; stamina fertilia 2, sterilia 2; antherae 2 loculares, longitudinaliter dehiscentes.

4. Gratiola L.

3. Tribus. *Antirrhineae* Chav. monogr. antirrh. p. 73. — Corolla campanulata vel tubulosa, irregularis, tubo basi saepe saccato vel calcarato; stamina 4, raro 2, omnia fertilia; capsula poris vel dentibus valvaeformibus dehiscens.

a. Corolla basi calcarata.

α. Corollae faux palato clausa.

5. Linaria Tourn.

β. Corollae faux palato depresso pervio.

6. Chaenorrhinum DC.

b. Corolla basi saccata.

7. Antirrhinum L.

B. Corollae labium superius praefloratione a labio inferiore inclusum.

4. Tribus. *Digitaleae* Benth. in DC. pr. X. p. 448. — Corolla tubulosa, campanulata vel rotata, irregularis, lobis planis vel reflexis; stamina 4 vel 2; antherarum loculi apice confluentes; capsula bivalvis.

a. Corolla campanulata vel tubulosa, limbo valde inaequali, subbilabio; stamina 4; capsula septicide dehiscens.

8. Digitalis L.

b. Corolla rotata, tubo brevissimo, limbo parum inaequali; capsula loculicide dehiscens.

α. Stamina 4.

9. Sibthorpia L.

β. Stamina 2.

10. Veronica L.

5. Tribus. *Rhinantheae* Spreng. aul. II. 1 p. 390. — Corolla irregularis bilabiata, labio superiore galeato, rarius plano; stamina 4 vel 2; antherarum loculi distincti; capsula (in nostris) loculicide dehiscens.

a. Ovarii loculi pluriovulati.

α. Calyx subregulariter 4 dentatus vel 4 fidus.

× Semina parva, non lenticulari-compressa.

○ Semina laevia.

11. Parentucellia Viv.

○○ Semina longitudinaliter elevato-costata.

. Capsula compressa.

; Lobi corollae labii inferioris integri.

12. Odontites Hall.

;; Lobi corollae labii inferioris emarginati.

13. Euphrasia L.

.. Capsula turgida.

14. Bellardia All.

XX Semina majuscula, lenticulari-compressa.

15. Alectorolophus Hall.

β. Calyx inaequaliter 5 fidus vel bilabiatus.

× Calyx bibracteolatus.

16. Siphonostegia Benth.

XX Calyx ebracteatus.

○ Calyx tubulosus, saepe inflatus.

17. Pedicularis L.

○○ Calyx a latere compressus.

18. Rhinanthus L.

b. Ovarii loculi biovulati.

19. Melampyrum L.

1. Tribus. VERBASCEAE Benth. in DC. pr. X. p. 224.

1. Verbascum L. gen. n. 245.

Dispositio specierum:

A. Herbae inermes.

1. Sectio. *Thapsus* Rchb. fl. germ. p. 383. — Flores fasciculati vel glomerati; staminum longiorum (2 inferiorum) antherae adnato-decurrentes.

a. Filamentorum lana albida vel lutescens.

α. Corolla concava.

1. V. thapsus L.

β. Corolla omnino explanata.

× Plantae tomento denso, albo vel lutescenti vestitae.

○ Folia caulina a folio ad folium late decurrentia.

2. V. densiflorum Bert. **3. V. macrurum** Ten.

○○ Folia caulina non ad folium proximum usque decurrentia.

. Flores brevissime pedicellati.

4. V. phlomoides L. **5. V. Sartorii** Bois. et Heldr.

.. Flores longiuscule pedicellati, pedicellis nempe aliis calyci aequilongis, aliis eo 2—3plo longioribus.

6. V. longifolium Ten.

XX Planta indumento tenui subdetersili obsitum, viridi-cinerascens; flores pedicellati, pedicellis calyce subaequilongis.

7. V. Guicciardii Bois. et Heldr.

b. Filamentorum lana purpurea vel violacea.

α. Flores subsessiles vel pedicellis calyce brevioribus suffulti.

× Caulis elatus, ramosus.

8. V. malacotrichum Bois. et Heldr. **9. V. foetidum** Bois. et Heldr.

XX Caulis simplex.

10. V. epixanthinum Bois. et Heldr.

β. Flores pedicellati, pedicellis calyce aequilongis vel longioribus.

× Folia radicalia et caulina inferiora in petiolum attenuata.

○ Flores in racemum simplicem vel basi parce breviterque ramulosum dispositi.

. Flores omnes fasciculati.

11. V. taygeteum Hal. **12. V. pindicolum** Freyn. et Sint.

13. V. epirotum Hal.

.. Flores solitarii et 2—3 fasciculati.

14. V. agrimonoides Deg. et Borb.

○○ Flores in paniculam amplam pyramidatam dispositi.

15. V. adenotrichum Hal.

×× Folia radicalia et caulina inferiora petiolata, basi cordata.

16. V. pelium Hal.

2. Sectio. *Lychnitis* Rchb. fl. germ. p. 380. — Flores fasciculati vel glomerati; antherae omnes reniformes, subaequales.

a. Calyx majusculus, 8—10 mm longus; corolla extus stellatim pilosa.

× Flores pedicellati, pedicellis calyce aequilongis; filamentorum lana lutescens.

17. V. Reiseri Hal.

×× Flores sessiles.

○ Filamentorum lana lutescens.

18. V. undulatum Lam. **19. V. pinnatifidum** Vahl.

○○ Filamentorum lana purpurea.

20. V. Boerhavii L.

b. Calyx parvus, 2 mm, ad maximum 6 mm longus.

α. Capsula globosa vel ellipsoidea, calyce ad $2^1/_2$plo longior; corolla extus stellatim pilosa.

× Filamentorum lana albida vel lutescens.

⊙ Planta tota pilis glanduliferis creberrimis viscosa.

21. V. gloeotrichum Haussk. et Heldr.

○○ Planta eglandulosa.

. Flores in racemum simplicem vel basi parce breviterque ramosum dispositi.

22. V. delphicum Bois. et Heldr. **23. V. acutifolium** Hal.

.. Flores in paniculam amplam dispositi.

; Folia (saltem juniora) utrinque vel saltem subtus dense tomentosa.

, Folia utrinque tomento denso persistente vestita.

24. V. mucronatum Lam. **25. V. megaphlomos** Bois. et Heldr.

,, Folia utrinque tomento floccoso tandem detersili obsita.

26. V. pulverulentum Vill.

,,, Folia viridia, supra fere glabra, subtus tomento tenuissimo obsita.

27. V. banaticum Schrad. **28. V. Heldreichii** Bois.

;; Folia (saltem juniora) indumento adpresso, tenuissime arachnoideo cinerascentia, subnitentia.

29. V. Haussknechtii Heldr. **30. V. leucophyllum** Griseb.

×× Filamentorum lana purpurea vel violacea.

○ Folia inferiora in petiolum attenuata.

. Planta eglandulosa.

; Folia crenulata vel integra.

31. V. Adeliae Heldr. **32. V. mallophorum** Bois. et Heldr.

;; Folia sinuato-pinnatifida.

33. V. sinuatum L.

.. Planta glandulosa.

34. V. meteoricum Haussk.

○○ Folia inferiora basi cordata vel rotundata.

35. V. thyrsoideum Host.

β. Capsula cylindrica, calyce 3—5plo longior; corolla glabra.

36. V. graecum Heldr. et Sart.

3. Sectio. *Blattaria* Rchb. fl. germ. p. 379. — Flores solitarii, racemosi vel spicati; antherae adnato-decurrentes vel reniformes.

a. Corolla lutea, antherae staminum longiorum adnato-decurrentes, breviorum reniformes.

37. V. blattaria L.

b. Corolla violacea, antherae omnes reniformes.

38. V. phoeniceum L.

B. Suffrutex erinaceus, ramis spinescentibus.

4. Sectio. *Spinosae* Bois. fl. or. IV. p. 300. — Flores solitarii; antherae reniformes.

39. V. spinosum L.

A. Herbae inermes.

1. Sectio. *Thapsus* Rchb. fl. germ. p. 383. — Flores fasciculati vel glomerati; staminum longiorum (2 inferiorum) antherae adnato-decurrentes.

a. Filamentorum lana albida vel lutescens.

α. Corolla concava.

1. **V. thapsus** L. sp. p. 177; S. et S. pr. I. p. 149; Mazz. in ant. ion. V. p. 210; Ch. et B. exp. p. 73, Fl. pelop. p. 15; Marg. et R. fl. Zante p. 70; Fraas fl. class. p. 191; Bois. fl. or. IV. p. 301; Hal. in z. b. G. 1898 p. 121. — *V. Schraderi* Mey. chlor. hanov. p. 326. — Icon: Fl. dan. t. 631.

Dense lutescenti-tomentosum; simplex vel parce ramosum; foliis crenulatis, radicalibus oblongis, obtusis, in petiolum attenuatis, caulinis oblongo-lanceolatis, acutiusculis, a folio ad folium late decurrentibus; floribus fasciculatis, brevissime pedicellatis, spicam compactam vel basi interruptam formantibus; antheris staminum longiorum glabrescentium breviter decurrentibus, filamento 4 plo brevioribus, breviorum albo-lanatorum reniformibus; capsula ovata, calyce sublongiore. — Corolla flava, parva, circa 15—20 mm. diametro, hac nota primo aspectu a sequentibus distinguendum.

In siccis, lapidosis regionis inferioris. In Graeciae provinciis et insulis frequens (Sibth.); Euboea et Phthiotis (Fraas); Arcadia: pr. Tripolis (Chaub.); Zante (Marg.); Corcyra: pr. Spartilla (Mazz.); an saepius cum aliis speciebus confusum. — Maio, Jul. ⊙ N. v.

β. Corolla omnino explanata.

× Plantae tomento denso, albo vel lutescenti vestitae.

○ Folia caulina a folio ad folium late decurrentia.

2. **V. densiflorum** Bert. rar. pl. dec. III. p. 52, Fl. ital. II. p. 574; Hal. in z. b. G. 1898 p. 122. — Exsicc.: Sint. it. thessal. n. 924.

Dense lutescenti-tomentosum, simplex vel rarius ramosum; foliis crenulatis, radicalibus oblongis, obtusis, in petiolum attenuatis, caulinis oblongis vel oblongo-lanceolatis, acutis vel acuminatis; floribus fasciculatis, brevissime pedicellatis, spicam compactam formantibus; antheris staminum longiorum glabrescentium longe decurrentibus, filamento subduplo brevioribus, breviorum albo-lanatorum reniformibus; capsula ovata, calyce sublongiore. — Corolla flava 3—5 cm. diametro.

β. **thapsiforme** Schrad. in comm. Goett. II. p. 13; Heldr. fl. cephal. p. 55; pro sp.; Vis. fl. dalm. II. p. 154; Hal. l. c. . — *V. thapsus* Dallap. prosp. p. 23, non L. . — Spica laxiuscula, basi saepe ramosa. — Exsicc.: N. v.

In lapidosis, sterilibus regionis inferioris et montanae. Thessalia: mt. Sina pr. Malakasi (Sint.); — *β.* Cephalonia: pr. Argostoli, Pessada, mt. Aenos (Heldr.). — Maio, Jul. ⊙

3. **V. macrurum** Ten. fl. neap. pr. app. V. p. 9, Fl. nap. III. p. 216 t. 214; Form. in D. bot. Mon. 1891 p. 26; Hal. Beitr. fl. Achaia p. 27, in z. b. G. 1898 p. 122; Bald. viagg. Creta p. 79. — *V. Schraderi* Raul. cret. p. 819, non Mey. . — *V. thapsiforme* Bois. fl. or. IV. p. 301, saltem quoad pl. graecam, non Schrad. . — Exsicc.: Orph. fl. gr. n. 733; Heldr. pl. fl. hellen. a. 1887.

Dense albido-tomentosum, simplex; foliis crenulatis, oblongo-lanceolatis, radicalibus obtusiusculis, in petiolum attenuatis, caulinis acutis vel acuminatis; floribus fasciculatis, brevissime pedicellatis, spicam longissimam et densissimam formantibus; antheris staminum longiorum glabrescentium breviter decurrentibus, filamento 4 plo brevioribus, breviorum albo-lanatorum reniformibus; capsula ovata, calyce sublongiore. — Certe species propria, a praecedente spica valde elongata, densissima, valde lanata et antheris breviter decurrentibus optime distincta.

In lapidosis praesertim montanis. Thessalia: mt. Ossa pr. Selicany et Spilia (Form.); Peloponnesus: in regione abietina mt. Panachaicon (Hal.), mt. Kyllene (Heldr.), mt. Malevo supra Kastanitza (Orph.); Creta: ad promontorium Meleka, pr. Lutro, Rhodakino, Avdhu (Raul.), mt. Ida (Bald.). Maio, Jul. ⊙

○○ Folia caulina non ad folium proximum usque decurrentia.

. Flores brevissime pedicellati.

4. **V. phlomoides** L. sp. p. 1194; Mazz. in ant. ion. V. p. 210; Bois. fl. or. IV. p. 301; Form. in D. bot. Monat. 1891 p. 26, in Ver. Brünn 1896 p. 62; Haussk. symb. p. 161; Hal. in z. b. G. 1898 p. 123. — *V. australe* Schrad. in comm. goett. II. p. 24 t. 2. — Exsicc.: Sint. et Bornm. it. thessal. n. 1391; Orph. fl. gr. n. 1008, mt. Korthiati.

Dense tomentosum, simplex vel superne ramosum; foliis crenulatis, radicalibus ellipticis vel oblongo-lanceolatis, in petiolum attenuatis, caulinis ovatis, ellipticis vel oblongo-lanceolatis, acutis vel acuminatis,

amplexicaulibus, saepe cordatis vel brevissime decurrentibus; floribus fasciculatis, brevissime pedicellatis, spicam longam inferne interruptam et saepius laxam formantibus: antheris staminum longiorum glabrescentium longe decurrentibus, filamento subduplo brevioribus, breviorum albo-lanatorum reniformibus; capsula ovata, capsula sesquilongiore. — Corolla magna 3—5 cm. diametro, flava.

In siccis montanis. Epirus: mt. Peristeri (Form.); Thessalia: pr. Karditza, mt. Pelion (Form.), pr. Katherina (Sint.); Euboea: mt. Tympanon, Dirphys (Heldr.): Argolis: mt. Chelone in peninsula Methana (Haussk.); Cycladum insula Tenos (Sart.); Corcyra: pr. Klonu, Pauliana (Mazz.). — Maio, Aug. ⊙

Obs. *V. phlomoides* S. et S. pr. I. p. 149 (circa Byzantium). Fl. gr. III. p. 19 t. 224; Ch. et B. fl. pelop. p. 15 (Kalamata Messeniae) ob filamentorum lanam purpuream ad speciem aliam mihi ignotam spectant.

5. **V. Sartorii** Bois. et Heldr. diagn. VII. p. 38: Hal. Beitr. fl. Achaia p. 27; Haussk. symb. p. 160. — *V. phlomoides β. Sartorii* Bois. fl. or. IV. p. 302. — Exsicc.: Dörfl. fl. aeg. n. 108.

Praecedenti valde affine, ab eo inflorescentia saepe densius tomentosa, filamentis omnibus albo-lanatis, antheris staminum longiorum breviter decurrentibus, filamento 3—4 plo brevioribus specifice differre videtur.

In siccis montanis et regionis inferioris. Thessalia: in Pindo dolopico, pr. Aivali, Volo (Haussk.); Attica: mt. Pentelicon (Heldr.): Achaia: mt. Omplo pr. Patras (Heldr.), mt. Kastro pr. Kalavryta (Hal.); Messenia: mt. Ithome (Heldr.); Cycladum insula Paros (Leon.): Zante (Marg.). — Maio, Jul. ⊙

5 × 18. **V. Sartorii × rigidum** Hal. in z. b. G. 1898 p. 124. — *V. phlomoides × rigidum* (*V. semirigidum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72. — *V. rigidum × Sartorii* (*V. semirigidum*) Haussk. symb. p. 166. — Habitu quasi quodam *V. Sartorii* ramoso, fasciculis remotis et hac nota ac calycis laciniis longius acuminatis ad *V. rigidum* accedens, sed indumentum densius, magis lutescens et non detersile; capsulae steriles. — In dumetis inter parentes pr. Volo et in ruderatis pr. Karditza Thessaliae (Haussk.). — N. v.

5 × 26. **V. Sartorii pulverulentum** Hal. in z. b. G. 1898 p. 124. — *V. floccosum × phlomoides* (*V. subphlomoides*) Haussk. in Mitth. thür. bot. Ver. V. 71. — *V. pulverulentum × Sartorii* (*V. subphlomoides*) Haussk. symb. p. 164. — Differt a *V. Sartorii* indumento dilutiore, facile detersile, caule magis ramoso, foliis minute crenulatis, floribus minoribus, manifeste pedicellatis et antheris staminum longiorum brevius decurrentibus: a *V. pulverulento* ramis crassioribus brevioribusque, florum fasciculis densioribus, floribus majoribus, antheris staminum longiorum breviter decurrentibus et indumento minus floccoso. — Ad monasterium Korona in Pindo (Haussk.). — N. v.

5 × 28. **V. Sartorii × Heldreichii** Hal. in z. b. G. 1898 p. 125. — *V. Heldreichii × phlomoides* (*V. coenobitarum* Haussk. et Heldr. in Mitth. thür. bot. Ver. V. p. 72. — *V. Heldreichii × Sartorii* (*V. coenobitarum*) Haussk. et Heldr. in Haussk. symb. p. 165. — A *V. Sartorii* inflorescentia paniculata, foliis inferioribus grosse crenatis, tenuiter tomentellis, floribus minoribus et antheris staminum longiorum brevius decurrentibus; a *V. Heldreichii* ramorum indumento magis tomentoso, florum fasciculis minus remotis, floribus fere duplo majoribus, brevius pedicellatis et antheris staminum longiorum subdecurrentibus, discedens; capsulae steriles. — Exsicc.: Heldr. it. IV. thessal. a. 1885. — In nemorosis quercinis regionis inferioris mt. Pindus circa monasterium Korona (Heldr.).

5 × 33. **V. Sartorii × sinuatum** (*V. pseudosinuatum*) Haussk. symb. p. 166; Hal. in z. b. G. 1898 p. 125. — *V. phlomoides × sinuatum* (*V. pseudosinuatum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72. — Quoad foliorum formam nunc ad unam nunc ad alteram stirpem accedens; a *V. Sartorii* foliis inferioribus irregulariter sinuatis, superioribus sinuato-crenatis et filamentorum lana purpurascente; a *V. sinuato* tomento crassiore, caulibus strictis, crassioribus, dense tomentellis, nec glabrescentibus, ramis crassioribus brevioribus, strictis, nec divaricatis, fasciculis florum irregulariter congestis, foliis inferioribus minus profunde sinuatis, superioribus basi rotundato sessilibus, floribus majoribus et antheris staminum longiorum breviter decurrentibus distinctum; capsulae steriles. — Thessalia: ad monasterium Korona in Pindo (Haussk.). — N. v.

.. Flores longiuscule pedicellati, pedicellis aliis nempe calyce aequilongis, aliis eo 2—3 plo longioribus.

6. **V. longifolium** Ten. fl. nap. I. p. 89, t. 21; Bald. riv. coll. bot. Alb. 1896 p. 80. — Exsicc.: Bald. it. alb. epir. IV. n. 135.

Dense lutescenti-tomentosum, spica simplici vel inferne parce et breviter ramulosa; foliis acutis, integris crenulatisve, radicalibus longe oblongo-lanceolatis, in petiolum attenuatis, caulinis oblongis, subcordato-amplexicaulibus; floribus fasciculatis, dense racemosis; antheris staminum longiorum glabrescentium decurrentibus, filamento 3—4 plo brevioribus, breviorum albo-lanatorum reniformibus; capsula ovata, calyce sublongiore. — Floribus longiuscule pedicellatis, calycibus lutescenti-tomentosis a praecedentibus distinctum.

In herbidis alpinis mt. Papignon distr. Zagorion in Epiro (Bald.). — Jul. Aug. ⊙

×× Planta indumento tenui subdetersili obsitum, viridicinerascens; flores pedicellati, pedicellis calyce subaequilongis.

7. **V. Guicciardii** Bois. et Heldr. diagn. ser. 2 VI. p. 127, Fl. or. IV. p. 303; Heldr. chlor. Parn. p. 24; Hal. Beitr. fl. Aetol. p. 9; in z. b. G. 1898 p. 125; Haussk. symb. p. 160. — Exsicc.: Heldr. herb. norm. n. 696; Hal. it. gr. II. a. 1893.

Caule simplici, glabriusculo, in racemum longum, angustum, laxiusculum, inferne interruptum abeunti; foliis repando-crenatis, subtus tenuiter tomentellis, supra subglabris, infimis oblongis, in petiolum attenuatis, caulinis cordato-amplexicaulibus; floribus fasciculatis; antheris staminum longiorum glabrescentium breviter decurrentibus, filamento multoties brevioribus, breviorum albo-lanatorum reniformibus; capsula sphaerica, calyce sublongiore. — Caulis 50—130 cm. altus, fuscus, subnitens, florum fasciculis praesertim infimis valde remotis, racemum ad 70 cm. longum formantibus; corolla flava, 25—30 mm. diametro; capsula magnitudine pisi.

In fruticetis, ad sepes regionis inferioris et montanae, rarissime. Thessalia: mt. Ghavellu in Pindo dolopico (Haussk.); Acarnania: in marginibus silvularum sempervirentium ad pedem mt. Makrinoros non procul a pago Menidion ad sinum Ambracicum (Hal.); ad radices mt. Parnassus l. d. Pagna (Guicc.). — Jun. Jul. ⊙

7 × **33. V. Guicciardii** × **sinuatum** Hal. Beitr. fl. Aetol. p. 9, in z. b. G. 1898 p. 126. — *V. ambracicum* Hal. l. c. . — Indumento tenui subdetersili obsitum, viridi-canescens; foliis radicalibus oblongo-spathulatis, sinuato-dentatis, in petiolum attenuatis, caulinis oblongis, sessilibus; floribus breviter pedicellatis, fasciculatis, in paniculam pyramidatam dispositis; corolla ampla, 3 cm. diametro; filamentis albo-lanatis; antheris omnibus reniformibus; capsula sterili. — Differt a *V. Guicciardii* indumento subdensiori, inflorescentia paniculata, foliis radicalibus sinuato-dentatis et antheris omnibus reniformibus; a *V. sinuato* praesertim corolla magna et filamentis albo-lanatis. — Acarnania: ad pedem mt. Makrinoros pr. Menidion ad sinum Ambracicum (Hal.).

b. Filamentorum lana purpurea vel violacea.

α. Flores subsessiles vel pedicellis calyce brevioribus suffulti.

× Caulis elatus, ramosus.

8. **V. malacotrichum** Bois. et Heldr. diagn. ser. 2 III. p. 142, Fl. or. IV. p. 305; Hal. in z. b. G. 1898 p. 126. — Exsicc.: Orph. fl. gr. n. 734 (mt. Corthiati); Sint. et Bornm. it. turc. n. 1390.

Totum lana alba, floccosa, tandem detersili obsitum; foliis subcrenatis, inferioribus oblongis, in petiolum attenuatis, ceteris basi rotundata vel cordata sessilibus; floribus fasciculatis, breviter pedicellatis, racemos longos laxiusculos formantibus; bracteis lanceolatis, glanduliferis, calyce aequilongis; calycis laciniis lanceolatis, glanduliferis; filamentis violaceo-barbatis, antheris longiorum longe adnato-decurrentibus, breviorum reniformibus; capsula subglobosa, calyce sublongiore. — Caulis elatus (observante Sintenis ad 3—4 m. altus), fuscus, inferne parum, superne densius floccosus et hic sicuti bracteae et calyces pilis glanduliferis brevibus obsitus; corolla flava.

In regione silvatica mt. Olympus prope coenobium Hagios Dionysios (Heldr.). — Jul. Aug. ⊙

9. **V. foetidum** Bois. et Heldr. diagn. ser. 2 III. p. 141, Fl. or. IV. p. 305; Heldr. chlor. Parn. p. 24; Hal. in z. b. G. 1898 p. 127. — Exsicc.: Orph. herb. a. 1854.

Tenuiter tomentosum, flavo-virens; foliis radicalibus amplis, repando-crenatis, ovatis vel oblongis, breviter petiolatis, caulinis basi rotundata vel cordata sessilibus; floribus fasciculatis, breviter pedicellatis, racemos longos laxos formantibus; bracteis linearibus, glanduliferis, calyce aequilongis; calycis laciniis lanceolatis, glanduliferis; filamentis violaceo-barbatis, antheris longiorum tota longitudine adnatis, breviorum reniformibus; capsula ovata, calycinis laciniis duplo longiore. — Species autoribus ipsis incerta videtur, mihi ex unico specimine incompletissimo tantum nota et ulterius observanda. — Odor totius plantae fere *Daturae stramonii*. Characteribus ad *V. malacotrichum* accedit, quod vero indumento albido, capsula duplo minori subglobosa differre dicitur.

In regione abietina mt. Parnassus ad Dipotamo (Heldr.). — Jul. ⊙

×× Caulis simplex.

10. **V. epixanthinum** Bois. et Heldr. diagn. VII. p. 39, Fl. or. IV. p. 304; Heldr. chlor. Parn. p. 24, in Sitzungsb. acad. Wiss. Berl. 1883 p. 4; Hal. in z. b. G. 1898 p. 127. — Exsicc.: Heldr. herb. norm. n. 695.

Caule erecto, breviter tomentello, dense glanduloso; foliis oblongis, obtusiusculis, obsolete crenulatis, utrinque dense albo-tomentosis, subtus dilutioribus, inferioribus in petiolum attenuatis, superioribus multo minoribus, basi attenuata sessilibus; floribus fasciculatis, subsessilibus, laxe spicatis; bracteis et calycis laciniis dense tomentosis, glandulosis; filamentis violaceo-barbatis, antheris longiorum adnato-decurrentibus, filamento 3 plo brevioribus, breviorum reniformibus. — Caulis 25—50 cm. altus; corolla 20—25 mm. diametro, lutea, basi saepe violacea. — A duobus praecedentibus caule nano, simplici, foliis infimis approximatis, tomento denso et inflorescentia crebre glandulosa egregie distinctum.

β. **Samaritanii** Heldr. in Bois. diagn. ser. 2 VI. p. 127; Hal. in z. b. G. 1888 p. 18; Heldr. chlor. Parn. p. 24, pro sp.; Bois. fl. or. IV. p. 304; Hal. in z. b. G. 1898 p. 128. — Caulis densissime glandulosus; folia superne aeruginoso-aureo, subtus cano-griseo tomentosis; flores alii subsessiles, alii pedicellati. — Exsicc.: Hal. it. gr. a. 1888.

In herbidis lapidosis regionis alpinae mt. Parnassus, Olenos et Taygetos (Heldr.); indicatur quoque in mt. Pelion Thessaliae (Heldr.), sed probabiliter ex confusione; — β. mt. Tymphrestus (Samar.), Kiona (Hal.). — Jul. Aug. ⊙

β. Flores pedicellati, pedicellis calyce aequilongis vel longioribus.

× Folia radicalia et caulina inferiora in petiolum attenuata.

○ Flores in racemum simplicem vel basi parce breviterque ramulosum dispositi.

. Flores omnes fasciculati.

11. **V. taygeteum** Hal. in z. b. G. 1898 p. 128. — Exsicc.: Pichl. pl. gr. a. 1876.

Caule erecto, tomentoso, parce glanduloso; foliis oblongo-lanceolatis, acuminatis, obsolete crenulatis, utrinque albotomentosis, subtus dilutioribus, infimis in petiolum brevissimum attenuatis, caulinis paucis, multo minoribus, basi attenuata sessilibus; floribus fasciculatis, laxiuscule racemosis; pedicellis calyce aequilongis vel duplo longioribus; bracteis et calycis laciniis tomentosis, parce glandulosis; filamentis violaceo-barbatis, antheris longiorum adnato-decurrentibus, filamento 4 plo brevioribus, breviorum reniformibus. — Caulis 25 cm. altus, folia 10 cm. longa, 3 cm. lata; corolla lutea 20—25 mm. diametro. — Praecedenti affine, quocum habitu et praesertim foliis infimis approximatis, caulinis subito decrescentibus congruit, sed ab eo tomento laxiore, foliis acuminatis, floribus longe pedicellatis et glandulositate parciore specifice differre videtur.

In regione alpina mt. Taygetos Laconiae (Pichl.). — Jul. Aug. ⊙

12. **V. pindicolum** Freyn. et Sint. in bull. herb. Bois. V. p. 793; Hal. in z. b. G. 1898 p. 129. — *V. Samaritanii* Haussk. symb. p. 160, vix Bois. — *V. tymphaeum* Freyn. et Sint. in bull. herb. Bois. V. p. 795; Hal. in z. b. G. 1898 p. 135; quod ex exsiccatis Sint. it. thessal. n. 923 iterum examinatis a *V. pindicolo* meo sensu non differt. — Exsicc.: it. thessal. IV. a. 1885; Sint. it. thessal. n. 922.

Caule erecto, tomentoso, parce glanduloso; foliis oblongis vel oblongo-lanceolatis, acutis vel acuminatis, obsolete crenulatis, superne aeruginoso-aureo, subtus cano-griseo tomentosis, inferioribus in petiolum longum attenuatis, superioribus sursum decrescentibus, basi subrotundata sessilibus; floribus fasciculatis, laxiuscule racemosis; pedicellis calyce aequilongis vel duplo longioribus; bracteis et calycis laciniis tomentosis, parce glandulosis; filamentis violaceo-barbatis, antheris longiorum breviter adnato-decurrentibus, filamento multoties brevioribus, breviorum reniformibus. — Caulis 40—70 cm. altus; folia inferiora cum petiolo ad 20 cm. longa, 7 cm. lata, corolla 30—35 mm. diametro. — *V. Samaritanii*, nec *V. sinuato* et *V. Adeliae*, ut voluerunt auctores, maxime affine et ab eo foliis acuminatis et floribus longius pedicellatis probabiliter vix specifice distinctum. Affine quoque *V. taygeteo*, foliis infimis brevissime petiolatis, caulinis paucis, antheris longius decurrentibus, diverso.

In regione subalpina Thessaliae: mt. Plaka pr. Chaliki, mt. Dokimi. (Sint.), mt. Ghavellu (Haussk.) et Karava (Heldr.) in Pindo. — Jun. Jul. ⊙

13. **V. epirotum** Hal. in z. b. G. 1898 p. 129. — *V. epixanthinum* Hal. Beitr. Fl. Epir. p. 32, non Bois. et Heldr. . — Exsicc.: Hal. it. gr. II. a. 1893.

Caule erecto, breviter tomentello, eglanduloso, aequaliter foliato; foliis lanceolatis oblongo-lanceolatisve, acutis vel acuminatis, obsolete crenulatis, superne tenuiter tomentosis, viridibus, subtus cano-tomentosis ab infimis parvis ad intermedia sursum auctis, in petiolum attenuatis,

superioribus basi subrodundata sessilibus; floribus fasciculatis, densiuscule racemosis; pedicellis calyce aequilongis; bracteis parce glandulosis calycis laciniisque lutescenti-tomentosis; filamentis violaceo-barbatis, antheris longiorum adnato-decurrentibus, filamento 4 plo brevioribus, breviorum reniformibus. — Caulis 40—50 cm. altus, simplicissimus, folia media 15 cm. longa, 3—5 cm lata; corolla lutea, 20—25 mm. diametro, basi saepe violacea. — A praecedente foliis supra tenuiter tomentosis, viridibus, ab infimis ad intermedia sursum auctis, inflorescentia flavido-tomentosa, glandulis fere omnino destituta et pedicellis brevioribus specifice distinctum.

.. Flores solitarii et 2—3 fasciculati.

14. **V. agrimoniodes** Deg. et Borb. in természettud. közl. pótfüz. XLI. p. 85; Form. in Ver. Brünn 1897 p. 46. — Exsicc.: Form. pl. thessal. a. 1896.

Totum lana detersili obsitum, superne glandulosum; caule rubescente, in racemum elongatum, angustum, simplicissimum, laxissimum abeunte; foliis oblongis vel oblongo-lanceolatis, acutis, obsolete crenulatis, inferioribus in petiolum longum attenuatis, superne demum viridescentibus, superioribus sursum diminutis, sessilibus; pedicellis calyce aequilongis vel longioribus; filamentis purpureo-barbatis, antheris longiorum adnato-decurrentibus, breviorum reniformibus. — Species ex unico specimine incompleto, in herbario Borbasiano asservato, descripta, *V. pindicolo* ut videtur maxima affinis, ulterius inquirenda et observanda.

Prope Phlambures et in mt. Mitriza in mt. Chassia Thessaliae (Form.). — Jul. Aug. ⊙

○○ Flores in paniculam amplam pyramidatam dispositi.

15. **V. adenotrichum** Hal. in z. b. G. 1898 p. 130. — *V. Samaritanii* Hal. Beitr. fl. Thessal. p. 16, non Heldr.. — *V. viscidulum* Freyn. et Sint. in bull. herb. Bois. V. p. 792, non Pers. syn. I. p. 215. — Exsicc.: Hal. it. gr. II. a. 1893; Sint. it. thessal. n. 1218.

Totum glanduloso-viscidum, parce tomentellum, viride; caule elato, in paniculam amplam, virgato-ramosissimam abeunte; foliis inferioribus amplis, late ellipticis, acutis, in petiolum attenuatis, superioribus sursum diminutis ellipticis vel lanceolatis, crenulatis vel serratis, acuminatis, sessilibus; floribus fasciculatis, in racemos laxiusculos dispositis; pedicellis calyce aequilongis vel longioribus; antheris staminum longiorum parce lanatorum breviter decurrentibus, breviorum omnino lanatorum reniformibus. — Species egregia, caule $1^1/_2$—2 m. alto, paniculae ramis ad 56 cm. longis, foliis radicalibus 25—30 cm. longis, 10—15 cm. latis, corolla 3 cm. diametro.

Ad silvarum oras regionis montanae, rarissime. Thessalia: in valle fluvii Prion (Sint.), mt. Oxya supra Kastania non procul a statione militari (Hal.). — Jul. ⊙

×× Folia radicalia et caulina inferiora petiolata, basi cordata.

16. **V. pelium** Hal. in z. b. G. 1898 p. 131. — Exsicc.: Hal. it. thess. a. 1882; Heldr. pl. fl. hell. a. 1887, quod prius incaute pro *V. mallophoro* habui.

Caule elato, ? simplici, parce floccoso, eglanduloso; foliis obtuse crenatis, superne tenuiter, subtus dense flavido-tomentosis, inferioribus cordato-ovatis ellipticisve, obtusis, petiolatis, superioribus ovatis vel ovato-oblongis, acuminatis, cordato-semiamplexicaulibus; floribus fasciculatis, in racemum densiusculum dispositis; bracteis et calycis laciniis lanceolatis, tomentoso-floccosis, parce glandulosis; antheris staminum longiorum glabrescentium adnato-decurrentibus, filamento 3 plo brevioribus, breviorum purpureo-lanatorum reniformibus. — Foliis inferioribus cordatis insigne.

In regione superiori mt. Pelion supra Portaria (Heldr.) et Zagora (Aphentulis). — Jul. Aug. ⊙

2. Sectio. *Lychnitis* Rchb. fl. germ. p. 380. — Flores fasciculati vel glomerati; antherae omnes reniformes, subaequales.

a. Calyx majusculus, 8—10 mm. longus; corolla extus stellatim pilosa.

× Flores pedicellati, pedicellis calyce aequilongis; filamentorum lana lutescens.

17. **V. Reiseri** Hal. in z. b. G. 1898 p. 132. — Exsicc.: Reis. fl. gr. a. 1894.

Totum dense albo-tomentosum; caule erecto, simplici; foliis obsolete crenulatis; inferioribus ellipticis vel elliptico-rotundatis, obtusis, breviter apiculatis, in petiolum longum attenuatis, superioribus ovatis, sessilibus sursum in bracteas abeuntibus; floribus fasciculatis, in racemum densum simplicem dispositis; calycis laciniis lanceolatis; corolla flava. — Ex unico specimine mihi tantum notum, caule 20 cm. alto, densissime floccoso-tomentoso, crassiusculo, foliis inferioribus cum petiolo 10—12 cm. longis, 4 cm. latis, calyce 8 mm. longo, corolla 2 cm. diametro. Habitu magis ad species sectionis praecedentis accedens, sed antherae omnes reniformes.

Aetolia: in mt. Vardusia pr. Granitza (Reis.). — Jul. ⊙

×× Flores sessiles.

○ Filamentorum lana lutescens.

18. **V. undulatum** Lam. dict. IV. p. 221 p. p.; Ch. et B. fl. pelop. p. 15; Friedr. Reise p. 274; Clem. sert. p. 71; Hal. in z. b. G. 1898 p. 132. — *V. plicatum* S. et S. pr. I. p. 150, Fl. gr. III. p. 21 t. 226; Link in Linnaea IX. p. 577; Fraas fl. class. p. 190; Bois. fl. or. IV. p. 312; Haussk. symb. p. 160; Heldr. fl. Aegina p. 381. — Exsicc.: Heldr. herb. norm. n. 399 et 1161; Orph. fl. gr. n. 1160; Dörfl. fl. gr. n. 225.

Totum adpresse albo-tomentosum; caule erecto, simplici vel ramoso; foliis inferioribus in petiolum brevem attenuatis, oblongo-spathulatis, undulato-plicatis, sinuato-pinnatifidis, lobis rotundatis, subimbricatis,

superioribus cordato-amplexicaulibus, ovatis, subintegris; floribus fasciculatis, racemos longos interruptos formantibus; calycis laciniis lanceolatis; corolla lutea; capsula ovato-globosa, calyce aequilonga. — Foliis undulato-plicatis insigne. Caulis 50—100 cm. altus, folia inferiora 10—25 cm. longa, calyx 10 mm. longus, corolla 25—35 mm. diametro.

β. **rigidum** Bois. et Heldr. diagn. ser. 2 II. p. 143; Hal. in ö. b. Z. 1892 p. 375; Haussk. symb. p. 161; pro sp.; Bois. fl. or. IV. p. 312; Hal. Beitr. fl. Aetol. p. 9, Beitr. fl. Achaia p. 27, in z. b. G. 1898 p. 132; Bald. riv. coll. bot. Alb. 1896 p. 81. — Indumentum tenuius, griseum; folia inferiora remote sinuata, parum undulato-plicata vel (*V. calaurium* Heldr. exs. a. 1854) plana, leviter sinuato-undulata. — Formis intermediis cum typo conjunctum. — Exsicc.: Orph. fl. gr. n. 732 (mt. Corthiati); Sint. et Bornm. it. turc. n. 394; Sint. it. thessal. n. 30.

In collibus arenosis, locis asperis regionis inferioris et submontanae. Attica: copiose, saepe gregarie pr. Athenas, in Acropoli, ad Phalerum, mt. Parnes, Pentelicon, Hymettus, pr. Sunium; Phocis: ad radices Parnassi pr. Rachova (Leon.); Achaia: pr. Patras (Link.); insula Aegina (Heldr.); Argolis: pr. Nauplia (Link.), Poros (Friedr.), insula Hydra (Sibth.); Cycladum insula Keos, Kythnos (Tourn. voy. p. 25); — *β.* Epirus: ad Vromonero distr. Ljaskovik (Bald.); Thessalia: monasterium Meteora, pr. Karditza, Pharsalus, Aivali, Pirsufli, Volo (Haussk.), mt. Pelion pr. Volo et Milies (Heldr.), pr. Katerina (Sint.); insula Scopelos (Leon.); Euboea: mt. Telethrion (Heldr.); Aetolia: mt. Taphiassos et Chalkis (Hal.); Achaia: pr. Patras (Hal.); Argolis: peninsula Methana (Haussk.), insula Poros (Haussk.). — Jun. Jul. ⊙

18 × 19. V. undulatum pinnatifidum Hal. in z. b. G. 1898 p. 133. — *V. pinnatifidum* × *plicatum* (*V. paradoxum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 167. — A *V. undulato* caule graciliori, foliis minoribus, dentatis vel pinnatifidis, non undulato-plicatis et indumento tenui; a *V. pinnatifido* caule firmiore, minus ramoso, foliis minus divisis; saepe tantum dentatis, fasciculis florum magis approximatis et corolla majori distinctum. — In arenosis maritimis ad Phaleron Atticae (Haussk.). — Exsicc.: Haussk. it. gr. a. 1885.

18 × 33. V. undulatum sinuatum Hal. in z. b. G. 1898 p. 133. — *V. plicatum* × *sinuatum* (*V. erraticum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 167; Hal. in z. b. G. 1888 p. 763. — Hybrida sat frequens, nunc (*f. undulatiforme*), ad *V. undulato* nunc (*f. sinuatiforme*) ad *V. sinuato* magis accedens, a priori caule graciliori, ramosiori, foliis minoribus, minus undulato-plicatis, superioribus saepius subdecurrentibus, floribus duplo minoribus, nonnullis areviter pedicellatis et filamentorum lana pallide purpurascente; ab bltero caule minus ramoso, ramis brevioribus, foliis undulato-plicatis et floribus brevius pedicellatis, nonnullis sessilibus distinctum. — Exsicc.: Haussk. it. gr. a. 1885; Heldr. herb. norm. n. 1064. — Attica: ad

Phalerum, Kephissia, ad monasterium Mendeli in mt. Pentelicon (Haussk.), pr. Tatoi ad radices mt. Parnes (Hal.); Euboea: mt. Telethrion (Heldr.).

19. **V. pinnatifidum** Vahl symb. II. p. 39; Urv. enum. p. 25; Ch. et B. fl. pelop. p. 15; Bois. fl. or. IV. p. 312; Haussk. symb. p. 161; Heldr. fl. Aegina p. 38; Hal. in z. b. G. 1898 p. 133. — Icon: S. et S. fl. gr. III. t. 228. — Exsicc.: Orph. fl. gr. n. 369; Heldr. herb. norm. n. 398 et 1162.

Tenuiter tomentosum, tandem plus minusve glabratum, caule rigido, a basi pyramidatim paniculato; foliis oblongo-lanceolatis, radicalibus petiolatis, pinnatifidis partitisque, segmentis obtuse lobatis, foliis caulinis sessilibus pinnatifidis vel dentatis; floribus fasciculatis, racemos virgatos interruptos formantibus; calycis laciniis lanceolatis; corolla lutea; capsula ovata, calyce aequilonga. — Caulis 30—45 cm. altus, a basi longe ramosus, ramis fere pungentibus; calyx 8 mm. longus; corolla 25 mm. diametro.

β. **dentatum** Haussk. l. c.; Hal. l. c. — Foliis inferioribus dentatis. — Exsicc.: Heldr. pl. fl. hell. a. 1890.

In arenosis maritimis. Thessalia (Aucher); insula Euboea, Aegina, Hydra (Sprun.); frequens ad Phalerum pr. Athenas; insula Syra (Urv.); — *β.* ad Phalerum (Haussk.); Euboea: pr. Oreus (Heldr.). — Jun. Jul. ⊙

19 × 33. V. pinnatifidum × sinuatum (*V. phalereum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 68. — Caule longe divaricatim ramoso; foliis eis *V. pinnatifidi* duplo majoribus, segmentis longioribus, acutis, dentatis et breviter lobatis; floribus sessilibus et breviter pedicellatis; filamentorum lana pallide purpurescente; capsula sterili. — In arenosis maritimis ad Phalerum Atticae (Haussk.). — N. v.

○○ Filamentorum lana purpurea.

20. **V. Boerhavii** L. mant. p. 45; Bois. fl. or. IV. p. 314; Hal. in z. b. G. 1898 p. 134. — Icon: Rchb. t. 33.

Tomento floccoso, tantum detersili obsitum; caule elato, in racemum longum, inferne interruptum abeunti; foliis crenato-dentatis, inferioribus ovatis, obtusis, petiolatis, superioribus cordato-amplexicaulibus, acutis vel acuminatis; floribus fasciculatis, subsessilibus; calycis laciniis lanceolatis, glandulosis; corolla lutea, ad faucem violaceo-maculata; capsula ellipsoidea, calyce tertia parte longiore.

Messenia: pr. Kalamata (Gittard), sed schedulis hujus collectoris pauca fides habenda (Bois. l. c.); nemini postea lectum. — Maio, Jul. ⊙ N. v.

b. Calyx parvus, 2 mm., ad maximum 6 mm. longus.

α. Capsula globosa vel ellipsoidea, calyce ad $2^1/_2$ plo longior; corolla extus stellatim pilosa.

× Filamentorum lana albida vel lutescens.

○ Planta tota pilis glanduliferis creberrimis viscosa.

21. **V. gloeotrichum***) Haussk. et Heldr. in Mitth. thür. bot. Ver. V. p. 87, symb. p. 162; Hal. in z. b. G. 1898 p. 134. — *V. perviscosum* Borb. ap. Form. in Ver. Brünn 1897 p. 46, solum nomen (ex specimine authentico Formanekiano). — Exsicc.: Heldr. it. thessal. IV. a. 1885; Sint. it. thessal. n. 1000.

Caule erecto, paulo supra basin in ramos virgatos diviso; foliis crenulatis, cinereo-tomentosis, radicalibus ovato- vel oblongo-spathulatis, obtusiusculis, in petiolum attenuatis, caulinis oblongo-lanceolatis, acutis vel acuminatis, semiamplexicaulibus; florum fasciculis dissitis, 1—6 floris; pedicellis calyce aequilongis vel subbrevioribus; calycis laciniis lanceolatis; filamentis longioribus ad tertiam partem, brevioribus tota longitudine sordide albo-lanatis; capsula ovata, calyce aequilonga. — Species egregia, siccando facile nigricans, caule pluripedali, corolla lutea, pellucido-punctata, 25—35 mm. diametro.

In collibus herbosis Thessaliae: pr. Meteora, Kalabaka, Trikala, Zarkos (Haussk.), Povelci, Kerasia Sina (Form.). — Jun. Jul. ⊙

○○ Planta eglandulosa.

. Flores in racemum simplicem vel basi parce breviterque ramosum dispositi.

22. **V. delphicum** Bois. et Heldr. diagn. ser. 2 III. p. 146, Fl. or. IV. p. 324; Hal. in z. b. G. 1898 p. 136. — Exsicc.: Heldr. herb. norm. n. 799.

Caule erecto, glabrescenti, in racemum simplicem vel inferne parce breviterque ramosum abeunti; foliis ovatis, crenatis, supra viridibus, subtus tomentoso-canescentibus, infimis petiolatis, basi rotundatis vel cordatis, superioribus diminutis, cordato-semiamplexicaulibus; florum glomerulis racemos densiusculos formantibus; pedicellis calyce longioribus; calycis tomentosi laciniis lanceolatis; filamentis albo-barbatis; capsula ovata, calyce duplo longiore. — Caulis 40—60 cm. altus, fuscus; foliorum infimorum lamina 15 cm. longa, 8 cm. lata, supra tenuiter tomentella; petiolus 5 cm. longus; calyx 2—3 mm. longus; corolla flava, 20 mm. diametro; capsula 4—5 mm. longa. Habitu *V. lanato* Schrad. accedens, quod autem filamentis purpureo-lanatis specifice discedit.

In regione silvatica et superiori mt. Dirphys, usque ad cacumen l. d. Fleba (Heldr.). — Jun. Jul. ⊙

23. **V. acutifolium** Hal. in z. b. G. 1898 p. 136. — *V. pulverulentum* Hal. Beitr. fl. Epir. p. 33 p. p. — Exsicc.: Hal. it. gr. II. a. 1893.

Caule erecto, glabrescenti, in racemum inferne parce breviterque ramosum abeunti; foliis integris, supra tenuiter tomentellis, viridibus, subtus cano-tomentosis, infimis oblongo-lanceolatis, acutissimis, in petiolum brevissimum attenuatis, superioribus subito decrescentibus, oblongis vel ovatis, acuminatis, basi semiamplexicauli sessilibus; floribus fasciculatis, in racemum elongatum, laxiusculum, basi ramulosum dispositis; pedicellis calyce aequilongis vel parum longioribus; calycis floccosi laciniis

* Nomen specificum deductum a γλοιός = viscositas et θρίξ = pilum, ideo *gloeotrichum* nec *glocotrichum*, ut vult Hansknecht, scribendum est.

lanceolatis; filamentis lutescenti-barbatis; capsula globosa, calyce aequilonga. — Caulis 70—80 cm. altus, fuscus, subnitens, parce floccosus; folia radicalia 10—13 cm. longa, 2—3 cm. lata; corolla lutea, 20—25 mm. diametro; capsula minuta. — A praecedente omnino diversum, characteribus saltem quoad indumentum, racemum laxum, florum magnitudinem magis ad *V. pulverulentum* accedens, a quo autem racemo subsimplici et praesentim forma foliorum radicalium egregie discedit.

In herbidis lapidosis regionis montanae pr. Kalentini ad septentrionem urbis Arta in Epiro (Hal.). — Jun. Jul. ⊙

.. Flores in paniculam amplam dispositi.

; Folia (saltem juniora) utrinque vel saltem subtus dense tomentosa.

, Folia utrinque tomento denso persistente vestita.

24. **V. mucronatum** Lam. dict. IV. p. 218; Raul. cret. p. 819; Bois. fl. or. IV. p. 315; Hal. in z. b. G. 1898 p. 137. — *V. auriculatum* S. et S. pr. I. p. 149, Fl. gr. III. p. 20 t. 225. — *V. candidissimum* DC. fl. fr. suppl. p. 413. — Exsicc.: Heldr. herb. a. 1845 (Pisidia).

Tomento denso niveo-pannosum; caule elato, ample paniculato; foliis crassis, radicalibus crenatis, interdum sublobatis, abrupte mucronatis; florum glomerulis globosis, valde dissitis, racemos virgatos formantibus; floribus subsessilibus; calyce lana bombycina occultato, laciniis late lanceolatis; capsula globosa, calyce non longiore. — Caulis 1—1½ altus, folia radicalia ampla, ad 25 cm. longa; calyx 4—5 mm. longus; capsula semine *Cannabeos* non major. — *V. pulverulento* affine, sed robustior caule crasso, foliis radicalibus amplis crenatis, caulinis decurrentibus.

Creta (Labill. ex Bois. l. c.), sine loci specialis indicatione; recenter non lectum. — Jul. Aug. ⊙

Obs. *V. auriculatum* Mazz. in ant. ion. V. p. 210 e Corcyra: pr. Granades, certe ad aliam speciem spectat.

25. **V. megaphlomos** Bois. et Heldr. diagn. ser. 2 III. p. 144; Hal. in z. b. G. 1898 p. 137. — *V. speciosum v. megaphlomos* Bois. et Heldr. l. c., Fl. or. IV. p. 325. — Exsicc.: Orph. fl. gr. n. 370; Dörfl. fl. gr. n. 222.

Tomento denso flavescenti-pannosum; caule elato, in paniculam pyramidatam amplam abeunti; foliis crassis, integris, radicalibus oblongo-lanceolatis, basi attenuatis, caulinis brevibus, cordato-amplexicaulibus; florum fasciculis laxiusculis, racemos floribundos formantibus; pedicellis calyce aequilongis vel longioribus; calyce tomentoso, laciniis anguste lanceolatis; capsula ovato-oblonga, calyce subduplo longiore. — Caulis orgyalis; folia radicalia 30—35 cm. longa, 5—7 cm. lata; calyx 3 mm. longus; corolla flava, 20—25 mm. diametro; capsula 5 mm. longa. Subspecies *V. speciosi* Schrad.. quod tomento tenuiori et pedicellis longioribus gaudet.

β. **macrophyllum** Hal. in z. b. G. 1898 p. 137. — Folia radicalia ovalia vel elliptica, amplissima, circa 30 cm. longa, 15 cm. lata. — Exsicc.: Heldr. pl. fl. hellen. a. 1896.

γ. **microcalyx** Hal. l. c. . — Calyx fere duplo minor, longiuscule pedicellatus; capsula minor. — Exsicc.: Heldr. it. thessal. n. 2.

In regione abietina mt. Parnassus et Kyllene (Heldr.); — β. mt. Korax (Tunt.), mt. Kiona pr. Segditza (Hal.); — γ. mt. Pelion Thessaliae (Heldr.). — Jul. Aug. ⊙

,, Folia utrinque tomento floccoso tandem detersili obsita.

26. **V. pulverulentum** Vill. hist. pl. dauph. II. p. 490; Pieri corc. fl. p. 32; Mazz. in ant. ion. V. p. 210; Bois. fl. or. IV. p. 322; Hal. Beitr. fl. Epir. p. 33 p. p., Beitr. fl. Aetol. p. 9; Haussk. symb. p. 161; Bald. riv. coll. bot. alb. 1895 p. 38. — *V. floccosum* W. et K. pl. rar. hung. I. p. 81 t. 79; Mazz. l. c. p. 212; Form. in Ver. Brünn 1895 p. 34, 1896 p. 62. — Exsicc.: Sint. it. thessal. n. 535.

Tomento denso, floccoso, niveo, tandem deciduo obsitum; caule erecto, in paniculam amplam abeunte; foliis integris vel obsolete crenulatis, inferioribus ellipticis vel oblongis, obtusis, in petiolum brevem attenuatis, superioribus ovatis vel rotundatis, amplexicaulibus, plus minus abruptè acuminatis; floribus fasciculatis, in racemos laxiusculos dispositis; pedicellis calyce sublongioribus; calycis laciniis lanceolatis; capsula ovato-globosa, obtusa, calyce longiore. — Caulis 40—100 cm. altus; corolla flava, 15—20 mm. diametro; capsula parva, 2—3 mm. longa.

Ad sepes, vias, in herbidis regionis inferioris in subalpinam adscendens. Epirus: pr. Muzakia (Bald.), Preveza, Han Mescho, Janina, Sadovica, Han Levka, Govrica (Form.), Vulgarelion, Arta (Hal.); Acarnania: pr. Arapis, Anina (Hal.); Thessalia: pr. Korona, Pezula (Haussk.), Kalabaka (Sint.), pr. Miluna in Olympo (Form.); Aetolia: pr. Sikya, Karpenisi (Heldr.); Corcyra: pr. Signes, Manducchio (Mazz.). Jun. Aug. ⊙

26 × 28. V. pulverulentum × Heldreichii Hal. in z. b. G. 1898 p. 138. — *V. Heldreichii* × *pulverulentum* (*V. sterile*) Haussk. symb. p. 170. — A *V. pulverulento* indumento tenuiore et foliis inaequaliter lobatis; a *V. Heldreichii* indumento foliorum tomentoso, inflorescentiae parce floccoso discedens; capsulae steriles. — In ruderatis ad monasterium Korona in Pindo dolopico (Haussk.). — N. v.

26 × 35. V. pulverulentum × thyrsoideum (*V. permixtum*) Hal. in z. b. G. 1898 p. 139. — *V. nigrum* × *pulverulentum* Haussk. symb. p. 164. — A *V. pulverulento* foliis inferioribus longiuscule petiolatis, grosse crenatis, floribus longius petiolatis et filamentis purpureo-lanatis; a *V. thyrsoideo* indumento denso, foliis inferioribus in petiolum attenuatis diversum. — In valle Aspropotami pr. Kodri in Pindo (Haussk.). — N. v.

,,, Folia viridia, supra fere glabra, subtus tomento tenuissimo obsita.

27. **V. banaticum** Roch. ap. Schrad. monogr. verb. II. p. 28, pl. Ban. rar. p. 55 t. 18; Form. in D. bot. Monat. 1891 p. 25, in Ver. Brünn 1895 p. 34; Bald. riv. coll. bot. alb. 1896 p. 82. — Exsicc.: Bald. it. alb. epir. IV. n. 191.

Indumento tenui, tandem evanescenti obsitum, viride; caule erecto, a medio paniculato, ramis tenuiter virgatis, in racemos elongatos, laxos abeuntibus; foliis inferioribus oblongis, sinuato-crenatis, in petiolum attenuatis, versus basin auriculatis vel pinnati-lobatis, superioribus ovatis vel ovato-oblongis, crenato-dentatis, basi rotundata vel subcordata sessilibus; florum glomerulis subdissitis; pedicellis calyce aequilongis; calycis tomentosi laciniis lanceolatis, apice glabrescentibus; corolla lutea, tomentella; capsula ovata, calyce subduplo longiore. — Caulis 50—100 cm. altus; corolla parva; capsula 3—4 mm. longa. — Habitu *V. thyrsoidei*, sed filamentorum lana non purpurea.

In arvis et in pratis in valle Dipotami (Form.), infra Vromonero et Liaskovik in Epiro (Bald.); Thessalia: pr. Malakasi, Kastania (Form.). — Jun. Jul. ⊙

28. **V. Heldreichii** Bois. diagn. ser. 2 III. p. 147, Fl. or. IV. p. 326; Hal. Beitr. fl. Thessal. p. 16, in z. b. G. 1898 p. 139; Haussk. symb. p. 162. — Exsicc.: Orph. fl. gr. n. 733 (Thessalonica); Sint. et Bornm. it. turc. n. 807 et 1852; Sint. it. thessal. n. 1504.

Indumento tenui, tandem evanescenti obsitum, viride; caule erecto, a medio paniculato, ramis tenuiter virgatis, in racemos elongatos, laxos abeuntibus; foliis inferioribus lyratis, in petiolum attenuatis, segmentis lateralibus utrinque 1—3, parvis, oblongis, irregularibus, terminali magno, oblongo, foliis superioribus ovatis, crenato-dentatis, basi subcordata sessilibus; florum glomerulis subdissitis; pedicellis inaequalibus, calyce aequilongis vel duplo longioribus; calycis tomentosi laciniis lanceolatis, apice glabrescentibus; corolla lutea, tomentella; capsula ovata calyce subduplo longiore. — *V. banatico* nimis affine, quod tantum diagnoscitur sec. Velenovsky (fl. bulg. suppl. p. 208), qui species ambas vivas observavit, segmentis foliorum inferiorum paucis (pro more unico), infra lobum terminalem auriculatim sitis et ut lobi folii reliqui obtusis, foliis caulinis minus profunde cordatis, floribus et capsula paulo minoribus.

In collibus apricis, fruticetis, ad vias regionis inferioris et montanae. Thessalia: pr. Tsungeri, Malakasi, Korona, Karditza, Aivali (Haussk.), Palaeocastro (Leon.), Orman Magula (Heldr.), ad Hagios Dionysios in Olympo, pr. Drakia in mt. Pelion (Sint.). — Jun. Jul. ⊙

Obs. *V. lychnitis* L. sp. p. 177. — A Mazziari in ant. ion. V. p. 210 prope San Marco Corcyrae probabiliter ex confusione quadam indicatur.

;; Folia (saltem juniora) indumento adpresso, tenuissime arachnoideo cinerascentia, subnitentia.

29. **V. Haussknechtii** Heldr. ap. Haussk. in Mitth. thür. bot. Ver. V. p. 71, symb. p. 161; Hal. in z. b. G. 1898 p. 139. — Exsicc.: Heldr. it. IV. Thessal. a. 1885.

Caule erecto, glabrato, a basi vel a medio in ramos elongatos, arcuato-adscendentes diviso; foliis crenulatis, inferioribus ovato-oblongis, acutis, in petiolum attenuatis, supra glabrescentibus, subtus leviter incanis, caulinis ovatis, subintegris, breviter acuminatis, basi subcordata sessilibus; floribus 3—5 fasciculatis, in racemos laxos dispositis; pedicellis calyce aequilongis vel brevioribus; calycis floccosi laciniis lanceolatis; capsula globosa, calyce aequilonga. -- Caulis pluripedalis, teres, fuscus, nitidus; folia basilaria 25 cm. longa, 7—10 cm. lata, caulina fere omnino glabra, obscure viridia; calyx 3—4 mm. longus, laciniis mox glabratis; corolla lutea 25—30 mm. diametro. — Ex affinitate praecedentis, quod colore flavo-virenti, caule stricto, densius foliato, foliis inciso-lyratis, racemis stricte erectis, magis confertis, corolla intensius stellato-tomentella specifice differt.

In collibus herbosis pr. Aivali (Heldr.), Orman Magula, Pharsalus, Karditza (Haussk.). — Jun. Jul. ⊙

18 × 29. V. rigidum × Haussknechtii Hal. in z. b. G. 1898 p. 140. — *V. Haussknechtii* × *rigidum* (*V. dervichorum*) Haussk. et Heldr. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 169. — Habitu et colore *V. Haussknechtii,* ab eo foliis inferioribus in petiolum longe attenuatis, densius tomentosis et calyce duplo majore; a *V. rigido* caulibus fuscis, subglabris, virgato-ramosis, foliis inferioribus majoribus, subintegris et calyce minore, discedens. — Exsicc.: Heldr. it. IV. Thessal. a. 1885.

30. **V. leucophyllum** Griseb. spic. II. p. 46; Bois. fl. or. IV. p. 320; Haussk. symb. p. 161; Hal. in z. b. G. 1898 p. 141. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Sint. it. thessal. n. 217.

Caule erecto, parce arachnoideo a basi vel a medio in ramos strictos, rigidulos diviso; foliis utrinque adpressissime canis, radicalibus confertis, oblongis, in petiolum attenuatis, profunde sinuato-lobatis, lobis rotundatis, foliis caulinis diminutis, sessilibus, obtuse-dentatis; floribus 1—5 fasciculatis, in racemos laxos dispositis; pedicellis calyce vix brevioribus; calycis tomentelli, tandem glabrati laciniis lanceolatis; capsula ellipsoidea, calyce duplo longiore. — Stirps gracilis, pulchra, caule 20—50 cm. alto, foliis ad 13 cm. longis, 3 cm. latis, interdum subundulatis, calyce 3 mm. longo, corolla flava, 10 mm. diametro.

γ. **integrifolium** Haussk. symb. p. 161; Hal. in z. b. G. 1898 p. 141. — Foliis inferioribus integris, planis. — Exsicc.: Sint. it. thessal. n. 217 cum typo.

In collibus herbosis Thessaliae: pr. Kalabaka (Sint.), ad Acropolin pr. Pharsalum (Heldr.), pr. Aivali (Haussk.). — Maio, Jun. ⊙

×× Filamentorum lana purpurea vel violacea.

○ Folia inferiora in petiolum attenuata.

. Planta eglandulosa.

; Folia crenulata vel integra.

31. **V. Adeliae** Heldr. in Bois. diagn. ser. 2 III. p. 145, in ö. b. Z. 1898 p. 184, chlor. Mykon. p. 248; Bois. fl. or. IV. p. 321; Hal. in z. b. G. 1898 p. 141. — Exsicc.: Heldr. herb. norm. n. 1063; Dörfl. fl. aeg. n. 121 et 255.

Totum tomento flavido, denso, persistente obsitum; caule erecto, crebre foliato, in paniculam confertam, densifloram abeunti; foliis minute crenulatis, inferioribus ovato-oblongis, obtusis, in petiolum attenuatis, superioribus ovatis, breviter acuminatis, cordato-auriculatis; florum fasciculis approximatis; pedicellis calyce aequilongis vel brevioribus; calycis tomentosi laciniis triangulari-lanceolatis; filamentis violaceo-barbatis; capsula ovata, obtusa, calyce duplo longiore. — Caulis 30—60 cm. altus, folia basilaria ad 20 cm. longa, 9 cm. lata, sed saepius minora; panicula nunc pauciramosa, nunc plus minus ampla, illae *V. megaphlomos* similis, calyx 2 mm. longus, corolla flava 15 mm. diametro, capsula 4 mm. longa.

In subhumidis regionis litoralis, nec non in montosis Cycladum: insula Naxos (Heldr.), Mykonos, Rhenea (Tunt.). — Jun. Jul. ⊙

32. **V. mallophorum** Bois. et Heldr. diagn. VII. p. 39, Fl. or. IV. p. 323; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 156, chlor. Parn. p. 24; Haussk. symb. p. 161; Hal. in z. b. G. 1898 p. 142. — Huc forsan e descriptione: *V. limnense* Fraas fl. class. p. 191 ex Euboea boreali pr. Limni. — Exsicc.: Orph. fl. gr. n. 1161; Sint. it. thessal. n. 921 b.

Totum tomento niveo, denso, floccoso, plus minus detersili obsitum; caule erecto, in paniculam amplam abeunti; foliis minute crenulatis vel integris, inferioribus ovato-oblongis oblongisve, acutiusculis, in petiolum brevem attenuatis, superioribus ovatis, acutis, basi rotundatis; florum fasciculis approximatis vel dissitis; pedicellis calyce aequilongis vel longioribus; calycis dense floccosi laciniis anguste linearibus; filamentis rufescenti-lanatis; capsula oblonga, mucronata, calyce triplo longiore. — Caulis 40—80 cm. altus, detersili-floccosus, saepissime ample paniculatus, rarius subsimplex, folia basilaria ad 35 cm. longa, 10 cm. lata, praesertim subtus dense niveo-tomentosa, pedicelli ad 5 mm. longi, calyx 2—3 mm. longus, corolla flava, 25 mm. diametro, capsula 5—6 mm. longa. — Indumento niveo-floccoso, foliis radicalibus brevissime petiolatis, pedicellis longioribus, filamentorum lana dilute purpurascenti et capsula ellipsoidea a praecedente egregie distinctum et magis *V. pulverulento* affine, quod autem filamentorum lana albida et capsula ovato-globosa differt.

In regione montana et subalpina. Thessalia: mt. Kokkino Lithari pr. Sermeniko (Sint.), mt. Ghavellu, Karava, Baba, Zygos (Haussk.); mt. Pelion (Heldr.); Euboea: mt. Dirphys (Heldr.); mt. Parnassus (Orph.); Peloponnesus: mt. Olenos, Taygetos (Heldr.), Malevo, Kyllene (Orph.). — Jun.

pinnatifida.

33. **V. sinuatum** L. sp. p. 178; S. et S. pr. I. p. 150, Fl. gr. III. p. 22 t. 227; Mazz. in ant. ion. V. p. 212; Ch. et B. exp. p. 73, Fl. pelop. p. 15; Marg. et R. fl. Zante p. 70; Friedr. Reise p. 271; Fraas fl. class. p. 191; Raul. cret. p. 819; Weiss in z. b. G. 1869 p. 745; Bois. fl. or. IV. p. 322; Heldr. fl. cephal. p. 55, chlor. Mykon. p. 248, chlor. Thera p. 18; Hal. in z. b. G. 1888 p. 760; Haussk. symb. p. 161; Bald. viagg. Creta p. 80; Form. in D. bot. Mon, 1891 p. 26, in Ver. Brünn 1895 p. 33, 1896 p. 62, 1897 p. 46. — Huc probabiliter: *V. undulatum* Sieb. avis p. 3, rem. p. 3; Pieri fl. corc. p. 31; Mazz. l. c. — Exsicc.: Orph. fl. gr. n. 731; Sint. et Bornm. it. turc. n. 1395.

Plus minus dense tomentosum, lutescens vel griseum; caule erecto, saepe jam a basi subdivaricatim ramoso; foliis radicalibus oblongo-spathulatis, obtusis, subsessilibus, sinuato-pinnatifidis, saepe undulatis, caulinis superioribus oblongis, acutis, integris, breviter decurrentibus; florum glomerulis dissitis, in racemos virgatos dispositis; pedicellis calyce aequilongis; calycis tomentelli laciniis breviter lanceolatis; filamentis purpureo-lanatis; capsula globosa, calycem subaequante. — Caulis 50—80 cm. altus, folia basilaria circa 10—20 cm. longa, caulina valde diminuta, calyx 2—3 mm. longus, corolla flava, 20 mm. diametro.

In sterilibus, ad vias regionis inferioris per totam Graeciam frequens. — Jun. Aug. ⊙

26 × 43. **V. pulverulentum × sinuatum** Hal. in z. b. G. 1898 p. 143. — *V. hybridum* Brot. fl. lusit. p. 210; Form. in Ver. Brünn 1895 p. 33. — A *V. pulverulento* indumento parum floccoso, foliis caulinis subdecurrentibus et filamenits purpureo-lanatis; a *V. sinuato* caule superne ramoso, foliis crenatis, nec sinuatis discedens. — Corcyra: pr. Kanali et Hagios Joannes (Form.). — N. v.

28 × 33. **V. Heldreichii × sinuatum** (*V. ambigens*) Haussk. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 169; Hal. in z. b. G. 1898 p. 143. — A *V. Heldreichii* indumento densiori, foliis inferioribus brevissime petiolatis, profunde dentatis, subundulatis et filamentis purpurascenti-lanatis; a *V. sinuato* indumento tenuiori, filamentorum lana pallide purpurascenti et habitu discedens; capsulae steriles. — Ad monasterium Korona in Pindo (Haussk.). — N. v.

29 × 33. **V. Haussknechtii × sinuatum** (*V. thessalum*) Haussk. in Mitth. thür. bot. Ver. V. p. 72, symb. p. 169; Hal. in z. b. G. 1898 p. 143. — A *V. Haussknechtii* caule tomentello, foliis basilaribus brevissime petiolatis, inaequaliter undulato-subsinuatis, superioribus crenatis et filamentis rufescenti-lanatis; a *V. sinuato* indumento tenuiori, foliis subsinuatis discedens; capsulae steriles. — In ruderatis pr. Karditza Thessaliae (Haussk.). — N. v.

.. Planta glandulosa.

34. **V. meteoricum** Haussk. symb. p. 163; Hal. in z. b. G. 1898 p. 143.

Caule tomento tenui detersili obsito, fere a basi paniculato-ramoso; foliis utrinque tomento brevi denso cinereis, radicalibus anguste oblongo-

lanceolatis, in petiolum longe decurrentibus, irregulariter lobulato-crenatis, caulinis elliptico-lanceolatis, levissime crenulatis, superioribus diminutis subsessilibus, integris, in apicem valde elongatum, uncinato-curvatum angustatis; florum glomerulis 2—5 floris, bractea lanceolata, falcata, dense glandulifera suffultis; calyce subsessili, glanduloso, laciniis ovato-lanceolatis, acuminatis; filamentis binis longioribus dimidia parte inferiori roseo-barbatis, reliquis glabris. — Caulis rubellus, bipedalis, folia radicalia ad 20 cm. longa, 5—10 mm. tantum lata, calyx 5 mm. longus, corolla lutea, pellucido-punctata, 2 cm. diametro.

In collibus herbosis supra Kalabaka pr. monasterium Hagios Stephanos in societate *V. gloeotrichi* (Haussk.). — Jul. ⊙ N. v.

○○ Folia inferiora basi cordata vel rotundata.

35. **V. thyrsoideum** Host fl. austr. I. p. 289; Hal. in z. b. G. 1898 p. 144. — *V. nigrum* Mazz. in ant. ion. V. p. 212; Bois. fl. or. IV. p. 328; Haussk. symb. p. 162; non L. . — *V. Chaixi* Hal. Beitr. fl. Epir. p. 33, Beitr. fl. Thessal. p. 16, non Vill. — Exsicc.: Orph. herb. n. 559; Sint. it. thessal. n. 925.

Caule erecto, tomentello, in paniculam strictam abeunti; foliis obtuse crenatis, supra virentibus, subtus tomentoso-canescentibus, inferioribus ovato-oblongis oblongisve, longe petiolatis, superioribus brevius petiolatis vel sessilibus; floribus fasciculatis, in racemos laxiusculos dispositis; pedicellis calyce longioribus; calycis tomentosi laciniis lanceolatis; filamentis violaceo-barbatis; capsula ellipsoidea, calyce longiore. — Caulis 60—100 cm. altus, superne angulatus, folia ad 25 cm. longa, 15 cm. lata, petiolis fere aequilongis, calyx 2 mm. longus, corolla lutea, basi saepe violacea, 10—15 mm. diametro. — *V. nigri* L. sp. p. 178 subspecies thyrsoidea, foliis subtus cano-tomentosis. Affine *V. Chaixi* Vill. hist. pl. Dauph. II. p. 491, caule tereti, foliis inferioribus lobulato-dentatis, basi inciso-lyratis et pedicellis calyce aequilongis diverso.

In silvaticis montanis. Epirus: pr. Kalarrytes, Chaliki (Hal.); Thessalia: pr. Tsungeri, Kodri, Neuropolis (Haussk.), Korona (Heldr.); Achaia: mt. Kyllene pr. Flamburitza (Orph.). — Jun. Jul. ⊙

? 32 × 35. **V. mallophorum × thyrsoideum** Hal. in z. b G. 1898 p. 145. — *V. hypoleucum* (*nigrum* × ? *mallophorum*) Bois. et Heldr. diagn. ser. 2 III. p. 147. — *V. hypoleucum* (*nigrum* × ? *sinuatum*) Bois. fl. or. IV. p. 329.

Caule erecto, crasso, glabrescenti, superne in paniculam strictam abeunti; foliis inferioribus petiolatis, amplis, supra glabrescentibus, subtus canis, oblongis, grosse lobatis, integris vel repando-crenatis, basi rotundatis vel breviter cuneatis, caulinis superioribus diminutis, ovato-rotundis, attenuato-acuminatis, basi cordata sessilibus; floribus in glomerulos paucifloros dispositis, racemos longos laxiusculos formantibus; pedicellis calyce 2—3 plo longioribus; filamentis purpurascentibus. — In faucibus umbrosis regionis abietinae mt. Kyllene Achaiae (Heldr.). — N. v.

β. Capsula cylindrica, calyce 3—5 plo longior; corolla glabra.

36. **V. graecum** Heldr. et Sart. in Bois. diagn. ser. 2 III. p. 148, Fl. or. IV. p. 336; Hal. in z. b. G. 1888 p. 760, 1898 p. 145. — Exsicc.: Orph. fl. gr. n. 371; Sint. it. thessal. n. 29.

Caule erecto, praeter basin foliosam floccoso-tomentosam glabro, in paniculam amplam abeunte; foliis utrinque dense albo-tomentosis, inferioribus oblongis vel obovato-oblongis, in petiolum attenuatis, integris vel subrepandis, interdum subundulatis, caulinis oblongis vel ovatis, sessilibus; floribus 4—5 fasciculatis in racemos virgatos, laxos dispositis; pedicellis calyce aequilongis vel longioribus; calyce floccoso vel glabriusculo, laciniis linearibus; filamentis albo-barbatis; capsula demum glabra. — Caulis 50—70 cm. altus; folia radicalia ad 30 cm. longa, calyx 15—25 mm. longus, corolla lutea, 15—20 mm. diametro, capsula 6—8 mm. longa. — Species pulchra, inflorescentia fere glabra et capsulis cylindricis eximia.

β. **aetolicum** Bois. fl. or. IV. p. 336; Haussk. symb. p. 162; Hal. in z. b. G. 1898 p. 145. — Folia majora, tenuius tomentosa, subvirentia; capsulae submajores. — Exsicc.: Sint. et Bornm. it. turc. n. 1389.

γ. **Zuccarinii** Bois. l. c.; Hal. l. c. — *Celsia tomentosa* Zucc. pl. nov. II. p. 22. — Folia radicalia obtusiora, spathulato-suborbiculata; fasciculi 1—2 flori; calycis laciniae obtusiores. — Exsicc.: N. v.

In dumosis regionis submontanae in abietinam adscendens. Thessalia: mt. Olympus, mt. Pelion pr. Agria (Sint.); Attica: mt. Parnes (Heldr.); mt. Parnassus pr. Rachova, pr. Amphissa (Hal.); Aetolia: mt. Arapokephala (Heldr.); — γ. pr. Zeitun Thessaliae (Berger). — Apr. Jun. ⊙

3. Sectio. *Blattaria* Rchb. fl. germ. p. 379. — Flores solitarii, racemosi vel spicati, antherae adnato-decurrentes vel reniformes.

a. Corolla lutea; antherae staminum longiorum adnato-decurrentes, breviorum reniformes.

37. **V. blattaria** L. sp. p. 178; S. et S. pr. I. p. 151; Pieri fl. corc. p. 31; Mazz. in ant. ion. V. p. 212; Ch. et B. exp. p. 73, Fl. pelop. p. 15; Marg. et R. fl. Zante p. 70; Bois. fl. or. IV. p. 308; Heldr. fl. cephal. p. 55; Gelmi in bull. soc. bot. ital. XXI. p. 450; Form. in D. bot. Mon. 1891 p. 26, in Ver. Brünn 1895 p. 33, 1896 p. 62, 1897 p. 46; Hal. Beitr. fl. Epir. p. 32, in z. b. G. 1898 p. 146; Haussk. symb. p. 160; Bald. riv. coll. bot. Alb. 1896 p. 82. — Icon: Rchb. t. 32. — Exsicc.: Heldr. pl. hellen. a. 1890 et 1899.

Caule erecto, simplici vel parce ramoso, glabro vel superne glanduloso-puberulo; foliis glabris, inferioribus oblongis vel obovato-oblongis, obtusis, in petiolum attenuatis, sinuato-pinnatifidis vel dentatis, superioribus oblongis, acutis, dentatis, cordato-amplexicaulibus; floribus in racemum laxum dispositis, pedicellis eis 2—3 plo longioribus suffultis; calyce glanduloso, laciniis lanceolatis; corolla flava, glabra; filamentis

violaceo-barbatis; capsula globosa, calyce longiore. — Caulis 50—150 cm. altus; corolla 25—30 mm. diametro.

β. **blattariforme** Griseb. in Wiegm. arch. XVIII. p. 321; Haussk. symb. p. 160, pro sp.; Hal. in z. b. G. 1898 p. 146. — *V. repandum* Griseb. spic. II. p. 41; Spreitz. in z. b. G. 1887 p. 665, non Willd. en. p. 226, quod pedicellis brevissimis discedit. — ? *V. viscidulum* Mazz. in ant. ion. V. p. 212, non Pers. syn. I. p. 215. — Pedicelli calyce 4—5 plo longiores; bracteae saepius minores, angustiores. — Intermediis ad typum transit. — Exsicc.: Sint. it. thessal. n. 1301.

In herbidis, dumosis regionis inferioris et montanae. Frequens in Epiro et Thessalia; Euboea: pr. Kastaniotissa (Heldr.), Steni (Pichl.); mt. Parnassus (Heldr.); Peloponnesus: pr. Lechaena Elidis (Heldr.), pr. Methone, Messene, mt. Kupe, ad fl. Pamisum, in Zarnate, pr. Sparta (Chaub.); Zante, Cephalonia, Corcyra. — Maio, Jun. ⊙

5 × 37. **V. Sartorii × blattaria** Hal. in z. b. G. 1898 p. 147. — *V. blattaria × phlomoides* (*V. flagriforme*) Haussk. in Mitth. thür. bot. Ver. V. p. 71, non Pfund. — *V. blattaria × Sartorii* (*V. pseudoflagriforme*) Haussk. symb. p. 164. — *V. blattariforme × Sartorii* (*V. parallelum*) Haussk. symb. p. 165. — A *V. Sartorii* caule graciliore, indumento tenuiore, glanduloso, floribus minoribus, longe pedicellatis; a *V. blattaria* pubescentia, foliis caulinis multo majoribus, subdecurrentibus, floribus approximatis, breviter pedicellatis; ab ambobus foliis inaequaliter sinuato-crenatis, floribus 1—3 fasciculatis et filamentorum lana pallide rubescenti discedens. — Prope Karditza Thessaliae (Haussk.). — Exsicc.: Haussk. it. thessal. a. 1885.

23 × 37. **V. acutifolium × blattaria** (*V. petrophilum*) Hal. in z. b. G. 1898 p. 147. — *V. macilentum* Hal. Beitr. fl. Epir. p. 32, non Franchet. — Habitu nunc ad unum nunc ad alterum parentium accedens: a *V. acutifolio* foliis viridibus, parce pubescentibus, foliis crenatis, fasciculis paucifloris, pedicellis longioribus, inflorescentia glandulosa et filamentis violaceo-barbatis; a *V. blattaria* foliis non sinuato-pinnatifidis, utrinque pubescentibus, floribus fasciculatis, brevius pedicellatis et antheris omnibus reniformibus discedens.

In montosis petrosis pr. Kalentini ad septentrionem urbis Arta in Epiro (Hal.).

b. Corolla violacea; antherae omnes reniformes.

38. **V. phoeniceum** L. sp. p. 178; Mazz. in ant. ion. V. p. 212; Form. in D. bot. Monat. 1891 p. 25. — *V. ferrugineum* Andr. bot. rep. t. 162; Pieri corc. fl. p. 32. — *V. triste* S. et S. pr. I. p. 151. — Icon: Jacq. fl. austr. t. 125.

Caule erecto, puberulo, superne glanduloso; foliis crenatis vel repandis, supra glabriusculis, subtus puberulis, inferioribus ovatis vel ellipticis, obtusis, basi rotundatis vel in petiolum brevem attenuatis, superioribus oblongis vel ovato-oblongis, acutis, valde decrescentibus, sessilibus; floribus in racemum laxum dispositis; pedicellis calyce 2—

3plo longioribus; calyce glanduloso, laciniis ellipticis, obtusiusculis; filamentis purpureo-lanatis; capsula ovata, calyce longiore. — Caulis 30—80 cm. altus, superne subnudus; corolla glabra, 25 mm. diametro.

In dumosis, rarissime. Corcyra (Pieri): pr. Kasopo (Mazz.); Thessalia: in valle Penei pr. Han Kuraneos (Form.). — Maio, Jun. ⊙

B. Suffrutex erinaceus, ramis spinescentibus.

4. Sectio. *Spinosa* Bois. fl. or. IV. p. 300. — Flores solitarii; antherae reniformes.

39. **V. spinosum** L. sp. ed. 2 p. 254; S. et S. pr. I. p. 151, Fl. gr. III. p. 23 t. 229; Sieb. Reise II. p. 318 t. 7, in Flora I. p. 275, avis p. 3; Raul. cret. p. 820; Spreitz. in z. b. G. 1890 p. 297; Bald. viagg. Creta p. 80. — Exsicc.: Rev. pl. cret. n. 118; Bald. it. cret. n. 83.

Frutex humilis, ramosissimus, ramis intricatis, spinescentibus, inferne foliosis, foliisque tenuiter tomentellis; foliis oblongis, in petiolum attenuatis, integris vel sinuato-lobatis; floribus solitariis, pedicello calyce aequilongo, demum elongato suffultis; calycis laciniis lanceolatis obtusiusculis; corolla flava, parva; filamentis sordide-lutescenti barbatis; capsula calyce duplo longiore.

In dumetis, rupestribus a mare ad alt. 1500 m. adscendens. Creta: mt. Lakkus (Rev.), Drakona, Hagios Paulos (Raul.), in oropedio Omalo, pr. Askyphos, mt. Hagios Theodoros (Spreitz.) in mt. Sphacioticis. — Jun. Jul. ♄

2. Celsia L. gen. n. 757.

1. Sectio. *Arcturus* Benth. in DC. pr. X. p. 244. — Staminorum longiorum antherae adnato-decurrentes.

a. Flores subsessiles.

1. **C. cretica** L. fil. suppl. p. 281; Sieb. avis rem. p. 5; Raul. cret. p. 820; Bois, fl. or. IV. p. 352. — Icon: Jacq. hort. Schoenbr. t. 480.

Pubescenti hirta, viridis, superne glandulosa; caule erecto, simplici, in racemum laxum abeunte; foliis inferioribus petiolatis, lyrato-pinnatifidis vel indivisis, ovatis vel oblongis, sublobatis et dentatis, superioribus sensim diminutis, cordato-auriculatis, ovatis oblongisve, dentatis; bracteis lanceolatis, serratis vel integris, calycem superantibus; calycis laciniis latis, serratis; corolla flava, purpureo-maculata; filamentis brevioribus violaceo-barbatis, longioribus glabris; capsula globosa, calyce breviore. — Caulis 30--80 cm. altus, corolla 25 mm. diametro.

Creta, sed a recentioribus non lecta. — Apr. Maio. ⊙ N. v.

b. Flores longe pedicellati.

α. Folia lanata, cana vel canescentia.

2. **C. arcturus** L. sp. p. 178 (*Verbascum*); Murr. syst. p. 469; S. et S. pr. I. p. 438; Sieb. avis. p. 4; Raul. cret. p. 821; Bois. fl.

or. IV. p. 352; Bald. viagg. Creta p. 80. — *C. sublanata* Jacq. fragm. p. 79 t. 126. — Exsicc.: Rev. pl. cret. n. 184; Bald. it. cret. n. 133, it. cret. alt. n. 264.

Caule erecto, simplici, glanduloso-pubescente, in racemum laxum abeunte; foliis infimis petiolatis, ovatis vel oblongis, indivisis vel lyratis, grosse crenatis, caulinis paucis, diminutis, semiamplexicaulibus, cito in bracteas abeuntibus; bracteis triangularibus, denticulatis, pedicello multo brevioribus; calycis glandulosi laciniis ellipticis, acutis; corolla flava; filamentis violaceo-barbatis; capsula globosa, calyce subtriplo longiore. — Valde variat quoad indumentum, magnitudinem et formam foliorum. — Caules 15—50 cm. alti, folia nunc parva indivisa, cum petiolo 4 cm. longa, nunc multo majora lyrata, interdum cum petiolo ad 25 cm. longa, lobo terminali maximo ad 10 cm. lato, corolla 25 mm. diametro.

Ad muros, rupes regionis inferioris et collinae Cretae: pr. Kissamos, Malaxa (Rev.), Palaeokastro, Epanochoria, Hagia Erini et Rumeli, Hagios Paulos, Aradhena, promontorium Meleka, ad Hagia Triada, Kritsa (Raul.), Hagios Joannes, pr. Anopolis, in fauce Nipros (Bald.). — Apr. Jun. ♃

β. Folia glabra vel glabriuscula.

× Pedicelli bractea subduplo longiores.

3. **C. speciosa** Fenzl. in Friedr. Reise p. 280; Bois. fl. or. IV. p. 353. — Exsicc.: Friedr. herb. n. 919.

Caule erecto, glanduloso-puberulo, basi parce ramoso, in racemum laxum abeunte; foliis glabriusculis, triangulari-ovatis, acutis, infimis petiolatis, inciso-serratis, reliquis subsessilibus sessilibusve, amplexicaulibus, in bracteas abeuntibus; bracteis cordato-ovatis, acuminatis, serratis; calycis glandulosi laciniis anguste lanceolatis; corolla flava; filamentis violaceo-barbatis, longioribus apice nudis; capsula ignota.

Arcadia: ad ripas fl. Alpheus pr. Carithena (Friedr.); a recentioribus non lecta. — Maio, Jun. ⊙

×× Pedicelli bractea multoties longiores.

4. **C. Daenzeri** Fauché et Chaub. in exp. mor. p. 342, Fl. pelop. p. 40; Bois. fl. or. IV. p. 353; Hal. Beitr. fl. Achaia p. 27. — Exsicc.: Orph. fl gr. n. 1114.

Caulibus solitariis vel e collo pluribus, simplicibus vel ramosis, pilosulis, superne parce glandulosis, in racemum laxum, elongatum abeuntibus; foliis pilosulis, infimis oblongo-lanceolatis, in segmenta oblongo-lanceolata, dentata pinnatisectis vel lyratis, superioribus triangulari-lanceolatis, inciso-dentatis, sessilibus, in bracteas abeuntibus; bracteis triangularibus, acuminatis, dentatis, superioribus linearibus, integris; calycis glandulosi laciniis oblongis, obtusiusculis; corolla flava; filamentis violaceo-barbatis, longioribus apice nudis; capsula globosa, obtusa, calyce duplo longiore. — Caules 50—150 cm. alti, folia radicalia 10—15 cm. longa, corolla 3 cm. diametro.

In silvaticis montanis. Achaia: mt. Kyllene pr. Trikala (Orph.), pr. Megaspilaeon (Hal.); Messenia: ad templum Apollinis pr. Phigalea

(Daenzer); Laconia: mt. Malevo (Orph.), mt. Taygetos supra Mistra (Bois.) et Anavryti (Heldr.). — Maio, Jun. ⊙

5. **C. Boissieri** Heldr. et Sart. in Bois. diagn. ser. 2 III. p. 151, Fl. or. IV. p. 353; Haussk. symb. p. 170. — *C. Daenzeri v. lyrata* Bois. et Heldr. diagn. l. c.. — Exsicc.: Heldr. herb. norm. n. 317 et 1267; Orph. fl. gr. n. 221; Sint. it. thessal. n. 632.

Caulibus solitariis vel e collo pluribus, simplicibus, glabris, superne parce glandulosis, in racemum laxum, elongatum abeuntibus; foliis glabris, oblongo-lanceolatis, lobato-dentatis vel pinnatifidis, lobis inciso-dentatis, superioribus diminutis, sessilibus, dentatis, in bracteas abeuntibus; bracteis minimis, lineari-lanceolatis, integris; calycis glandulosi laciniis lanceolatis, acuminatis; corolla flava; filamentis violaceo-barbatis, longioribus apice nudis; capsula ovata, acutiuscula, mucronata, calyce duplo longiore. — Differt ab affini praecedente caulibus simplicibus, foliis brevioribus, minus divisis, bracteis angustis integris et capsula acutiuscula.

In dumosis regionis inferioris et montanae. Thessalia: pr. Malakasi in Pindo (Sint.); Attica: mt. Parnes (Heldr.), Kerata pr. Eleusis (Haussk.). -- Maio, Jun. ♃

2. Sectio. *Nefflea* Benth. in DC. pr. X. p. 246. — Antherae omnes reniformes.

a. Folia crenato-dentata.

6. **C. cyllenea** Bois. et Heldr. fl. or. IV. p. 355. — Icon: Rouy ill. t. 93. — Exsicc.: Heldr. pl. fl. hellen. a. 1887.

Cauligera, glanduloso-puberula, viridis; caulibus erectis, humilibus, simplicibus, aphyllis, in racemum laxum abeuntibus; foliis radicalibus, oblongis, crenato-dentatis, in petiolum aequilongum vel longiorem attenuatis; bracteis minutis, linearibus, pedicello multo brevioribus; pedicellis calyce 3—4 plo longioribus; calycis laciniis linearibus, obtusiusculis; corolla lutea; filamentis inferne flavido, superne violaceo-barbatis; capsula ignota. — Caulis 10—20 cm. altus, foliorum lamina 20—35 mm. longa, corolla 20—25 mm. diametro.

In regione alpina mt. Kyllene Achaiae rarissime (Heldr.). — Jun. Jul. ♃

7. **C. acaulis** Bory et Chaub. exp. p. 177 t. 18, Fl. pelop. p. 40 t. 19; Bois. fl. or. IV. p. 354; Hal. Beitr. fl. Achaia p. 27. — Exsicc.: Heldr. herb. norm. n. 963 et 1460; Orph. fl. gr. 1115; Dörfl. fl. gr. n. 399.

Subacaulis, glanduloso-puberula, viridis; caulibus subnullis, ad racemum brevissimum radicalem reductis; foliis radicalibus, oblongis, crenato-dentatis, in petiolum aequilongum vel longiorem attenuatis; bracteis minutis, linearibus, pedicello multo brevioribus; pedicellis calyce multoties longioribus; calycis laciniis linearibus, obtusiusculis; corolla lutea; filamentis aurantiaco-barbatis; capsula ovato-globosa, acuta, mucronata. — Species pulchella, egregia, a praecedente caulibus subnullis,

caespitosis, pedicellis longioribus, interdum valde elongatis, ad 8 cm. longis et filamentis aurantiaco-lanatis specifice diversa.

In regione alpina Peloponnesi: mt. Chelmos supra fontem Stygis (Heldr.) et supra Sudena (Hal.), mt. Taygetos (Chaub). — Jun. Jul. ♃

b. Folia bipinnatisecta.

8. **C. orientalis** L. sp. p. 621; S. et S. pr. I. p. 438, Fl. gr. t. 605; Ch. et B. exp. p. 177, Fl. pelop. p. 40; Bois. fl. or. IV. p. 360.

Minute tomentella, viridis; caule erecto, simplici vel parce ramoso, folioso, in racemum laxum abeunti; foliis ambitu ovato-oblongis, in lacinias divergentes bipinnatisectis, inferioribus petiolatis, segmentis oblongis, superioribus sessilibus, laciniis anguste linearibus; floribus pedicello calyce breviori suffultis; bracteis inferioribus pinnatisectis, ceteris oblongo-lanceolatis, calyce longioribus; calyce glanduloso-puberulo, laciniis lanceolatis; corolla parva, lutea; filamentis flavido-barbatis; capsula ovata, calyce breviore.

In collibus saxosis regionis inferioris. Argolis (Sibth.); Cyclades (Chaub.). — Apr. Maio. ⊙ N. v.

2. Tribus. CHELONEAE Benth. in DC. pr. X. 298.

3. Scrofularia L. gen. n. 312.

Dispositio specierum:

1. Sectio. *Scorodonia* Don gard. dict. IV. p. 508. — Staminodium squamaeforme; flores pedicellis saepissime calyce longioribus suffulti; folia indivisa.

a. Cymae axillares, 2—9 flori; calycis laciniae non marginatae.

1. **S. peregrina** L. 2. **S. aestivalis** Griseb.

b. Cymae in paniculam terminalem aphyllam vel basi subfoliatam dispositi; calycis laciniae membranaceo-marginatae.

α. Folia plus minus pubescentia.

3. **S. Scopolii** Hoppe. 4. **S. cretica** Bois. et Heldr.

β. Folia glabra.

× Rhizoma tuberculoso-nodosum; calycis laciniae obsolete marginatae.

5. **S. nodosa** L.

×× Rhizoma non tuberculoso-nodosum; calycis laciniae late membranaceo-marginatae.

6. **S. Nesii** Wirtg. 7. **S. Samaritanii** Bois. et Heldr.

2. Sectio. *Tomoiophyllum* Benth. in DC. pr. X. p. 310. — Staminodium squamaeforme vel lineare; flores pedicellis saepissime calyce aequilongis vel brevioribus suffulti; folia varie partita, rarius indivisa et tunc pedicelli calyce breviores.

a. Staminodium squamaeforme.

α. Biennes.

8. **S. laxa** Bois. et Heldr. 9. **S. lucida** L.

β. Perennes, rhizomate lignescente.

× Plantae glabrae, inflorescentia tantum glanduloso-puberulae.

10. **S. heterophylla** Willd. 11. **S. laciniata** W. et K.

XX Planta tota pube patula crispa glandulosa obsita.

12. S. taygetea Bois.

b. Staminodium anguste lineare.

13. S. canina L.

1. Sectio. *Scorodonia* Don gard. dict. IV. p. 508.

a. Cymae axillares, 2—9 flori; calycis laciniae non marginatae.

1. **S. peregrina** L. sp. p. 621; S. et S. pr. I. p. 435, Fl. gr. t. 597; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 176, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 70; Friedr. Reise p. 269; Ung. Reise p. 129; Raul. cret. p. 822; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 395; Heldr. fl. cephal. p. 55, Fl. Aegina p. 382, in ö. b. Z. 1898 p. 184; Haussk. symb. p. 170, cum *v. trachytica* (f. a basi ramosa, foliis minoribus, cymis uni-paucifloris, floribus et capsulis minoribus). — Huc forsan: *S. scorodonia* Pieri corc. fl. p. 84, non L. sp. p. 620, quae species occidentalis et discedit: pubescentia, caule non fistuloso, cymis plurifloris, calycis laciniis obtusis, late membranaceo-marginatis. — Exsicc.: Baen. herb. europ. n. 9339.

Caule erecto, fistuloso, quadrangulo, simplici vel ramoso, brevissime puberulo, superne pedicellisque glanduloso; foliis petiolatis, ovato-cordatis, grosse dentatis, glabris, inferioribus oppositis, superioribus sparsis; cymis axillaribus, laxe 2—5 floris; pedicellis calyce 2—3 plo longioribus; calycis glabri laciniis ovato-lanceolatis, acutis, non marginatis; corolla fusco-purpurea; staminodio obovato-orbiculari, obtuso; capsula subglobosa, obtusa, mucronata.

In ruderatis, ad vias, sepes regionis inferioris. Aetolia: pr. Krioneri (Reis.); Attica: pr. Athenas (Heldr.), insula Aegina (Friedr.); Peloponnesus (Chaub.): pr. Nauplia, in peninsula Methana (Haussk.); Cycladum insula: Cythnos (Tunt.), Amorgos (Heldr.); Creta (Sieb.): pr. Canea, Malaxa (Raul.); Zante (Marg.); Cephalonia: pr. Pessada (Heldr.); Ithaca: pr. Vathy (Spreitz.); Corcyra (Ung.): pr. Castrades (Spreitz.). — Apr. Jun. ⊙

2. **S. aestivalis** Griseb. spic. II. p. 36; Hal. in ö. b. Z. 1892 p. 375. — *S. silvatica* Bois. et Heldr. diagn. ser. 2 III. p. 153, Fl. or. IV. p. 398; quae meo sensu a specie Griseb. nullo modo discedit. — Exsicc.: Orph. fl. gr. n. 730; Sint. et Bornm. it. turc. n. 1387.

Caule erecto, farcto, quadrangulo, simplici vel ramoso, glanduloso-pubescente; foliis oppositis, petiolatis, ovatis, grosse inaequaliter duplicato-serratis, basi cordatis vel subtruncatis, utrinque villosiusculis, summis sparsis; cymis axillaribus, laxe 3—9 floris; pedicellis calyce aequilongis vel longioribus; calycis glanduloso-puberuli laciniis ovatis, obtusis, non marginatis; corolla flavo-virenti; staminodio subrotundo; capsula ovata, acutissima, calyce subduplo longiore. — A praecedente pubescentia viscosa, radice perenni, foliis majoribus, calycis glandulosi laciniis obtusis, corolla virenti, capsula acutissima et toto habitu diversissima.

In silvaticis regionis montanae, rarissime. Thessalia: mt. Olympus pr. Hagios Dionysios (Heldr.) et Malathria (Sint.). — Jul. Aug. ♃

b. Cymae in paniculam terminalem aphyllam vel basi subfoliatam dispositi; calycis laciniae membranaceo-marginatae.

α. Folia plus minus pubescentia.

3. **S. Scopolii** Hoppe in Pers. syn. II. p. 160; Bois. fl. or. IV. p. 395; Form. in D. bot Monat. 1891 p. 26, in Ver. Brünn 1896 p. 63; Hal. Beitr. fl. Thessal. p. 16; Haussk. symb. p. 170; Boissieu in bull. soc. bot. Fr. 1896 p. 287. — *S. auriculata* Scop. fl. carn. ed. 2 I. p. 446 t. 32; ? Pieri corc. fl. p. 84; non L. sp. p. 620, quae statura elatiori, foliis saepius auriculato-appendiculatis, cymis breviter pedunculatis, pedicellis brevissimis, calycis laciniis latius marginatis discedit. — *S. glandulosa* W. et K. pl. rar. III. p. 238 t. 214. — Exsicc.: Orph. herb. n. 3519.

Caule erecto, obtuse quadrangulo, simplici vel ramoso, pubescente, superne pedicellisque glanduloso; foliis oppositis, petiolatis, ovato-oblongis, cordatis, crenato-dentatis, utrinque plus minus pubescentibus; panicula oblonga, cymis longiuscule pedunculatis; pedicellis calyce subduplo longioribus; calycis glabriusculi laciniis ovato-orbiculatis, membranaceo-marginatis; corolla virenti-purpurascenti; staminodio transverse latiore; capsula ovato-globosa, mucronata, calyce subduplo longiore.

β. **grandidentata** Ten. fl. nap. I. prodr. suppl. II. p. 69, pro sp.; Bois. fl. or. IV. p. 396; Haussk. symb. p. 170. — Foliorum saepe acutiorum et magis oblongorum dentes majores, acutiores, saepe denticulis 1—2 aucti. — Specimina typica, cum planta italica (Tod. fl. sic. n. 783; Lojac. pl. sic. rar. n. 459) congruentia e Graecia nondum vidi, omnia quoad foliorum serraturam immo potius formam intermediam sistunt inter plantam hungaricam et italicam, talia sunt e. g. a Heldr. it. thessal. IV. a. 1885 et a Sint. it. thessal. n. 757 distributa.

γ. **oligantha** Bois. et Heldr. diagn. XII. p. 33 pro sp.; Bois. fl. or. IV. p. 396; Haussk. symb. p. 170. — Humilior, folia minora, panicula depauperata; sec. Boissier forma alpina. Exsicc.: N. v.

In silvaticis regionis montanae et subalpinae. Thessalia: mt. Oxya (Hal.), pr. Vitomo (Sint.), mt. Zygos, pr. Sermeniko, mt. Ghavellu, Karava (Haussk.) in Pindo, pr. Selicany et Spilia in mt. Ossa, ad Nezeros in Olympo (Form.); Euboea: mt. Dirphys (Heldr.); Aetolia: mt. Tymphrestus (Heldr.); Boeotia: pr. Larmes (Boissieu); Laconia: mt. Malevo (Orph.); Corcyra si S. auriculata Pieri huc pertineat; — γ. in summis mt. Taygetos (Heldr.) et sec. Haussknecht in Thessalia in societate typi. — Maio, Jul. ⊙

4. **S. cretica** Bois. et Heldr. in Bois. fl. or. IV. p. 400. — *S. auriculata* Raul. cret. p. 822, non L. sp. p. 620, quae foliis saepius auriculato-appendiculatis, pedicellis brevissimis et calyce glabro differt. Exsicc.: — Rev. pl. cret. n. 123.

Caule erecto, acute quadrangulo, simplici vel ramoso, glabro, superne pedicellisque sparse glanduloso; foliis oppositis, petiolatis, oblongis, basi

cordatis vel subtruncatis, obtuse dentatis, utrinque sparse pubescentibus vel supra glabriusculis; panicula elongata, valde interrupta, cymis breviter pedunculatis; pedicellis calyce aequilongis; calycis glabri vel papilloso-puberuli laciniis ovatis, membranaceo-marginatis; corolla fusco-purpurascente; staminodio ovato-orbiculari; capsula globosa, mucronata, calyce sesquilongiore. — Differt a praecedente indumento sparso, foliis oblongis, minutius dentatis, cymis breviter pedunculatis, remotis, pedicellis brevioribus; magis accedit sequentibus, a quibus praesertim foliis pubescentibus, panicula valde interrupta et pedicellis brevibus discedit. — Huc forsan spectat: *S. salicifolia* Sieb. in Flora I p. 271 e fauce Perivoglia Cretae.

In locis humidis Cretae: pr. Canea (Heldr.). — Maio, Jun. ♃

β. Folia glabra.

× Rhizoma tuberculoso-nodosum, calycis laciniae obsolete marginatae.

5. **S. nodosa** L. sp. p. 619; S. et S. pr. I. p. 435; Dallap. prosp. p. 86; Ch. et B. exp. p. 176, Fl. pelop. p. 39; Bois. fl. or. IV. p. 399; Haussk. symb. p. 170. — Icon: Fl. dan. t. 1167.

Caule erecto, argute quadrangulo, simplici vel ramoso, glabro, superne pedicellisque glanduloso; foliis oppositis, petiolatis, ovato-oblongis, basi cordatis vel truncatis, subduplicatim et argute serratis, glabris; panicula angusta, ramis erecto-patulis; pedicellis calyce 2—3plo longioribus; calycis glabri laciniis ovatis, obsolete marginatis; corolla brunneo-virenti; staminodio transverse latiore, truncato vel retuso; capsula ovata, acuta, calyce 3—4 plo longiore.

In nemorosis subhumidis. Thessalia: in oropedio Neuropolis (Haussk.); Laconia (Sibth.); Cephalonia (Dallap.). — Jun. Jul. ♃ N. v.

×× Rhizoma non tuberculoso-nodosum; calycis laciniis late membranaceo-marginatae.

6. **S. Neesii** Wirtg. in Verh. naturhist. Ver. I. p. 29. — Exsicc.: Heldr. it. thessal. n. 44.

Caule erecto, quadrangulo, anguste alato, simplici vel ramoso, glabro, superne pedicellisque glanduloso; foliis oppositis, petiolatis, oblongis, basi truncatis vel subcordatis, obtusiuscule serratis, glabris; panicula elongata, ramis patulis; pedicellis calyce 2—3 plo longioribus; calycis glabri laciniis ovatis vel subrotundis, late membranaceo-marginatis; corolla viridi-purpurea; staminodio transverse latiore, truncato vel retuso; capsula subglobosa, acuta, calyce 2—3 plo longiore. — *S. alatae* Gilib. fl. lith. II. p. 127 = *S. Ehrharti* Stev. in ann. nat. hist. V. 3 t. 1 sec. Bab. bret. bot. p. 218 = *S. aquaticae* L. sp. p. 620 pro p., valde affinis, sed omnibus partibus tenerior, rami caulis et paniculae magis, fere rectangule patentes, corolla minor, staminodium non obcordato-bilobum. — A praecedente rhizomate non nodoso, caule cum petiolis alato et calycis laciniis late membranaceo-marginatis distincta.

Ad rivulos pr. Portaria in regione media mt. Pelion Thessaliae (Heldr.). — Jul. Aug. ♃

7. **S. Samaritanii** Bois. et Heldr. diagn. ser. 2 III. p. 153. — *S. Ehrharti v. Samaritani* Bois. et Heldr. ibidem. — *S. alata* Bois. fl. or. IV. p. 399, quoad pl. parnass.; Heldr. chlor. Parn. p. 24; non Gilib. . — Exsicc.: Heldr. herb. norm. n. 83; Orph. fl. gr. n. 443.

Caule erecto, quadrangulo, anguste alato, simplici vel ramoso, glabro superne pedicellisque glanduloso; foliis oppositis, petiolatis, ovato-oblongis oblongisve, crenatis, basi truncatis, in petiolum late alatum decurrentibus; panicula elongata, ramis erecto-patulis; pedicellis calyce 2—3plo longioribus; calycis glabri laciniis ovatis, late membranaceo-marginatis; corolla livido-flavescenti; staminodio transverse latiore, obcordato-bilobo; capsula subglobosa, acuta, calyce 2—3plo longiore. — Differt a praecedente foliis obtuse crenatis, in petiolum latum decurrentibus, paniculae ramis strictiusculis, corolla majore et staminodii forma; a *S. alata* quacum a Boissier postea conjuncta fuit, caule angustius alata, foliis obtuse crenatis, in petiolum latum decurrentibus, corolla majori.

In scaturiginosis regionis abietinae mt. Parnassus l. d Gurna pr. Rachova (Heldr.); probabiliter huc pertinet quoque planta e mt. Malevo Laconiae, a Boissier prope *S. alatam* commemorata. — Jul. Aug. ♃

2. Sectio. *Tomoiophyllum* Benth. in DC. pr. X. p. 310.

a. Staminodium squamaeforme.

α. Biennes.

8. **S. laxa** Bois. et Heldr. diagn. ser. 2 III. p. 154, Fl. or. IV. p. 404. — Exsicc : Orph. fl. gr. n. 243.

A collo saepius pluricaulis; caulibus erectis, obtuse tetragonis, simplicibus vel a basi parce ramosis, inferne tantum foliatis, glabris vel superne pedicellisque glandulosis; foliis oppositis, glabris, omnibus tenuiter bipinnatisectis, segmentis minutis, oblongis, acute dentatis; panicula elongata, aphylla, cymis inferioribus 5 floris, superioribus 3—1 floris; pedicellis calyce brevioribus; calycis glabri laciniis orbiculatis, membrana scariosa lata, denticulato-lacera marginatis; corollae lividae lobis atropurpureis; staminibus apice breviter exsertis; staminodio semicirculari; capsula globosa, mucronata, calyce duplo longiore. — Sequenti maxime affinis, ab ea foliis omnibus tenuissime sectis et calycis margine denticulato-lacera diversa.

In rupestribus regionis alpinae mt. Kyllene et Taygetos Peloponnesi (Heldr.). — Jun. Jul. ⊙

9. **S. lucida** L. sp. ed. 2 p. 865; S. et S. pr. I. p. 436; Urv. enum p. 75; Ch. et B. exp. p. 177, Fl. pelop. p. 40; Weiss in z. b. G. 1869 p. 745; Bois. fl. or. IV. p. 403. — Exsicc.: Rev. pl. cret. n. 122; Bald. it. cret. alt. n. 89; Dörfl. fl. aeg. n. 130, pl. cret. n. 25, 25 a et 58.

Caule saepissime solitario, erecto, obtuse tetragono, simplici vel parce ramoso, glabro vel superne pedicellisque glanduloso; foliis oppositis, glabris, glaucescentibus, imis parvis lobatis, ceteris semel vel bis pinnatisectis, segmentis oblongis, inciso-dentatis; panicula elongata,

aphylla, cymis inferioribus plurifloris, superioribus sursum decrescentibus, summis 2—1 floris; pedicellis calyce brevioribus; calycis glabri laciniis orbiculatis, membrana scariosa lata, integra marginatis; corollae lividae lobis atropurpureis; staminibus majoribus exsertis; staminodio late orbiculato; capsula globosa, mucronata, calyce duplo longiore. — Stirps variabilis, caulis 15—50 cm. altus, paniculae rami nunc (interdum omnes) breves et pauciflori, nunc praesertim inferiores elongati et pluriflori, ideo panicula nunc angusta, nunc plus minus pyramidata, folia saepius prope basin caulis approximata, nunc tenuiter nun late dissecta, flores semper magni, eis *S. caninae* duplo majores.

α. **glauca** S. et S. fl. gr. VI. p. 78 t. 599 pr. sp. — *S. lucida v. genuina* Bois. fl. or. IV. p. 403. — Saepius humilior, foliorum laciniae interdum angustiores.

β. **filicifolia** Mill. dict. n. 10; S. et S. pr. I. p. 436, Fl. gr. IV. p. 79 t. 600; Sieb. avis p. 4, rem. p. 5; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 822; Spreitz. in z. b. G. 1877 p. 716; Haussk. symb. p. 171; Heldr. fl. Aegina p. 383; pro sp.; Benth. in DC. pr. X. p. 312; Bois. fl. or. IV. p. 403; Spreitz. in z. b. G. 1890 p. 297; Heldr. chlor. Thera p. 18, chlor. Mykon. p. 248. — Saepius elatior, foliorum laciniae latiores, saepe minus dissectae. — Formis intermediis sine limitibus ad α. transit.

In collibus saxosis, rupestribus regionis inferioris et montanae. Thessalia: mt. Pelion pr. Lechonia (Heldr.); mt. Parnassus (Guicc.); Attica: mt. Parnes, Pentelicon (Sprun.), Kerata, pr. Laurion (Haussk.), mt. Hymettus; insula Aegina, Hydra (Heldr.); Cycladum insula: Andros (Sart.), Tenos (Chaub.), Delos, Mykonos (Tunt.), Naxos (Tourn. voy. p. 85), Melos, Thera, Mikra Kameni (Urv.); Creta: pr. Askyphos (Spreitz.), ad fl. Jardanus pr. Canea (Weiss), Akroteri, Malaxa, Rhamni, Nerokuru, Prosnero, insula Dia (Raul.), ad Hagios Myron distr. Mallevisi (Bald.), pr. Kritsa distr. Mirabello, pr. Parsas distr. Viano, pr. Michti distr. Hierapetra (Leon.); Zante: mt. Scopo (Weiss); Corcyra: mt. Pantocrator (Spreitz.). — Mart. Jun. ⊙

Obs. Ad formas *S. lucidae* quoque pertinet ex descriptione *S. methanaea* Haussk. symb. p. 170, e peninsula Methana pr. Vromolimni, quae nulla nota graviori, nisi panicula multiflora, squarroso-rigidissima, ramis elongatis, horizontaliter patentibus a typo differre videtur.

β. Perennes, rhizomate lignescente.

×× Plantae glabrae, in inflorescentia tantum glanduloso-puberulae.

10. **S. heterophylla** Willd. sp. III. p. 274; S. et S. fl. gr. p. 2 t. 603; Urv. enum. p. 75; Raul. cret. p. 822; Bois. fl. or. IV. p. 408; Heldr. chlor. Parn. p. 24, in ö. b. Z. 1898 p. 184, chlor. Thera p. 18; Hal. in ö. b. Z. 1896 p. 16; Haussk. symb. p. 172. — *S. frutescens* S. et S. pr. I. p. 437, non L. — *S. caesia* S. et S. pr.

I. p. 438, Fl. gr. VII. p. 3 t. 604; Ch. et B. fl. pelop. p. 40; Ung. Reise p. 129; Raul. cret. p. 822; Weiss in z. b. G. 1869 p. 745 pro var. *S. heterophyllae*; (f. foliis glaucescentibus, lyrato-pinnatifidis, incisis, caulibus numerosis, panicula abbreviata pauciflora). — *S. micrantha* Urv. enum. p. 39 (f. foliis oblongis, basi auriculatis, grosse dentatis, caulibus pedalibus sesquipedalibusque, floribus et capsulis minoribus). — *S. Sibthorpiana* Spreng. syst. II. p. 786. — *S. Oliveriana* Wydl. monogr. scrof. p. 39 (f. foliis ovatis, inciso-dentato-crenatis, basi plus minus pinnatisectis, segmentis petiolulatis subrotundis). — *S. Urvilleana* Wydl. l. c. p. 40; Weiss in z. b. G. 1869 p. 745 pro var. *S. heterophyllae* (f. foliis subrotundo-cordatis, sublobatis, rarius crenulatis). — *S. Urvillei* Wydl. l. c. t. 2 et 3. — *S. chysanthenifolia* Ch. et B. exp. p. 176, Fl. pelop. p. 40; Fraas fl. class. p. 189; sec. Bois. fl. or. IV. p. 408; non Willd.; (f. foliis pinnatisectis). — *S. laciniata v. obtusiloba* Marg. et R. fl. Zante p. 70. — *S. heterophylla v. pinnatisecta* Bois. l. c.; Heldr. fl. cephal. p. 55; Haussk. symb. p. 172 *f. viridis et caesia*; (f. foliis profundius divisis, pinnatisectis, segmento terminali non majore, caulibus saepe procerioribus). — *S. laciniata* Spreitz. in z. b. G. 1877 p. 716; Gelmi in bull. soc. bot. ital. 1889 p. 450; Form. in Ver. Brünn 1896 p. 63; Bald. riv. coll. bot. Alb. 1895 p. 59; non W. et K. — Huc spectare videtur etiam ex descriptione: *S. tenuis* Haussk. symb. p. 173 e peninsula Methana Argolidis. — Exsicc.: Orph. fl. gr. n. 244 et 728; Heldr. herb. norm. n. 383 et 1461; Sint. et Bornm. it. turc. n. 1386; Sint. it. thessal. n. 26; Baen. herb. europ. n. 9338; Dörfl. fl. aeg. n. 11 et 66.

A collo pluricaulis; caulibus erectis vel adscendentibus, fragilibus, tetragonis, simplicibus vel parce ramosis, glabris, superne pedicellisque glanduloso-puberulis; foliis oppositis, petiolatis, glabris, carnosulis, saepius glaucescentibus, ovatis vel oblongis, obtusis, basi subcuneatis, truncatis vel subcordatis, inciso-crenatis, indivisis vel lyrato-pinnatifidis sectisve, segmentis lateralibus 1—2 multo minoribus; panicula pyramidata, aphylla, cymis laxe 5—9 floris; pedicellis calyce subaequilongis; calycis glabri laciniis ovatis, latiuscule albo-marginatis; corolla luride purpurea; antheris subexsertis; staminodio transverse latiori, reniformi, interdum subretuso; capsula globosa, apiculata, calyce subtriplo longiore. — Stirps, uti jam ex synonymia patet, quoad foliorum formam praesertim valde variabilis; cum autem folia diversissima in unoeodem specimine saepe occurrunt, varietates perfecte describere mihi impossibile videtur.

Ad muros, in rupestribus regionis inferioris et submontanae. Epirus: ad Camarina distr. Prevesa (Bald.); Thessalia: pr. Kalabaka, Pharsalus, Han Tripa, Larissa, Nezeros, Rapsani, mt. Pelion (Form.), ad Lechonia pr. Volo, pr. Litochori (Sint.); Acarnania: in pago Katochi; mt. Parnassus (Heldr.); Boeotia: pr. Livadia (Orph.), ad lacum Hylike (Ung.); Attica: in Acropoli, Lycabetto, Turcovuni, mt. Hymettus (Heldr.), mt. Kerata (Haussk.); Argolis: pr. Nauplia (Orph.); Messenia: pr. Kalamata (Zahn.), Pylos, insula Sapienza (Chaub.); Laconia (Sibth.); Cyclades: insula Keos (Heldr.), Syra (Fraas), Paros (Leon.), Evreokastron (Reis.).

Naxos (Bois.), Amorgos (Heldr.), Melos, Thera, Therasia (Urv.); Creta: ad promontorium Grabusa (Raul.); Zante: pr. Litakia (Marg.), mt. Scopo (Weiss); Cephalonia: pr. Poros, mt. Atro et Gerolaki, ad coenobium Phanentes (Heldr.); Corcyra: pr. Castrades (Baen.), mt. Deca (Spreitz.). — Febr. Maio. ♃

11. **S. laciniata** W. et K. pl. rar. hung. II. p. 185 t. 170; Bois. fl. or. IV. p. 408; Hal. Beitr. fl. Epir. p. 33; Form. in Ver. Brünn 1895 p. 34, 1896 p. 63, cum. v. *pindicola*; Haussk. symb. p. 173; Boissieu in bull. soc. bot. Fr. 1896 p. 287; Bald. riv. coll. bot. Alb. 1896 p. 83. — *S. multifida* Willd. hort. berol. t. 58; Hal. in z. b. G. 1888 p. 761; (f. foliis in lacinias angustas dissecta, ad typum manifeste transiens). — *S. laciniata v. multifida* Bois. fl. or. IV. p. 409; Hal. Beitr. fl. Epir. p. 33 f. *puberula*, in ö. b. Z. 1897, p. 287; Form. in Ver. Brünn 1896 p. 63, 1897 p. 46. — Exsicc.: Orph. fl. gr. n. 729; Heldr. herb. norm. n. 682; Sint. it. thessal. n. 673; Dörfl. fl. gr. n. 226.

A collo pluricaulis; caulibus erectis vel adscendentibus, tetragonis, simplicibus vel parce ramosis, glabris, superne pedicellisque glanduloso-puberulis; foliis oppositis, petiolatis, glabris, tenuibus, laete viridibus, lyratis vel pinnatipartitis sectisve, segmentis acute dentatis vel incisis; panicula subpyramidata, subaphylla, cymis inferioribus plurifloris; pedicellis calyce brevioribus vel aequilongis; calycis glabri laciniis orbiculatis, late albo-marginatis; corolla luride purpurea; antheris inclusis; staminodio late orbiculato-reniforme, interdum retuso; capsula globosa, apiculata, calyce vix duplo longiore. — „Valde affinis formis pinnatisectis *S. heterophyllae*, pro cujus stirpe subalpina et alpina haberi posset, differt tamen colore laete viridi, foliis tenuius membranaceis, nec carnosis, lobis et dentibus saepius acutis, nec obtusis, caulibus non adeo fragilibus, capsula calycis proportione breviore“ (Bois. fl. or. IV. p. 409), attamen in herbario interdum difficile diagnoscenda.

In rupestribus regionis alpinae et subalpinae. Epirus: mt. Papignon, Smolika (Bald.), pr. Janina (Form.), mt. Peristeri (Hal.); Thessalia: mt. Salatura pr. Chaliki (Sint.), pr. Koturi, Velitsena, mt. Baba pr. Krania (Form.), Ghavellu, Karava (Haussk.), Agrapha, Othrys, pr. Neraida (Form.), mt. Olympus (Orph.), Oeta (Heldr.); Aetolia: mt. Tymphrestus (Samar.), Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Boeotia: pr. Larmes (Boissieu); Peloponnesus: mt. Olenos, pr. Megaspilaeon, mt. Chelmos, Kyllene, Taygetos (Heldr.), mt. Arachneon Argolidis (Sprun.). — Jun. Aug. ♃

×× Planta tota pube patula crispa glandulosa obsita.

12. **S. taygetea** Bois. diagn. IV. p. 68, Fl. or. IV. p. 406.

A collo pluricaulis; caulibus teretiusculis, fragillimis; foliis oppositis, petiolatis, triangulari-ovatis, cuneato-truncatis vel subcordatis, acutis, acute et subduplicatim dentatis, saepe lobatis; panicula abbreviata, aphylla, cymis breviter pedunculatis, 7—3 floris, summis unifloris; pedicellis calyce subaequilongis; calycis glabri laciniis orbiculatis, late

membranaceo-marginatis; corolla virescenti-flavida; antheris inclusis; staminodio transverse latiore, reniforme; capsula globosa, apiculata, calyce subduplo longiore. — A praecedentibus indumento viscido et floribus flavidis distincta.

In fissuris rupium regionis inferioris mt. Taygetos pr. Mistra (Bois.). — Apr. ♃ N. v.

b. Staminodium anguste lineare.

13. **S. canina** L. sp. p. 621; S. et S. pr. I. p. 436, non Fl. gr. t. 598, quae a Rchb. fl. germ. exc. certe erronee ad *S. lucidam* ducitur et sec. Bois. fl. or. IV. p. 403 ad *S. sphaerocarpam* probabiliter referenda est.; Pieri corc. fl. p. 84; Ch. et B. exp. p. 177, Fl. pelop. p. 40; Friedr. Reise p. 265, 269, 275 et 283; Ung. Reise p. 129; Raul. cret. p. 822; Bois. l. c. p. 419; Heldr. fl. ceph. p. 55, Fl. Aegina p. 382; Spreitz. in z. b. G. 1887 p. 665; Form. in D. bot. Mon. 1891 p. 26, in Ver. Brünn 1895 p. 34; Hal. Beitr. fl. Epir. p. 33, Beitr. fl. Thessal. p. 16; Haussk. symb. p. 171, cum v. *pallida* (f. colore subcaesio-viridi, inflorescentia vix glandulosa, corolla late flavido-marginata) et v. *dolopica* (f. foliorum segmentis brevibus, latiusculis, subobtusis, panicula angusta, magis glandulosa, cymis paucifloris, abbreviatis, capsula minore). — *S. bicolor* S. et S. pr. I. p. 437, Fl. gr. VI. p. 1 t. 602; Weiss in z. b. G. 1869 p. 745. — *S. chrysanthemifolia* Willd. hort. berol. t. 59. — *S. ramosissima* Urv. enum. p. 75; Haussk. symb. p. 171 pr. var. *S. caninae*; non Lois. fl. gall. II. p. 381; (f. caulibus a basi ramosissimis, ramis elongatis, intricatis). = *S. pyramidalis* Wydl. mon. scrof. p. 45; Marg. et R. fl. Zante p. 71. = *S. scoparia* Marg. et R. l. c. p. 24 et 71. — *S. pindicola* Haussk. symb. p. 172 (f. a basi intricato-ramosa, ramis tenuibus, foliorum infimorum bipinnatorum segmentis paucidentatis, foliis superioribus plerisque trisectis, segmento terminali elongato, foliis summis integris, corolla late flavido-marginata, capsula minore). — Exsicc.: Orph. fl. gr. n. 727; Sint. et Bornm. it. turc. n. 1384; Sint. it. thessal. n. 174, 674 et 1186; Baen. herb. europ. n. 9335; Dörfl. fl. aeg. n. 68.

Pluricaulis; caulibus erectis vel adscendentibus, rigidis, obtusangulis, simplicibus vel ramosis, glabris vel superne pedicellisque glandulosis; foliis plerisque oppositis, pinnatifidis vel pinnati-vel bipinnatisectis, inferioribus petiolatis, segmentis oblongis vel lanceolatis, dentatis vel integris; panicula elongata vel pyramidata, cymis semel vel pluries bifidis, laxe 9—3 floris; pedicellis calyce brevioribus; calycis glabri laciniis orbiculatis, late scariosis; corolla minuta, purpurea, lobis superioribus margine albidis; staminibus exsertis; staminodio anguste-lineari, rarius nullo; capsula globosa, apiculata. — Stirps quoad ramificationem et praesertim foliorum divisionem et laciniarum forman maxime polymorpha, sed inter omnes formas transitus observantur; ab omnibus praecedentibus floribus minutis et staminibus longe exsertis insignis.

In siccis, collibus, lapidosis regionis inferioris et montanae, frequens. Epirus: pr. Ephemiades, Safikbi, Pruvilastro, Sadovica, mt. Micikeli,

Han Kanberga, in valle Dipotami, mt. Prosgoli (Form.), pr. Theodoriana, Kalarrytes (Hal.).; Thessalia: pr. Chaliki (Hal.), Malakasi, mt. Zygos, Ghavellu, pr. Sermeniko (Haussk.), Klinovo, ad monasterium Korona (Heldr.), Kastania, Vendista, Velitsena, Koturi, Spilia et Selicany in mt. Ossa, pr. Portaria, Volo (Form.), Lithochori in Olympo (Sint.); pr. Rachova ad radices mt. Parnassus (Hal.); Euboea (Ung).; Attica: pr. Athenas (Friedr.), Menidi (Pichl.), Piraeum (Heldr.), mt. Kerata, Pentelicon (Haussk.); insula Aegina, Poros (Friedr.), Syra (Urv.), Paros (Leon.); Peloponnesus (Sibth.): in Arcadia, Messenia, mt. Taygetos (Chaub.), pr. Misitra Laconiae (Friedr.); Creta (Sibth.); Zante: mt. Scopo (Marg.); Cephalonia: pr. Argostoli, Drapano, Pessada, Omala (Heldr.); Leucas: ad monasterium Hagios Georgios (Spreitz.); Corcyra (Pieri): pr. Spartilla (Baen.). — Mart. Jul. ♃

4. Gratiola L. gen. 29.

1. **G. officinalis** L. sp. p. 17; Dallap. prosp. p. 9; Ch. et B. fl. pelop. p. 1; Heldr. fl. ceph. p. 55; Hal. in ö b. Z. 1896 p. 16; Bald. riv. coll. bot. Alb. 1896 p. 83; Form. in Ver. Brünn 1897 p. 47; Haussk. symb. p. 174. — Icon: Fl. dan. t. 363. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Glabra; rhizomate repente; caule erecto vel adscendente, simplici vel ramoso; foliis sessilibus, lanceolatis, integris vel serrulatis; floribus axillaribus, pedunculatis; calycis laciniis lanceolatis; corollae calyce 5plo longioris tubo luteolo, limbo roseo, fauce intus barbata; capsula ovata, acuta, calyce aequilonga.

Ad fossas, in humidis rarissime. Epirus: pr. Kastriza, Lapsista (Bald.).; Thessalia: ad monasterium Korona in Pindo (Heldr.), pr. Palaeokastro (Leon.); Peloponnesus: pr. Philiatra (Gittard); Cephalonia (Dall.). — Jun. Aug. ♃

3. Tribus. ANTIRRHINEAE Chav. monogr. antirrh. p. 73.

5. Linaria Tourn. inst. t. 76.

Dispositio specierum.

1. Sectio. *Linariastrum* Chav. monogr. antirrh. p. 114. — Folia sessilia, penninervia, integra; capsula 4—10 valvulis vel dentibus apice dehiscens; semina oblonga vel reniformia vel angulosa, immarginata, aut discoidea, membranaceo-marginata.

a. Semina angulato-triquetra, immarginata.

α. Corolla lutea, interdum coerulea variegata.

× Perennes.

○ Calycis laciniae lanceolatae, acutae, immarginatae.

1. L. dalmatica (L.). **2. L. genistifolia** (L.).

○○ Calycis laciniae ovatae vel oblongae, obtusae, membranaceo-marginatae.

3. L. peloponnesiaca Bois. et Heldr.

XX Annua.

4. L. triphylla (L.).

β. Corolla alba vel purpurea.

5. L. chalepensis (L.). **6. L. purpurea** (L.).

b. Semina oblonga, immarginata.

7. L. reflexa (L.).

c. Semina discoidea, plana, membranaceo-marginata.

α. Perennis; corolla majuscula, cum calcare 25—30 mm longa, flava.

8. L. vulgaris Mill.

β. Annuae; corolla parva, cum calcare 4—15 mm longa, violacea vel coerulescens, raro flava.

× Corolla cum calcare 12—15 mm. longa, intense violacea; seminum ala fimbriato-ciliata.

9. L. Pelisseriana (L.).

XX Corolla minuta, cum calcare 4—6 mm. longa, coerulescens vel flava; seminum ala non ciliata.

○ Corolla minuta, tubo calyce breviore, calcare subulato, corolla subaequilongo.

10. L. arvensis (L.). **11. L. parviflora** (Jacq.)

○○ Corolla minima, calycem parum excedente, calcare brevissimo, conico.

12. L. micrantha (Cav.).

2. Sectio *Elatinoides* Chav. monogr. antirrh. p. 103. — Folia penninervia, saepius breve petiolata, dentata lobatave, interdum integra; capsula operculis binis circumscissis latere dehiscens; semina ovata vel oblonga.

a. Semina reticulato-foveolata vel lacunosa; capsula membranacea.

13. L. spuria (L.). **14. L. Sieberi** Rchb.

b. Semina tuberculata; capsula coriacea.

15. L. commutata Bernh. **16. L. cirrhosa** (L).

3. Sectio. *Cymbalaria* Chav. monogr. antirrh. p. 98. — Folia palminervia, saepissime lobata, longe petiolata; capsula 6 valvulis apice dehiscens; semina oblonga.

a. Planta glaberrima.

17. L. cymbalaria (L.). **18. L. longipes** Bois. et Heldr.

b. Planta lana longa alba crispa villosa.

19. L. microcalyx Bois.

1. Sectio. *Linariastrum* Chav. mon. antirrh. p. 114.

a. Semina angulato-triquetra, immarginata.

α. Corolla lutea, interdum coeruleo-variegata.

× Perennes.

○ Calycis laciniae lanceolatae, acutae, immarginatae.

1. **L. dalmatica** L. sp. p. 616 (*Antirrhinum*); Mill. dict. n. 13; Raul. cret. p. 821; Bois. fl. or. IV. p. 376; Form. in Ver. Brünn 1895 p. 34. — *L. Smithii* Bois. et Orph. in Orph. fl. gr. n. 711. — Icon: Rchb. t. 65.

Glabra, simplex vel saepius paniculatim-ramosa, surculis sterilibus nullis; foliis alternis, ovato-lanceolatis lanceolatisve, semiamplexicaulibus

racemis tandem elongatis, laxis; bracteis ovato-lanceolatis, pedicello brevioribus; calycis laciniis ovato-lanceolatis, corollae tubo subbrevioribus; corolla magna, sine calcare ad 1 cm longa, lutea, palato villoso, calcare recto, corolla breviore; capsula ovata, calyce subbreviore; seminibus ovatis, triquetris, muricato-rugosis.

β. **macedonica** Griseb. spic. II. p. 19; Heldr. in Sitzungsb. acad. wiss. Berl. 1883 p. 4; pro sp.; Fenzl in Tchih. as. min. II. p. 25. — *L. dalmatica* Haussk. symb. p. 174; Form. in Ver. Brünn 1895 p. 64; 1897 p. 47 v. *thessala*. — Calycis laciniae corollae tubo manifeste breviores; pedicelli longiores; corolla minor. — Exsicc: Heldr. it thessal. n. 51; Form. pl. gr. a. 1896; Sint. it. thessal. n. 998 et 998b.

In vinetis, apricis montanis. Epirus: mt. Micikeli (Form.); Euboea: mt. Dirphys (Sprun); Laconia: mt. Malevo pr. Platanos (Orph.); Creta (Linné). — *β.* Thessalia: pr. Kalabaka (Sint.), Pharsalus, Aivali (Haussk.), ad Phlambures et mt. Mitriza in mt. Chassia, pr. Patsios et Godaman in Olympo (Form.), mt. Pelion pr. Portaria (Heldr.). — Jun. Jul. ♃

2. **L. genistifolia** L. sp. p. 616 (*Antirrhinum*); Mill. dict. n. 14; Form. in D. bot. Mon. 1891 p. 26. — *L. dolopica* Form in Ver. Brünn. 1897 p. 47. — ? *L. dalmatica* v. *ramosissima* Haussk. symb. p. 174. — Icon: Jacq. fl. austr. t. 244. — Exsicc.: Orph. fl. gr. n. 714 (Thessalonica, f. *angustifolia*); Heldr. it. thessal. IVa. 1885 (f. *latifolia*); Sint. it. thessal. n. 1300 (f. *brevicalcarata*).

Glabra, paniculatim-ramosa, surculis sterilibus nullis; foliis alternis, lanceolatis vel oblongo-lanceolatis, sessilibus; racemis laxiusculis, virgatis; bracteis lanceolatis, pedicello longioribus; calycis laciniis lanceolatis, corollae tubum aequantibus; corolla mediocri, sine calcare ad 1 cm longa, lutea, palato barbato, calcare rectiusculo, corolla aequilonga vel breviore; capsula ovata, calyce subbreviore; seminibus ovatis, triquetris, muricato-rugosis. — Foliis angustioribus, pedicellis brevioribus et floribus fere duplo minoribus a praecedente distincta.

In vineis, dumosis regionis inferioris et montanae. Thessalia: ad monasterium Korona (Held.), pr. Vendista Kalabaka (Form.), pr. Portaria in mt. Pelion (Sint.). — Jun. Aug. ♃

○○ Calycis laciniae ovatae vel oblongae, obtusae, membranaceo-marginatae.

3. **L. peloponnesiaca** Bois. et Heldr. diagn. ser. 2. III. p. 163; Hal. Beitr. fl. Epir. p. 33, in ö. b. Z. 1897 p. 324. — *L. Sibthorpiana α. peloponnesiaca* Bois. fl. or. IV. p. 378. — Exsicc.: Orph. fl. gr. n. 137; Heldr. herb. norm. n. 1462.

Glabra, pluricaulis, a collo surculos steriles edente; caulibus erectis vel adscendentibus, foliosis, simplicibus vel paniculato-ramosis; foliis alternis, anguste-linearibus, sessilibus; racemis confertifloris, oblongis, fructiferis parum elongatis; bracteis squamaeformibus; pedicellis glabris, calyce aequilongis vel sublongioribus; calycis glabri laciniis ovatis; corolla mediocri, pallide lutea, palato barbato, calcare rectiusculo, corollae

aequilongo; capsula ovata, calyce 2—3plo longiore; seminibus triquetris, minute tuberculatis. — Stirps ad 60 cm. alta et tunc saepius ramosa, sed in alpinis ad 10 cm. tantum alta et simplex. Surculi steriles, qui a Boissier negantur, saepe adsunt.

β. **parnassica** Bois. et Heldr. diagn. ser. 2. III. p. 164; Hal. in z. b. G. 1888 p. 761; pro sp.; Hal. in ö. b. Z. 1897 p. 324. — *Antirrhinum strictum* S. et S. pr. I p. 433 p. p. (cum et plantam orientalem et siculam = *L. strictam* Guss. pl. rar. p. 250, a nostra calycis laciniis lineari-lanceolatis longioribus, corolla majori palato intense luteo diversam, amplectit), Fl. gr. t. 594. — *L. Sibthorpiana β parnassica* Bois. fl. or. IV. p. 378; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 6, chlor. Parn. p. 24; Form. in D. bot. Mon. 1891 p. 26; Bald. riv. coll. bot. Alb. 1896 p. 82; Haussk. symb. p. 175. — *L. Sibthorpiana* Form. in Ver. Brünn 1896 p. 63. — *L. olympica* Form. l. c. — Calycis laciniae oblongo-lineares pedicellisque pubescenti-glandulosi. — Exsicc.: Orph. fl. gr. n. 715 et 1009; Heldr. herb. norm. n. 655; Bald. it. alb. epir. IV. n. 192; Sint. it. thessal. n. 920.

In herbidis lapidosis regionis montanae et alpinae. Epirus: in summis mt. Peristeri (Hal.); Achaia: mt. Kyllene (Heldr.); Arcadia: pr. Zatuna (Orph.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.); — β. Epirus: mt. Gamila distr. Zagorion (Bald.); Thessalia: pr. Malakasi (Sint.), mt. Zygos (Haussk.), Olympus (Orph.), Godaman (Form.), Ossa (Heldr.), Pelion (Form.); Eurytania: mt. Chelidoni (Heldr.); mt. Kiona (Hal.), Parnassus (Sprun.). — Jul. Aug. ♃

×× Annua.

4. **L. triphylla** L. sp. ed. 2. p. 852; Ch. et B. exp. p. 341, Fl. pelop. p. 39 (*Antirrhinum*); Mill. dict. n. 2; Urv. enum. p. 74; Friedr. Reise p. 269; Clem. sert. p. 71; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 821; Bois. fl. or. IV. p. 380; Haussk. symb. p. 175; Heldr. fl. Aegina p. 382. — Icon: Rchb. t. 63. — Exsicc.: Orph. fl. gr. n. 779; Heldr. herb. norm. n. 471, herb. dimorph. n. 39; Rev. pl. cret. n. 121; Dörfl. fl. gr. n. 467.

Glabra, 1 — pluricaulis, a collo surculos steriles edente; caulibus erectis, simplicibus vel parce breviterque ramulosis; foliis sessilibus, ovatis vel ellipticis, obtusis, surculorum oppositis, caulium ternatis; spica densiuscula, floribus sessilibus; calycis laciniis ovato-oblongis, obtusis; corolla lutea, lilacino vel coeruleo variegata, calcare leviter arcuato, corollae subaequilongo; capsula ovata, calyce aequilonga; seminibus triquetris, grosse reticulatis.

In arvis, vinetis, inter segetes regionis calidae. Attica: ubique cica Athenas; Corinthus, Acrocorinthus (Haussk.); Argolis: pr. Argos, Nauplia (Daenzer); insula Aegina (Friedr.), Melos (Urv.); Creta: pr. Canea (Raul.). — Febr. Maio. ⊙

β. Corolla alba vel purpurea.

5. **L. chalepensis** L. sp. p. 617; S. et S. pr. I. p. 433; Ch. et B. exp. p. 175, Fl. pelop. p. 39; (*Antirrhinum*); Mill. dict. n. 12;

Marg. et R. fl. Zante p. 71; Friedr. Reise p. 269; Raul. cret. p. 821; Bois. fl. or. IV. p. 381; Heldr. fl. cephal. p. 55, Fl. Aegina p. 382; Bald. riv. coll. bot. alb. 1896 p. 62; Haussk. symb. p. 175. — Icon: Fl. gr. t. 592. — Exsicc.: Orph. fl. gr. n. 394; Dörfl. fl. gr. n. 475.

Glabra, 1 — pluricaulis, a collo surculos steriles edente; caulibus erectis, simplicibus vel parce ramosis; foliis sessilibus surculorum et caulinis inferioribus oblongis, oppositis, ceteris linearibus, alternis; racemo laxo; pedicellis calyce multo brevioribus; bracteis linearibus, pedicello longioribus; calycis laciniis linearibus, acutis, corollam aequantibus vel superantibus; corolla alba, calcare gracillimo, arcuato, corolla sesqui vel duplo longiore; capsula ovata, calyce acuto valde superata; seminibus triquetris, rugoso-scrobiculatis.

In arenosis, arvis, cultis regionis inferioris. Epirus: pr. Dodona, Kurenta (Bald.); Thessalia: pr. Volo (Heldr.); Euboea (Sprun.); Attica: pr. Athenas (Heldr.), Oropos (Orph.); Peloponnesus (Sibth.): pr. Nauplia (Haussk.), Phigalea (Chaub.); insula Aegina (Friedr.), Cyclades (Oliv.); Creta: pr. Canea (Raul.); Zante (Marg.); Cephalonia: pr. Drapano (Heldr.). — Mart. Maio. ⊙

6. **L. purpurea** L. sp. p. 613; S. et S. pr. I. p. 432 (*Antirrhinum*); Mill. dict. n. 5; Bois. fl. or. IV. p. 380. — Icon: Fl. gr. t. 589. —

Glabra, 1 — pluricaulis, a collo surculos steriles edente; caulibus erectis, foliosis, simplicibus vel fastigiatim paniculatis; foliis sessilibus, lineari-lanceolatis linearibusve, inferioribus et surculorum verticillatis, ceteris alternis; racemis densiusculis; pedicellis calyce aequilongis vel sublongioribus; bracteis linearibus, pedicello aequilongis vel brevioribus; calycis laciniis anguste lanceolatis, acutiusculis, corollae tubo brevioribus; corolla violacea, calcare arcuato, corollae tubo longiore; capsula globosa, retusa, calyce subtriplo longiore; seminibus triquetris, rugosis.

In Graecia (Sibth.), loco non notato. — Jun. Jul. ♃ N. v.

b. Semina oblonga, immarginata.

7. **L. reflexa** L. sp. ed. 2 p. 857; S et S. pr. I. p. 433; Ch. et B. exp. p. 176, Fl. pelop. p. 39; (*Antirrhinum*); Desf. fl. atl. II. p. 42; Bois. fl. or. IV. p. 380. — Icon: Fl. gr. t. 593. —

Glabra, multicaulis, a collo surculos steriles edente; caulibus diffusis, adscendentibus simplicibus vel parce ramosis; foliis sessilibus, ovatis vel oblongis, acutis, surculorum et inferioribus ternis, ceteris alternis; floribus ad axillas racemum pauciflorum, laxum, foliatum formantibus; pedicellis folio brevioribus vel aequilongis, demum arcuato-reflexis, elongatis, folio longioribus; calycis laciniis lanceolatis, acutis, corollae tubum subaequantibus; corolla coerulea vel albida, calcare gracili rectiusculo, corollae subaequilongo; capsula globosa, calyce breviore; seminibus subincurvis, profunde rugoso-foveolatis.

In agro Argolico (Sibth.). — Jan. Apr. ⊙ N. v.

c. Semina discoidea, plana, membranaceo-marginata.

α. Perennis; corolla majuscula, cum calcare 25—30 mm. longa, flava.

8. **L. vulgaris** Mill. dict. n. 1; Bois. fl. or. IV. p. 372; Heldr. fl. cephal. p. 55. — *Antirrhinum linaria* L. sp. p. 616; S. et S. pr. I. p. 434; Dallap. prosp. p. 86; Ch. et B. exp. p. 176, Fl. pelop. p. 39. — Icon: Fl. dan. t. 982.

Glabra vel inflorescentia pubescenti-glandulosa, 1 — pluricaulis, a collo surculos steriles edente; caulibus erectis, foliosis, simplicibus vel superne ramosis; foliis sessilibus, anguste lineari-lanceolatis, acutis; racemis densis; pedicellis calyce brevioribus; bracteis breviter lanceolatis, pedicellum superantibus; calycis laciniis triangulari-ovatis, corollae tubo multo brevioribus; corolla flava, palato aurantiaco, calcare rectiusculo subaequilongo; capsula ovata, calyce duplo longiore; seminibus marginatis, disco saepius tuberculatis.

In arvis, cultis, rarissime. Attica: pr. Ampelokipos (Heldr.); in agro Laconico (Sibth.); Cephalonia (Dallap.). — Jun. Aug. ♃ N. v.

β. Annuae; corolla parva cum calcare 4—15 mm. longa, violacea vel coerulescens, raro flava.

× Corolla cum calcare 12—15 mm. longa, intense violacea; seminum ala fimbriato-ciliata.

9. **L. Pelisseriana** L. sp. p. 615; S. et S. pr. I. p. 432; Fl. gr. t. 591; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 176, Fl. pelop. p. 39 (*Antirrhinum*); Mill dict. n. 11; Urv. enum. p. 74; Friedr. Reise p. 276; Ung. Reise p. 129; Weiss in z. b. G. 1869 p. 745; Raul. cret. p. 821; Bois. fl. or. IV. p. 375; Gelmi in bull. soc. bot. it. 1889 p. 451; Hal. Beitr. fl. Epir. p. 33, in ö. b. Z. 1897 p. 77; Bald. riv. coll. bot. alb. 1896 p. 82; Haussk. symb. p. 175; Heldr. fl. Aegina p. 382, chlor. Mykon. p. 248. — *Antirrhinum saphirinum* Sieb. in Flora V. II. p. 639. — Exsicc.: Orph. fl. gr. n. 713; Sint. it. thessal. n. 506; Dörfl. pl. cret. n. 54.

Glabra 1—pluricaulis, a collo surculos steriles edente; caulibus gracilibus, erectis, simplicibus vel ramosis; foliis sessilibus, surculorum oblongis, verticillatis, caulinis remotis, alternis, anguste linearibus, acutis; floribus paucis, capitatis, tandem laxe racemosis; pedicellis calyce sublongioribus; bracteis linearibus, pedicello brevioribus; corolla violacea, calcari rectiusculo subaequilonga; capsula globosa, calyce breviore; seminibus marginatis, radiatim ciliatis, disco laeviusculis.

In herbidis regionis inferioris et montanae. Epirus: pr. Kalentini supra Arta (Hal.), mt. Cuka, pr. Janina (Bald.); Thessalia: pr. Vutades in Pindo tymphaeo (Sint.), monasterium Korona, Kalabaka (Haussk.); Sporadum insula Scopelos (Leon.); Euboea: pr. Kurbatzi (Willd.), mt. Dirphys (Ung.); Attica: pr. Athenas, mt. Pentelicon, insula Aegina (Heldr.); Argolis: peninsula Methana (Haussk.), pr. Poros (Friedr.), Astros (Orph.); Elis (Sibth.): in silva Kapellis pr. Lala (Reis.);

Messenia (Sibth.): pr. Pylos, Arcadia usque Messene (Chaub.); Cycladum insula: Tenos, Kythnos, Mykonos, Kimolos (Heldr.), Melos (Urv.); Creta: pr. Canea (Weiss), Nerokuru (Raul.), Males distr. Hierapetra (Leon.); Corcyra: pr. Signes (Gelmi). — Apr. Jul. ⊙

×× Corolla minuta, cum calcare 4—6 mm. longa, coerulescens vel flava; seminum ala non ciliata.

○ Corolla minuta, tubo calyce breviore, calcare subulato, corolla subaequilongo.

10. **L. arvensis** L. sp. p. 614; S. et S. pr. I. p. 432; Ch. et B. exp. p. 175, Fl. pelop. p. 39 (*Antirrhinum*); Desf. fl. alt. II. p. 45; Bois. fl. or. IV. p. 375. — Icon: Rchb. t. 62.

Glabra, in inflorescentia glandulosa, 1—pluricaulis, a collo surculos steriles edente; caulibus gracilibus, erectis, simplicibus vel ramosis; foliis sessilibus, linearibus, acutis, inferioribus verticillatis, superioribus alternis; floribus paucis, capitatis, tandem laxe racemosis; pedicellis calyce brevioribus; bracteis linearibus, pedicello sublongioribus; corolla lilacina, violaceo-striata, calcare subulato, curvato, fere angulum rectum cum tubo formante, corolla subaequilongo; capsula subglobosa, calycem excedente; seminibus marginatis, disco laevi vel minute punctulato.

In agris, campis incultis, arenosis regionis inferioris Peloponnesi (Sibth.): pr. Navarin (Chaub.). — Apr. Jun. ⊙ N. v.

11. **L. parviflora** Jacq. ic. rar. III. t. 499, collect. IV. p. 204 (*Antirrhinum*). — *A. simplex* Willd. sp. III. p. 243. — *L. simplex* DC. fl. fr. III. p. 588; Weiss. in z. b. G. 1869 p. 744; Bald. riv. coll. bot. alb. 1896 p. 82; Haussk. symb. p. 175. — *L. arvensis β. flaviflora* Bois. fl. or. IV. p. 375. — Icon: Fl. gr. t. 590. — Exsicc.: Sint. it. thessal. n. 25.

Differt a praecedente corolla paullo majore, lutea, interdum violaceo-striata, calcare recto, seminibus saepius minute tuberculatis.

In herbidis, rupestribus regionis inferioris et montanae. Epirus: mt. Maria distr. Ljaskovik (Bald.); Thessalia: ad radices mt. Pelion pr. Volo (Heldr.), Agria (Sint.); Attica: ad Phalerum (Haussk.); Arcadia: ad radices mt. Chelmos pr. Planideri (Hald.); Cycladum insula Syra (Weiss). — Apr. Jun. ⊙

○○ Corolla minima, calycem parum excedente, calcare brevissimo, conico.

12. **L. micrantha** Cav. ic. I. p. 51 t. 59 (*Antirrhinum*); Spreng. syst. II. p. 793; Clem. sert. p. 7; Bois. fl. or. IV. p. 375; Haussk. symb. p. 175; Heldr. fl. Aegina p. 382. — *L. parviflora* Desf. fl. atl. II. p. 44, non (Jacq.). — *A. parviflorum* Willd. sp. III. p. 245, non Jacq. — Icon: Fl. gr. t. 587. — Exsicc.: Orph. fl. gr. n. 46; Heldr. herb. norm. n. 358; Dörfl. fl. gr. n. 468.

Glabra vel in inflorescentia plus minus glandulosa, 1—pluricaulis, a collo surculos steriles edente; caulibus erectis, simplicibus vel ramosis; foliis sessilibus, surculorum linearibus, caulinis oblongo-lanceolatis, acutis;

floribus paucis, capitatis, tandem laxe racemosis; pedicellis brevissimis; bracteis linearibus, pedicello multo longioribus; corolla coerulea, calcare conico brevissimo, fere angulum rectum cum tubo formante; capsula subglobosa, calycem excedente; seminibus marginatis, disco minute tuberculatis. — A praecedentibus forma foliorum, floribus minoribus brevissime pedicellatis, calcare gibbiformi distincta.

In arvis, Attica: pr. Athenas (Orph.), insula Aegina (Heldr.); Acrocorinthus (Haussk.); Euboea (Sprun.); insula Syra (Heldr.). — Febr. Apr. ⊙

2. Sectio. *Elatinoides* Chav. mon. antirrh. p. 103.

a. Semina reticulato-foveolata vel lacunosa; capsula membranacea.

13. **L. spuria** L. sp. p. 613; S. et S. pr. I. p. 430; Pieri corc. fl. p. 83; Ch. et B. fl. pelop. p. 39 (*Antirrhinum*); Mill. dict. n. 15; Marg. et R. fl. Zante p. 71; Friedr. Reise p. 275; Clem. sert. p. 71; Raul. cret. p. 821; Bois. fl. or. IV. p. 366; Heldr. fl. cephal. p. 55; Bald. viagg. Creta p. 81; Form. in Ver. Brünn 1895 p. 34, 1896 p. 63, 1897 p. 47; Haussk. symb. p. 175. — Icon: Fl. dan. t. 913. — Exsicc.: Heldr. herb. norm. n. 656 et 866.

Villosa, multicaulis; caulibus prostratis, elongatis, subsimplicibus; foliis sessilibus vel brevissime petiolatis, ovatis, obtusis vel acutiusculis, integris vel subdentatis, basi rotundatis rarius cordatis; inferioribus oppositis, ceteris alternis; pedicellis axillaribus, villosis, patentibus, inferioribus folio brevioribus, superioribus eo longioribus; calycis villosuli laciniis ovato-oblongis, acutis, basi subcordatis; corollae hirtulae, flavae, labio superiore atroviolaceo, calcare incurvo, corollae aequilongo; capsula globosa, minutissime puberula; seminibus reticulato-foveolatis.

In arvis, vinetis, olivetis regionis inferioris. Epirus: pr. Prevesa, Ephemiades, Safikbi, mt. Prosgoli (Form.); Thessalia: pr. Han Tripa in valle Penei, Trikala, Pharsalus, Agria, Lechonia, Volo (Form.); Euboea: pr. Kastaniotissa (Heldr.); Attica: pr. Athenas, Phalerum (Heldr.); Peloponnesus (Chaub.): pr. Poros (Friedr.); Archipelagus (Sibth.); Creta: pr. Hagios Deka, Istovai in promontorio Sidero (Raul.), Gonia, Rustika (Bald.); Zante (Marg.); Cephalonia: pr. Drapano, Heno, Akoli (Heldr.); Corcyra (Pieri): pr. Alipu, Kanali, Stratia, Analypsis, Hagios Elias, Potamos, Kontokali (Form.). — Jun. Sept. ⊙

14. **L. Sieberi** Rchb. fl. exc. p. 374; Haussk. symb. p. 175; Heldr. fl. Aegina p. 382, in ö. b. Z. 1898 p. 184, chlor. Thera p. 18, chlor. Mykon. p. 248. — *Antirrhinum elatine* S. et S. pr. I. p. 430; Sieb. avis p. 4, rem. p. 6; non L. sp. p. 612, quod statura graciliore, caulibus minus ramosis, indumento multo parciore, pedicellis glabris, [illegible] incedit. — *L. lanigera* Link in Linn. IX. [illegible] II. p. 38, quae foliis floralibus orbiculari- [illegible] nec hastatis et inflorescentia paniculata [illegible] B. fl. pelop. p. 39, non Willd. sp.

III. p. 235. — *L. elatine* Friedr. Reise p. 263, non Mill. dict. n. 16. — *L. elatine v. lasiopoda* Vis. fl. dalm. II. p. 161; Weiss in z. b. G. 1869 p. 744 — *L. Prestandreae* Tin. in Guss. Syn. II. p. 842. — *L. elatine β. villosa* Bois. fl. or. IV. p. 367; Bald. in nuov. giorn. bot. ital. 1894 p. 99, viagg. Creta p. 81. — Exsicc.: Heldr. herb. norm. n. 656, 867 a et b.

Dense villosa, multicaulis; caulibus prostratis, elongatis, ramosis, rarius subsimplicibus; foliis breviter petiolatis, ovatis, obtusis vel acutiusculis, integris vel remote dentatis, saepe hastatis vel sagittatis, inferioribus oppositis, ceteris alternis; pedicellis axillaribus, glabris vel hirsutis, patentibus, folio axillari saepius multo longioribus; calycis dense villosi laciniis lanceolatis, acutis; corolla hirtula, albida, labio superiore violaceo, calcare subincurvo, corollae subaequilongo; capsula globosa, minutissime puberula; seminibus foveolato-lacunosis. — Indumento copiosiori, foliis hastatis, calycis laciniis angustioribus, basi non cordatis a praecedente distincta.

β. **bombycina** Bois. et Blanche diagn. ser. 2 III. p. 161, pro sp. — Tota lana crispa bombycina obducta, pedicellis saepe folio axillari brevioribus. — Intermediis ad typum transit. — Exsicc.: Rev. pl. cret. n. 120.

In locis aridis, campis, cultis regionis inferioris. Epirus: pr. Prevesa (Bald.); Thessalia: pr. Pharsalus (Haussk.); Attica: pr. Athenas Tatoi, Phaleron; insula Aegina, Corinthus (Heldr.); Argolis: peninsula Methana (Haussk.); Achaia: pr. Diakophto (Heldr.); Cycladum insula: Rhenea (Tunt.), Syra (Weiss), Naxos, Amorgos, Thera (Heldr.); Creta: pr. Kissamos, Canea (Rev.), Kladiso (Raul.), Armeni (Bald.); Corcyra (Friedr.). — Maio, Sept. ⊙

b. Semina tuberculata; capsula coriacea.

15. **L. commutata** Bernh. in Rchb. ic. IX. p. 6 t. 815, Fl. exc. p. 373; Haussk. symb. p. 175. — *Antirrhinum graecum* Ch. et B. exp. p. 175 t. 21, Fl. pelop. p. 39 t. 22. — *L. graeca* Chav. monogr. antirrh. p. 108; Vis. ill. alc. piante p. 11; Fraas fl. class. p. 189; Raul. cret. p. 821; Bois. fl. or. IV. p. 367; Hal. Beitr. fl. Epir. p. 33, Beitr. fl. Aetol. p. 9; Bald. in nuov. giorn. bot. ital. 1894 p. 99; Heldr. fl. Aegina p. 382, *f. gracilior*, foliis superioribus remotis, valde diminutis hastatis, pedunculis sublongioribus. — Huc forsan: *A. aegyptiacum* S. et S. pr. I. p. 431 non L. — Exsicc.: Heldr. pl. fl. hellen. a. 1885 et 1899.

Pilosa, multicaulis; caulibus prostratis, elongatis, simplicibus vel ramosis; foliis breviter petiolatis, inferioribus ovato-oblongis, oppositis, superioribus sensim diminutis, alternis, ovatis vel ovato-lanceolatis, hastatis; pedicellis axillaribus, glabris, patentibus, folio multo longioribus, apice plerumque inflexis; calycis hirti laciniis lanceolatis, acutis; corolla cum calcare 12—15 mm. longa, flavida vel coerulescenti, hirtula, calcare rectiusculo, corollae subaequilongo; capsula globosa, glabra.

β. **polygonoides** Rev. pl. cret. n. 119 pro var. *L. elatines.* — Glabra vel minutissime puberula.

In collibus herbidis, montosis, aridis, fruticetis. Epirus: pr. Kalentini supra Arta (Hal.); Thessalia: ad monasterium Korona, pr. Malakasi, Kalabaka, Pharsalus, Sophates, Karditza (Haussk.); Acarnania: pr. Agrinion (Hal.); Aetolia (Nied.); Attica: mt. Parnes, insula Aegina (Heldr.); Argolis (Fraas); peninsula Methana (Haussk.); Elis: pr. Lintzi (Heldr.); Laconia: pr. Kardamyle, Chimova (Chaub.); Creta: pr. Theriso, Platania, Akroteri (Raul.); Cycladum insula Rhenea (Parolini); Corcyra: pr. Spartilla (Bald.); — β. Creta: pr. Platania, Kissamos (Rev.). — Maio, Aug. ⊙

16. **L. cirrhosa** L. mant. p. 249; S. et S. pr. I. p. 431; Ch. et B. fl. pelop. p. 39 (*Antirrhinum*); Dum. cours. bot. cult. II. p. 92; Raul. cret. p. 829; Bois. fl. or. IV. p. 368. — Icon: Jacq. hort. vind. t. 82.

Inferne pubescens, superne glabrescens, multicaulis; caulibus gracillimis, prostratis, elongatis, simplicibus vel ramosis; foliis parvis, petiolatis, inferioribus ovato-oblongis, oppositis, ceteris valde diminutis, alternis, triangulari- et lineari-lanceolatis, acutis, sagittatis; pedicellis axillaribus, capillaribus, glabris, patentibus, folio multo longioribus, apice cernuis; calycis glabri laciniis lineari-lanceolatis, acutis; corolla cum calcare 4—5 mm. longa, coerulescente, glabriuscula, calcare recto, corollae aequilongo; capsula globosa, glabra. — Habitu gracillimo, foliis lanceolato-sagittatis, corolla duplo minori, capsula minuta a praecedente distincta.

In planitie pr. Korone Messeniae (Chaub.) et in vineis Cretae (Sibth.). — Jun. Aug. ⊙ N. v.

3. Sectio. *Cymbalaria* Chav. monogr. antirrh. p. 98.

a. Planta glaberrima.

17. **L. cymbalaria** L. sp. p. 612; S. et S. pr. I. p. 430; (*Antirrhinum*); Mill. dict. n. 17; Raul. cret. p. 821. — Icon: Fl. dan. t. 1220. — Exsicc.: Heldr. pl. fl. hellen. a. 1899.

Multicaulis; caulibus elongatis, filiformibus, diffusis, intricatis, ramosis; foliis longe petiolatis, cordato-reniformibus, 5—7 lobis, lobis late ovatis, obtusis vel acutiusculis, mucronatis; pedicellis axillaribus, folio aequilongis vel longioribus; calycis laciniis lineari-lanceolatis, acutis, corollae subtriplo brevioribus; corolla violacea, palato flavo, calcare subcurvato, tubo corollae breviore; capsula subglobosa; seminibus rugosis.

In muris, inter saxa. Arcadia: pr. Andrizena (Heldr.); Laconia (Sibth.); Creta: inter Platania et Canea, pr. Dramya, Retymo (Raul.); loca nonnula forsan ad sequentem pertinent. — Maio, Oct. ♃

18. **L. longipes** Bois. et Heldr. diagn. XII. p. 40, Fl. or. IV. p. 365; Heldr. in ö. b. Z. 1877 p. 157; Haussk. symb. p. 175. — *L. cymbalaria v. Sieberi* Chav. monogr. antirrh. p. 99. — *L. cym-*

balaria Friedr. Reise p. 273, non L. — Exsicc.: Heldr. pl. fl. hell. a. 1876 et 1878, herb. norm. n. 964.

Differt a praecedente foliis infimis integris, ceteris minus profunde 3—5 lobatis, lobis obtusissimis, calycis laciniis minutis, oblongis, obtusis, corollae tubo multo brevioribus, calcare corollae saepius parum breviore; sed formae intermediae, uti specimina in Heldr. herb. norm. n. 964 distributa docent, adesse videntur.

In rupibus maritimis. Attica: insula Salamis et Mikra Kyra Pharmacusarum (Heldr.); Argolis: pr. Vromolimni in peninsula Methana (Heldr.); Creta (Sieb.). — Febr. Maio. ♃

b. Planta lana longa alba crispa villosa.

19. **L. microcalyx** Bois. diagn. IV. p. 72, Fl. or. IV. p. 365; Hal. in ö. b. Z. 1897 p. 324. — *Antirrhinum pilosum* Ch. et B. exp. p. 175, Fl. pelop. p. 39, sec. Bois. l. c.; non L., quod indumento saepius breviori, foliorum lobis numerosioribus, mucronatis, calycis minuti pilosi laciniis lanceolatis, duplo longioribus, corolla minore puberula distinguitur. — Huc forsan: *L. pilosa* Spreitz. in z. b. G. 1887 p. 665, vix DC. — Orph. fl. gr. n. 712; Dörfl. fl. aeg. n. 39.

Multicaulis; caulibus diffusis, filiformibus, valde intricatis, ramosis; foliis longe petiolatis, cordato-orbiculatis, 3—5 lobis, lobis obtusissimis; pedicellis axillaribus, folio longioribus; calycis laciniis minutis, ovatis, obtusis, corollae tubo multo brevioribus; corolla glabra, violacea, palato aurantiaco, calcare recto, corolla dimidio breviore.

In fissuris rupium regionis montanae inferioris et alpinae. Elis: mt. Olenos (Heldr.); Laconia: mt. Malevo pr. Platanos (Orph.), mt. Taygetos pr. Mistra, pr. Gortys (Chaub.); Cyladum insula: Naxos in mt. Dia (Chaub.), Amorgos in mt. Hagios Elias (Leon.); Cephalonia: mt. Aenos (Griseb.); Leucas: cap Zuana (Spreitz.). — Mart. Jul. ♃

Obs. Quid sit *L. monspeliensis* Friedr. Reise p. 266 (pr. Athenas) nescio.

6. Chaenorrhinum DC. fl. fr. V. p. 410.

a. Semina costata costis muricatis.

1. **C. rubrifolium** Rob. et Cast. in DC. fl. fr. V. p. 410; Bois. fl. or. IV. p. 383 (*Linaria*); Willk. et Lange pr. fl. hisp. II. p. 578. — Icon: Mor. fl. sard. t. 101. — Exsicc.: Heldr. pl. fl. hellen. a. 1878.

Glanduloso-pilosum; caule erecto, fere a basi ramoso; foliis subtus purpurascentibus, infimis oppositis, breviter petiolatis, ovatis, ceteris alternis, sessilibus, oblongis vel oblongo-linearibus; racemis laxissimis, paucifloris; pedicellis, calyce 3—4 plo longioribus, erecto-patulis; calycis laciniis lineari-spathulatis, obtusis, capsula longioribus; corolla pubescente, violascente, calcare recto, acuto, corolla duplo breviore; capsula ovato-globosa.

In collibus aridis, saxosis regionis inferioris et montanae, rarissime. Attica: mt. Pentelicon (Sprun.), pr. Pikermi, mt. Hymettus, insula Salamis (Heldr.). — Apr. Maio. ⊙

b. Semina costata, costis laevibus.

2. **C. minus** L. sp. p. 617; S. et S. pr. I. p. 431 (*Antirrhinum*); Willk. et Lange pr. fl. hisp. II. p. 577. — *L. minor* Desf. fl. atl. II. p. 46; Bois. fl. or. IV. p. 383; Hal. in ö. b. Z. 1892 p. 375, Beitr. fl. Epir. p. 33, Beitr. fl. Thessal. p. 16, in ö. b. Z. 1897 p. 324; Form. in Ver. Brünn 1895 p. 34; Bald. riv. coll. bot. Alb. 1895 p. 59, 1896 p. 82; Haussk. symb. p. 175. — Icon: Fl. dan. t. 502. — Exsicc.: Sint. et Bornm. it. turc. n. 1389.

Glanduloso-pilosum; caule erecto, saepe a basi ramoso; foliis sessilibus, lanceolatis linearibusve, obtusiusculis, fere omnibus alternis; racemis laxissimis, paucifloris; pedicellis calyce 2—3 plo longioribus, patulis; calycis laciniis lineari-spathulatis, capsula sublongioribus; corollae pubescentis, violaceae lobis labii superioris divaricatis, calcare conico, corolla duplo breviore; capsula ovata.

In arenosis, cultis, herbidis praesertim montanis. Epirus: pr. Panagia, Safikbi, in valle Dipotami (Form.), mt. Murga, pr. Cepelovon distr. Zagorion (Bald.), mt. Tsumerka, Peristeri (Hal.); Thessalia: mt. Oxya (Hal.), ad monasterium Korona (Haussk.), Hagios Dionysios in Olympo (Sint.); Aetolia: mt. Korax (Tunt.); mt. Parnassus (Sprun.). — Jun. Sept. ⊙

3. **C. litorale** Bernh. in DC. fl. pr. V. p. 410 (*Antirrhinum*); Fritsch exc. p. 494. — *Linaria litoralis* Willd. en. hort. berol. p. 641; Haussk. symb. p. 175. — Exsicc.: Hal. it. gr. sec. a. 1893.

Praecedenti proximum, sed robustius, caulibus crassioribus, etiam pedalibus et paulo ultra, ramis magis strictis, pedunculis abbreviatis, et saepe calyce ipso brevioribus, rarius sublongioribus, corolla submajore, pallidiore, lobis labii superioribus parallelis.

In incultis, herbidis montanis. Epirus: pr. Kalarrytes (Hal.); Argolis: in peninsula Methana (Haussk.). — Jun. Sept. ⊙

7. Antirrhinum L. gen. n. 750.

a. Annua; capsula calyce brevior; semina compressa, marginata, dorso laevia, carinata, ventre sulcata et ad marginem sulci crenulata.

1. **A. orontium** L. p. p. 617; S. et. S. pr. I. p. 435; Pieri corc. fl. p. 83; Friedr. Reise p. 273; Fraas fl. class. p. 188; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 385; Heldr. fl. cephal. p. 55, chlor. Thera p. 18, chlor. Mykon. p. 248; Bald. riv. coll. bot. alb. 1896 p. 83; Haussk. symb. p. 174; Hal. in ö. b. Z. 1897 p. 97. — Icon: Fl. dan. t. 941. — Exsicc.: Sint. it. thessal. n. 675.

Glabriusculum, superne glanduloso-pilosum; caule erecto, simplici vel parce ramoso; foliis subpetiolatis, margine revolutis, inferioribus

oppositis, oblongo-lanceolatis lanceolatisve, superioribus alternis, linearibus; floribus axillaribus, in racemum spiciformem laxissimum dispositis; pedicellis brevissimis; calycis longe albo-pilosi laciniis linearibus, corollam purpuream glabriusculam aequantibus vel superantibus; capsula ovata, pilosa. — Variat foliis angustioribus et latioribus, corollae magnitudine, sed *A. calycinum* Lam. enc. IV. p. 365 = *A. orontium β. grandiflorum* Chav. mon. p. 89 t. 4 a Raul. cret. p. 821 in Creta et a Heldr. fl. Aegina p. 382 in insula Aegina indicatum, typice (corolla magna, speciosa, calycem manifeste excedente) in ditione non provenit.

In collibus aridis, rupestribus, cultis, vinetis regionis inferioris. Epirus: ad Kurenta distr. Janina (Bald.); Thessalia: pr. Malakasi (Sint.), Klinovo, Korona (Haussk.); Aetolia: mt. Chalkis ad sinum Patranum (Hal.); Attica: pr. Athenas, in colle Lycabettus, insula Aegina (Heldr.); insula Poros (Friedr.); Sporadum insula Skopelos (Leon.); Cycladum insula: Tenos (Weiss), Rhenea (Tunt.), Kimolos, Thera (Heldr.); Cephalonia: pr. Drapano, in oropedio Omala (Heldr.); Corcyra (Pieri); Creta: pr. Khalepa, Nerokuru, Kephala, Istovai (Raul.). — Mart. Jun. ⊙

b. Perennia; capsula calyce longior; semina non compressa, nec marginata, undique foveolato-cristata.

2. **A. majus** L. sp. p. 617; S. et S. pr. I. p. 434; Pieri corc. fl. p. 83; Ch. et B. exp. p. 176, Fl. pelop. p. 39; Fraas fl. class. p. 188; Raul. cret. p. 821; Bois. fl. or. IV. p. 385. — Icon: Rchb. t. 58. — Exsicc.: Heldr. et Hal. fl. aeg. a. 1889.

Basi tortuoso-ramosum, saepe lignescens, caulibus erectis, crebre foliatis, simplicibus vel parce ramosis, inferne glabris, superne pedicellis calycibusque glanduloso-pubescentibus; foliis oppositis, subternis vel alternis, oblongo-lanceolatis linearibusque, glabris; racemo densifloro, fructifero laxe elongato; pedicellis calyce brevioribus; calycis laciniis ovatis, obtusis, corollae tubo multo brevioribus; corolla violaceo-purpurea, ampla, pubescente, palato flavescente; capsula oblique ovato, glanduloso-pubescente.

β. **angustifolium** Chav. monogr. antirrh. p. 86; Bois. fl. or. IV. p. 385; Haussk. symb. p. 174. — Folia anguste linearia, margine interdum revoluta. — Exsicc.: N. v.

Ad muros vetustos, rupes regionis inferioris. Attica (Fraas); Laconia (Sibth.); in Maina (Chaub.); insula Hydra: ad muros coenobii Profetae Elias subsponte (Heldr.); Cyladum insula Tenos (Chaub.); Corcyra (Pieri); — *β.* Argolis: pr. Nauplia (Haussk.); Cycladum insula Andros (Sart.); Creta: in faucibus mt. Icari pr. Candia (Heldr.). — Mart. Sept. ♃

3. **A. siculum** Ucria pl. ad Linn. op. add. in Roem. arch. bot. I. p. 69; Raul. cret. p. 822.

Basi tortuoso-ramosum, lignescens; caulibus erectis, crebre foliatis, simplicibus vel ramosis, inferne glabris, superne pedicellis calycibusque

glanduloso-pubescentibus; foliis oppositis ternisve, lineari-lanceolatis linearibusque, glabris; racemo laxo, fructifero valde elongato; pedicellis calyce brevioribus vel subaequilongis, fructiferis elongatis, capsula subaequilongis; calycis laciniis ovato-lanceolatis, acutiusculis, corollae tubo triplo brevioribus; corolla albida vel lutescente, mediocri, pubescente; capsula oblique ovata, glanduloso-pubescente. — Praecedenti simile, ab eo racemo laxo, floribus fere dimidio minoribus, flavidis, suaveolentibus et calycis laciniis acutiusculis discedit.

In collibus saxosis Cretae: pr. Samaria (Heldr.). — Mart. Sept. ♃

Obs. *A. asarina* L. sp. p. 612, a Fraas fl. claas. p. 189 ex confusione quadam pr. Lamia indicatur. — *A. altissimum* Sieb. avis p. 5 e Creta, species ignota est. — *A. bellidifolium* L. sp. p. 617; Ch. et B. fl. pelop. p. 39; = *Anarrhinum bellidifolium* Desf. fl. atl. II. p. 51, „ex schedula mendaci Gittardi in Peloponneso erronee indicatur," cf. Bois. fl. or. IV. p. 363.

4. Tribus. **DIGITALEAE** Benth. in DC. pr. X. p. 448, sensu ampl.

8. Digitalis L. gen. n. 758.

a. Corollae tubus abrupte inflatus, subglobosus, lobo medio labii inferioris aequilongus vel sublongior.

α. Calycis laciniae oblongae, obtusae.

1. **D. ferruginea** L. sp. p. 622; S. et S. pr. I. p. 439, Fl. gr. t. 606; Fraas fl. class. p. 189; Bois. fl. or. IV. p. 429; Heldr. chlor. Parn. p. 24; Form. in D. bot. Monat. 1891 p. 26, in Ver. Brünn 1896 p. 64, 1897 p. 47; Hal. Beitr. fl. Thessal. p. 16, Beitr. fl. Achaia p. 27; Bald. riv. coll. bot. Alb. 1895 p. 59; Haussk. symb. p. 174. — Exsicc.: Orph. fl. gr. n. 40 et 1164; Heldr. herb. norm. n. 239; Sint. it. thessal. n. 1189.

Caule erecto, glabro, simplici, in racemum densiflorum vel laxiusculum abeunti; foliis oblongo-lanceolatis vel lanceolatis, integris vel obscure repandis, glabris vel ad margines et subtus parce puberulis, inferioribus in petiolum attenuatis, superioribus sessilibus; calycis glabri laciniis oblongis, obtusis, late scarioso-marginatis; corollae glanduloso-pubescentis, fulvae, ferrugineo-reticulatae lobis superioribus abbreviatis, rotundatis, lateralibus brevibus, acutis, inferiore elongato, oblongo, integro vel obscure trilobo, intus villoso.

In silvaticis regionis montanae et subalpinae. Epirus: pr. Syraku ad radices mt. Peristeri (Bald.); Thessalia: pr. Chaliki, mt. Baba (Haussk.), Oxya (Hal.), Koturi, Velitsena, Vendista, mt. Zygos, Ghavellu in Pindo, mt. Godaman in Olympo (Form.); Phthiotis: in valle fl. Sperchius (Fraas); mt. Parnassus (Sibth.); Euboea: mt. Dirphys (Heldr.); Peloponnesus: mt. Olenos, Kyllene (Hal.), Malevo (Orph.), Taygetos (Fraas). — Jul. Aug. ♃

β. Calycis laciniae lanceolatae, acutae.

2. **D. laevigata** W. et K. pl. rar. hung. II. p. 171 t. 158; Ch. et B. exp. p. 178, Fl. pelop. p. 40; Bois. fl. or. IV. p. 430; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 4, chlor. Parn. p. 24; Hal. Beitr. fl. Epir. p. 33; Bald. in nuov. giorn. bot. ital. 1894 p. 99, riv. coll. bot. Alb. 1896 p. 83; Haussk. symb. p. 174; Form. in Ver. Brünn 1897 p. 47. — Exsicc.: Orph. fl. gr. n. 41; Heldr. herb. norm. n. 445, it. thessal. n. 25; Sint. et Bornm. it. turc. n. 1382.

Caule erecto, glabro, simplici, in racemum elongatum laxum abeunti; foliis oblongo-lanceolatis vel lanceolatis, integris, glabris, inferioribus in petiolum attenuatis, superioribus sessilibus; calycis glabri laciniis lanceolatis, acutis vel acuminatis, margine subscariosis; corollae glanduloso-puberulae, fulvae, purpureo-venosae lobis superioribus et lateralibus brevibus, inferiore elongato, oblongo, repando, intus barbato.

In silvaticis regionis montanae et subalpinae. Epirus: mt. Vratedon distr. Zagorion, pr. Vutzindro (Bald.), Kalarrytes (Hal.); Thessalia: mt. Oxya, Karava, in oropedio Neuropolis (Haussk.), mt. Agrapha, Othrys (Form.), mt. Olympus, Pelion (Heldr.); Phocis: mt. Parnassus (Heldr.); Peloponnesus: mt. Kyllene, Malevo (Orph.), Taygetos (Chaub.). — Jun. Jul. ♃

3. **D. lanata** Ehrh. Beitr. VII. p. 153; Bois. fl. or. IV. p. 430; Heldr. chlor. Parn. p. 24; Hal. Beitr. fl. Epir. p. 33; Form. in Ver. Brünn 1895 p. 34; Bald. riv. coll. bot. Alb. 1896 p. 83; Haussk. symb. p. 174, *v. abbreviata* (f. inflorescentiae indumento breviori, corrollae lobo inferiore abbreviato). — Exsicc.: W. et K. pl. rar. hung. t. 74. — Exsicc.: Orph. fl. gr. n. 726 (Macedonia); Hal. it. gr. II. a. 1893.

Caule erecto, glabro, apice lanato, in racemum densum vel laxum, interdum ramulosum abeunti; foliis oblongo-lanceolatis vel lanceolatis, integris vel subrepandis, glabris vel ciliatis, inferioribus in petiolum attenuatis, superioribus sessilibus; calycis lanati laciniis lanceolatis, acutis, margine non scariosis; corollae glanduloso-villosae, pallide fusco-coerulescentis, ferrugineo-reticulatae lobis superioribus et lateralibus brevibus, inferiore elongato, ovato vel oblongo, albido.

In dumosis regionis inferioris et montanae. Epirus: pr. Prevesa (Form.), Kalentini supra Arta (Hal.), mt. Gamila distr. Zagorion (Bald.); Thessalia: pr. Karditza (Haussk.); mt. Parnassus (Sprun.). — Jun. Jul. ♃

b. Corollae tubus late campanulatus, lobis omnibus brevibus.

4. **D. ambigua** Murr. pr. stirp. gotting. p. 62; Bald. riv. coll. bot. Alb. 1895 p. 59; Form. in Ver. Brünn 1896 p. 64, 1897 p. 47; Haussk. symb. p. 174, *v. obtusioba* Neilr. fl. Niederöst. p. 547 = *D. grandiflora v. obtusiflora* Koch. syn. p. 518 (f. lobis labii inferioris obtusis). — *D. ochroleuca* Jacq. fl. austr. I. p. 36, Fl. austr. t. 57 (f. lobis labii inferioris acutis). — *D. grandiflora* Lam. fl. fr. II. p. 332; Bois. fl. or. IV. p. 432; Heldr. in Sitzungsb. acad. Wiss.

Berlin 1883 p. 4; Hal. Beitr. fl. Epir. p. 33. — Exsicc.: Orph. fl. gr. n. 754; Heldr. fl. thessal. n. 31.

Caule erecto, simplici, puberulo, superne glanduloso, in racemum laxum, secundum abeunte; foliis oblongis oblongo-lanceolatisve, denticulatis, supra glabris, subtus parce puberulis, inferioribus in petiolum attenuatis, superioribus sessilibus; calycis glanduloso-puberuli laciniis lanceolatis, acutiusculis, non scarioso-marginatis; corolla ochroleuca, glanduloso-puberula, labio superiore brevissimo, late emarginato, lobis labii inferioribus acutis vel obtusis, intermedio majore.

In silvaticis, dumosis regionis montanae et subalpinae. Epirus: mt. Strungula (Hal.); Thessalia: ad monasterium Korona, in oropedio Neuropolis in Pindo dolopico (Haussk.); mt. Oxya, Chassia (Form.), mt. Olympus (Orph.), Pelion (Heldr.). — Jun. Jul. ♃

c. Corollae tubus tenuiter cylindricus, lobis omnibus brevibus.

5. **D. lutea** L. sp. p. 622; S. et S. pr. IV. p. 439. — Icon: Jacq. hort. vindob. II. t. 105.

Caule erecto, simplici, glabro vel apice parce glanduloso-puberulo, in racemum densiusculum secundum abeunte; foliis oblongis vel oblongo-lanceolatis, denticulatis, glabris vel ciliatis, subtus interdum parce puberulis, inferioribus in petiolum attenuatis, superioribus sessilibus; bracteis inferioribus corolla, superioribus calyce brevioribus; calycis laciniis lanceolatis, acutis, emarginatis, glanduloso-ciliatis; corollae ochroleucae, glabriusculae, tubo subventricoso, lobo infimo ceteris dimidio longiore.

In Arcadia (Sibth.), sine loci specialis indicatione; a recentioribus non lecta. — Jun. Jul. ♃

6. **D. viridiflora** Lindl. mon. p. 21 t. 18. — Exsicc.: Sint. it. thessal. n. 963; Orph. fl. gr. n. 725 (mt. Corfiati Macedoniae).

Tota glanduloso-pubescens; caule erecto, simplici, in racemum densiusculum abeunte; foliis oblongis vel ovato-lanceolatis, denticulatis, inferioribus in petiolum attenuatis, superioribus sessilibus; bracteis inferioribus corolla, superioribus calyce longioribus; calycis laciniis lanceolatis, acutis, emarginatis; corollae lutescenti-virentis, viscoso-pubescentis, tubo vix ventricoso, lobis patentibus, parum inaequalibus. — Species a praecedente optime distincta.

In mt. Tragopetra pr. Malakasi Thessaliae (Sint.). — Jul. Aug. ♃

9. Sibthorpia L. gen. n. 775.

1. **S. africana** L. sp. p. 631; Raul. cret. p. 822; Bois. fl. or. IV. p. 428; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 3. — *S. europaea* S. et S. pr. I. p. 439, II. p. 359; Sieb. in Flora I. p. 276; non L., quae differt floribus minoribus, corolla carneola, calycem vix excedente, ovario cum stylo basi barbatis. — *Disandra africana* Camb. enum. pl. bal. p. 117 t. 9. — Exsicc.: Heldr. it. thessal. n. 32.

Caespitosa, hispidula; ramis teneris, filiformibus, prostratis radicantibus; foliis longe petiolatis; orbiculatis, 7—9 crenatis; pedicellis

axillaribus, petiolum aequantibus vel superantibus; calycis laciniis triangularibus, acutis; corollae flavae laciniis ovatis, calyce subduplo longioribus; antheris corollam subaequantibus; stylo glabro; capsula depresse globosa, calyce breviore.

Ad rivulos, in rupibus irrigatis, inter muscos regionis montanae, rare. Thessalia: mt. Pelion (Hawkins); Laconia: ad Hagios Petros (Bois.); Creta (Hawkins): in castanetis pr. Enneachoria (Heldr.), Vrisinas (Raul.), Serisso (Sieb.). — Jun. Sept. ♃

10. Veronica L. gen. n. 25.

Dispositio specierum:

A. Radix perennis.

1. Sectio. *Pseudolysimachium* Koch syn. p. 527. — Racemi terminales; corollae tubus latitudine sua longior.

1. V. longifolia L. **2. V. spicata** L.

2. Sectio. *Chamaedrys* Koch syn. p. 524. — Racemi axillares, oppositi vel rarius alterni; corollae tubus brevissimus.

a. Capsulae valvulae columnae placentiferae arcte cohaerentes.

α. Caulis pilosus; capsula cuneata, ovata vel orbiculata.

× Caulis elongatus.

○ Capsula basi cuneata.

. Folia parva, circa 1 cm. longa, oblonga vel linearia, integra, revoluta.

3. V. thymifolia S. et S.

.. Folia majuscula, ovata vel elliptica, dentata, margine non revoluta.

4. V. officinalis L. **5. V. chamaedrys** L.

○○ Capsula basi rotundata.

. Capsula calyce duplo longior.

6. V. latifolia L.

.. Capsula calyce aequilonga.

, Folia indivisa.

7. V. teucrium L. **8. V. austriaca** L.

,, Folia pinnatisecta.

9. V. multifida L.

×× Caulis brevissimus, fere subnullus.

10. V. aphylla L.

β. Caulis glaberrimus; capsula transverse latior.

11. V. scutellata L.

b. Capsulae valvulae demum bipartitae, altera vel utraque a columna placentifera secedeus.

α. Folia sessilia, acuta.

12. V. anagallis L. **13. V. anagalloides Guss.**

β. Folia petiolata, obtusa.

14. V. beccabunga L.

3. Sectio. *Veronicastrum* Benth. in DC. pr. X. p. 479. — Racemi terminales; corollae tubus brevissimus.

a. Caules herbacei; folia integra vel crenulata; capsula basi rotundata.

15. V. balcanica Vel. **16. V. serpyllifolia** L.

b. Caules basi lignosi; folia acute dentata; capsula cuneata.

17. V. thessalica Benth.

B. Radix annua.

4. Sectio. *Alsinebe* Griseb. spic. II. p. 23. — Flores in racemum terminalem dispositi vel axillares solitarii.

a. Semina plano-biconvexa, facie plus minus manifeste umbilicata.

α. Pedicelli calyce 3—4 plo longiores.

18. V. acinifolia L.

β. Pedicelli calyce breviores.

× Folia pinnatifida.

19. V. verna L.

×× Folia indivisa, crenata.

20. V. arvensis L. **21. V. Sartoriana** Bois. et Heldr.

b. Semina cyathiformia.

α. Folia floralia saltem superiora bracteiformia.

× Folia dentata.

○ Pedicelli calyce 2 — multo longiores.

. Planta crispule et breviter pubescens.

22. V. glauca S. et S. **23. V. Chaubardi** Bois. et Reut.

. . Planta patule glanduloso-pilosa.

24. V. peloponnesiaca Bois. et Orph.

○○ Pedicelli calyce breviores vel parum longiores.

25. V. praecox All.

×× Folia palmatisecta.

26. V. triphyllos L.

β. Folia floralia omnia caulinis conformia.

× Folia ovata, crenato-dentata; capsulae loculi 4—10 spermi; semina minora ovalia.

○ Capsula elevatim reticulato-venosa, lobis divaricatis.

27. V. persica Poir.

○○ Capsula laevis vel tenuiter reticulata, lobis non divaricatis.

28. V. agrestis L. **29. V. didyma** Ten.

×× Folia suborbiculata, 3—9 loba; capsulae loculi 1—2 spermi; semina magna, subglobosa.

30. V. cymbalaria Bodard. **31. V. hederifolia** L.

A. Radix perennis.

1. Sectio. *Pseudolysimachium* Koch syn. p. 527. — Racemi terminales; corollae tubus latitudine sua longior.

1. **V. longifolia** L. sp. p. 10; Bois. fl. or. IV. p. 455. — *V. spuria* S. et S. pr. I. p. 5, non L. (teste Bois. l. c.). — Icon: Rchb. t. 93. —

Breviter pubescens; caulibus elatis, strictis, racemis terminalibus, densiusculis, paniculatis; foliis breviter petiolatis, oppositis, ternis quaternisve, a basi ovata vel cordata lanceolatis, acuminatis, argute duplicato-

serratis; calycis laciniis acutis; corolla coerulea; capsula obcordato-rotundata, turgida glabra.

In locis humidiusculis Arcadiae (Bois.) et Laconiae (Sibth.); a recentioribus non lecta et ulterius inquirenda. — Jul. Aug. ♃ N. v.

2. **V. spicata** L. sp. p. 10; S. et S. pr. I. p. 6; Pieri corc. fl. p. 2; Ch. et B. exp. p. 15, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 444; Bois. fl. or. IV. p. 455. — ? *V. australis* Schrad. comm. p. 24; Mazz. l. c. p. 442. — Icon: Fl. dan. t. 52. —

Plus minus tomentella, canescens; caulibus erectis vel adscendentibus, racemis solitariis vel pluribus, densis; foliis oppositis, petiolatis, crenatis, apice integris, inferioribus ovatis, obtusis, superioribus lanceolatis; calycis laciniis oblongis, obtusiusculis; corolla coerulea; capsula obcordato-rotundata, turgida, puberula.

In pascuis siccis. Laconia (Sibth.); Messenia: inter Gargaliano et Philatra (Chaub.); Corcyra (Pieri): pr. Afiona et Messongi (Mazz.); a recentioribus non lecta. — Jul. Aug. ♃ N. v.

2. Sectio. *Chamaedrys* Koch syn. p. 524. — Racemi axillares, oppositi vel rarius alterni; corollae tubus brevissimus.

a. Capsulae valvulae columnae placentiferae arcte cohaerentes.

α. Caulis pilosus, capsula cuneata, ovata vel orbiculata.

× Caulis elongatus.

○ Capsula basi cuneata.

. Folia parva, circa 1 cm. longa, oblonga vel linearia, integra, revoluta.

3. **V. thymifolia** S. et S. pr. I. p. 6, Fl. gr. I p. 5 t. 6; Sieb. in Flora I. p. 275; Raul. cret. p. 823; Bois. fl. or. IV. p. 444; Heldr. chlor. Parn. p. 24; Hal. Beitr. fl. Achaia p. 27; Bald. viagg. Creta p. 81. — *V. cretica* Pall. in Link Jahrb. I. p. 41; Raul. l. c. — *V. tymphrestea* Bois. et Sprun. diag. IV. p. 77; cf. Hal. l. c. — Exsicc.: Heldr. pl. cret. n. 1488; Hal. it gr. II. a. 1893.

Breviter et crispule velutina, cinerea, caespitosa, basi suffrutescens; caudiculis prostratis, radicantibus, ramis adscendentibus, crebre foliatis; foliis sessilibus, lineari-vel rarius oblongo-spathulatis, integris, margine revolutis; racemis breviter pedunculatis, dense capitatis, 1—2 in axillis supremis, interdum gemmae terminalis abortu pseudo-terminalibus; pedicellis calyce brevioribus; calycis laciniis quaternis, oblongis, subinaequalibus; corolla rosea vel coerulea, calyce parum longiore; capsula obtriangulari, hirta, calyce longiore. — Species egregia, rara.

In rupestribus regionis alpinae. ? Aetolia: mt. Tymphrestus (Sprun.); mt. Parnassus (Sprun.); Achaia: in cacumine mt. Kyllene in pulvinaribus *Astragali angustifolii* caespites fere inextricabiles formans (Hal.); Creta: inter Prosnero et Askyphos (Raul.) et mt. Hagion Pneuma (Bald.) montium Sphacioticorum (Sibth.), mt. Ida, Lassiti (Heldr.). — Maio, Jul. ♃

.. Folia majuscula, ovata vel elliptica, dentata, margine non revoluta.

4. **V. officinalis** L sp. p. 11; Dallap. prosp. p. 5, sed. sec. Heldr. fl. cephal. p. 56 confusa; Form. in D. bot. Monat. 1891 p. 26, in Ver. Brünn 1897 p. 48 v. *glandulosa*; Haussk. symb. p. 175. — Icon: Fl. dan. t. 248. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Pubescens, in inflorescentia plus minus glandulosa; caulibus prostratis, radicantibus, circumcirca pilosis; foliis ovatis vel ellipticis, serratis, breviter petiolatis; racemis ex axillis superioribus ortis, densiusculis; pedicellis calyce brevioribus; calycis laciniis quaternis, lanceolatis, subinaequalibus; corolla pallide coerulea, calyce sublongiore; capsula obtriangulari, glandulosa, calyce dimidio longiore.

In silvaticis montanis et subalpinis. Thessalia: mt. Zygos supra Metzovo (Heldr.), pr. Vlachava, Hepdominta Aderfia in mt. Oxya (Form.), in oropedio Neuropolis (Haussk.); ? Cephalonia (Dallap.). — Jun. Jul. ♃

5. **V. chamaedrys** L. sp. p. 13; S. et S. pr. I. p. 8; Mazz. in ant. ion. II. p. 442; Friedr. Reise p. 280; Bois. fl. or. IV. p. 446; Gelmi in bull. soc. bot. ital. p. 451; Form. in D. bot. Monat. 1890 p. 26, in Ver. Brünn 1895 p. 34, 1896 p. 65, 1897 p. 48; Haussk. symb. p. 175 v. *pindica* (f. racemis densis, breviter pedunculatis, capsulis brevius et crasse pedicellatis, calycis laciniis brevioribus). — *V. chamaedryoides* Ch. et B. exp. p. 15 t. 1, Fl. pelop. p. 1 t. 1; Spreitz in z. b. G. 1887 p. 665; (f. caulibus undique pilosis; nota in speciminibus unius et ejusdem loci saepe inconstans); = *V. chamaedrys v. pilosa* Benth. in DC. pr. X. p. 475; Bois. fl. or. IV. p. 447; Heldr. chlor. Parn. p. 24. — Exsicc.: Orph. fl. gr. n. 717; Heldr. herb. norm. n. 287, 697 et 1568; Sint. it. thessal. n. 351; Dörfl. fl. gr. 334.

Pubescens; caulibus adscendentibus, basi prostrata radicantibus, bifariam vel superne circumcirca pilosis; foliis ovatis vel cordato-ovatis, inciso-crenato-serratis, inferioribus breviter petiolatis, superioribus sessilibus; racemis ex axillis supremis ortis, laxis; pedicellis calyce sublongioribus; calycis laciniis quaternis, lanceolatis, subinaequalibus; corolla calyce subduplo longiore, coerulea, lobo inferiore albido; capsula obtriangulari, hirta, calyce breviore.

In dumosis montanis et subalpinis. Epirus: mt. Prosgoli (Bald.); Thessalia: pr. Vitomo in Pindo tymphaeo (Sint.), mt. Karava, Ghavellu, monasterium Korona (Haussk.), Klinovo, Mandra Hodza, mt. Agrapha, Oxya, Chassia (Form.), mt. Olympus, Pelion, mt. Telethrion et Dirphys Euboeae (Heldr.); Aetolia: mt. Arapocephala (Heldr.), Korax (Leon.), mt. Parnassus (Heldr.); Achaia (Bois.); Arcadia: mt. Diaforti pr. Andrizena (Friedr.); Messenia: pr. Phigalea (Chaub.); Laconia (Sibth.): pr. Hagios Petros (Orph.), Megali Anastasova (Zahn), mt. Taygetos (Chaub.); Corcyra: pr. Castagna (Mazz.), mt. Deca (Spreitz.). — Apr. Maio. ♃

○○ Capsula basi rotundata.

. Capsula calyce duplo longior.

6. **V. latifolia** L. sp. p. 13. — *V. urticaefolia* Jacq. fl. austr. I. p. 37 t. 59; Bois. fl. or. IV. p. 448.

Parce hirta; caulibus erectis; foliis sessilibus, ovatis, argute serratis, superioribus acuminatis, basi subcordatis; racemis axillaribus, oppositis, laxis; pedicellis erecto-patulis, calyce 3—4 plo longioribus; calycis laciniis quaternis, minutis, lanceolatis, obtusiusculis; corolla griseo-coerulea, calyce longiore; capsula glabriuscula, orbiculari, calyce duplo longiore.

In silvaticis subalpinis mt. Olympus Thessaliae (Heldr.). — Jun. Jul. ♃ N. v.

.. Capsula calyce aequilonga.

, Folia indivisa.

7. **V. teucrium** L. sp. ed 2 p. 16; Kern. in ö. b. Z. 1873 p. 371. — *V latifolia* Jacq. obs. I. p. 41; Ch. et B. exp. p. 15, Fl. pelop. p. 1; Bois. fl. or. IV. p. 449; non L. — *V. pseudochamaedrys* Jacq. fl. austr. I. p. 38 t. 60. —

Pubescens; caulibus erectis, elatis; foliis sessilibus, ovatis vel ovato-oblongis, inciso-crenatis; racemis 2—4 ex axillis supremis ortis, tandem elongatis; pedicellis strictis, calyce longioribus; calycis laciniis quinis, lanceolatis, inaequalibus; corolla coerulea; capsula puberula, obovato-obcordata, calyce vix longiore.

β. **glandulosa** Form. in Ver. Brünn 1896 p. 64. — Inflorescentia glanduloso-hirta.

In montosis pr. Skardamula Laconiae rare (Chaub.); — *β.* mt. Dokimi in Pindo (Form.) — Maio, Jun. ♃ N. v.

8. **V. austriaca** L. sp. ed 2 p. 17; Kern. in ö. b. Z. 1873 p. 372; Bald. riv. coll. bot. Alb. 1896 p. 84. — *V. dentata* Schmidt fl. boem. I. p. 20; Form. in Ver. Brünn 1896 p. 65. — *V. teucrium* Jacq. obs. I. p. 41; Ch. et B. fl. pelop. p. 1; Bois. fl. or. IV. p. 448; Heldr. chlor. Parn. p. 24; Haussk. symb. p. 175. — *V. Schmidtii* Roem. et Schult. syst. I. p. 115. — Exsicc.: Sint. it. thessal. n. 631; Bald. it. alb. epir. IV. n. 193.

Pubescens; caulibus adscendentibus; foliis oblongo- vel lineari-lanceolatis, obtuse inciso-crenatis vel remote dentatis, infimis subpetiolatis, ceteris sessilibus; racemis 2—4, ex axillis supremis ortis, tandem elongatis; pedicellis strictis, calyce aequilongis vel longioribus; calycis laciniis quinis rarius quaternis, oblongo-linearibus, inaequalibus; corolla coerulea; capsula puberula, obovato-obcordata, calyce subaequilonga. — A praecedente foliis angustis distincta.

β. **teucrioides** Bois. et Heldr. diagn. ser. 2 III. p. 169 pro sp. — *V. prostrata* v. *filicaulis* Hal. in z. b. G. 1888 p. 761. — Caulibus tenuiter filiformibus, diffusis, basi procumbentibus; racemis solitariis binisque; calycis laciniis quaternis. — Sine dubio huc, nec ad *V thymifoliam,* ut vult Bois. in fl. or. IV. p. 441, pertinet. — Exsicc.: Heldr. (pl. fl. hellen. n. 2505 et 3312.

In herbidis rupestribus regionis montanae et alpinae. Epirus: mt. Gamila distr. Zagorion (Bald.); Thessalia: mt. Said Pascha, Dokimi Form.), Ghavellu, Karava (Haussk.); Peloponnesus: pr. Messene (Chaub.).

— β. Aetolia: mt. Tymphrestus (Samar.); mt. Kiona (Hal.), mt. Parnassus et mt. Olympus (Held.). — Jun. Aug. ♃

„ Folia pinnatisecta.

9. **V. multifida** L. sp. p. 13; Jacq. austr. t. 327; Mazz. in ant. ion. II. p. 442; Kern. in ö. b. Z. 1873 p. 373; Bald. riv. coll. bot. alb. 1896 p. 84. — *V. Jacquini* Baumg. en. trans. I. p. 26; Schott in Roem. et Schult. syst. I. p. 108. — *V. austriaca* Bois. fl. or. IV. p. 449; non L. — Exsicc.: Bald. it. alb. a. 1892 n. 292.

Pubescens; caulibus erectis vel adscendentibus; foliis sessilibus, ambitu ovatis, in lacinias oblongas vel lineares, integras vel incisas pinnatisectis; racemis 1—4, ex axillis supremis ortis, tandem elongatis; pedicellis strictis, calyce longioribus; calycis laciniis quinis, rarius quaternis, linearibus, inaequalibus; corolla coerulea; capsula glabra vel puberula, obovato-obcordata vel orbiculata, emarginata, calyce aequilonga vel breviore. — Foliis pinnatisectis insignis.

In herbidis, saxosis montanis. Epirus: mt. Olycika distr. Janina, mt. Sasica distr. Vallona (Bald.); Aetolia: mt. Tymphrestus (Sprun.); Corcyra: pr. Spartilla (Mazz.). — Maio, Jul. ♃

×× Caulis brevissimus, fere subnullus.

10. **V. aphylla** L. sp. p. 11; Bois. fl. or. IV. p. 450; Hal. in z. b. G. 1888 p. 761; Heldr. chlor. Parn. p. 24. — *V. depauperata* W. et K. pl. rar. hung. III. p. 272 t. 245. — Exsicc.: Hal. it. gr. a. 1888.

Pubescens, superne glandulosa, caespitosa; caudiculis tenuibus, radicantibus; caulibus brevissimis, rosulatim foliosis; foliis ovatis vel rotundatis, integris vel crenatis, breviter petiolatis; racemis ex axilla superiore longe pedunculatis, corymbose 3—5 floris; pedicellis calyce longioribus; calycis laciniis quaternis, oblongis; corolla coerulea; capsula glandulosa, ovato-obcordata, calyce duplo longiore. — Plantula 3—8 cm. alta; habitu egregia.

In rupium fissuris regionis alpinae, rarissime. Doris: mt. Kiona (Hal.), mt. Parnassus (Heldr.). — Jun. Aug. ♃

β. Caulis glaberrimus; capsula transverse latior.

11. **V. scutellata** L. sp. p. 12; Mazz. in. ant. ion. II. p. 444. — Icon: Fl. dan. t. 209.

Glaberrima; caulibus tenuibus, debilibus, basi radicantibus; foliis lineari-lanceolatis, integris vel remote denticulatis, acuminatis, basi semiamplexicauli sessilibus; racemis axillaribus, plerumque alternis, longe pedunculatis, laxifloris; pedicellis calyce 3—5plo longioribus, patulis vel reflexis; calycis laciniis quaternis, ellipticis, acutiusculis; corolla parva, coerulescente; capsula obreniformi, subdidyma, calyce longiore.

In locis humidiusculis. Corcyra: ad Rodostamo pr. Castagna (Mazz.). — Jun. Sept. ♃ N. v.

b. Capsulae valvulae demum bipartitae, altera vel utraque a columna placentifera secedens.

α. Folia sessilia, acuta.

12. **V. anagallis** L. sp. p. 12; S. et S. pr. I. p. 7; Pieri corc. fl. p. 4; Sieb. avis rem. p. 2; Ch. et B. exp. p. 15; Mazz. in ant. ion. p. 442; Marg. et R. fl. Zante p. 71; Friedr. Reise p. 272; Weiss in z. b. G. 1896 p. 744; Raul. cret. p. 823; Bois. fl. or. IV. p. 437; Heldr. fl. cephal. p. 55, Fl. Aegina p. 382; Gelmi in bull. soc. bot. it. 1889 p. 451; Form. in D. bot. Mon. 1891 p. 26, in Ver. Brünn 1896 p. 64, 1897 p. 47; Hal. Beitr. fl. Epir. p. 33; Bald. riv. coll. bot. Alb. 1895 p. 59, 1896 p. 83; Haussk. symb. 175. — Icon: Fl. dan. t. 903. — Exsicc.: Heldr. herb. norm. n. 1065.

Glabra; caulibus erectis, fistulosis; foliis sessilibus, integris vel serratis, a basi saepius cordato-amplexicauli ovatis vel lanceolatis; racemis axillaribus, oppositis, elongatis, laxifloris; pedicellis erecto-patulis, calyce longioribus; calycis laciniis quaternis, ellipticis, acutis; corolla pallide coerulea, parva; capsula orbiculata vel ovata, calyce saepius breviore. — Caules saepius crassi, usque 1 m. alti, capsulae $2^1/_2$—$3^1/_3$ mm. longae, saepius orbiculatae et calyce breviores, occurrunt tamen interdum in eodem specimine capsulae ovatae calyce breviores.

β. **anagalliformis** Bor. fl. centr. ed. 3 II. p. 489. Inflorescentia capsulisque glandulosa. — Exsicc.: Heldr. pl. fl. hellen. a. 1886, 1887, 1888, 1898, 1899; Dörfl. pl. cret. n. 7.

In humidis, ad fossas, rivos per totam Graeciam. — Apr. Aug. ♃

13. **V. anagalloides** Guss. pl. rar. I. p. 5 t. 3; Ch. et B. fl. pelop. p. 1; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 437; Bald. riv. coll. bot. Alb. 1896 p. 83; Haussk. symb. p. 175. — Huc forsan: *V. graminea* Mazz. in ant. ion. II. p. 442. — Exsicc.: Heldr. herb. norm. n. 1066; Bald. it. alb. epir. IV. n. 88.

Differt a praecedente caulibus subsolidis, racemis numerosioribus, saepe compositis, capsula minore 2—$2^1/_2$ mm. longa, elliptica, latitudine fere duplo longiore, calyce saepissime longiore. Inflorescentia ut videtur semper glandulosa.

In iisdem locis quibus praecedens et saepe promiscue cum ea, ubi tunc formae intermediae (hybridae?) occurrere videntur. Epirus: pr. Kastrica, Lapsista (Bald.); Thessalia: pr. Karditza, Sophades (Haussk.); pr. Kastaniotissa Euboeae, ad rivulum Podoniphti Cephissi affluentem et ad Phalerum Atticae, in Boeotia (Heldr.); Messenia: mt. Kupe (Chaub.); insula Tenos (Weiss); Corcyra: ad Porta reale (Mazz.); et certe alibi, sed saepe cum praecedente confusa. — Apr. Aug. ♃.

β. Folia petiolata, obtusa.

14. **V. beccabunga** L. sp. p. 12; S. et S. pr. I. p. 7; Dallap. prosp. p. 5; Ch. et B. exp. p. 15, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 442; Clem. sert. p. 72; Bois. fl. or. IV. p. 438; Heldr. fl. cephal. p. 55; Hal. Beitr. fl. Epir. p. 33; Bald. riv. coll. bot. alb. 1896 p. 83. — Exsicc.: Heldr. fl. thessal. a. 1883; Hal. it. gr. secund. a. 1893.

Glabra; caulibus basi radicantibus, fistulosis; foliis breviter petiolatis, oblongo-ellipticis, crenato-serratis; racemis axillaribus; saepius oppositis, laxifloris; pedicellis patulis, calyce longioribus; calycis laciniis quaternis, ovato-lanceolatis, acutis; corolla coerulea; capsula orbiculata, calyce aequilonga.

Ad aquas regionis submontanae in alpinam adscendens. Epirus: mt. Papignon (Bald.), Peristeri (Hal.); Thessalia: mt. Pelion pr. Portaria (Heldr.); Attica: pr. Athenas, mt. Pentelicon (Clem.), pr. Chelidonu (Heldr.); Achaia: mt. Panachaicon, Kyllene (Hal.); Arcadia, Elis (Sibth.); Messenia (Chaub.); Cephalonia (Dallap.); Corcyra: pr. Chrisidas, Canale (Mazz.). — Apr. Jul. ♃

3. Sectio. *Veronicastrum* Benth. in DC. pr. X. p. 479. — Racemi terminales; corollae tubus brevissimus.

a. Caules herbacei; folia integra vel crenulata; capsula basi rotundata.

15. **V. balcanica** Vel. fl. bulg. p. 431; Hal. Beitr. fl. Epir. p. 33; Bald. riv. coll. bot. alb. 1895 p. 59, 1896 p. 84. — *V. serpyllifolia v. glandulosa* Haussk. symb. p. 175, non Wirtg. fl. Rheinprov. p. 333. — Exsicc.: Hal. it. gr. sec. a. 1893; Bald. it. alb. epir. IV. n. 194; Sint. it. thessal. n. 676.

Caulibus pumilis, tenuibus glanduloso-pilosis, basi radicantibus; foliis glabris vel superioribus glanduloso-puberulis, integris, inferioribus suborbiculatis vel ovatis, breviter petiolatis, superioribus ellipticis, sessilibus subsessilibusve; racemo terminali, sub 10 floro, laxo, dense glanduloso, demum elongato; pedicellis calyce 3—4plo longioribus; calycis laciniis oblongis, corolla azurea multo brevioribus; capsula orbiculato-obcordata, glandulosa, calyce sublongiore. — Nana, 3—6 cm. alta, sequenti affinis, sed ab ea meo sensu, racemo paucifloro dense glanduloso-piloso, calyce minore et praesertim corolla majore, 5—7 mm. diametro, saturate azurea specifice differt.

Ad rivulos, in scaturiginosis regionis subalpinae et alpinae. Epirus: mt. Papignon, Gamila, Smolika, Tsumerka (Bald.), Peristeri (Hal.), Turnara pr. Chaliki (Sint.); Thessalia: mt. Karava, Ghavellu, in oropedio Neuropolis (Haussk.). — Jun. Jul. ♃

16. **V. serpyllifolia** L. sp. p. 12; Mazz. in ant. ion. II. p. 444; Form. in Ver. Brünn 1897 p. 48, *v. oxya*, quae ex descriptione a typo non differre videtur.

Caulibus saepius elongatis, basi radicantibus, minute puberulis, superne saepe parce glandulosis; foliis glabris vel parce puberulis, integris vel crenulatis, inferioribus breviter petiolatis, ovatis, superioribus oblongis oblongo-lanceolatisve; racemo terminali, multifloro, laxo, demum valde elongato; pedicellis calyce 2—3 plo longioribus; calycis laciniis oblongis, corolla albida vel pallide coerulea parum brevioribus; capsula orbiculato-obcordata, parce glandulosa, calyce sublongiore.

In umbrosis praesertim montanis. Indicatur in Corcyra: pr. Viro, Castagna, Missostrati, Combizzi (Mazz.); Thessalia: ad Hepdominta Aderfia in mt. Oxya (Form.). — Maio, Jul. ♃ N. v.

b. Caules basi lignosi, folia acute dentata; capsula cuneata.

17. **V. thessalica** Benth. in DC. pr. X. p. 480; Bois. fl. or. IV. p. 453. — *V. erinoides* Bois. et Sprun. diagn. IV. p. 78. — Exsicc.: Heldr. pl. fl. hellen. a. 1887.

Pilis crispulis parce velutina, superne glandulosa, caespitosa; caudiculis prostratis radicantibus, ramis adscendentibus, crebriuscule foliatis; foliis sessilibus, ovato-rhombeis, acute dentatis; floribus 3—5, ad ramorum apicem approximatis; pedicellis calyce brevioribus; calycis laciniis oblongis, corolla violacea duplo brevioribus; capsula obtriangulari, glandulosa, calyce sublongiore. — Species egregia, caulibus nanis, floribus subcapitatis, corolla et capsula majuscula.

In rupestribus regionis alpinae, rarissime. Thessalia: mt. Olympus (Auch.); Phocis: mt. Parnassus (Sprun.); Achaia: mt. Kyllene, ad cacumen (Heldr.). — Jun. Jul. ♃

B. Radix annua.

4. Sectio. *Alsinebe* Griseb. spic. II. p. 23. — Flores in racemum terminalem dispositi vel axillares solitarii.

a. Semina plano-biconnexa, facie plus minus manifeste umbilicata.

α. Pedicelli calyce 3—4plo longiores.

18. **V. acinifolia** L. sp. ed. 2 p. 19; Pieri corc. fl. p. 3; Mazz. in ant. ion. II. p. 442; Ch. et B. fl. pelop. p. 1; Raul. cret. p. 823; Bois. fl. or. IV. p. 458. — Icon: Rchb. t. 38. — Exsicc.: Heldr. fl. thessala a. 1883.

Glanduloso-hirta, pumila, simplex vel a basi ramosa, caulibus lateralibus adscendentibus; foliis ovatis, subcrenatis, inferioribus petiolatis, superioribus sessilibus; racemo tandem elongato, laxo; calycis laciniis, ovali-ellipticis, corolla coerulea subbrevioribus; capsula obreniforme, profunde emarginato-biloba, glanduloso-ciliata, calyce submajore; seminibus ovatis.

In cultis, herbidis, praesertim montanis et subalpinis. Thessalia: mt. Pelion pr. Portaria (Heldr.); Attica: in cacumine mt. Parnes (Heldr.); pr. Messene (Gittard); Creta: pr. Nerokuru (Raul.); Corcyra (Pieri): ad muros urbis (Mazz.). — Apr. Maio. ⊙

β. Pedicelli calyce breviores.

× Folia pinnatifida.

19. **V. verna** L. sp. p. 14; Pieri corc. fl. p. 3; Bois. fl. or. IV. p. 456; Form. in Ver. Brünn 1897 p. 48. — Icon: Fl. dan. t. 252. — Exsicc.: Orph. herb. n. 2971; Hal. it. gr. II. a. 1893.

Erecta, inferne pubescens, superne glanduloso-puberula; caule simplici vel ramoso, ramis erectis; foliis ovatis, inferioribus breviter petio-

latis, crenatis; ceteris sessilibus, profunde 3—7 sectis, floralibus lanceolato-linearibus, integris; pedicello longioribus; racemis elongatis, densifloris vel demum laxiusculis; calycis laciniis lineari-lanceolatis, corollam parvam coeruleam superantibus; capsula glandulosa, obcordata, biloba; seminibus ovatis, minute umbilicatis.

In herbidis, rupestribus regionis montanae et alpinae. Corcyra (Pieri); Epirus: in cacumine mt. Peristeri (Hal.); Thessalia: ad Hepdominta Aderfia in mt. Oxya (Hal.); Laconia: in cacumine mt. Malevo (Orph.). — Apr. Jul. ⊙

×× Folia indivisa, crenata.

20. **V. arvensis** L. sp. p. 13; S. et S. pr. I. p. 9; Pieri corc. fl. 3; Ch. et B. exp. p. 16, Fl. pelop. p. 1; Mazz. in ant. ion. p. 442; Friedr. Reise p. 266; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 823; Bois. fl. or. IV. p. 457; Heldr. fl. cephal. p. 55, Fl. Aegina p. 382; Hal. Beitr. fl. Epir. p. 34, Beitr. fl. Achaia p. 28, in ö. b. Z. 1897 p. 97; Form. in Ver. Brünn 1896 p. 65, 1897 p. 48; Haussk. symb. p. 176. — Icon: Fl. dan. t. 515. — Exsicc.: Heldr. herb. norm. n. 747 et 1569; Dörfl. fl. gr. n. 377.

Erecta vel adscendens, pilosiuscula, superne saepius glandulosopuberula; caule simplici vel ramoso, ramis crenato-adscendentibus; foliis ovatis, crenatis, inferioribus breviter petiolatis, superioribus sessilibus, floralibus lanceolato-linearibus, integris, pedicello multo longioribus; racemis demum elongatis, laxiusculis; calycis laciniis lineari-lanceolatis, corollam parvam, coeruleam superantibus; capsula glandulosa, obcordata, biloba; seminibus ovatis, minute umbilicatis.

In herbidis, rupestribus praesertim montanis, in alpinam usque adscendens. Epirus: mt. Strungula, Peristeri (Hal.); Thessalia: pr. Klinovo, Vlachava (Form.), mt. Karava, pr. Velestinos (Heldr.), Pharsalus, Orman Magula (Haussk.); Sporadum insula Peristeri (Leon.); Aetolia: mt. Korax (Leon.); mt. Parnassus pr. Rachova (Heldr.); Attica: ad Ilyssum (Friedr.), pr. Athenas, mt. Parnes (Heldr.), pr. Eleusis, Geraka (Haussk.); insula Aegina (Heldr.), Melos (Leon.); Peloponnesus (Sibth.): mt. Panachaicon pr. Patras (Hal.), pr. Nauplia Argolidis (Haussk.), pr. Gargaliano (Chaub.) et Kalamata (Zahn) Messeniae; Creta: pr. Canea (Weiss); Cephalonia: pr. Francata (Heldr.) Corcyra: pr. Botumia (Mazz.). — Apr. Jul. ⊙

21. **V. Sartoriana** Bois. et Heldr. diagn. ser. 2 III. p. 171, Fl. or. IV. p. 457; Heldr. chlor. Parn. p. 24. — Exsicc.: Heldr. herb. norm. n. 748.

Praecedenti maxime affinis, ab ea racemis paucifloris (5—9), foliis floralibus pedicello aequilongis vel sublongioribus, pedicellis calyce aequilongis vel sublongioribus discedit. — Semina ovata vel suborbiculata, uti in praecedente.

In pascuis saxosis mt. Parnes Atticae et mt. Parnassus pr. Rachova in consortio *V. arvensis*, rara (Heldr.). — Maio, Jun. ⊙

cyathiformia.

α. Folia floralia saltem superiora bracteiformia.

× Folia dentata.

○ Pedicelli calyce 2—multo longiores.

. Planta crispule et breviter pubescens.

22. V. **glauca** S. et S. pr. I. p. 9, Fl. gr. I. p. 6 t. 7; Friedr. Reise p. 267; Clem. sert. p. 72; Bois. fl. or. IV. p. 461; Haussk. symb. p. 176. — *V. graeca* Sprun. in schedis. — Exsicc.: Orph. fl. gr. n. 45; Heldr. herb. norm. n. 400 sub *V. amoena* (non Stev. quae indumento glanduloso discedit) et 1163, herb. fl. hellen. a. 1889, in Baen. herb. europ. n. 3612; Rev. pl. cret. n. 258; Dörfl. fl. gr. n. 466.

Glauca, patentim ramosa, ramis adscendentibus; foliis ovatis, dentatis, inferioribus petiolatis, superioribus sessilibus, floralibus diminutis, praeter infimas dentatas integris, linearibus; racemis laxis; pedicellis filiformibus, calyce et bractea 3—5 plo longioribus, fructiferis patentibus, apice incurvis; calycis laciniis oblongo-linearibus, corolla ampla, coerulea, basi albido-maculata triplo brevioribus; capsula orbiculata, turgida, glabra, truncato-subretusa, calyce subbreviore. — Corolla magna, ad 15 mm. diametro, pulcherrime azurea insignis.

In campis, arvis Atticae ubique circa Athenas in montibus ad 1000 m. adscendens. — Febr. Apr. ⊙

23. V. **Chaubardi** Bois. et Reut. diagn. ser. 2 III. p. 174, Fl. or. IV. p. 461; Haussk. symb. p. 176. — *V. peduncularis* Ch. et B. exp. p. 15, Fl. pelop. p. 1; non M. a B., quae planta perennis, diversissima, ex affinitate *V. chamaedrys*.

Praecedenti affinis, sed non glauca, pilis longioribus obducta, pedicelli arcuato-deflexi, tandem longissimi capsula fere ad medium usque biloba, semina majora.

In herbidis, arenosis. Messenia: a Phigalea usque Pylos (Chaub.), pr. Methone, Kalamata, Armyros (Bois.); indicatur quoque a Haussknecht in mt. Hymettus Atticae. — Febr. Apr. ⊙ N. v.

. . Planta patule glanduloso-pilosa.

24. V. **peloponnesiaca** Bois. et Orph. fl. or. IV. p. 462; Heldr. fl. cephal. p. 55; Gelmi in bull. soc. bot. it. 1889 p. 451; Hal. Beitr. fl Achaia p. 28; Bald. riv. coll. bot. alb. 1895 p. 60, 1896 p. 84, Boissieu in bull. soc. bot. Fr. 1896 p. 287; Haussk. symb. p. 176. — *V. glauca* Ung. Reise p. 129, non S. et S. — *V. Chaubardi* Spreitz. in z. b. G. 1879 p. 716, non Bois. et Reut. — Exsicc.: Orph. fl. gr. n. 716; Bald. it. alb. epir. III. n. 77, IV. n. 89; Sint. it. thessal. n. 31.

Subsimplex vel saepius patentim ramosa, ramis adscendentibus; foliis ovatis, dentatis, inferioribus petiolatis, superioribus sessilibus, floralibus diminutis, praeter infimas dentatas integris, linearibus; racemis laxis; pedicellis filiformibus, calyce et bractea 2—4 plo longioribus, fructiferis recurvis; calycis laciniis oblongo-linearibus, corolla coerulea 2—3 plo brevioribus; capsula orbiculata, turgida, glandulosa, truncato-

subretusa, calyce subbreviore. — Duabus praecedentibus affinis, ab eis differt indumento glanduloso et ut videtur floribus minoribus, a *V. Chaubardi* insuper capsula non biloba.

In arvis, rupestribus regionis montanae et subalpinae. Epirus: mt. Micikeli, Ciuka, Olycika distr. Janina (Bald.); Thessalia: pr. Kalabaka (Boissieu); mt. Helicon (Orph.); Achaia: mt. Panachaicon, Kastro pr. Kalavryta (Hal.); Argolis (Sprun.): mt. Palamidi pr. Nauplia (Haussk.): Laconia: mt. Malevo supra Hagios Joannes (Orph.); Cephalonia: mt. Aenos (Heldr.); Corcyra: mt. Deca (Spreitz.) — Mart. Jun. ⊙

○○ Pedicelli calyce breviores vel parum longiores.

25. **V. praecox** All. auct. ad fl. ped. p. 5 t. 1; Mazz. in ant. ion. II. p. 444; Raul. cret. p. 823; Bois. fl. or. IV. p. 463. — *V. thessala* Form. in D. bot. Monat. 1897 p. 197, in Ver. Brünn 1897 p. 48, = *V. Formanekii* Heldr. in D. bot. Monat. 1897 p. 321, quae ex descriptione nullo modo discrepat. —

Glanduloso-pilosa, simplex vel saepius a basi ramosa, ramis adscendentibus; foliis ovatis, dentatis, inferioribus breviter petiolatis, superioribus sessilibus, floralibus praeter infimas dentatas integris, linearibus; racemis laxis; pedicellis crassiusculis, calyce et bractea brevioribus vel sublongioribus, fructiferis erecto-patulis; calycis laciniis oblongo-linearibus, corolla coerulea subbrevioribus; capsula suborbiculata, turgidula, glandulosa, emarginata, calyce breviore.

In herbidis, rupestribus montanis. Corcyra: pr. Canale, Evangelistra (Mazz.); Thessalia: pr. Konisko, Asproklisia (Form.); Creta: mt. Ida, Lazaro in mt. Lassiti (Raul.). — Apr. Maio. ⊙ N. v.

×× Folia palmatisecta.

26. **V. triphyllos** L. sp. p. 14; Mazz. in ant. ion. II. p. 442; Bois. fl. or. IV. p. 463. — Icon: Fl. dan. t. 627. — Exsicc.: Heldr. fl. thessala a. 1883.

Glanduloso-pilosa, simplex vel saepius a basi ramosa, ramis adscendentibus; foliis imis ovatis, petiolatis, integris vel crenatis, intermediis in 3—5 lacinias oblongo-spathulatas palmatisectis, floralibus sessilibus, trisectis vel summis indivisis; racemis laxis; pedicellis crassiusculis, calyce et bractea aequilongis vel longioribus, fructiferis erecto-patulis; calycis laciniis oblongis, corolla coerulea longioribus; capsula suborbiculata, turgidula, glandulosa, emarginata, calyce aequilonga.

In arvis, arenosis Graeciae (Bois.): ad radices mt. Pelion pr. Volo (Heldr.); Corcyra: pr. Castellanus (Mazz.). — Mart. Maio. ⊙

β. Folia floralia omnia caulinis conformia.

× Folia ovata, crenato-dentata; capsulae loculi 4—10 spermi; semina minora ovalia.

○ Capsula elevatim reticulato-venosa, lobis divaricatis.

27. **V. persica** Poir. dict. VIII. p. 542. — *V. Tournefortii* Gmel. fl. bad. I. p. 36, non Vill. — *V. agrestis v. byzantina* S. et S. fl. gr. I. p. 6 t. 8. — *V. Buxbaumii* Ten. fl. nap. I. p. VI. et 7 t. 1;

Mazz. in ant. ion. II. p. 444; Friedr. Reise p. 268; Ung. Reise p. 128; Bois. fl. or. IV. p. 465. — *V. filiformis* DC. fl. fr. suppl. p. 388; Ch. et B. exp. p. 16, Fl. pelop. p. 1; non Sm. — ? *V. byzantina* Mazz. in ant. ion. II. p. 444. — Exsicc.: Heldr. herb. norm. n. 747; Sint. it. thessal. n. 964.

Pubescens; caulibus elongatis, prostratis, inferne ramosis et saepe radicantibus; foliis petiolatis, ovatis vel cordato-ovatis, crenato-serratis; pedicellis filiformibus, folio longioribus, fructiferis patulo-recurvis; calycis laciniis ovato-oblongis, fructiferis valde divaricatis, capsula longioribus; corolla cyanea, calycem superante; capsula obreniforme, late et obtusangule emarginata, puberula.

In cultis regionis inferioris. Thessalia: pr. Malakasi (Sint.); Attica: pr. Athenas, Patissia (Held.), Marussi (Friedr.); Achaia: pr. Patras (Chaub.); Corcyra: pr. Castrades, Garizza, San Rocco (Mazz.). — Febr. Jul. ○

○○ Capsula laevis vel tenuiter reticulata, lobis non divaricatis.

28. **V. agrestis** L. sp. p. 13; Ch. et B. exp. p. 16, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 442; Raul. cret. p. 823; Bois. fl. or. IV. p. 466; Heldr. fl. cephal. p. 56. — *V. alba* Mazz. l. c. — Icon: Rchb. t. 79. —

Parce et breviter papillari-hirta; caulibus elongatis, prostratis, a basi ramosis; foliis breviter petiolatis, ovatis, crenato-dentatis; pedicellis folio aequilongis vel brevioribus, fructiferis recurvis; calycis laciniis oblongis, obtusis, corollam coerulestem vel albidam, striatam superantibus; capsula calyce subbreviore, glanduloso-pilosa, profunde emarginata, stylo emarginatura subbreviore, loculis 4—5 spermis.

In cultis. Corcyra: pr. Ascensione (Mazz.); Cephalonia: mt. Aenos (Heldr.); Achaia: pr. Patras (Chaub.); Creta: pr. Murnies, Canea (Raul.). — Febr. Apr. ⊙ N. v.

29. **V. polita** Fr. nov. fl. succ. p. 63; Haussk. symb. p. 176. — *V. didyma* Ten. in Spreng. syst. I. p. 75, Fl. nap. II. 2 fl. neap. syll. p. 7 (1830). — Icon: Rchb. t. 77. — Exsicc.: Heldr. herb. norm. n. !164.

Parce pubescens; caulibus elongatis, prostratis, a basi ramosis; foliis breviter petiolatis, ovatis, crenato-dentatis; pedicellis folio aequilongis vel brevioribus, fructiferis recurvis; calycis laciniis ovatis, acutiusculis, corollam coeruleam aequantibus; capsula calyce subbreviore, breviter tomentella, emarginata, stylo emarginatura longiore, loculis 8—10 spermis.

In cultis. Thessalia: ad monasterium Korona (Haussk.), pr. Velestinos (Heldr.); Attica: pr. Athenas (Heldr.); Acrocorinthus (Haussk.); insula Melos (Leon.). — Dec. Apr. ⊙

×× Folia suborbiculata, 3—9 loba; capsulae loculi 1—2 spermi, semina magna subglobosa.

30. **V. cymbalaria** Bodard diss. p. 3; S. et S. pr. I. p. 10, Fl. gr. I. p. 7 t. 9; Sieb. avis p. 2, rem. p. 2; Ch. et B. exp. p. 16, Fl.

pelop. p. 1; Mazz. in ant. ion. II. p. 444; Marg. et R. fl. Zante p. 71; Ung. Reise p. 128; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 823; Bois. fl. or. IV. p. 467; Heldr. fl. cephal. p. 56, Fl. Aegina p. 382; Haussk. symb. p. 176. — Exsicc.: Orph. fl. gr. n. 718 (Chios); Heldr. herb. norm. n. 1165.

Parce hirsuta; caulibus flaccidis, decumbentibus; a basi ramosis foliis subcordato-semiorbiculatis vel cuneatis, petiolatis, 5—9 lobis; pedicellis folio longioribus, fructiferis patentibus; calycis laciniis ovatis vel ellipticis, obtusis, ciliatis, basi attenuatis, fructiferis patulis vel reflexis; corolla alba, calyce vix majore; capsula inflata, didyma, hispida.

β. **panormitana** Tin. in Guss. pr. suppl. p. 4; Hal. Beitr. fl. Achaia p. 28, in ö. b. Z. 1897 p. 97. — Capsula glabra. — Exsicc.: Heldr. pl. fl. hellen. a. 1872 et 1900.

In cultis, muris, ruderatis, rupestribus regionis inferioris et montanae per totam Graeciam. — Febr. Maio. ⊙

31. **V hederifolia** L. sp. p. 13; S. et S. pr. I. p. 9; Pieri corc. fl. p. 3; Ch. et B. exp. p. 16, Fl. pelop. p. 1; Mazz. in ant. ion. II. p. 444; Friedr. Reise p. 265; Bois. fl. or. IV. p. 468; Haussk. symb. p. 176. — Icon: Fl. dan. t. 428. — Exsicc.: Heldr. in Baen. herb. europ. n. 2463; Dörfl. fl. gr. n. 478.

Plus minus hirsuta; caulibus decumbentibus, a basi ramosis; foliis suborbiculatis, petiolatis, infimis subintegris, ceteris 5—3 lobis; pedicellis folio subbrevioribus vel longioribus, fructiferis patenti-recurvis; calycis laciniis late cordatis, acutis, ciliatis, fructiferis erectis; corolla lilacina vel albida, calyce breviore; capsula globoso-quadriloba, glabra.

β. **triloba** Opiz hesper. a. 1815 pro sp. — Folia saepius 3 loba, pedicelli calyce aequilongi vel breviores. — Exsicc.: Heldr. fl. thessala a. 1883.

Inter segetes, in cultis, ruderatis, regionis inferioris in alpinam usque adscendens, (ubi f. *nana*, 2—4 cm. alta, simplex tantum occurrit) per totam ut videtur Graeciam. — Febr. Jun. ⊙

Obs. Ex confusione quadam indicantur: *V. pectinata* L. mant. p. 24, a S. et S. pr. I. p. 8, in summis jugis mt. Sphacioticorum Cretae et *V. syriaca* Roem. et Sch. syst. I. p. 116, a Mazz. ant. ion. II. p. 444, pr. Synadares Corcyrae.

5. Tribus. **RHINANTHEAE** Spreng. aul. II. 1 p. 390.

11. Parentucellia Viv. fl. lyb. spec. p. 31.

1. **P. latifolia** L. sp. p. 604; Sieb. avis p. 4, rem. p. 5; Urv. enum. p. 73; Ung. Reise p. 129; (*Euphrasia*); Car. in Parl. fl. ital. VI. p. 480. — *Bartsia latifolia* S. et S. pr. I. p. 428, Fl. gr. VI. p. 69 t. 586; Ch. et B. exp. p. 174, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 72; Friedr. Reise p. 269 et 271. — *Trixago latifolia* [illegible] p. 360; Weiss in z. b. G. 1869 p. 745; Haussk.

symb. p. 176; Form. in Ver. Brünn 1897 p. 48. — *Eufragia latifolia* Griseb. spic. II. p. 14; Raul. cret. p. 823; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 473; Heldr. fl. cephal. p. 56, Fl. Aegina p. 382; Hal. in ö. b. Z. 1897 p. 97. — Exsicc.: Orph. fl. gr. n. 710; Heldr. herb. norm. n. 1268 (f. *major*), 1268b, 1570 (*v. pycnantha*, f. spica etiam fructifera densa); Sint. it. thessal. n. 350; Dörfl. pl. cret. n. 51.

Glanduloso-pilosa; caule erecto, 3—20 cm. alto, simplici vel basi ramoso; foliis sessilibus, ovatis, crenato-pinnatilobis; bracteis palmatifidis, flores vix superantibus; spica compacta, subcapitata, demum elongata, inferne saepius interrupta; calyce vix ad tertiam partem in lacinias oblongo-lanceolatas, erectas diviso; corollae purpureae, tubo albae, rarius omnino albae (v. *albiflora* Raul. cret. p. 823) labio inferiore superiori aequilongo; antheris styloque glabris, capsula glabra vel breviter pilosa.

In collibus siccis, graminosis, pascuis regionis inferioris, in subalpinam adscendens. Thessalia: ad monasterium Meteora (Sint.), mt. Othrys (Form.), Pelion pr. Portaria (Heldr.); Sporadum insula Scopelos et Sciathos (Leon.); Attica: pr. Eleusis (Haussk.), Chelidonu, Kalyvia, Hagia Glycaria, mt. Parnes (Heldr.), insula Aegina (Friedr.); Elis (Sibth.): mt. Olenos (Hal.); Messenia (Sibth.): pr. Pylos (Chaub.); Argolis (Sibth.): pr. Nauplia (Haussk.), Poros (Friedr.); Cycladum insula: Cythnos (Tunt.), Melos (Urv.); Creta (Sieb.): pr. Nerokuru, Suda (Raul.), Mesalakonia distr. Mirabello (Leon.); Zante: ad paludem Makri (Marg.); Cephalonia: mt. Atro (Heldr.); Corcyra: pr. urbem (Sagb.), Bragagniotica (Ung.), mt. Deca (Spreitz.) — Mart. Jun. ⊙

2. **P. viscosa** L. sp. p. 602; S. et S. pr. I. p. 427; Sieb. in Flora V. 2. p. 639; Ch. et B. exp. p. 174, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 71; Friedr. Reise p. 283 (*Bartsia*); Caruel in Parl. fl. ital. IV. p. 482. — *Rhinanthus viscosus* Lam. fl. fr. II. p. 354; Pieri corc fl. p. 82. — *Eufrasia viscosa* Sieb. avis p. 4, non L. . — *Trixago viscosa* Rchb. fl. germ. exc. p. 360; Ung. Reise p. 129; Weiss in z. b. G. 1869 p. 745; Gelmi in bull. soc. bot. ital. 1889 p. 451; Haussk. symb. p. 176. — *Eufragia viscosa* Benth. in DC. pr. X. p. 543; Raul. cret. p. 823; Spreitz. in z. b. G. 1877 p. 717, 1887 p. 665; Bois. fl. or. IV. p. 474; Heldr. fl. cephal. p. 56. — Icon.: Rchb. t. 105. — Exsicc.: Orph. fl. gr. n. 709; Baen. herb. europ. n. 9362; Dörfl. fl. aeg. n. 35.

Glanduloso-pilosa; caule erecto, 10—50 cm. alto, simplici vel rarius ramuloso; foliis sessilibus, oblongis, crenato-dentatis; bracteis dentatis, inferioribus flores superantibus; spica elongata, inferne interrupta; calyce ad medium in lacinias lineari-lanceolatas, erecto-patulas diviso; corollae flavae labio inferiore superiori longiori; antheris barbatis; stylo hispido; capsula pilosa. — Praecedenti omnibus partibus major.

In pratis, locis arenosis regionis inferioris et montanae. Aetolia: pr. Aetolikon (Reis.); Euboea (Ung.): mt. Dirphys (Heldr.); Peloponnesus: pr. Patras (Hal.), pr. Carithena, ad fl. Alpheus, Sarandapotamos

(Chaub.), Kalamata (Friedr.), Tirynthum Argolidis (Orph.); insula Naxos (Leon.); Creta (Sieb.); pr. Canea, Suda (Weiss); Zante (Marg.); Cephalonia: pr. Lixuri (Heldr.); Corcyra (Pieri): pr. Kastrades (Baen.), Manducchio, Potamo (Spreitz.). — Apr. Jun. ⊙

12. Odontites Hall. hist. stirp. helv. p. 133.

a. Corolla lutea.

α. Planta suffruticosa, glabra.

1. **O. frutescens** Sieb. avis p. 4 (*Euphrasia*). — *E. fruticosa* Sieb. avis rem. p. 5. — *O. Bocconi* Walp. rep. III. p. 400; Benth. in DC. pr. X. p. 551; quoad pl. creticam; Raul. cret. p. 824; an *E. Bocconi* Guss. cat. hort. Baccad. p. 76. — *O. cretica* Bois. fl. or. IV. p. 477. — Exsicc.: Sieb. pl. cret. —

Caudice ramos steriles, abbreviatos, foliosos et floriferos elongatos, simplices vel apice in spicas 4—5 breviter paniculatas abeuntes, edente; foliis anguste lineari-lanceolatis, uninerviis; bracteis lineari-lanceolatis, calyce subaequilongis; calycis dentibus triangularibus, tubo triplo brevioribus; corolla glabra, tubo calyce sublongiore; antheris glabris, breviter mucronatis, capsula glabra, calyce sublongiore. — Tota planta siccitate nigrescit. — Species mihi ex unico specimine tantum nota et ulterius observanda, an revera ab *O. Bocconi* (Guss.), quae ex Boissier bracteis longioribus apice recurvis et capsulis puberulis differre dicitur, distincta sit. — Planta sicula praeterea ut videtur robustior, foliis latioribus.

Creta: ad Omalo in mt. Sphacioticis (Sieb.) — Aug. Sept. ♄

2. **O. Linkii** Heldr. et Sart. in Bois. diagn. ser. 2 III. p. 177; Fl. or. IV. p. 476; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 4, chlor. Parn. p. 24; Haussk. symb. p. 176. — Exsicc.: Heldr. herb. n. 67 et 1571, it. thessal. n. 68.

Caudice ramos steriles, abbreviatos, foliosos et floriferos elatos, saepius jam a medio in ramulos florales, elongatos, virgatos divisos, paniculam amplam formantes, edente; foliis anguste lineari-lanceolatis, uninerviis; bracteis lineari-lanceolatis, calyce subaequilongis; calycis dentibus triangularibus, tubo quadruplo brevioribus; corolla glabra, tubo calyce sublongiore; antheris breviter mucronatis, ad insertionem parce lanatis; capsula glabra, calyce sublongiore. — Praecedenti et *O. Bocconi* (Guss.) maxime affinis, ab illa calycis dentibus brevioribus, ab altera foliis angustioribus, bracteis brevioribus, apice non recurvis, ab ambabus caulibus ample paniculatis et antheris sublanatis differre videtur; characteres certiores invenire nequeo.

In petrosis regionis inferioris et montanae. Thessalia: mt. Pelion pr. Makrinitza (Heldr.) et Volo (Haussk.); mt. Parnassus ad Pagna et Gurna pr. Rachova (Heldr.); Laconia: in districtu Alagonia, l. d. Strasti pr. Megali-Anastasova (Zahn.). — Aug. Sept. ♄

β. Planta annua, glanduloso-pubescens.

3. **O. glutinosa** M. a B. fl. taur. cauc. II. p. 70 (*Euphrasia*); Benth. in DC. pr. X. p. 549; Bois. fl. or. IV. p. 475; Rouy ill. p. 55; t. 164. — *O. ixodes* Bois. et Heldr. diagn. VII. p. 45. —

Caule erecto, simplici vel stricte ramoso; foliis linearibus, floralibus abbreviatis, calyce longioribus; spicis angustis, demum laxiusculis; calycis dentibus linearibus, tubo sublongioribus; corollae tubo glanduloso, calyce 2—3plo longiore, limbo margine glabro; antheris mucronatis, ad insertionem villosis; capsula glandulosa, calyce subbreviore.

In regione alpina mt. Taygetos (Heldr.). — Aug. Sept. ⊙ N. v.

b. Corolla purpurea.

4. **O. serotina** Lam. fl. fr. ed. 2 II. p. 350 (*Euphrasia*); Rchb. fl. germ. p. 359; Bois. fl. or. IV. p. 476. — *E. odontites* L. sp. p. 604 var. *β*; Ch. et B. fl. pelop. p. 39. — *O. divergens* Jord. arch. fl. fr. et all. p. 191. — Exsicc.: Orph. herb. n. 564; Sint. et Bornm. it. turc. n. 1381; Heldr. herb. norm. n. 1463.

Pubescens, caule procero, divaricatim ramoso; foliis sessilibus, lanceolatis remote serratis; bracteis oblongo-linearibus, integris, flore brevioribus; spicis elongatis, laxiusculis; calycis dentibus lanceolatis, tubo aequilongis; corollae pubescentis tubo non exserto; antheris apiculatis, basi barbulatis; capsula calyci aequilonga, apice hispida. — Planta graeca ad var. *canescens* Rchb. fl. germ. p. 359 pertinet.

In cultis, herbidis regionis inferioris et montanae, rare. Thessalia: ad Litochori in Olympo (Sint.); Laconia: pr. Megali Anastasova (Zahn), mt. Taygetos (Orph.), pr. Kalamata (Gittard.). — Jul. Sept. ⊙

13. Euphrasia L. gen. n. 741.

1. **E. pectinata** Ten. fl. nap. I. pr. p. 36; Wettst. monogr. p. 82; Haussk. symb. p. 176. — *E. puberula* Hal. Beitr. fl. Epir. p. 34; Form. in Ver. Brünn 1896 p. 65; non Jord. pug. pl. nov. p. 133; quae = *E. tatarica* Form. in Ver. Brünn 1897 p. 48, non Fisch. in Spreng. syst. II. p. 777, et quae indumento magis setuloso et bracteis basi rotundatis parum discedit. — Huc forsan: *E. officinalis* S. et S. pr. I. p. 429; Dallap. prosp. p. 85; Ch. et B. exp. p. 175, Fl. pelop. p. 39; non L. . — Icon: Wettst. monogr. t. 7. — Exsicc.: Heldr. it. thessal. IV. a. 1885; Sint. it. thessal. n. 677.

Caule stricto, simplici vel rarius inferne ramo uno alterove praedito, pilis crispulis, brevibus, eglandulosis obsito; foliis glabris vel plus minus puberulis, inferioribus cuneatis, obtusis, utrinque obtusiuscule 1—3 dentatis, superioribus ovatis, acutis, utrinque aristato 4—6 dentatis; bracteis ovatis, rhomboideis vel ovato-lanceolatis, medio latissimis, basi cuneatis, utrinque elongato-aristato 3—5 dentatis; spica densiuscula, demum elongata; calycis setulosi dentibus lanceolatis; corolla 1 cm. longa, pallide lilalicina, violaceo-striata, puberula; capsula truncata, pilosa.

In pascuis subalpinis alpinisque, rare. Epirus: mt. Tsumerka (Hal.); Thessalia: in valle Negerli pr. Chaliki (Sint.), mt. Zygos, Karava, Gha-

velii (Haussk.), in oropedio Neropolis Helikis mt. Dokimi pr. Malakasi, Kerasia Sina, Mitrika mt. Oxya Form.; Aetolia (Wettst.); ? in Laconia Sibth. et Cephalonia Dall. — Jun. Jul. ♃

2. **E. salisburgensis** Funk in Hoppe bot. Taschenb. 1799 p. 184 et 190; Bois. fl. or. IV. p. 473; Form. in D. bot. Mon. 1891 p. 27 cum ?; Wettst. monogr. p. 278; Baldi viaggio Creta p. 51. — Icon: Wettst. monogr. t. 10. — Exsicc.: N. v.

Caule erecto, valde ramoso, ramis simplicibus pilis crispulis, brevibus eglandulosis obsito; foliis glabris vel parcissime setulosis, lanceolatis, acutis, utrinque aristato 2—3-dentatis, bracteis lanceolatis, utrinque et integro-aristatis 2—5, plerumque 3-dentatis; spica densa, demum elongata; calycis glabri vel parcissime setulosi lentiformi-lanceolatis; corolla 6—8 mm. longa, albida vel coerulea, puberula; capsula truncato-emarginata, glabra.

3. **olympica** Hal. in Sitz. k. ac. a. 1889 u. 1894. — *E. Olympica* Hal. et Sint. in S. b. Z. 1894 p. 99. — *E. Sintenisii* Hal. sp. Wettst. in Engl. et Prantl pflanzenfam. IV. 3 b. p. 101. — Folia et bracteae anguste linearia vel linear-lanceolata, dentibus eglandulosis, longe setaceo-aristatis, laminae latitudine saepe plus triplo longioribus, antrorsum curvatis. —

In rupestribus regionis subalpinae et alpinae rar. Thessalia: mt. Dokimi in Pindo Form.; mt. Parnassus Heldr.; Creta: supra Hellenochori Ilem. Sphakia Baldi; — 5. vel Hardie Phrygiae in Olympo (Sint.). — Aug. Sept. ♃

14. Bellardia All. fl. ped. I. p. 91.

1. **B. trixago** L. sp. p. 602; S. et S. L. p. 427; Fl. gr. t. 555; Ch. et B. exp. p. 174; Fl. pelop. p. 89; Marg. et R. fl. Zante p. 71; Freyn. Beiße p. 274 et 288; *Barsia*: All. l. c. — *Rhinanthus trixago* L. sp. ed. 2 p. 849; Pard. cont. fl. p. 82. — *R. maximus* Willd. sp. III. p. 189; Sieb. arb. p. 4, rem. p. 5. — *Barsia versicolor* Pers. syn. II. p. 151; Urv. enum. p. 75. — *Trixago apula* Stev. in mem. mosq. VI. p. 4; Urv. Beiße p. 29; Rand. cret. p. 524; Spreitz. in a. b. G. 1877 p. 717; Bois. fl. or. IV. p. 474; Heldr. fl. cephal. p. 56; Fl. Aegina p. 552; Gelm. in bull. soc. bot. ital. 1899 p. 431; Haussk. symb. p. 176; Hal. in S. b. Z. 1897 p. 97. — *T. carnea* Gmelin spic. II. p. 11; Weiss in a. b. G. 1899 p. 745. — Exsicc.: Baenitz herb. europ. n. 9161; Dörfl. fl. sup. n. 56; pl. cret. n. 161.

Pubescentia-hirsuta, superne glandulosa; caule erecto, simplici rarius ramoso; foliis ovalibus, linearibus-lanceolatis, dense serrato-dentatis; spica brevi, densiflora, tandem subelongata; calycis dentibus ovatis, obtusis, corolla tubo longioribus; corolla alba, purpureo-variegata vel luteovirid. puberula; capsula villosa, calyce sequilonga.

In pratis praesertim humidis, locis arenosis regionis inferioris. Thessalia: ad lacum Karla (Reiser); Sporadum insula Scopelos (Leon.); Aetolia: pr. Antirrhion (Hal.); Attica: pr. Athenas (Heldr.), Eleusis, Phaleron (Haussk.); Achaia: pr. Patras (Hal.); Messenia: pr. Arcadia, Methone, Messene, mt. Diaforti (Chaub.), Kalamata (Friedr.), Selitza (Zahn); Argolis: pr. Nauplia (Haussk.), Poros (Friedr.); insula Aegina (Heldr.), Cythnos (Tunt.), Tenos, Syra (Weiss), Melos (Urv.), Naxos (Leon.); Creta: pr. Canea, Akroteri, Tokadapora, Retymo, mt. Karadagh pr. Candia (Raul.), Males distr. Hierapetra (Leon.); Zante (Marg): pr. urbem (Reis.); Cephalonia: pr. Argostoli (Spreitz.); Corcyra (Pieri): pr. urbem (Baen.), Manducchio (Spreitz.). — Apr. Maio. ⊙

15. Alectorolophus Hal. hist. I. p. 137.

a. Calyx undique dense glanduloso-pubescens.

1. **A. pubescens** Bois. et Heldr. in Orph. fl. gr. n. 442 (*Rhinanthus*); Stern. in ö. b. Z. 1895 p. 126 t. 7, monogr. p. 49. — *R. major β. parnassicus* Bois. fl. or. IV. p. 479.

Caule erecto, 10—12 cm. alto, glanduloso-piloso, simplici vel rarissime parce ramoso, ramis brevibus; foliis caulinis internodio subaequilongis, ovatis oblongisve, obtusis, crenato-dentatis; bracteis dense glanduloso-pilosis, triangularibus, calycem paulo superantibus, in dentes breviter triangulares incisis; calyce undique dense glanduloso-piloso; corolla flava, 18 mm. longa, tubo paulum sursum curvato, dente labii superioris breviter triangulari, obtuso, labio inferiore recto, superius aequante et saepius dentem obtegente, eo dense adpresso, faucem occludente.

In regione superiori mt. Parnassus pr. Ses rarissime (Orph.). — Jun. Jul. ⊙

b. Calyx margine excepto glaberrimus.

2. **A. pindicus** Stern. in ö. b. Z. 1902 p. 177 t. 6. — Huc probabiliter: *A. alpinus* Fraas fl. class. p. 188, non (Baumg.) Stern. — *A. glandulosus* Haussk. symb. p. 176, (non Simonk.) Stern. — *A. minor* Stern. monogr. p. 103, quoad pl. graec., non (Ehrh.) Wim. et Grab. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Caule erecto, 30—40 cm. alto, subglabro, ramis virgatis, pari foliorum intercalarium unico instructis; foliis caulinis internodio brevioribus, oblongo-lanceolatis, crenato-dentatis; bracteis minute scabridis, triangularibus, calycem aequantibus, in dentes lanceolatos, apicem bracteae versus sursum decrescentes incisis; calyce margine scabrido, ceterum glaberrimo; corolla flava, 18 mm. longa, tubo paulum sursum curvato, dente labii superioris breviter triangulari, obtuso, labio inferiore recto, superius ad $^3/_4$ aequante, eo fere adpresso, faucem fere occludente. — Differt a praecedente caule elato, virgato-ramoso, foliis angustioribus, corollae labio inferiore subbreviori, superiori minus adpresso et praesertim indumento eglanduloso.

In nemorosis oropedii Neuropolis in Pindo dolopico (Heldr.). — Jun. Jul. ⊙

3. **A. Sintenisii** Stern. in ö. b. Z. 1902 p. 179 t. 6. — Exsicc.: Sint. it. thessal. n. 630.

Caule erecto, 25—40 cm. alto, sparsim piloso, simplici vel parce ramoso, ramis brevibus; foliis caulinis internodio subduplo brevioribus, ovato- vel oblongo-lanceolatis, crenato-dentatis; bracteis minute scabridis, triangularibus, calyce brevioribus, in dentes elongatos, lanceolatos, apicem bracteae versus sursum decrescentes incisis; calyce margine glanduloso-scabrido, ceterum glaberrimo; corolla flava, 20 mm. longa, tubo paulum sursum curvato, dente labii superioris breviter triangulari, obtuso, labio inferiore recto, superius ad $^3/_4$ aequante, eo adpresso, faucem occludente. — A praecedente caule subsimplici, foliis intercalaribus nullis, internodiis longioribus, bracteis brevioribus, calyce margine glanduloso, corolla majori, tubo magis curvato discedit.

In pratis montanis pr. Malakasi in Pindo tymphaeo (Sint.); memini me vidisse speciem *Alectrolophi*, forsan huc spectantem, in rupibus inadscensibilibus regionis alpinae inferioris mt. Tsumerka inter pagum Tsumerka et Melisurgi in Epiro. — Jun. Jul. ⊙

Obs. Alectrolophi species a Pieri corc. fl. p. 82 sub nomine *R. crista galli* L. in Corcyra indicata, ibi inquirenda.

16. Siphonostegia Benth. Hook. et Arn. bot. Beech. p. 203.

1. **S. syriaca** Bois. et Reut. diagn. XII. p. 44 (*Lesquereuxia*); Bois. fl. or. IV. p. 471; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 3; Hal. in ö. b. Z. 1892 p. 375. — Exsicc.: Heldr. it. thessal. n. 30; Sint. et Bornm. it. turc. n. 1378.

Glanduloso-hirsuta; caulibus erectis, virgatis, simplicibus vel parce ramosis, in spicam foliosam, laxiusculam abeuntibus; foliis sessilibus, oblongo-lanceolatis, floralibus calycem aequantibus vel superantibus; bracteolis linearibus; floribus subsessilibus; calycis glandulosissimi, elevato-costati laciniis oblongo-lanceolatis, tubo triplo brevioribus; corollae purpureae, glanduloso-puberulae tubo exserto, limbo parvo. — Caulis 30—50 cm. altus, crebre foliatus, folia 3—4 cm. longa, circa 5 mm. lata, flores 30—35 mm. longi. — Species egregia, rarissima. — Planta graeca ab asiatica indumento copiosiori et bracteolis brevioribus latioribusque differre videtur.

In fruticetis vallis Megarema pr. Bitos in Olympo (Sint.) et pr. Anilios in mt. Pelion (Heldr.). — Jul. Aug. ♃

17. Pedicularis L. gen. n. 746.

1. **P. graeca** Bunge in bull. phys. math. Petersb. I. p. 10; Bois. fl. or. IV. p. 490; Hal. in z. b. G. 1888 p. 761; Beitr. fl. Epir. p. 34; Heldr. chlor. Parn. p. 24; Bald. riv. coll. bot. alb. 1895 p. 60; Haussk.

symb. p. 176. — *P. rupestris* Bois. et Orph. diagn. ser. 2. III. p. 175; Heldr. herb. norm. n. 669.

Radicis fibris cylindricis, longis; caulibus erectis, parte superiore praesertim arachnoideo lanatis; foliis petiolatis, glabriusculis, ambitu ovato-oblongis, in lacinias breviter lineares bipinnatisectis, caulinis superioribus subsessilibus; spica densa, floriflera ovata, dein elongata; bracteis linearibus, pinnatifidis, inferioribus calycem superantibus; calycis albolanati dentibus lanceolatis, subdenticulatis, dimidium tubum aequantibus; corollae ochroleucae, glabrae galea incurva, erostri, apice truncata, subulato-bidentata, labii inferioris lobis denticulatis, cristis binis longis parallelis; filamentis duobus puberulis; capsula oblonga, acuta, subobliqua, rostrata, calyce triente longiore.

β. **inermis** Bald. riv. coll. bot. alb. 1896 p. 84. — Corollae galea sub apice truncato non subulato-bidentata. — Exsicc.: N. v.

In pascuis alpinis. Epirus: mt. Tsumerka, Peristeri (Hal.), Olycika (Bald.); Thessalia: mt. Karava (Haussk.); Aetolia: mt. Tymphrestus (Fraas), Korax (Heldr.); mt. Kiona (Hal.); Parnassus (Heldr.); Achaia: mt. Chelmos, Kyllene (Orph.); — β. mt. Smolika supra Kerasovo in Epiro boreali (Bald.). — Jun. Jul. ♃

2. **P. brachyodonta** Schloss. et Vuk. syll. fl. croat. p. 89, Fl. croat. p. 682; Haussk. symb. p. 176. — *P. ochroleuca* Schloss. in ö. b. Wochenbl. VII. p. 248; Rchb. icon. XXX. t. 134. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Radicis fibris fusiformibus interdum napuliformibus; caulibus erectis, pubescentibus; foliis petiolatis, glabris, ambitu oblongo-lanceolatis, in lacinias lanceolatas, pinnatipartitas, calcareo-denticulatas pinnatisectis, caulinis superioribus sessilibus; spica oblonga, densa; bracteis lanceolatis, pinnatifidis et integris, calycem aequantibus; calycis parce pilosi dentibus triangularibus, latitudini aequilongis, tubo multo brevioribus; corollae ochroleucae, glabrae galea incurva, subrostrata, apice truncata, bidentata, labii inferioris lobis denticulatis, cristis ad gibbos ovatos binos sub lobo intermedio reductis; filamentis duobus superne puberulis; capsula oblongo-lanceolata, falcato-rostrata, calyce subduplo longiore.

In pratis oropedii Neuropolis in Pindo dolopico (Heldr.). — Jun. ♃.

18. Rhinanthus L. gen. n. 740.

1. **R. elephas** L. sp. p. 603. — *Elephas Columnae* Guss. fl. sic. pr. II. p. 155. — *Rhynchocorys elephas* Griseb. spic. II. p. 12; Haussk. symb. p. 176. — Exsicc.: Heldr. it. gr. septentr. a. 1879, it. thessal. IV. a. 1885.

Parce pilosa, superne glandulosa; caulibus flaccidis, simplicibus vel ramosis; foliis brevissime petiolatis, ovatis, obtusis, dentato-crenatis, floralibus sensim diminutis, ellipticis; floribus in axillis solitariis, pedicellatis, racemum laxum formantibus; calycis labiis inaequalibus, inferiore longiore, profundius bifido; corolla lutea, glabra, calyce subduplo longiore, galea

in rostrum subulatum, supra basin dentibus binis dependentibus instructum abeunti, labio inferiori amplo, obtuse trilobo; capsula hirta, globosa, calyce breviore.

Ad rivulos, locis humidiusculis regionis subalpinae, rarissime. Thessalia: in oropedio Neuropolis pr. Bezula et Sermeniko in Pindo dolopico (Haussk.); in faucibus Ruphias mt. Oeta Phthiotidis (Heldr.). — Jun. Jul. ♃

19. Melampyrum L. gen. n. 305.

1. **M. ciliatum** Bois. et Heldr. diagn. ser. 2. III. p. 176. — *M. barbatum* Bois. fl. or. IV. p. 481, non W. et K. pl. rar. Hung. I. p. 89, quod differt foliis saepius latioribus, spicis latioribus, minus elongatis, laciniis bractearum lateralibus margine scabris vel breviter ciliatis, calycis tubo hispido. — Exsicc.: Orph. fl. gr. n. 722.

Breviter asperulo-hirtum; caule erecto, patule ramoso; foliis anguste linearibus; spicis terminalibus, densifloris; bracteis ovatis, in lacinias e basi lanceolata subulatas, margine valde albo-ciliatas partitis, lacinia intermedia aliis multo longiori et latiori; calycis tubo glabro, dentibus e basi lanceolata setaceis, ciliatis subbreviore; corollae luteae, tomentellae tubo calyce subduplo longiore.

Inter frutices sempervirentes ad radices mt. Olympus pr. Skala Thessaliae (Orph.). — Jul. Aug. ⊙

LXXXIII. Ordo. Orobanchaceae*).

Lindl. Nat. syst. ed. II. 287; Beck in Engl. Prantl Nat. Pflanzenfam. IV. 3b, 123.

Orobanche

(Tourn., Inst., 175 t. 181); L. gen. ed. VI. 321 nr. 779 p. p. rectius Beck Monogr. Orobanche (1890) 73 et in Nat. Pflanzenfam. l. c. 129.

1. Sectio. *Trionychon* Wallr. Sched. crit. I. p. 314 (1822); Beck Orob. p. 86. — *Kopsia* Dum. Comm. p. 16 (1822). — *Phelipaea* sect. E. A. Mey. in Led. Fl. alt. II. p. 459 (1830). — Flores bractea et bracteolis duabus minoribus, pedunculo vel calycis basi affixis suffulti; calyx gamosepalus, 4-rarius 5 dentatus; corolla (in nostris) coeruleo-violacea, supra staminum insertionem constricta; capsula bivalvis (Spec. 1—7).

a. Scapus ramosus (solum in plantis minoribus vel rarius simplex), saepius gracilis. (Spec. 1—5.)

α. Flores parvi, 10 summum 17 mm. longi. (Spec. 1—2).

1. **Orobanche ramosa** L. spec. pl. p. 633; S. et S. pr. p. 440; Urv. enum. p. 76; Ch. et B. exp. p. 179; Fl. pelop. p. 40

*) Auctore: G. Beck de Mannagetta.

Marg. et R. fl. Zante p. 72; Beck, Orob. 87. — *Phelipaea ramosa* C. A. Mey. Enum. pl. Cauc. p. 104 (1831); Raul. cret. p. 824. — Icon: Fl. Graec. VII. p. 7 t. 608; Reichenb. Iconogr. VII. f. 933—934 et Ic. fl. germ. XX. t. 152.

Spica laxi- et multiflora, demum elongata; floribus paulo pronus curvatis; dentibus calycis triangularibus, acuminatis, tubo suo brevioribus; corolla plurimum coerulea, laciniis rotundato-obtusis; filamentis basi puberulis; antheris plurimum glabris; stylo parce glanduloso-piloso; stigmate obsolete lobato, plurimum albido.

Plantae nutrices: *Cannabis sativa L.*, *Nicotiana tabacum L.*, *Lycopersicum esculentum* Mill. et plantae cultae aliae.

In cultis. Laconia: mt. Taygetos, pr. Androvista, in Maina pr. Vitylos (Chaub.); Messenia: (Sibth.), pr. Kubeh (Chaub.); Melos (Urv.); Creta (Sibth.); Zante (Marg.) — Aug., Sept. ⊙ N. v. Certo sequente contusa.

2. **O. nana** Noë in Reich. herb. norm. n. 1352; Beck Orob., 92; Hal. Beitr. fl. Epir. 34. — *Phelipaea Muteli β. nana* Reut. in DC. Prodr. XI. p. 9. — *Ph. nana* Reich. f. Icon. XX. 88. — *Ph. ramosa γ. nana* Boiss. Fl. or. IV. 499. — *O. Muteli* Hal. in ö. b. Z. 1897 p. 97, non Schultz. — Icon: Reich. f., l. c., t. 151. — Exsicc.: Heldr. herb. fl. hell. a. 1877 et 1879. —

Scapo simplici, gracili, rarius ramoso; spica brevi rarius elongata; floribus pronus curvatis; dentibus calycis subulatis, apicem versus filiformibus, tubum suum longitudine aequantibus; corolla intense violaceo-coerulea, laciniis ellipticis, subacutis; filamentis basi villosulis; antheris comosis vel glabris.

Plantae nutrices: Plantae annuae perplures (*Trifolium-*, *Vicia-*, *Coronilla-*, *Thlaspi-*, *Glechoma* spec. etc.).

In graminosis et herbidis. Epirus: mt. Tsumerka pr. Vulgarelion (Hal.); Attica: mt. Parnes p. Tatoi, in Pharmacusarum insula Megáli Kyrá (Heldr.!); Euboea: prope Kumi, Kurbatzi!; Argolis: pr. Nauplia!; Laconia: pr. Vitylos!; Sporadum ins. Skopelos (Leonis!); Cycladum insula Amorgos; Corcyra: pr. urbem (Sagb.) — April. Julio. ⊙?

β. Flores majores, 15—37 mm. longi (Spec. 3—4).

3. **O. Muteli** F. Schultz in Mutel Fl. franç. II. p. 353; Beck Orob. p. 95; Weiss in z. b. G. 1869 p. 744. — *Phelipaea Muteli* Reut. in DC. Prodr. XI. p. 8; Raul. cret. p. 824; Heldr. fl. Aegina p. 382, chlor. Mykon. p. 248. — *Ph. ramosa β. Muteli* Boiss. Fl. orient. IV. p. 499 p. p. — Icon: Mutel l. c. Atlas t. 43 f. 314 et Suppl. t. 2 f. 5; Reich. Icon. fl. Germ. XX. t. 150.

Spica plurimum densi- et multiflora; floribus pronus curvatis, 15—22 plurimum 20 mm. longis; calycis dentibus subulato-lanceolatis, tubum suum longitudine aequantibus; corollae plurimum dilute coeruleae laciniis rotundatis; filamentis basi pilosulis; antheris glabris v. comosis.

Plantae nutrices: *Anthemis cotula* L., *Pinardia coronaria* Less., *Endoptera Diosoridis DC.*, *Filago spathulata* Presl, *Orlaya maritima L.* etc.

α. **typica** Beck Orob. p. 97. — Scapus ramosus; spica multiflora plurimum densiflora et apice acuminata; corollae 2 cm. longae, rarius minores, paulo pronus curvatae, laciniis latis praeditae. — Exsicc.: Heldr. pl. fl. hell. a. 1875.

β. **spissa** Beck Orob. p. 96; Haussk. symb. p. 177. — *Phelipaea caesia* Gris. Spic. fl. rumel. II. 59 (1844) p. p. — *O. lavandulacea* Hal. in ö. b. Z. 1897 p. 97, non Rchb. — Icon. Buxb. Cent. III. t. I f. 2. — Scapus plurimum simplex; spica densiflora, apice rotundata; corollae circa 2 cm. longae, laciniis latis praeditae. — Exsicc.: Heldr. et Hal. fl. sporad. a. 1896.

In graminosis et herbidis. Attica!, ins. Aegina (Heldr.); Euboea!; Aetolia!; Argolis!; Cyclades: ins. Syros! (Weiss), Cythnos (Tuntas!), Mykonos (Heldr.); Creta: pr. Lutro, Damasta, in mt. Aphendi Kavutsi (Raul); Cephalonia, Zante. — β. Attica: pr. Peristeri (Heldr.!), in mt. Hymetto et pr. Kamaritza (Haussk.); Sporades: ins. Skopelos (Leonis!). April. Maio. ♃

4. **O. aegyptiaca** Pers. Ench. II. p. 181; Beck Orob. 100. — *Phelipaea aegyptiaca* Walp. Rep. III. 463; Reut. in DC. Prodr. XI. p. 9; Boiss. fl. orient. IV. p. 499; ?Heldr. fl. cephal. p. 56. —

Spica multi- saepe laxiflora, serius elongata; floribus paulo curvatis, subrectis, 20—37 plurimum 30 mm. longis; calycis dentibus lanceolato-subulatis, tubum suum longitudine aequantibus vel superantibus; corollae cyaneae laciniis latis, rotundatis, saepe suborbicularibus; filamentis basi pilosulis, supra saepe parce glanduloso-pilosis; antheris copiose lanuginoso-pilosis.

Plantae nutrices: *Gossypium herbaceum* L., *Solanum melongena* L., *Cucumis sativa* L.

Cephalonia: pr. Drapano (Heldr.), certe commutata. — Aug. Oct. ⊙ N. v.

5. **O. lavandulacea** Reich. Iconogr. VII. p. 48 (1829); Marg. et R. fl. Zante p. 72; Heldr. fl. cephal. p. 56; Beck Orob. p. 105. — *Phelipaea lavandulacea* Reuter in DC. Prodr. XI. p. 7 et Schultz Archiv. p. 101; ?Bois. fl. or. IV. p. 497. — *O. Fraasii* F. Schultz in Flora XXVI. (1893) p. 127 = *Phel. Fraasii* Walp. Repert. III. p. 460. — Icon: Reich. l. c. f. 935 (optime) et Icon. fl. Germ., XX. t. 147 (mediocriter).

Spica multi- et in apice densiflora, demum elongata laxa; floribus pronus curvatis, subhorizontaliter patentibus, 17—20 mm. longis; bracteis ovato-lanceolatis; calycis dentibus e basi triangulari subulatis, tubum suum longitudine subaequantibus vel eo paulo longioribus; laciniis corollae azureae, siccae pullae, copiose pilis intense azureis praeditis, rotundato-obtusis, rarius acutis; filamentis basi saepe pilosulis, supra parce glanduloso-pilosis; antheris circa basim ciliato-lanuginosis, rarius subcalvis.

Planta nutriens: *Psoralea bituminosa L.*

In herbidis. Attica: pr. Athenae (Heldr.!), ins. Salamis (Heldr.!); Argolis: pr. Naupliam (Heldr.!); Arcadia: pr. Tripolitsam (Fraas);

Cephalonia: pr. Argostoli, Livatho (Heldr.!); Zakynthos (Mazz.!); Corcyra!: pr. Pelleca (Gelmi). — April. Majo. ♃

b. Scapus plurimum simplex (rarius sub spica breviter ramosus) (spec. 6—7).

α. Scapus plurimum gracilis, saepe tener; calycis tubus inter dentes albo-membranaceus, paucis nervis praeditus (spec. 6).

6. **O. Schultzii** Mutel fl. franç. II. p. 352 (1835), add. p. 429; Beck Orob. 111. — *Ph. Schultzii* F. Schultz apud Mutel l. c. — *O. stricta* Moris Pl. sard. exs. (1840)!; Bert. Fl. ital. VI. p. 450 = *Phelipaea stricta* Moris ap. Reut. in DC. Prodr. XI. p. 7. — Icon: Mutel l. c., suppl. t. II. f. 4 (optime); Moris Fl. Sard. III. t. 102 (p. p. bene). — Exsicc.: Heldr. reliqu. Orph. a. 1886.

Spica multi- et densiflora, demum laxa, in apice plurimum pyramidaliter acuminata et subcomosa; floribus summo 20 mm. longis; calycis dentibus lineari-lanceolatis, apicem versus subsetaceis, semper saepe triplo tubo suo longioribus; corollae pronus curvatae coeruleo-violaceae laciniis ellipticis, acuminatis rarius rotundatis; filamentis basi pilosulis, supra parce glanduloso-pilosis; antheris comosis v. glabris.

Plantae nutrientes: *Ferula nodiflora* L., *Rubia peregrina* L., *Rumices.*

α. **typica** Beck, Orob., 112. — Scapus simplex v. ramosus; spica densi- v. laxiflora saepius pyramidaliter elongata; corollae laciniae acutae vel ellipticae.

? β. **pyramidalis** Beck Orob. 112; Haussk. symb. p. 177. — *Phelipaea pyramidalis* Reut. in Boiss., Fl. orient., IV, 498 (an Reuter in DC. Prodr. XI. 7?). Scapus plurimum simplex; spica densiflora; flores inferiores conspicue pedicellati, saepe verticillato-conferti; laciniae corollae ovatae, saepe suborbiculares, obtusae.

In herbidis. Thessalia: in mt. Pindo pr. Korona et Neuropolis (Haussk.!); Laconia: mt. Malevo pr. Hagios Petros (Orph.!), mt. Taygetos pr. Androuvista (Heldr.!) et pr. Gaïtzes (Psarides!); — β. Argolis: in mt. Kyllene pr. Trikalam (Orph.). — April. Junio. ♃

β. Scapus firmus, saepe crassior et altior; calycis tubus firmus, nervis copiose praeditus (Spec. 7).

7. **O. purpurea** Jacq. Enum. agr. Vind. p. 108 et observ. p. 252 (1762); Beck Orob. p. 123; Haussk. symb. p. 177. — *O. coerulea* Vill. Pl. Dauph. II. p. 406 (1787); — ?S. et S. pr. I. p. 440; Ch. et B. exp. p. 179, Fl. pelop. p. 40; Friedr. Reise p. 283; Schulz in Flora XXVI. p. 127; Ung. Reise p. 129. — *Phelipaea purpurea* Asch. Fl. Brand. p. 499 (1864). — *Phel. coerulea* C. A. Mey. Enum. pl. cauc. p. 104 (1831); Reut. in DC. Prodr. XI. p. 5; Bois. Fl. orient. IV. p. 495. — Icon: Jacqu. Fl. austr. III. t. 276; Reich. Iconogr. VII. f. 928 (bene), XX. t. 149 (male).

Scapo cum calyce dense furfuraceo- vel farinoso-glanduloso, parce squamato; spica in apice rotundata, plurimum laxiflora; floribus paulo

pronus curvatis, 25—30 mm. longis, rarius brevioribus; calycis dentibus saepe tubo suo brevioribus; corolla coeruleo-violascente, limbum versus intensius colorata; labii inferi laciniis ellipticis, plurimum rotundato-acuminatis; filamentis basi glabris vel pilosulis; antheris plurimum calvis vel comose pilosis.

Plantae nutrientes: *Achillea setacea* W. K., *A. nobilis* L., *Artemisia vulgaris* L., *Anthemis*.

In herbidis. Thessalia: pr. Klinovo in Pindo tymphaeo (Haussk.!); Attica: mt. Pentelicon (Sartori!); Peloponnesus (Sibth.): mt. Ithome Messeniae (Friedr.), mt. Kupe (Chaub.); Cephalonia (Unger); Creta (Sibth.). — Junio, Julio. ♃

2. Sectio. *Osproleon* Wallr. Sched. I. p. 307 (1822); Beck Orob. p. 132. — *Orobanche* C. A. Mey. in Ledeb. Fl. alt. II. p. 450 (1830); Reut. in DC. Prodr. XI. p. 15; Boiss. Fl. orient. IV. p. 502. — Flores bractea modo suffulti; calyx plurimum postice et antice fissus, segmenta 1—2dentata formans, rarius connatus; capsula saepius birimosa. (Spec. 8—24 et adnexae.)

1. Subsectio. *Inflatae* Beck in Hal. Braun Nachtr. Fl. Nied. Oest. p. 124 (1882) et Orob. p. 136. — Corolla sub insertione staminum inflata, in media parte constricta, limbum versus (in nostris) coeruleo-amethystina; stigma dilute coloratum (spec. 8).

8. **O. cernua** Loefl. It hisp. p. 152 (1758); Boiss. Fl. or. IV. p. 514; Beck Orob. p. 141.

Tota subfurfuraceo breviter glandulosa, saepe glabrescens; calycis partibus in apice rarius profundius bidentatis v. integris; corolla 10—20 mm. longa, acriter pronus curvata, cum fauce saepe deflexo, coerulescente, parce glandulosa v. glabrescente, laciniis in margine glabris; antheris plurimum, filamentis et stylo saepius glabris.

Indicatur modo *β.* **cumana** Beck Orob. p. 143. — *O. cumana* Wallr. Orob. gen. *δίασκ.* p. 58 (1825); Reut. in DC. Prodr. XI. p. 33; ?Raul. cret. p. 825. — *O. arenaria* M. Bieb. Fl. taur. cauc. II. p. 83 (1808) et III. Suppl. 418, non Borkh. — *O. bicolor* C A. Mey. in Ledeb. Fl. alt. II. p. 458 (1830). — Icon: Reich., Iconogr., VII. 926 (optime), XX. t. 189; Ledeb., Icon. t. 390. — Scapus saepius gracilis; spica laxiflora rarius in parte supera densior; corolla 15—18 mm. longa, acriter pronus curvata, supra tertiam infimam genuflexa et fauce iterum deflexa, limbo parvulo.

Plantae nutrientes: *Artemisiae* species, *Nicotiana rustica* L., *Lycium europaeum* L.

In herbidis Cretae (s. Raul.); certissime commutata. April. Julio. ♃

2. Subsectio. *Angustatae* Beck in Hal. Braun Nachtr. Fl. Nied. Oest. p. 125 (1882) et Orob. p. 152. Corolla sub insertione staminum angustior, supra insertionem sensim ampliata v. inflato-ventricosa; stigma saepissime coloratum (Spec. 9—14 et adnexae).

a. Corolla major 20—35 mm. longa (rarius minor), late tubulosa vel ampla (Spec. 9—14).

α. Calycis segmenta lateralia plurimum bidentata (Spec. 9—12).

× Stigma typice fuscopurpureum v. rubrum (Spec. 9).

9. **O. caryophyllacea** Sm. in Transact. Linn. soc. IV. (1797) p. 169; S. et S. pr. I. p. 440; Urv. enum. p. 76; Ch. et B. exp. p. 178, Fl. pelop. p. 40; Beck Orob. p. 155. — *O. Galii* Duby Bot. Gall. I. p. 349 (1828); Marg. et R. fl. Zante p. 72; Schultz in Flora XXVI. p. 129; Reut. in DC. Prodr. XI. p. 20; Boiss. Fl. or. IV. p. 508. — Icones: formae typicae et aliarum: Reich. Iconogr. V. f. 662, VII. f. 890—895, 905—907, 910, 913; IX. f. 1127—1128; XX. t. 162, 217; Dietr., Fl. boruss. t. 147, 253, 437—441; Sturm Deutschl. Fl. Heft 94.

Calycis partibus liberis v. antice coalitis, conspicue nervosis, vix dimidium corollae tubi aequantibus; corolla 20—35 mm. longa rarius minore, normaliter fusco-lilacina, sicca obscure fusca v. fusco-purpurea, cum linea dorsali e basi curvata, in medio dorso subrecta v. concava, in labio supero late lobato acriter curvata et declivi, laciniis margine glanduloso-pilosis; filamentis ad basim (rarius — 5 mm. supra basim corollae) insertis, infra usque ad mediam copiose pilosis, supra cum stylo plurimum copiose glanduloso-pilosis; stigmate normaliter fusco-purpureo. — Scapo saepe firmiore, squamis oblongo-lanceolatis, ad 2,5 cm. longis numerosis praedito.

Plantae nutrices: *Galii*, *Asperulae* et *Sherardiae* species.

In herbidis, inter frutices in Graecia et insulis (Sibth.); Euboea: (Fraas); Achaia: pr. Patras (Heldr.!); Laconia: pr. Vitylos, Chimova (Chaub.); Messenia inf. (Chaub.); Zacynthos: in mt. Scopo (Marg.); Cycladum ins. Melos (Urv.). — Julio ♃

×× Stigma luteum (Spec. 10—11).

10. **O. major** L. spec. pl. p. 632; Pieri corc. fl. p. 85; Beck Orob. p. 169. — *O. elatior* Sutt. in Trans. Linn. Soc. IV. p. 178 (1797); Friedr. Reise p. 275 cum?; Reut. in DC. Prodr. XI. p. 25. — *O. stigmatodes* Wimm. Fl. siles. p. 280 (1840). — *O. Kochii* F. Schultz in Flora (1847) I. p. 66. — Icones: Sutton l. c. t. 17 f. 4; Reich., Icon. fl. Germ., XX. t. 173 (male), t. 215 fig. II.; Schultz, l. c. t. II, f. 1.

Scapo plurimum firmo, squamis copiosis, basi imbricatis, oblongo-lanceolatis, glanduloso-pilosis; spica densi- et multiflora; corolla 15—26 plurimum 20 mm. longa, supra staminum insertionem non inflata, in dorso aequaliter plus minus curvata, rosea, demum pallide luteola v. sulphurea, labio superiore integro vel emarginato, lobis patentibus, laciniis in margine inaequaliter denticulato-crenato rarius repando subglabris; staminibus 4—6 mm. supra basim corollae insertis; filamentis supra glanduloso-pilosis, infra copiose pilosis; antheris oblongis, sensim acuminatis; stylo glanduloso piloso; stigmate luteo.

Plantae nutrices; *Centaureae* et *Echinopis* species.

In herbidis. Argolis: insula Poros (Friedr.); Corcyra (Pieri). — Junio, Aug. ♃ N. v.

11. **O. gracilis** Smith in Trans. Linn. soc. IV. (1797) p. 172: Beck Orob. p. 196. — *O. cruenta* Bert. rar. it. pl. dec. III. p. 56 (1810) et Fl. ital. VI. p. 430 (1844); Marg. et R. fl. Zante p. 72: Fraas fl. class. p. 188; Reut. in DC. Prodr. XI. p. 15; Bois. fl. or. IV. p. 505. — Icones: Reich. Iconogr. VII. f. 896, 898—899, 903; XX. t. 159; Coss. Germ. Fl. Paris. Atlas t. XIX. B. (optime).

Scapo gracili v. firmiore, infra dense supra parcius squamato; spica primum densi- serius saepe laxiflora; calycis segmentis bidentatis (rarius integris), corollae tubo brevioribus; corolla ampla, 20—25 mm. longa rarius minore, supra staminum insertionem valde ventricosa, in dorso aequaliter curvata, typice lutea, limbum versus rubra — fusco-purpurea. sicca fusco-purpurea v. subatra, intus intensius colorata et nitente, extus plurimum cum bracteis glanduloso-pilosa, labio supero integro cum laciniis brevibus, laciniis labii inferi saepe brevibus, medio paulo majore, omnibus inaequaliter denticulatis et glanduliferis; staminibus in basi corollae insertis; filamentis infra pilosis, supra sicut stylus glanduloso-pilosis (v. glabrescentibus); stigmate luteo, purpureo cincto; germine in basi antica 3—5gibboso. — Odor gratus caryophyllaceus.

Plantae nutrientes: *Leguminosae* variae, imprimis *Dorycnii, Trifolii, Loti, Coronillae, Spartii, Genistae, Cytisi* species.

α. **typica** Beck Orob. p. 196. — *O. gracilis* Sm. l. c. etc. — Corolla 15—25 plurimum 20 mm. longa; lobi labii superi breves; filamenta basi pilosa, supra glanduloso-pilosa. Scapus firmus vel gracilis, spica densi- v. latiflora. — Plantae nutrices s. c.

β. **Spruneri** F. G. Schultz in Flora XXVI. (1843) p. 130; Reuter in DC. Prodr. XI. p. 16; Boiss. Fl. Or. IV. p. 506; Hal. Beitr. fl. Achaia p. 28; Haussk. symb. p. 177; Heldr. fl Aegina p. 383; pro sp.; Beck Orob. p. 198. — Forma corolla tam longa quam lata, amplissima est *O. breviflora* Schultz in Flora XXVI. (1843) p. 130; Bois. fl. or. IV. p. 517. — Icones: Beck, l. c. f. 59 (2). — Exsicc. Heldr. herb. graec. norm. nr. 75! et Exsicc. nr. 1904! — Corolla magna, 20—25 mm. longa; lobi labii superioris saepe latiores, nervis obscuris subreticulatis perducti; filamenta supra parce glanduloso-pilosa, ea labii superioris ibidem saepe subglabra; scapus firmus, crassus, saepe copiose et late squamatus; spica cylindracea, multiflora, plurimum subdensa. Saepe transit ad *α.*

Plantae nutrices: sec. aut. *Cisti* spec. et *Spartium junceum* L.

In herbidis, graminosis, inter frutices. Euboea: pr. Kumi (Heldr.!); Attica: mt. Hymettus, Parnes pr. Dekeleia (Heldr.!); Boeotia: pr. Haliartus (Fraas); Argolis: pr. Nauplia (Heldr.!), Mykene (Boiss.!); Creta: pr. Zulatika Gortyniae (Heldr.!); Zante (Marg.); — *β.* Attica: mt. Pentelikon (Heldr.!), ins. Aegina (Heldr.!); Argolis: pr. Naupliam.

Mykenas (Heldr.!); Achaia: ad mt. Chelmos pr. Kalavryta (Hal.!) — April. Junio. ♃

×× Stigma pallidum (Spec. 12).

12. **O. crenata** Forsk. Fl. Aeg. p. LXVIII. et 113 (1775); Beck Orob. p. 225; Haussk. symb. p. 177. — *O. speciosa* DC. fl. franç. VI. p. 393 (1815); Reut. in DC. Prodr. XI. p. 19; Ung. Reise p. 129; Raul. cret. p. 824; Spreitz. in z. b. G. 1877 p. 717; Boiss. Fl. or. IV. p. 506, suppl. p. 357. — *O. pruinosa* Lapeyr. Abr. Pyr. suppl. p. 87 (1818); Marg. et R. fl. Zante p. 72; Weiss in z. b. G. 1869 p. 744; Heldr. fl. cephal. p. 56, Fl. Aegina p. 383; Gelmi in bull. soc. bot. ital. 1889 p. 451. — *O. grandiflora* Ch. et B. Exp. p. 178, Fl. pelop. p. 40, non Presl. — *O. segetum* Sprun. in Flora (1843) p. 129 et *O. canescens* Sprun. l. c. non Presl. — *O. angustisepala* Schultz in Flora (1843) 130 (ex mt. Hymetto) est forma calycis segmentis bifidis v. integris et dentibus angustissimis subulatis v. subfiliformibus, corolla saepe minore, c. 2 cm. longa. Icones: Reich. Icon, VII. f. 911 (optime) et XX. t. 161; Ch. et B. l. c. t. 22, Fl. Pelop. t. 23; Buxb., Cent. Byzanth III. t. II. — Exs.: Leonis Fl. Aeg. n. 95; Reverch. pl. creta n. 259.

Scapo firmo, spica cylindracea, multi- et plurimum densiflora; calycis partibus profunde bidentatis, cum dentibus angustissimis, plurinerviis, corollam subaequantibus; corolla 20—30 mm. longa, campanulata, ampla, supra staminum insertionem ampliata, cum linea dorsali in dorso subrecta, et in labio supero subdeclivi et ad marginem surrecta, alba v. ochroleuca, limbum versus imprimis in nervis rubra, coerulea v. purpurea, sicca tenui, papyracea, extus glabrescente, laciniis omnibus maximis, in margine crenulato-dentato glabris; filamentis 2—3 rarius 5 mm. supra basim corollae insertis, infra pilosis supra glanduloso-pilosis v. glabris; stylo glanduloso-piloso; stigmate pallide colorato.

Plantae nutrices: *Leguminosae* variae imprimis cultae sicut *Vicia faba* L., *Pisum sativum* L., *Lathyrus cicera* L., *Ervum lens* L., *E. ervilia* L. etc.

In cultis totius Graeciae et insularum frequens. Aprili, Majo. ♃

β. Calycis segmenta lateralia plurimum integra, lanceolato-acuminata (Spec. 13—14).

13. **O. alba** Steph. in Willd. spec. pl. III. p. 350 (1800); Reut. in DC. Prodr. XI. p. 20; Beck Orob. 209; Hal. in ö. b. Z. 1897 p. 97; Haussk. symb. p. 177. — *O. Epithymum* DC. Fl. franç. III. p. 490 (1805); Ch. et B. exp. p. 179, Fl. pelop. 40; Reuter l. c. p. 21; Ung. Reise p. 129; Raul. cret. p. 824; Bois. fl. or. IV. p. 509; Heldr. fl. cephal. p. 56. — *O. diaphana, O. punctata, O. crassistyla* Schultz in Flora (1843) I. p. 130—131; Boiss. Fl. or. IV. p. 506, 509. — *O. gracilis* Hal. in ö. b. Z. 1897 p. 97, non Sm. — Icones: Reich., Icon., VII. f. 895—889, 912, 914—915; XX. t. 215 f. 1, t. 163.

Scapo gracili paucifloro v. firmiore multifloro, spica saepe laxiuscula; calycis partibus plurimum indivisis (rarius bidentatis), cum dentibus 1—3nerviis, in siccitate fuscescentibus; corolla ampla, 10—28 mm. longa, cum linea dorsali plurimum leviter curvata, in labio superiore integro v. plicato emarginato declivi, albida v. ochroleuca limbum versus imprimis in nervis rarius tota rubra, amethystea vel purpurea et pilis glanduliferis coloratis saepe tuberculo brevi insidentibus praedita, sicca saepe purpurascente, laciniis magnis, in margine crenulato-denticulatis glanduloso-pilosis; filamentis fere in basi corollae insertis, infra pilosis (rarissime glabris) supra sicut stylus copiose (rare parce) glanduloso-pilosis; stigmate typice purpureo, rubro v. coccineo.

Plantae nutrices: *Labiatae* variae, imprimis *Thymi*, *Thymbrae*, *Salviae*, *Origani*, *Calaminthae*, *Clinopodii*, *Melissae*-Species.

Formae sequentes observatae sunt. 1. *communis* (Beck Orob. p. 209). Caulis gracilis, floribus paucis, laxifloris rarius appropinquatis, summo 20 mm. longis. — 2. *subalpina* (Beck Orob. p. 210; Haussk. symb. p. 177). Caulis subfirmus, spica pluriflora saepe densiore, floribus ultra 20 mm. longis, corollis purpureo pictis. — 3. *minutiflora* (Beck l. c. p. 210). Corollis minimis, 10—13 mm. longis. — 4. *microsepala* (Beck l. c. p. 211). Calycis partes integrae, brevissimae, suboccultae, vix sextam v. septimam corollae longitudine aequantes. — 5. *bidentata* (Beck l. c.). Calycis partes separatae, in omnibus v. pluribus floribus plus minus profunde saepe inaequaliter bidentatae. — 6. *glabrata* C. A. Meyer in Eichw. Pl. Casp. Cauc. I. p. 16. t. XVII (1831); Reuter in DC. Prodr. XI. p. 33 p. sp.; Beck Orob., p. 212; Haussk. symb. p. 177. Filamenta in basi glaberrima supra sicut stylus parce glandulosa v. eglandulosa. — 7. *leptocalamus* (Beck Orob., p. 210: Hausk. symb. p. 177). Scapus elongatus, tenuissimus; floribus remotissimis.

In herbidis et graminosis sat frequens. Attica: circa Athenas, mt. Hymettus, Parnassus; Lycabettus, Pentelikon, pr. Laurion etc. (Heldr.! et al.); Thessalia: mt. Zygos, pr. Klinovo (Haussk.), pr. Chaliki (Sint. s. *O. crinita*); Sporadum insula Skopelos (Leonis!); Phthiotis (Bois.); Argolis: pr. Naupliam (Heldr.!); Corinthus!, Messenia: pr. Kubeh et Messene (Chaub.); Cephalonia (Ung.); pr. Omala (Heldr.); Zante (Boiss.); Creta: pr. Retymo, Candia (Raul); f. 1—2 pervulgatae; f. 3—5 in Sporadum ins. Skopelos (Leonis); f. 6 Attica: pr. Laurion (Heldr.!), pr. Kamariza (Haussk.); f. 7. Argolis: in peninsula Methana (Hausskn.). — April. Majo. ♃

14. **O. reticulata** Wallr. Orob. gen. *διασκ.* p. 42 (1825); Beck Orob. p. 217.

Scapo plurimum firmo: spica cylindracea, multi- et plurimum densi-
flora; calyci [illegible] **indivisis** (rarius bidentatis), cum dentibus
uniner [illegible] **corolla tubulosa**, 20—22 mm. longa
rari [illegible] **dorso subrecta concava** v. leviter
c [illegible] **alba** v. ochroleuca, in labium

superum amethystina v. purpurascente et pilis glanduliferis purpureis saepe e tuberculo ortis tecta, sicca saepe nigrescente, laciniis magnis, in margine denticulato plurimum subglabris; filamentis 2—4 mm. supra basim corollae insertis, infra parce pilosis, supra parce glanduloso-pilosis vel utroque subglabris; stylo plurimum parce glanduloso-piloso; stigma typice rubro v. purpureo.

Formas sequentes ostendit:

α) **typica** Beck Orob. p. 219. — *O. reticulata* Wallr. l. c. — *O. platystigma* Reich. Iconogr. VII. 31 (1829); Hal. in z. b. G. 1888 p. 761. — *O. scabiosae* Koch Deutschl. Fl. IV. p. 440 (1833); Reut. in DC. Prodr. XI. p. 22. — Icones: Reich. Ic. fl. Germ., XX. t. 164. — Exsicc.: Hal. it. gr. a. 1888. — Corolla pilis glanduliferis obscuris copiose praedita. — Plantae nutrices: *Cirsii, Cardui, Knautiae* et *Scabiosae*-species.

β. **pallidiflora** Wimm. Grab. Fl. Sil., II. 1 p. 233 (1829); Reut. in DC. Prodr. XI. 22; Beck Orob. 218. — *O. Cirsii* Fries Mant. III p. 58 (1842). — *O. leucantha* Gris. Spic. fl. Rum. II. p. 57 (1844); Hal. in ö. b. Z. 1897 p. 97. — Icon.: Reich., Icon. XX. t. 168 et 211 (male). — Corolla pilis glanduliferis obscuris parcius obsita. Plantae nutrices: *Cirsii* et *Cardui*-Species.

In herbidis. Doris: α. in regione alpina montis Kionae (Hal.!). — Julio. — β. Phocis: mt. Parnassus (Guicc.!); Argolis: pr. Naupliam (Heldr.!); Sporadum ins. Skopelos (Hal.) non vidi. — Majo. ♃

b. Corolla minor, 10—20 post anthesim modo —23 mm. longa, anguste tubulosa, albida v. ochroleuca, limbum versus saepius coerulea, violacea v. purpurascens, sicca papyracea, laciniis in margine glabris; stamina in quadrante vel tertio infimo corollae inserta (Spec. 15—23).

α. Corolla supra staminum insertionem faucem versus sensim ampliata nec sub limbo constricta (Spec. 15—22).

× Stigma luteum (Spec. 15—16).

15. **O. attica** Reut. in DC. Prodr. XI. p. 28; Hal. in z. b. G. 1888 p. 761 et 763; Haussk. symb. p. 177. — *O. amethystea* Thuill. γ. *attica* Beck Orob. p. 230. — *O. amethystea* Boiss. Fl. or. IV. p. 512 (p. p.). — *O. major* Hal. Beitr. fl. Epir. p. 34, non L. — Icon. Beck l. c. f. 70 (2). — Exsicc.: Sint. It. thess. nr. 759 s. *O. crinita*.

Scapo subfirmo; spica cylindracea, multi- et densiflora, bracteis flores 15—23 mm. longos superantibus; calycis segmentis plurimum bidentatis, dentibus angustis, corollae tubum aequantibus; corolla a basi curvata pronus curvata serius saepius genuflexa vel deflexa, alba, limbum versus in nervis saepe amethystea, extus subglabra, sicca dilute fusca, basi alba, labio superiore profunde bilobo, laciniis crenulato-lobatis, in margine glaberrimis; filamentis 3—6 mm. supra basim corollae insertis, infra ad medium pilosis, supra parce glandulosis v. glabris; stylo parce glanduloso-piloso v. glabro; stigmate luteo.

O. amethystea Thuill. Fl. Paris, ed. 2 I. p. 317 (1797) typica glabritie corollae, filamentis basi parce puberulis, stigmate rubro fusco mox nigrescente paulo differt.

Planta nutriens adhuc nota: *Ballota acetabulosa* Bth.

In herbidis, pascuis, rupestribus. Epirus: pr. Kalarrytes (Hal.!); Thessalia: pr. Chaliki (Sint. s. *O. crinita!*), monasterium Korona (Haussk.!); Attica: mt. Lycabettus, pr. Oropos, Phaleron (Heldr.!); Doris: in reg. alpina mt. Kionae alt. 6000 (Hal.!), pr. Itea, Amphissa (Heldr.!), mt. Parnassus (Hal.!). — Junio, Julio. ♃

16. **O. canescens** J. et C. Presl Del. Prag. I. p. 72 (1822); F. W. Schultz in Flora XXVI. p. 129; Raul. cret. p. 825; Gelmi in bull. soc. bot. it. 1889 p. 451; Beck Orob. p. 235; Haussk. symb. p. 177. — *O. australis* Moris ap. Bert. Fl. ital. VI. p. 440 (1844); Reut. in DC. Prodr. XI. p. 29. — *O. tridentata* = *O. grandisepala* F. G. Schultz in Flora (1843) p. 128 (f. segmentis calycis tridentatis); Bois. fl. or. IV. p. 516. — Icones: Reich., Icon., XX. t. 210. — Exsicc: Sint. It. thess. n. 552 s. *O. procera.*

Scapo firmo; spica cylindracea, saepe longissima, densi- in basi modo laxiflora; bracteis flores plurimum superantibus; calycis segmentis bidentatis, dentibus angustis, corollae tubum longitudine aequantibus; corolla 13—16 mm. longa, erecto-patente cum linea dorsali e basi curvata porrecta, sordide lutea vel rubescente, sicca dilute in limbo intensius fusca, extus glanduloso-pilosa, glabrescente, labio supero emarginato v. bilobo, laciniis in margine crenulato-dentatis, glabris; filamentis 3—4 mm. alte insertis, basi ad medium copiose pilosis, supra glabris; stylo parce glanduloso-piloso; stigmate luteo. — Ab *O. versicolore* spica densi- et multiflora, saepe longissima, corollis extus non villosis, porrectis, indumento staminum, stigmate luteo, ab *O. attica* corollis minoribus, gracilibus, earum tubo multo angustiore distinguitur.

Plantae nutrientes incognitae.

In herbidis. Thessalia: pr. Kalabaka (Sint.! s. *O. procera*); Attica: ad Phalerum pr. Athenas (Heldr.!), pr. Kamariza (Haussk.); Argolis: pr. Naupliam, Tolon (Haussk.), ibidem *O. grandisepala* (Fraas); Corfu: pr. Pelleca (Gelmi); Creta (Friv. et Raul). — Majo, Junio. ♃

×× Stigma violaceum v. fusco-purpureum (Spec. 17—22).

17. **O. versicolor** F. G. Schultz in Flora (1843) I. p. 129; Beck Orob. p. 237. — *O. pubescens* Urv. Enum. pl. or. p. 76; Ch. et B. fl. pelop. p. 40 (1822); Reut. in DC. Prodr. XI. p. 27; Ung. Reise p. 129; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 825; Spreitz. in z. b. G. 1877 p. 717; Boiss. Fl. Or. IV. p. 507, suppl. p. 357; Heldr. fl. ceph. p. 56, Fl. Aegina p. 383, chlor. Thera p. 18; Hal. in ö. b. Z. 1897 p. 97; Haussk. symb. p. 177. — *O. villosa* Schultz in litt. — *O. villiflora* Koch in litt. = *villosiflora* Schultz in Flora (1845) p. 737 et 740. — *O. arachnoidea* Schultz in Linnaea XIX.

p. 26 et XXII. (1849) p. 669. — *O. fragrans* Gris. Spic. fl. Rum. II. p. 58, non Koch. — *O. livida* Hal. in ö. b. z. 1897 p. 97, non Sendt. — Icones: Beck Orob. t. IV. f. 75 (flos.). — Exsicc.: Heldr. herb. norm. nr. 668!; Reverch. pl. cret. (1883) nr. 124!; Orphan., Fl. Graec., nr. 21!

Scapo longe saepe copiose glanduloso-piloso, gracili paucifloro v. firmiore et spica cylindrica, multiflora, laxa v. densa praedita; bracteis subvillosis, flores plurimum superantibus; calycis segmentis bidentatis, dentibus angustis, corollae tubum saepe superantibus; corolla 10—20 mm. longa, e basi curvata porrecta cum dorso recto vel acrius pronus curvato saepe subgenuflexa, ochroleuca limbum versus lilacina v. violascente ibidem sicca saepe fusco-purpureo- v. nigrescente, extus imprimis in labio supero longe lanuginoso-pilosa, labio supero integro, emarginato v. bilobo, laciniis in margine glabro crenulato-dentatis; filamentis 3—4 mm. alte insertis, ad mediam v. ultra longe pilosis, supra sicut stylus glanduloso-pilosis; stigmate violaceo-coeruleo.

Plantae nutrientes: *Compositae*, *Labiatae*, *Umbelliferae* variae: *Pinardia coronaria* Less., *Orlaya maritima* L. etc.

Formae permultae observantur: (cfr. Beck Orob., 238—239). 1. *monochroa* (Beck = f. *pallida* Reut. l. c. non Schultz). Corollae limbus non intensius colorata. — 2. *typica* (Beck). — Corolla in limbo lilacina v. fusco-purpurea, sicca pulla v. nigrescens. — 3. *pseudobarbata* (Beck). Bracteae flores conspicue superantes, in spicae apice comosae. — 4. *macrophyllon* (Beck. — *O. minor v. adenostyla* Hal. Beitr. fl. Achaia p. 28, non Vis.) — Bracteae flores superantes; calycis dentes angustissimi, in apice filiformes, corollae longitudinem aequantes. — 5. *lanifera* (Beck). In omnibus partibus copiosissime albido-arachnoideo-villosa. — 6. *campuliflora* (Lo Jacono Crit. Orob. in Natural. Sic. 1882—1883 p. 57 nr. 35 t. IX. fig. 1 a. p. sp.; Beck l. c. p. 239). Corolla exigue pronus curvata, dilute colorata, 10—12 mm. longa (in f. 1—5, 12—15 post anthesim — 20 mm. longa).

Pervulgata imprimis in incultis, arenosis maritimis per totam Graeciam et in insulis. — Aprili, Majo in calidioribus Martio. ♃

Obs. *O. arcuata* F. G. Schultz in Flora (1843) p. 129; Boiss., Fl. Or., IV. 517; Beck, Orob., p. 240, ex Attica (Fraas) forma quaedam parviflora, corollis magis pronus curvatis esse videtur. Non vidi.

18. **O. Grisebachii** Reut. in DC. Prodr. XI. p. 28 (1847); Beck Orob. p. 240; Hausk. symb. p. 177. — *O. condensata* Gris. Spic. fl. Rum. II. p. 56, non Moris. — *O. striata* Reut. l. c. (p. max. p.). — *O. palaestina* Boiss. Fl. Or. IV. p. 506 p. p. — Icon.: Beck Orob. t. IV. f. 76 (flos). — Exsicc.: Reverch. Pl. Crète nr. 124!

Scapo firmiore, breviter glanduloso-piloso, spica cylindracea, densi- et multiflora, basi modo laxiflora; bracteis flores aequantibus v. superantibus; calycis segmentis bidentatis, dentibus longe acuminatis, tubum corollae aequantibus; corolla 12—15 mm. longa, erecto-patente, cum linea dorsali in dorso subrecta, limbum versus complanata vel arcuato-declivi, albida v. violascente, extus glanduloso-pilosa, labio supero integro

v. plicato-emarginato, laciniis in margine glabro inciso-crenulato-dentatis; filamentis 3—4 mm. alte insertis, ultra medium longe pilosis, supra glabris; stylo parce glanduloso-piloso; stigmate violaceo-purpureo. Difficillime ab *O. versicolore* indumento caulis et corollae, ab *O. Picridis* scapo brevius piloso, squamis minoribus, floribus minoribus, labio superiore vix emarginato, ab *O. attica* corolla minore, erecto patente, labio supero subintegro, stigmate violaceo-purpurea distinguitur.

Planta nutriens: *Coronilla parviflora* W.

In campis, arenosis et incultis. Attica: Athenis in Acropoli (Haussk.!), pr. Phaleron (Heldr.!); Laconia: pr. Spartam (Heldr.!); Creta: pr. Canea (Reverch.!). — Aprili, Majo. ♃?

19. **O. loricata** Reich. Iconogr., VII. p. 41 (1829); Marg. et R. fl. Zante p. 72; Reut. in DC. Prodr. XI. p. 27; Beck Orob. p. 243. — *O. Artemisiae campestris* Vauch. in Gaud. Fl. Helv. IV. p. 179 (1829). — Icon.: Reichenb. l. c. VII. f. 917; XX. t. 176, 216 f. 1; Dietr. Fl. boruss. IX. t. 580.

Scapo plurimum firmiore; spica cylindracea, multi- et plurimum densiflora, serius basi laxiflora, bracteis apice comosa; calycis segmentis profunde bidentatis rarius integris, dentibus angustis, corollae tubum longitudine aequantibus; corolla 14—22 mm. longa, erecto-patente, cum linea dorsali in dorso subrecta in labio supero curvata declivi, ochroleuca, limbum versus violacea, sicca dilute fusca, extus glanduloso-pilosa, labio supero plicato-emarginato v. bilobo, laciniis in margine glabro crenulato-lobatis; filamentis 3—4 mm. alte insertis, usque ad mediam pilosis supra plurimum sicut stylus glanduloso-pilosis; stigmate rubro-violaceo.

Planta nutriens: *Artemisia campestris*. L.

In herbidis. Zante in mt. Scopo (Marg.) non vidi et valde in dubium revoco.

20. **O. picridis** Schultz in Flora (1830) Litt. Bl. p. 500; Reut. in DC. Prodr. XI. p. 26; Beck Orob. p. 245. — *O. Reichardiae* Freyn in z. b. G. (1877) p. 392.

Scapo gracili, paucifloro v. firmiore, spica cylindracea multiflora ornato, glanduloso-piloso; bracteis flores vix superantibus; calycis segmentis ad mediam bidentatis (rarius integris), dentibus saepe subulatis, tubum corollae aequantibus; corolla 15—20 mm. longa, erecto-patente v. patula, cum linea dorsali in dorso subrecta limbum versus declivi, alba v. ochroleuca, limbum versus saepe violacea, sicca dilute fusco-lutea, extus glanduloso-pilosa v. subglabra, labio supero plicato-emarginato v. bilobo, laciniis in margine glabro crenulatis; filamentis 3—5 mm. alte insertis, usque ad mediam pilosis, supra plurimum glabris; stylo glandulosopiloso; stigmate violaceo-purpureo.

Plantae nutrices: *Compositae* nonnullae, *Picridis, Crepidis, Tragopogonis* species, porro *Dauci* et *Orlayae* species.

Variat: α. **typica** Beck Orob. p. 245; Synon. l. c. — Corolla sicut tota copiose glanduloso-pilosa. — Icon.: Reich. Icon. XX. t. 175 (α). — Exsicc. Heldr. pl. Cephal. nr. 3565!

β. **carotae** Des Moul. in Ann. sc. nat. III. (1835) p. 78 pro spec.; Beck Orob. p. 246. — *O. ambigua* Moris diagn. sard. p. 2 et fl. sard. III. p. 251. — *O. Tommasinii* Reich. f. Icon. XX. p. 92 = *O. elatior* b. ibidem 118. — *O. lilacina* Schultz in Flora (1843) I. p. 129; Bois. fl. or. IV. p. 517. — *O. Freynii* Nym. consp. p. 561. — Icon. Reich. l. c. t. 209, 216. — Corolla sicut tota parce glanduloso-pilosa, subglabra; spica plurimum multi- et densiflora, bracteis rubro violaceis, siccis nigrescentibus.

O. picridis ab *O. loricata* calycis segmentis minus profunde bidentatis, filamentis supra subglabris, ab *O. attica* floribus erecto-patentibus, stigmatis colore, ab *O. minore* corollis majoribus, laciniis labii superi patulis majoribus, filamentis infra copiose pilosis distinguitur.

In herbidis et rupestribus. In insulis jonicis!; Cephalonia: pr. Argostolis (Heldr.!); Aetolia: in mt. Taphiassos ad sinum Patranum (Hal.!); Attica: mt. Lycabettus, Hymettus, pr. Laurion (Heldr.!), Kamariza (Haussk.); Argolis: pr. Naupliam! — β. Attica: in mt. Lycabetto (Heldr.!). — Majo, Junio. ♃

21. **O. fuliginosa** Reut. in DC. Prodr. XI. p. 23; Boiss. Fl. or. IV. p. 510; Beck Orob. p. 248. — Icon. Jord. observ. III. t. 9 fig. B. (male); Beck l. c., t. IV, f. 80 (flos.)

Scapo subfirmo, breviter glanduloso-piloso; spica ovata v. cylindrica, multi- et densiflora in basi saepe laxiflora; calycis segmentis integris (rarius bidentatis), dentibus angustis, tubo corollae subbrevioribus; corolla 15—18 mm. longa, pronus curvata, cum linea dorsali in dorso subrecta, in labio supero subbilobo complanata v. declivi, in margine arrecta, ochroleuca, limbum versus rubescente v. violaceo-fusca, extus parce glanduloso-pilosa, glabrescente; filamentis 2—3 mm. alte insertis, infra pilosulis, supra glabris rarius sicut stylus glanduloso-pilosis; antheris brevissime apiculatis; stigmate violaceo-aurantiaco.

Plantae nutrientes sec. aut. *Anthemis maritima* L., *Cephalaria ambrosioides* Boiss., *Cineraria maritima* L.

Variat: α. **typica** Beck Orob. p. 249. — Laciniae corollae rotundatae; filamenta supra glabra (v. subglabra).

β. **oxycheilos** Beck l. c. — Flores erecto-patentes; laciniae corollae omnes acuminatae, profundius crenato-denticulatae; corolla in labio supero insigniter galeata et biloba.

In herbidis. α. Attica: in mt. Hymetto (Heldr.!); Laconia: in mt. Taygetos pr. Anavryti (Heldr.!); Cycladum insula Syra (Weiss!). — β. in mt. Hymetto Atticae (Heldr.!). — Majo, Junio. ♃

22. **O. minor** Sutton in Trans. Linn. soc. IV. (1797) p. 179; Reut. in DC. Prodr. XI. p. 29 p. p.; Ung. Reise p. 129; Spreitz. in z. b. G. 1877 p. 717; Bois. fl. or. IV. p. 512; Heldr. fl. ceph. p. 56;

Beck Orob. p. 251; Haussk. symb. p. 177. — *O. nudiflora* Wallr. Sched. crit. I. p. 310 (1822) ex p. — *O. apiculata* Wallr. l. c. — *O. hyalina* Sprun. apud Reut. l. c. p. 24; Bois. fl. or. IV. p. 508, suppl. p. 357; Heldr. fl. ceph. p. 56, Fl. Aegina p. 383; = *O. Reuteri* Schultz in litt. fide Reut. — Icones: Reich., Icon., VII, f. 876—880, 883—884, 916, XX. t. 183.

Scapo gracili rarius firmiore; spica cylindrica, plurimum multiflora, basi v. tota laxiflora; bracteis flores plurimum aequantibus; calycis segmentis integris v. bidentatis, dentibus longe subulatis, corollam longitudine plurimum aequantibus; corolla 10—18 mm. longa, erecto-patula v. pronus curvata, cum linea dorsali aequaliter curvata, ad limbum complanata, ochroleuca, limbum versus saepius coerulea v. violascente, extus glanduloso-pilosa (v. subglabra), laciniis labii superi emarginati v. bilobi porrectis, in margine glabro crenulato-dentatis; filamentis 2—3 mm. alte insertis, infra parce pilosis, supra sicut stylus glabris v. subglabris; stigmate violaceo-purpureo.

Plantae nutrientes: imprimis Leguminosae, Compositae (*Trifolii*, *Ornithopi*, *Helichrysi*-Species) sed etiam permultae aliae, atque cultae.

Formae sequentes observantur. 1. *minima* (Beck Orob. p. 252). Corolla minutissima, c. 10 mm. longa. — 2. *typica* (Beck l. c.). Spica densi- v. laxiflora, corolla c. 15 mm. longa, extus glanduloso-pilosa, ochroleuca, limbum versus violascens. — 3. *pumila* (Koch in Noë exs.; Reich. f., Icon., XX, 104 t. 165 f. II. p. sp.; Beck Orob. p. 252. — *O. livida* Sendt. (fide Freyn.) v. *adenostyla* Vis. Fl. Dalm. II 179). — Spica mox laxiflora; corolla primum parce glanduloso-pilosa, mox glabrescens.

In herbidis et graminosis. Attica: in mt. Lycabetto (Unger), ad Phalerum (Boiss.), ins. Aegina (Boiss.), Angistri (Heldr.); Argolis: pr. Naupliam (f. 2!); Sporadum ins. Skopelos (Leonis!); Aetolia: pr. Mesolongion (Heldr.!); Cephalonia: pr. Thaphion (Heldr.), Argostoli (Spreitz.); Corcyra: pr. Potamos (f. 3, Grimb.!); Thessalia: in mt. Karava Pindi (Haussk.!). — Aprili, Julio. ♃

β. Corolla sub limbo conspicue constricta, stigmate luteo (Spec. 23).

23. **O. hederae** Duby Bot. Gall. I. p. 350 (1828); Reut. in DC. Prodr. XI. p. 28; Beck Orob. p. 259. — *O. medicaginis* Reich. Iconogr. VII. p. 41; Griseb. Spic. fl. Rum. II, 57; non Duby. — *O. laurina* Reich. f., Icon., XX. 104, non Ch. Bonap.

Scapo subfirmo; spica cylindracea, multiflora, plurimum densi- rarius laxiflora; calycis segmentis integris v. bidentatis, dentibus saepe filiformibus, plurimum tubo corollae brevioribus; corolla 10—20 mm. longa, erecto-patula v. pronus curvata, sub limbo constricta, ochroleuca v. flavescens, in labium superum integrum v. plicato-emarginatum rubescens v. lilacea, laciniis in margine glabro crenato-denticulatis; filamentis 3—4 mm. alte insertis, infra breviter pilosis supra glabris; stylo plurimum parce glandu[illegible] stigmate luteo v. aurantiaco.

Planta nutriens: *Hedera helix* L.

Observata est forma spica elongata subtota laxiflora, floribus infimis remotissimis = f. *stenantha* Lo Jacono Crit. Orob. in Nat. Sic. I—II. 1882—1883, 53 p. sp.; Beck Orob. 260; Haussk. symb. p. 177. —

Attica: pr. Athenas (Heldr.!). — Majo, Julio. ♃

Species insufficienter descriptae et incertae sedis:

O. lycica F. G. Schultz in Flora (1843) 128; Boiss., Fl. or., IV, 516; Beck, Orob., 264. — Sepalis plurinerviis, ovatis, bifidis rarissime cum dente altero truncato praeditis, dentibus lanceolato-subulatis, tubum corollae aequantibus v. longioribus; corolla tubuloso-campanulata, in dorso curvata, in fauce ampliata, labio supero bilobo, laciniis inaequaliter et acute dentatis; staminibus sub medio tubi insertis, a basi ad medium subvillosis. — Attica: ad Lycos (Fraas), non vidi. E minoribus, *O. atticae* affinis.

O. subverticillata F. G. Schultz in Flora (1843), 128; Boiss., Fl. or., IV. 516; Beck, Orob., 264. — Sepalis obsolete 5 nerviis, ovatis; inaequaliter bifidis v. indivisis, tubum corollae aequantibus; corolla tubuloso-campanulata, arcuata, labio supero vix bilobo, laciniis inaequaliter denticulatis; filamentis in curvatura corollae insertis, a basi ad medium subvillosis. — Boeotia: in *Vicia* sp. ad Thebas (Fraas), non vidi.

O. parviloba F. G. Schultz in Flora (1843) 128; Boiss., Fl. or., IV. 516; Beck, Orob., 265. — Planta, praesertim inflorescentia pilis albidis subvillosa, exsiccata griseo-ochroleuca; sepalis obsolete 3—6 nerviis, e basi ovata lanceolato-subulatis, bifidis, rarissime indivisis, corollae tubum superantibus; corolla tubulosa, subarcuata, laciniis parvis, obsolete denticulatis; staminibus infra medium tubi insertis, sparsim villoso-pilosis. — Argolis: ad Naupliam (Fraas). *O. canescenti* affinis ?.

O. chassia Form. in Deutsch. bot. Mon., 1897 p. 197, in Ver. Brünn, 1897, 45. — Scapo breviter et remote squamoso; spica longa, laxiuscula, bracteis corollae tubo longioribus; calycis segmentis bidentatis, dentibus lineari-subulatis, corollae tubo sublongioribus; corolla dorso leviter arcuata, sicca rubella, labio snpero bilobo, lobis retusis, laciniis non ciliatis, subdenticulatis; filamentis paulo supra basim corollae insertis, inferne puberulis; stigmate e sicco rubro; ovario superne parce glanduloso. — Thessalia: pr. Mitrica in mt. Chassia (Form.) non vidi. — *O. amethysteae* Thuill., nec *O. cupreae* Boiss. ex descriptione affinis.

O. prosgolica Form. in Ver. Brünn 1895 p. 52 = *O. pruinosa* Form. non Lapeyr. l. c. p. 33. — Caule crassiusculo minute et tenuissime pruinoso; spica laxiuscula, bracteis pruinoso-puberulis, corollae tubo longioribus; calycis segmentis bidentatis, dentibus lineari-subulatis tubum corollae aequantibus v. subsuperantibus; corolla flavo-rubella, glabra, dorso laeviter arcuata, labio supero bilobo, laciniis labii inferi subrotundis subaequalibus, omnibus denticulatis non ciliatis; filamentis supra tertiam tubi insertis, basi puberulis; stigmate e sicco fusco rubro. —

Epirus: in mt. Prosgoli, 1000 m. s. m. (Form.) non vidi. — Ex descriptione forma *O. crenatae* Forsk., certe negligenter descripta.

LXXXIV. Ordo. **Acanthaceae** Juss. gen. p. 102.

1. **Acanthus** L. gen. n. 893.

a. Foliorum laciniae ovatae, dentatae, inermes.

1. **A. longifolius** Host. fl. austr. II. p. 217. — Exsicc.: Sint. it. thessal. n. 691. —

Caule erecto, simplici, puberulo, in spicam crassam abeunte; foliis glabriusculis, pinnatipartitis, laciniis divaricatis, ovatis, lobato-dentatis, dentibus inermibus; bractea ovato-lanceolata, lateraliter longe dentato-spinosa, in cuspidem subrecurvam, pungentem abeunti; bracteolis lineari-subulatis, puberulis, acerosis, bractea brevioribus; calycis glabriusculi lacinia superiore obovato-oblonga, violacea, apice denticulata, inferiore breviore et angustiore, apice denticulata, lateralibus multo brevioribus, obovato-orbiculatis, scariosis, glabris; corollae albidae labio glabro, obtuse trilobo.

In declivibus vallis Negerli pr. Chaliki in Pindo tymphaeo (Sint.). — Jun. Jul. ♃

b. Foliorum laciniae longe lanceolatae, spinoso-dentatae.

2. **A. Caroli Alexandri** Haussk. in Mitth. thür. bot. Ver. 1886 p. 62, in Reg. et Engl. Gartenfl. XXXV. p. 626 f. 73—75, in symb. p. 178. — Exsicc.: Sint. it. thessal. n. 442.

Caule erecto, simplici, puberulo, in spicam crassam abeunte; foliis glabriusculis, pinnatipartitis, laciniis divaricatis, longe lanceolatis, lobato-dentatis, dentibus spina subpungente terminatis; bractea ovato-lanceolata, lateraliter longe dentato-spinosa, in cuspidem subrecurvam, spinosam abeunti; bracteolis lanceolatis, puberulis, acerosis, bracteae subaequilongis; calycis puberuli lacinia superiore oblongo-spathulata, violacea, apice spinuloso-denticulata, inferiore breviore et angustiore, apice spinuloso-denticulata, lateralibus multo minoribus, a basi ovati lanceolatis, sensim in spinam excurrentibus, hirtulis; corollae albidae labio glabro, obtuse trilobo. — Differt a praecedente foliis spinoso-dentatis, a sequente spinis foliorum mollioribus, ab ambobus floribus majoribus et praesertim calycis laciniarum lateralium forma.

β. **tenuissimus** Haussk. in Reg. et Engl. l. c. p. 633. — Foliis brevioribus, in lacinias angustas pinnatipartitis.

γ. **minor** Haussk. l. c. . — Caule spithameo vel dodrantali, densius villoso, spica subcapitata, pauciflora.

In regione montana Pindi Agraphae in nemorosis quercinis, praecipue ad monasterium Korona et in toto tractu ad Pindum tymphaeum usque (Haussk.). — Maio, Jul. ♃

3. **A. spinosus** L. sp. p. 939; S. et S. pr. I. p. 141, Fl. gr. VII. p. 10 t. 611; Pieri corc. fl. p. 86; Sieb. avis rem. p. 5; Ch. et

B. exp. p. 179, Fl. pelop. p. 40; Marg. et R. fl. Zante p. 77; Fraas fl. class. p. 185; Weiss in z. b. G. p. 782; Raul. cret. p. 842; Bois. fl. or. IV. p. 522; Heldr. fl. cephal. p. 56, Fl. Aegina p. 383, chlor. Mykon. p. 248; Spreitz. in z. b. G. 1887 p. 665; Form. in D. bot. Monat. 1890 p. 25, in Ver. Brünn 1895 p. 33; Hal. Beitr. fl. Epir. p. 34, Beitr. fl. Aetol. p. 9, Beitr. fl. Thessal. p. 16; Haussk. symb. p. 178. — *A. spinosissimus* Pers. syn. II. p. 179; Sieb. avis p. 4; Raul. cret. p. 842; Gelmi in bull. soc. bot. ital. 1899 p. 451; qui ab *A. spinoso* vix differre videtur. — *A. mollis* Dallap. prosp. p. 87, non L. sp. p. 939, qui foliorum dentibus inermibus et calycis laciniis lateralibus ovato-acutis discedit et in ditione nostra vix occurrit. — *A. spinulosus* Friedr. Reise p. 277, non Host fl. austr. II. p. 217, qui probabiliter hybrida inter *A. mollem* et *A. spinosum* et ab ulteriore dentibus foliorum brevibus, triangularibus nec lanceolato-acuminatis, molliter spinosis discedit (cf. Vis. fl. dalm. II. p. 227). — Exsicc.: Heldr. herb. norm. n. 301 et 1166; Orph. fl. gr. n. 869; Rev. pl. cret. n. 145; Sint. et Bornm. it. turc. n. 1436. —

Caule erecto, simplici, puberulo, in spicam crassam abeunte; foliis glabriusculis, elevatim nervosis, pinnati- vel bipinnati-partitis, laciniis primariis divaricatis, longe lanceolatis, secundariis triangulari-lanceolatis, apice et lateraliter in spinas pungentes abeuntibus, bractea ovato- vel oblongo-lanceolata, lateraliter longe dentato-spinosa, in spinam validam subrecurvam abeunte; bracteolis lanceolato-subulatis, pubescentibus, acerosis, bracteae subaequilongis; calycis puberuli lacinia superiore oblongo-spathulata, violacea, integra vel apice spinuloso-denticulata, inferiore breviore et angustiore, apice spinuloso-denticulata, lateralibus multo brevioribus, obovato-orbiculatis, scariosis, glabriusculis; corollae albidae labio glabro, apice obtuse trilobo.

β. **thessalus** Form. in Ver. Brünn 1896 p. 62, 1897 p. 45. — Caulis, folia, bracteae, bracteolae et calycis laciniae dense hirsutae, corollae labium parce puberulum. — Exsicc.: Form. pl. gr. a. 1889.

In collibus apricis, cultis derelictis regionis inferioris et submontanae, per totam Graeciam; — β. frequens in Thessalia (Form.). — Maio, Jul. ♃

LXXXV. Ordo. **Verbenaceae** Juss. ann. mus. VII. p. 63.

1. Vitex L. gen. n. 790.

1. **V. agnus castus** L. sp. p. 938; S. et S. pr. I. p. 441, Fl. gr. t. 541; Pieri corc. fl. p. 85; Sieb. avis p. 4, rem. p. 5; Dallap. prosp. p. 87; Urv. enum. p. 73; Ch. et B. exp. p. 179, Fl. pelop. p. 40; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 264 et 275; Fraas fl. class. p. 185; Clem. sert. p. 79; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 842; Bois. fl. or. IV. p. 535; Heldr. fl. cephal. p. 57, in Sitzungsb. acad. Wiss. Berl. 1883 p. 7, Fl. Aegina p. 383,

chlor. Thera p. 18; Chloros Waldverh. p. 38; Spreitz. in z. b. G. 1877 p. 665, 1890 p. 297; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Epir. p. 34, Beitr. fl. Thessal. p. 16; Form. in D. bot. Mon. 1891 p. 27, in Ver. Brünn 1895 p. 34, 1896 p. 65, 1897 p. 49; Haussk. symb. p. 178. — Exsicc.: Rev. pl. cret. n. 185.

Frutex erectus, 1—3 m. altus; ramis tomentellis; foliis longe petiolatis, digitatim 5—7 foliolatis; foliolis lanceolatis, supra obscure viridibus, subtus incano-tomentellis; floribus in cymas compactas, subsessiles, verticillastros formantes dispositis, verticillastris in racemos longos terminales interruptos aggregatis; bracteis parvis, linearibus; calyce tomentoso, dentibus triangularibus, obtusis; corolla campanulato-bilabiata, lilacina, violacea vel raro alba, calyce triplo longiore; drupa sphaerica.

In dumetis, alveis, ad ripas regionis inferioris totius ditionis. — Jun. Aug. ♄

2. Verbena L. gen. n. 32.

1. **V. officinalis** L. sp. p. 20; S. et S. pr. I. p. 402; Pieri corc. fl. p. 4; Dallap. prosp. p. 6; Ch. et B. exp. p. 164, Fl. pelop. p. 36; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 263 et 273; Fraas fl. class. p. 186; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 841: Bois. fl. or. IV. p. 534; Heldr. fl. cephal. p. 56, chlor. Mykon. p. 248; Spreitz. in z. b. G. 1890 p. 297; Form. in D. bot. Mon. 1891 p. 27, in Ver. Brünn 1895 p. 34, 1896 p. 65, 1897 p. 49; Hal. Beitr. fl. Epir. p. 34, Beitr. fl. Thessal. p. 16; Haussk. symb. p. 178, cum var. *grandiflora* (f. floribus fere duplo majoribus). — Icon: Fl. dan. t. 628. — Exsicc.: Sint. it. thessal. n. 981.

Caule erecto, ad angulos scabridulo, a medio paniculato-ramoso; foliis oblongis vel oblongo-lanceolatis, strigulosis, incisis vel laciniatis vel lyrato-pinnatifidis, in petiolum brevem contractis; floribus parvis, subsessilibus, bractea oblongo-lanceolata, calyce breviore suffultis; calycis strigosi dentibus acutis; corolla lilacina.

In ruderatis, ad vias, muros passim per totam Graeciam. — Maio. Sept. ⊙

2. **V. supina** L. sp. p. 21; Raul. cret. p. 842; Bois. fl. or. IV p. 534; Heldr. chlor. Parn. p. 25; Haussk. symb. p. 178. — Icon: Fl. gr. t. 554. — Exsicc.: Heldr. it. thessal. a. 1886.

Caule procumbente, rarius erecto, undique strigoso-hispido, valde ramoso; foliis strigosis, canescentibus, in petiolum attenuatis, infimis oblongis, incisis, ceteris in lacinias ovato-oblongas, integras vel dentatas bipinnatisectis; floribus minimis, subsessilibus, bractea lanceolata, calyce breviore suffultis; calycis strigosi dentibus brevibus, corolla lilacina. — Foliis mollioribus, tenuius dissectis, spicis brevioribus et floribus minoribus a praecedente distincta.

In ruderatis, arenosis hyeme inundatis, rare. Thessalia: pr. Trikala, Sophades, Karditza (Haussk.); mt. Parnassus pr. Livadi (Heldr.); Creta: pr. Aradhena (Raul.). — Jun, Jul. ⊙

1 × 2. **V. officinalis × supina.** — *V. Haussknechtii* Heldr. it. IV. pr. Thessal. a. 1885. — *V. adulterina* Haussk. symb. p. 178. — Differt a. *V. officinali* foliis minoribus, magis dissectis et indumento canescente; a *V. supina* foliis minus dissectis et indumento tenuiore. — Thessalia: pr. Karditza (Heldr.).

3. Lippia L. gen. n. 781.

1. **L. nodiflora** L. sp. p. 20; S. et S. pr. I. p. 401, Fl. gr. t. 553; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 164, Fl. pelop. p. 36; (*Verbena*); Rich. in Michx. fl. bor. amer. II. p. 15; Raul. cret. p. 842; Bois. fl. or. IV. p. 532. — *Zapania nodiflora* Lam. ill. p. 248 t. 17. — *L. repens* Spreng. syst. II. p. 752; Haussk. symb. p. 178; f. bracteis obtusis, haud vel obsolete mucronatis. — Exsicc.: Pichl. pl. gr. a. 1876.

Adpresse strigulosa; caulibus procumbentibus, radicantibus; foliis cuneato-spathulatis, breviter petiolatis, a medio ad apicem usque argute serratis; floribus parvis, in capitula ovoidea, demum cylindrica, pedunculo axillari, folio longiore insidentia congestis; bracteis imbricatis, obovatis, concavis, mucronatis vel muticis, erosulis; calyce bipartito; corolla exigua, rosea; fructu ovoideo.

In arenosis, inundatis, maritimis, paludosis regionis inferioris. Acarnania: pr. Actium (Bald.); Attica: pr. Tatoi (Haussk.); Messenia: pr. Skala (Chaub.); Laconia: in planitie Helos (Heldr.), pr. Marathonisi (Pichl.); Creta: pr. Platania (Sibth.), Canea, Kladiso (Raul.); insula Syra (Chaub.). — Jun. Oct. ♃

LXXXVI. Ordo. Labiatae Juss. gen. p. 110.

Dispositio tribuum generumque:

A. Corolla ob labium superius brevissimum subuuilabiata.

1. Tribus. *Ajugoideae* Benth. lab. p. 657. — Stamina (in nostris) 4, adscendentia, approximata, inferiora longiora; nuculae siccae.

a. Corollae tubus intus saepe piloso-annulatus; labium superius brevissimum, emarginatum vel fissum, inferius elongatum, patens trifidum.

1. Ajuga L.

b. Corollae tubus intus exannulatus; labium superius bipartitum, laciniis margini labii inferioris impositis, inferius trilobum, lobo medio maximo.

2. Teucrium L.

B. Corolla bilabiata.

I. Nuculae carnosae, subbaccatae.

2. Tribus. *Prasieae* Benth. lab. p. 646. — Stamina 4, adscendentia, approximata, inferiora longiora.

3. Prasium L.

II. Nuculae siccae.

1. Stamina adscendentia approximata vel recta divergentia.

§ Stamina 2.

3. Tribus. *Monardeae* Benth. lab. p. 190. — Stamina adscendentia.

a. Connectivum elongatum, filiforme, transverse in filamentum articulatum postice adscendens et loculum fertilem ferens, antice porrectum vel deflexum, loculum cassum, rarius polliniferum ferens vel nudum.

4. Salvia L.

b. Connectivum parvum; antherae loculi divaricati, confluentes.

α. Calyx ovato-campanulatus, bilabiatus, fauce nudus; corollae labium inferius trilobum, lobis lateralibus oblongis erectis, medio maximo concavo dependente.

5. Rosmarinus L.

β. Calyx tubulosus, bilabiatus, fauce villosus; corollae labium inferius trifidum, lobis rotundatis, medio longiore emarginato.

6. Zizyphora L.

§§ Stamina 4.

4. Tribus. *Stachydeae* Benth. in DC. pr. XII. p. 407. — Stamina sub labio superiore adscendentia, approximata, inferiora longiora.

1. Subtribus. *Scutellarieae* Benth. in DC. pr. XII. p. 407. — Calyx bilabiatus, non inflatus, labiis in fructu clausis; stamina exserta.

a. Calycis labia integra, superius squamigerum.

7. Scutellaria L.

b. Calycis labia dentata, superius non squamigerum.

8. Prunella L.

2. Subtribus. *Marrubieae* Endl. gen. p. 627. — Calyx bilabiatus vel 5 dentatus, non inflatus, fructifer non clausus; stamina inclusa.

a. Nuculae apice rotundatae.

9. Sideritis L.

b. Nuculae apice truncatae.

10. Marrubium L.

3. Subtribus. *Melitteae* Willk. et Lange pr. fl. hisp. II. p. 460. — Calyx bilabiatus, inflatus, fructifer apertus; stamina exserta.

a. Calyx obconicus, labio superiore breviter 2—3 dentato, inferiore bifido; corollae tubus longe exsertus; nuculae trigonae, apice rotundatae.

11. Melittis L.

b. Calyx maximus, campanulatus, labio superiore integro, inferiore denticulato; nuculae triquetrae, apice truncatae.

12. Moluccella L.

4. Subtribus. *Lamieae* Endl. gen. p. 624. — Calyx 5 dentatus, non inflatus, fructifer apertus; stamina exserta.

a. Labium corollae superius valde concavum, galeatum.

α. Galea lateraliter compressa, dorso carinata, extus plus minus tomentosa.

13. Phlomis L.

β. Galea dorso non carinata, extus glabra aut margine solum ciliata.

14. Lamium L.

b. Labium corollae superius concaviusculum vel subplanum, erectum.

× Nuculae apice rotundatae.

○ Antherae loculi paralleli vel subparalleli; stamina inferiora demum ad latere non recurvata.

15. Betonica L.

○○ Antherae loculi divergentes vel divaricati.

. Stamina inferiora demum ad latera recurvata; antherae loculi rima longitudinali dehiscentes.

16. Stachys L.

.. Stamina inferiora demum ad latera non recurvata; antherae loculi rima longitudinali propria dehiscentes.

17. Ballota L.

×× Nuculae apice plano triangulari truncatae.

18. Leonurus L.

5. Tribus. *Nepeteae* Benth. lab. p. 462. — Stamina sub labio superiore adscendentia, approximata, superiora longiora.

a. Antherae per paria in crucem dispositae.

19. Glechoma L.

b. Antherae non in crucem dispositae.

20. Nepeta L.

6. Tribus. *Satureieae* Benth. lab. p. 324, s. ampl. — Stamina distantia, atque vel superne divergentia vel sub labio superiore conniventia, inferiora longiora.

a. Stamina superne plus minus conniventia.

α. Calyx bilabiatus.

× Antherae loculi divergentes, apice connexi.

21. Melissa L.

×× Antherae loculi divergentes, apice sejuncti.

22. Calamintha Tourn.

β. Calyx 5dentatus.

× Calycis tubus anguste cylindricus, 13—15nervius.

23. Micromeria Benth.

×× Calycis tubus campanulatus, 10nervius.

24. Satureia L.

b. Stamina superne patula vel divergentia.

α. Calyx campanulatus vel turbinatus.

25. Origanum L.

β. Calyx a dorso plano-compressus, subanceps.

26. Thymbra L.

γ. Calyx ovoideus.

27. Thymus L.

2. Stamina declinata.

7. Tribus. *Ocymoideae* Benth. lab. p. 1. — Stamina 4, inferiora longiora.

28. Lavandula L.

C. Corolla subregularis, limbo 4—5fido.

8. Tribus. *Menthoideae* Benth. lab. p. 152. — Stamina 2—4, distantia, recta, superne divergentia.

a. Stamina 2.

29. Lycopus L.

b. Stamina 4.

30. Mentha L.

1. Tribus. **AJUGOIDEAE** Benth. lab. p. 657.

1. **Ajuga** L. gen. n. 705.

1. Sectio. *Bugula* Benth. lab. p. 692. — Verticillastri multiflori; corolla saepissime coerulea, raro rosea vel alba, nunquam flava.

1. **A. reptans** L. sp. p. 561; S. et S. pr. I. p. 389; Pieri corc. fl. p. 73; Ch. et B. exp. p. 161, Fl. pelop. p. 35; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 282; Ung. Reise p. 127; Bois. fl. or. IV. p. 798; Gelmi in bull. soc. bot. it. 1889 p. 451; Hal. in ö. b. Z. 1896 p. 16, 1895 p. 325; Form. in Ver. Brünn 1896 p. 72; Haussk. symb. p. 33. — Icon: Fl. dan. t. 925. — Exsicc.: Sint. it. thessal. n. 35; Dörfl. fl. gr. n. 295.

Rhizomate truncato, stolones elongatos edente; caule erecto, bifariam piloso; foliis obovato-oblongis oblongisve, crenatis vel dentatis, glabriusculis, inferioribus in petiolum attenuatis, ceteris sessilibus, sensim in bracteas abeuntibus; calycis laciniis lanceolatis, villosis, tubo aequilongis; corollae tubo exserto, recto; staminibus exsertis; nuculis reticulato-rugosis.

In herbidis, nemorosis, ad rivulos regionis inferioris et subalpinae. Thessalia: pr. Klinovo (Form.), Bezula, Sermeniko, in oropedio Neuropolis (Gaussk.) in Pindo, pr. Kissos (Heldr.) et Lechonia (Sint.) in mt. Pelion; Aetolia: pr. Krioneri, Naupactos (Reis.), mt. Korax pr. Musinitza (Leon.); Peloponnesus: mt. Kyllene (Orph.), mt. Alvena et Smirna pr. Krysthena Arcadiae (Friedr.), mt. Kupe, pr. Messene (Chaub.), mt. Taygetos (Heldr.); Zante: pr. Argassi (Marg.); Corcyra (Pieri.): pr. urbem (Sagb.). — Mart. Maio. ♃

2. **A. orientalis** L. sp. p. 561; Ung. Reise p. 127; Spreitz. in z. b. G. 1877 p. 715; Bois. fl. or. IV. p. 800; Gelmi in bull. soc. bot. it. 1889 p. 451; Hal. Beitr. fl. Achaia p. 28, in ö. b. Z. 1897 p. 97; Haussk. symb. p. 33. — Icon: Dill. hort. elth. t. 53. — Exsicc.: Orph. fl. gr. n. 1113; Heldr. herb. dimorph. n. 41; Baen. herb. europ. n. 4198.

Rhizomate truncato, estolonoso; caule erecto, foliisque villoso; foliis oblongis, sinuato-crenatis vel inciso-lobulatis, inferioribus in petiolum attenuatis, superioribus sessilibus, sensim in bracteas ovatas, lobulatas abeuntibus; calycis laciniis lanceolatis, villosis, tubo duplo longioribus: corollae tubo breviter exserto, torto, limbo resupinato; staminibus inclusis; nuculis reticulato-rugosis.

β. **aenesia** Heldr. fl. cephal. p. 59; Spreitz. in z. b. G. 1877 p. 715. — Bracteis pulchre violaceo-purpurascentibus, profunde paucidentatis; corolla intensius coerulea, immaculata. — Exsicc.: N. v.

In pascuis, herbidis, dumosis regionis inferioris et subalpinae. Thessalia: mt. Pelion pr. Milies (Heldr.); Sporadum insula Skiathos (Leon.); Attica: mt. Pentelicon (Orph.), Parnes, Cithaeron (Heldr.); Boeotia, mt. Parnassus (Sprun.); Acrocorinthus (Haussk.); Laconia: pr. Megali Anastasova (Zahn); Elis: mt. Olenos (Hal.); Corcyra: mt. Panto-

crator (Spreitz.); — β. Cephalonia: mt. Aenos et Rhudi (Heldr.). — Mart. Jun. ♃

2. Sectio. *Chamaepitys* Benth. lab. p. 697. — Verticillastri biflori, raro quadriflori; corolla flava vel rosea.

a. Corollae labium superius emarginatum.

α. Folia indivisa.

3. **A. iva** L. sp. p. 563; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 161, Fl. pelop. p. 35; (*Teucrium*); Schreb. unilab. p. 25; S. et S. pr. I. p. 389; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 277; Fraas fl. class. p. 172; Ung. Reise p. 127; Weiss in z. b. G. 1869 p 743; Raul. cret. p. 841; Bois. fl. or. IV. p. 802; Heldr. fl. cephal. p. 59; Gelmi in bull. soc. bot. ital. 1889 p. 451; Haussk. symb. p. 33. — Icon: Rchb. t. 34 f. 3. — Exsicc.: Heldr. pl. fl. hellen. a. 1899.

Canescens vel villosa, a basi ramosa, caespitosa; caulibus prostratis vel adscendentibus, foliosis; foliis sessilibus, lineari-oblongis linearibusve, integris vel utrinque remote 1—3 dentatis; floribus in axillis foliorum superiorum sessilibus; calycis dentibus lanceolatis, tubo brevioribus; corolla rosea, calyce triplo longiore; nuculis reticulato-rugosis. — Planta tota moschum redolens, occurrit saepe floribus apetalis (*f. cleistogama* Heldr. fl. Aegina p. 383).

β. **pseudoiva** Rob. et Cast. ap. DC. fl. fr. suppl. p. 395; Heldr. fl. Aegina p. 383, chlor. Thera p. 19; pro sp.; Benth. in DC. pr. XII. p. 600; Weiss in z. b. G. 1869 p. 743. — Icon: S. et S. fl. gr. t. 525. — Flores lutei. — Exsicc.: Heldr. herb. norm. n. 1465.

In arvis, olivetis, locis aridis regionis inferioris et submontanae. Attica: pr. Athenas, Liosia, mt. Pentelicon (Heldr.), Eleusis, Kamariza Laurii (Haussk.); insula Aegina (Heldr.); Argolis: pr. Argos (Heldr.), Poros (Friedr.); Messenia (Chaub.); Elis: pr. Bartholomeo (Heldr.); Cycladum insula Syra (Weiss), Thera (Heldr.); Creta (Sibth.): pr. Candia (Raul.); Zante (Sibth.); Cephalonia: pr. Kundurata, Pessada (Heldr.); Corcyra (Gelmi). — Maio, Oct. ♃

β. Folia in lacinias lineares tripartita.

4. **A. chamaepitys** L. sp. p. 562; Dallap. prosp. p. 80; Ch. et B. exp. p. 161, Fl. pelop. p. 35; (*Teucrium*); Schreb. unilab. p. 24; S. et S. pr. I. p. 389; Friedr. Reise p. 282; Clem. sert. p. 78; Bois. fl. or. IV. p. 802; Heldr. fl. cephal. p. 59; Form. in D. bot. Monat. 1891 p. 31; Bald. riv. coll. bot. alb. 1895 p. 66, 1896 p. 88. — Icon: Fl. dan. t. 733.

Pilosa; caulibus diffusis, filiosis; foliis in lacinias anguste lineares tripartitis, floralibus flores superantibus; calycis dentibus lanceolatis, tubo sublongioribus; corolla lutea, calyce triplo longiore; nucularum areolis plerisque pentagonis.

In arvis, cultis regionis inferioris et montanae. Epirus: pr. Janina, mt. Micikeli (Bald.); Thessalia: pr. Han Kuraneos in valle Penei, pr. Kalabaka, Koryza, Volo (Form.); Attica: mt. Pentelicon (Clem.); Peloponnesus (Sibth.); mt. Alvena et Smirna pr. Krysthena Arcadiae (Friedr.); Archipelagus (Sibth.); Cephalonia (Dall.); loca plurima, forsitan omnia probabiliter ad sequentem spectant. — Maio, Oct. ⊙ N. v.

5. **A. chia** Poir. dict. suppl. II. p. 772 (*Teucrium*); Schreb. unilab. p. 25; S. et S. pr. I. p. 389, Fl. gr. VI. p. 21 t. 524; Fraas fl. class. p. 172; Bois fl. or. IV. p. 802; Form. in Ver. Brünn 1897 p. 54; Haussk. symb. p. 33. — Exsicc.: Orph. fl. gr. n. 533; Heldr. herb. norm. n. 202 et 869, herb. dimorph. n. 40.

Albo-hispida; caulibus diffusis, foliosis; foliis in lacinias latiuscule lineares tripartitis, floralibus floribus saepius subbrevioribus; calycis dentibus lanceolatis, tubo brevioribus; corolla lutea purpureo-striata vel omnino purpurea, calyce quintuplo longiore; nuculis praeter areolas paucas terminales pentagonas transverse corrugatis. — A praecedente vix differre videtur, nisi corollis duplo majoribus et indumento densiori.

β. **intermedia** Bois. et Orph. in Bois. fl. or. IV. p. 802, solum nomen; Heldr. herb. norm. n. 1466. — Corolla parva, calyce triplo longiore. — Medium tenet inter *A. chiam* et *chamaepityn*.

In campis, cultis, ad vias regionis inferioris et submontanae. Thessalia: pr. Malakasi, Klinovo, Korona, Meteora, Pharsalus (Haussk.), Neraida (Form.); Attica: ubique circa Athenas; Aetolia: ad radices mt. Chalkis ad sinum Patranum (Hal.); Peloponnesus: mt. Xerovuni Alagoniae, pr. Kalamata (Zahn); Archipelagus (Sibth.); et probabiliter loca omnia ad speciem praecedentem enumerata. — Mart. Jun. ⊙, sec. Bois. ♃

b. Corollae labium superius fissum.

6. **A. Piskoi** Deg. et Bald. egy uj ajuga faj. 1896 p. 1, in Bald. riv. coll. bot. Alb. 1896 p. 88. — *A. Halácsyana* Zahlbr. in z. b. G. 1896 p. 447. — Exsicc.: Bald. it. alb. epir. n. 136.

Viridis; caulibus pluribus, erectis, ramosis, parce hispidulis; foliis oppositis breviter petiolatis, ovato-oblongis, acutis, parce hispidulis, margine ciliatis, inferioribus grosse remoteque dentatis, ceteris integris, floralibus flores multoties superantibus; verticillastris bifloris; floribus breviter pedicellatis; calycis laciniis lineari-lanceolatis, hispidulis, tubo multo longioribus; corolla rosea, purpureo-venosa, calyce subtriplo longiore. — Species egregia, foliis 6—7 cm. longis, 3—4 cm. latis, corolla 2—3 cm. longa.

In humidis pr. Diovisda distr. Ljaskovik Albaniae australis (Bald.), jam extra ditionem nostram, in Epiro boreali inquirenda. — Jul. ♃

2. Teucrium L. gen. n. 706.

Dispositio specierum:

1. Sectio. *Teucris* Benth. lab. p. 665. — Pedunculi oppositi, 1—3 flori, racemosi vel paniculati; calyx aequaliter 5 dentatus.

1. T. aroanium Orph. **2. T. creticum** L. **3. T. brevifolium** Schreb.

2. Sectio. *Stachyobotrys* Benth. lab. p. 672. — Flores sparsi vel irregulariter subverticillati, in racemum simplicem dispositi; calyx 5 dentatus, dente supremo ovato, latiore.

4. T. Halácsyanum Heldr. **5. T. cuspidatum** Celak.

3. Sectio. *Scorodonia* Benth. lab. p. 674. — Pedunculi oppositi, uniflori, secundi, racemosi; calyx 5 dentatus, dente supremo ovato, latiore.

6. T. massiliense L. **7. T. scorodonia** L.

4. Sectio. *Scordium* Benth. lab. p. 678. — Flores in verticillastros 2—6 floros, axillares, distantes dispositi; calyx subaequaliter 5 dentatus, basi infra gibbus.

8. T. scordioides Schreb.

5. Sectio. *Chamaedrys* Benth. lab. p. 680. — Flores in verticillastros 2—6 floros, racemum laxum formantes dispositi; calyx subaequaliter 5 dentatus..

a. Herbaceum.

9. T. chamaedrys L.

b. Suffruticosa.

α. Corolla flava.

10. T. flavum L.

β. Corolla purpurea.

11. T. divaricatum Sieb. **12. T. microphyllum** Desf.

6. Sectio. *Polium* Benth. lab. p. 684. — Flores in verticillastros, capitula globosa vel ovata terminalia formantes, dispositi; calyx dentibus subaequalibus vel inferioribus subangustioribus et acutioribus.

a. Folia integra; corolla calyce subduplo longior.

13. T. montanum L.

b. Folia crenata (cf. T. polium v. capitatum).

α. Capitula laxiflora; corolla calyce subduplo longior.

14. T. cuneifolium S. et S. **15. T. alpestre** S. et S.

β. Capitula densiflora; corolla calyce parum longior.

16. T. polium L.

1. Sectio. *Teucris* Benth. lab. p. 665.

1. **T. aroanium** Orph. in Bois. diagn. ser. 2 IV. p. 55, Fl. or. IV. p. 807. — Exsicc.: Orph. fl. gr. n. 102.

Suffruticosum, ramosissimum, caespitosum; caudiculis duris, nudis, radicantibus; ramis novellis brevibus, herbaceis, adscendentibus, foliosis, adpresse canis; foliis parvis, obovatis vel ovato-oblongis, integris, in petiolum brevem attenuatis, utrinque adpresse sed subtus intensius canis; floribus oppositis, pedicellatis, in racemum ovatum, pauciflorum dispositis, bracteis ellipticis, acutis, calyce brevioribus suffultis; calycis campanulati, minute glandulosi dentibus ovato-triangularibus, tubo aequilongis; corolla coerulescente, calyce multo longiore, laciniis oblongis, lateralibus glabriusculis, inferiori multo majori, subtus papillis crispulis hirta; staminibus longissime exsertis. — Stirps pulcherrima, foliis 5—15 mm. longis, corolla circa 15 mm. longa.

In regione superiori mt. Chelmos Achaiae pr. Stygem, ubi rupes laetis vestit tapetibus (Orph.). — Jul. ♄

2. **T. creticum** L. sp. p. 563; Raul. cret. p. 839. — *T. hyssopifolium* Schreb. unilab. p. 28. — *T. rosmarinifolium* Lam. dict II. p. 693. — Icon: Fl. gr. t. 529. — Exsicc.: Sint. et Rigo. it. cypr. n. 565 (Cyprus).

Fruticosum; ramis virgatis, elongatis, demum glabrescentibus; foliis supra viridibus, subtus adpresse canis, margine revolutis, integris, inferioribus crebris, elliptico-linearibus, inferne attenuatis, ceteris anguste-linearibus, remotis; floribus solitariis ternisve, pedicellatis, in racemum elongatum, interruptum dispositis, bracteis calyce aequilongis vel longioribus suffultis; calycis campanulati, adpresse cani dentibus triangulari-lanceolatis, acutis, tubo brevioribus; corolla purpurascenti-coerulea, pubescente, calyce subduplo longiore, laciniis lateralibus oblongis, inferiori multo majori; staminibus exsertis, corolla subbrevioribus.

In saxosis regionis inferioris. Indicatur a Linnaeo in Creta, ubi tamen sec. Bois. fl. or. IV. p. 806 non crescit. — Maio, Jul. ♃ N. v.

3. **T. brevifolium** Schreb. unilab. p. 27; S. et S. pr. I. p. 397, Fl. gr. VI. p. 24 t. 528; Urv. enum. p. 63; Ch. et B. fl. pelop. p. 36; Raul. cret. p. 839; Heldr. in Gartenfl. 1873 p. 323; Bois. fl. or. IV. p. 807; Haussk. symb. p. 31. — *T. creticum* Lam. dict. II. p. 592; Sieb. avis. p. 4, rem. p. 5; non L. — Exsicc.: Heldr. herb. norm. n. 965; Rev. pl. cret. n. 205, in Baen. herb. europ. n. 5241.

Fruticosum, ramosissimum, ramis foliosis, saepe intricatis; foliis adpresse canescentibus, subsericeis, oblongis linearibusve, margine revolutis, integris, floralibus conformibus, calyces superantibus; floribus oppositis, pedicellatis, racemos breves formantibus; calycis campanulati adpresse tomentelli dentibus oblongis, obtusiusculis, tubo subbrevioribus; corolla coerulea, purpureo-striata, pubescente, calyce triplo longiore, laciniis oblongis, inferiori multe majore; staminibus exsertis, corolla brevioribus. — Frutex 1—2 pedalis.

In aridis, collibus saxosis, maritimis, rare. Attica: pr. Laurion Sunium (Heldr.); insula Poros, Naxos (Wiedem.), Melos (Urv.); Creta (Schreb.): ad promontorium Meleka (Raul.), pr. Topolia distr. Kissamos (Rev.). — Febr. Maio. ♃

2. Sectio. *Stachyobotrys* Benth. lab. p. 672.

4. **T. Halácsyanum** Heldr. in ö. b. Z. 1879 p. 241; Bois. fl. or. suppl. p. 363; Hal. Beitr. fl. Aetol. p. 9 t. 1. — Exsicc.: Heldr. pl. fl. hellen. a. 1878; Hal. it. gr. sec. a. 1893; Bickn. pl. Corc. a. 1891.

Suffruticosum, caespitosum, molliter villoso-tomentosum; ramis e caudice brevi, crasso, lignoso numerosis, flexuosis, fragillimis; foliis petiolatis, e basi truncata vel in superioribus breviter attenuata ovatis vel subrhombeis, obtusis, crenatis; floribus sparsis, pedicellatis, racemos subsecundos, densos formantibus; bracteis linearibus vel lanceolatis, calycem aequantibus vel eo brevioribus; pedicellis calyce aequilongis vel paulo brevioribus; calycis campanulati dentibus triangulari-lanceolatis, acutis, superiore latiore; corolla violacea, pubescente, calyce duplo

longiore, laciniis lateralibus lineari-oblongis, inferiori multo majore; staminibus exsertis, corolla brevioribus. — Species distinctissima, typice caespites humiles, ramis 3—5 pollicaribus, foliis cum petiolo 10—15 mm. longis, racemis 1—2 pollicaribus formans, vidi tamen in umbrosis specimina (*f. laxa* Hal. it. gr. secund. a. 1893) caulibus ad 50 cm. longis valde intricatis.

In rupestribus regionis inferioris. Aetolia: ad radices mt. Taphiassos (Heldr.) et Chalkis (Hal.) ad sinum Patranum; Achaia: in faucibus Diakophto (Heldr.); Corcyra: pr. Palaeokastrizza (Bicknell). — Maio, Jun. ♄

5. **T. cuspidatum** Celak. in bot. Centralbl. 1883 p. 222; Bois. fl. or. suppl. p. 362. — Huc forsan: *T. coloratum* Scheele in Flora 1843 p. 568.

Suffruticosum, ramosissimum, hirsutum, superne subvillosum; ramis adscendentibus; foliis petiolatis, e basi truncata vel subcuneata ovatis, obtusis, crenatis, floralibus conformibus, minoribus, superioribus bracteaeformibus, ovato-lanceolatis, cuspidatis in petiolum brevem spathulato-attenuatis, calyce brevioribus; floribus sparsis, pedicello calyce dimidio breviore suffultis; calycis campanulati dentibus triangulari-lanceolatis, cuspidatis, supremo late breviterque ovali; corolla parva, rosea, calyce parum longiore; staminibus parum exsertis. — Species mihi ignota, e descriptione in hac sectione ut videtur collocanda.

Creta: pr. Mirabello (Sieb.). — Flor. ?. ♄ N. v.

Abs. *Scutellaria cretica* L. sp. p. 600, sec. Bois. fl. or. IV. p. 811 = *Teucrium lamiifolium* Urv. enum. p. 64 (non L.) = *T. Arduini* S. et S. fl. gr. VI. p. 26 t. 581 et Raul. cret. p. 839. — Erronee ut videtur in Creta indicatur.

3. Sectio. *Scorodonia* Benth. lab. p. 674.

6. **T. massiliense** L. sp. ed. 2 p. 789; S. et S. pr. I. p. 392; Sieb. avis. p. 4, rem. p. 5; Raul. cret. p. 840; Bois. fl. or. IV. p. 812; Bald. viagg. Creta p. 87. — Icon: Jacq. hort. vind. t. 94.

Pubescens, e rhizomate indurato pluricaule; caulibus erectis vel adscendentibus, ramosis; foliis petiolatis, ovato-oblongis, crenatis, basi rotundatis vel cordatis, floribus oppositis, pedicellis calyce subaequilongis suffultis, racemum laxiusculum, secundum formantibus; bracteis ovatis lanceolatisve, calyce brevioribus; calycis glanduloso-puberuli breviter campanulati dentibus ovatis, cuspidatis, supremo majore cordato-ovato, mucronato; corollae purpureae tubo incluso; staminibus exsertis, corolla aequilongis.

In rupestribus regionis inferioris. Creta: pr. Enneachoria, Rumata, mt. Lassiti (Raul.). — Jun. Jul. ♃ N. v.

7. **T. scorodonia** L. sp. p. 564; S. et S. pr. I. p. 392; Ch. et B. exp. p. 161, Fl. pelop. p. 31. — *T. silvestre* Lam. fl. fr. II. p. 412. — *Scorodonia heteromalla* Moench. meth. p. 384. — Icon: Rchb. t. 36.

Pubescens, e rhizomate stolonifero pluricaule; caulibus erectis, simplicibus vel ramosis; foliis petiolatis, ovato-oblongis, crenatis, basi rotundatis vel cordatis; floribus oppositis, pedicellis calyce subaequilongis suffultis, racemum laxiusculum, secundum formantibus; bracteis ovalibus vel suspathulatis, pedicello aequilongis; calycis glabriusculi, breviter campanulati dentibus ovatis, cuspidatis, supremo majore, orbiculato, mucronato; corollae lutescentis tubo calyce duplo longiore; staminibus exsertis, corolla aequilongis. — Differt ab affini praecedente praesertim calyce glabriusculo, corolla lutescenti, tubo calyce duplo longiore.

In rupestribus regionis inferioris. Laconia (Sibth.); Messenia: in montosis ad fl. Neda (Chaub.); Naxos (Chaub.). — Jun. Jul. ♃ N. v.

4. Sectio. *Scordium* Benth. lab. p. 678.

8. **T. scordioides** Schreb. unilab. p. 37; S. et S. pr. I. p. 393; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 161, Fl. pelop. p. 35; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 284; Clem. sert. p. 79; Raul. cret. p. 840; Bois. fl. or. IV. p. 813; Heldr. fl. cephal. p. 59; Form. in D. bot. Monat. 1891 p. 31, in Ver. Brünn 1895 p. 37, 1896 p. 72, 1897 p. 54 *f. acuminata* („calyce lanuginoso, eglanduloso, dentibus acuminatis"); Haussk. symb. p. 31. — *T. scordium* Pieri corc. fl. p. 74; Dallap. prosp. p. 80; Fraas fl. class. p. 172; Form. in Ver. Brünn 1895 p. 37; non L. sp. p. 565, quae indumento tenuiori, stolonibus saepissime foliiferis, foliis caulinis non amplexicaulibus, ramealibus attenuatis, a basi ad medium integris, discedit. — *T. lanuginosum* Hoffm. et Link. fl. port. p. 84 t. 3. — Exsicc.: Orph fl. gr. n. 540; Sint. et. Bornm. it. turc. n. 1414; Sint. it. thessal. n. 1253.

Lanuginosum; rhizomate repente, stolones saepissime aphyllos edente; caule adscendente vel erecto, ramoso, folioso; foliis ovato-oblongis, crenatis, caulinis basi cordato-amplexicaulibus, ramealibus basi rotundatis, floralibus flores superantibus; floribus 2—4 axillaribus, pedicellatis, paniculam longam formantibus; calycis campanulati, hispidi dentibus lanceolatis, acutis, tubo duplo brevioribus; corolla purpurascente, calyce duplo longiore; staminibus subexsertis. — Subspecies australis *T. scordii* L.

In humidis regionis inferioris et montanae. Epirus: pr. Prevesa, Janina (Form.); Thessalia: frequens a Malakasi per planitiem totam usque Litochoron ad mare Aegaeum; Boeotia (Fraas); Attica: mt. Pentelicon (Clem.), ad Phaleron (Haussk.); Argolis: pr. Nauplia, Argos, Astros (Chaub.); Creta: pr. Kladiso, Candia (Raul.); Zante (Sibth.); Cephalonia: pr. Lixuri, Cranea (Dallap.), Samoli, Livadi, Pessada (Heldr.); Corcyra (Pieri): pr. Kanali, Kontokali (Form.). — Maio, Aug. ♃

5. Sectio. *Chamaedrys* Benth. lab. p. 680.

a. Herbaceum.

9. **T. chamaedrys** L. sp. p. 565; S. et S. pr. I. p. 393; Pieri corc. fl. p. 74; Dallap. p. 80; Ch. et B. exp. p. 161, Fl. pelop.

p. 35; Bois. fl. or. IV. p. 816; Heldr. fl. cephal. p. 60; Form. in D. bot. Monat. 1891 p. 31, in Ver. Brünn 1895 p. 37 *v. canescens*, 1896 p. 72 *v. canum et revolutum*, 1897 p. 54; Hal. Beitr. fl. Epir. p. 34; Haussk. symb. p. 31. — Icon: Rchb. t. 38. — Exsicc.: Sint. it. thessal. 1177.

Pubescens vel patule hirsutum; caulibus adscendentibus vel prostratis, ramosis, basi sublignosis, ramis foliosis, in racemum oblongum, foliatum abeuntibus; foliis ovatis vel oblongis, inciso-crenatis, inferioribus in petiolum brevem attenuatis, superioribus subsessilibus, floralibus minoribus, summis subintegris; verticillastris 2—6 floris; pedicellis calyce subbrevioribus; calycis rubelli dentibus lanceolatis, acuminatis, tubo subbrevioribus; corolla purpurea, calyce duplo longiore; staminibus exsertis, corolla brevioribus. — Stirps praesertim quoad indumentum satis polymorpha.

β. **glanduliferum** Haussk. symb. p. 31. — Caules pilis patentibus ad $1^1/_2$ mm. longis, brevioribus intermixtis, omnibus glanduliferis obsiti; folia ad nervos marginesque sicuti pedunculi et calyces glanduloso-pilosa. — Exsicc.: N. v.

γ. **oxyodon** Heldr. et Hal. in ö. b. Z. 1897 p. 324. — Caules breviter pubescentes; folia oblongo-lanceolata, glabriuscula, argute profundeque serrata; calycis glabri dentes ciliati. — Varietas egregia. — Exsicc.: Heldr. pl. fl. hellen. a. 1896.

In dumosis, silvaticis regionis montanae et subalpinae. Epirus: in valle Dipotami, mt. Peristeri (Form.), pr. Kalentini, Mazuki (Hal.); frequens in Thessalia a Pindó per planitiem usque ad mt. Olympum et Pelion; Aetolia: mt. Korax (Tunt.); Achaia: mt. Panachaicon, pr. Megaspilaeon (Hal.), pr. Hiero Argolidis, mt. Diaforti Arcadiae, pr. Skardamula Laconiae (Chaub.); Cycladum insula Tenos, Naxos (Chaub.); Cephalonia (Dall.); Corcyra (Pieri); — *β.* ad monasterium Korona, pr. Aivali, Pharsalus (Haussk.); — *γ.* mt. Malevo pr. Kastanitza (Leon.). Maio, Aug. ♃

b. Suffruticosa.

α. Corolla flava.

10. **T. flavum** L. sp. p. 565; S. et S. pr. I. p, 394, Fl. gr. t. 533; Ch. et B. exp. p. 162, Fl. pelop. p. 36; Marg. et R. fl. Zante p. 76; Fraas fl. class. p. 173; Clem. sert. p. 79; Raul. cret. p. 840; Bois. fl. or. IV. p. 816; Heldr. fl. cephal. p. 60; Bald. riv. coll. bot. alb. 1896 p. 88; Form. in Ver. Brünn 1897 p. 54, 1898 p. 79. — Exsicc.: Orph. fl. gr. n. 541.

Basi ramosum; ramis erectis, crispule-pubescentibus, in racemum densiusculum abeuntibus; foliis subcoriaceis, petiolatis, a basi truncata vel cuneata ovatis, crenato-dentatis, utrinque velutino-pubescentibus, subtus pallidioribus, floralibus lanceolatis, integris, acutis, calyce subbrevioribus; verticillastris 4—6 floris; calycis campanulati, pedicello eo subbreviore suffulti, hirsuti dentibus lanceolatis, acutis, tubo subbrevi-

oribus; corolla flava, pubescente, calyce $2^1/_2$ plo longiore; staminibus exsertis, corolla longioribus.

β. **leiophyllum** Celak. in bot. Centralbl. 1883 p. 219. — ? *T. flavum v. glaucum* Haussk. symb. p. 31. — Folia glabra, supra laete viridia, subtus, glaucescentia. — Exsicc.: Heldr. pl. fl. hellen. a. 1878.

In rupestribus regionis inferioris et montanae. Epirus: pr. Vromonero distr. Ljaskovik (Bald.); Thessalia: pr. Pharsalus (Heldr.), Neraida (Form.); Euboea: mt. Telethrion (Heldr.), Hagios Elias (Fraas); Phocis (Fraas): mt. Parnassus (Orph.); Attica: mt. Pentelicon (Heldr.); Peloponnesus: mt. Malevo (Fraas), Taygetos (Heldr.), pr. Selitza (Zahn), Androvista, Turia, Phigalea (Chaub.); Archipelagus (Sibth.); Creta: pr. Enneachoria, in fauce Laki, pr. Mirabello (Raul.); Zante (Sibth.); Cephalonia: pr. Argostoli, Pessada, mt. Rhudi, pr. Lixuri, Kontogianuata, Kipuria (Heldr.); Corcyra: mt. Pantocrator (Clem.); — *β.* Attica: mt. Parnes, Pentelicon (Heldr.); ? Thessalia: pr. Pharsalus (Haussk.). — Maio, Jul. ♃

β. Corolla purpurea.

11. **T. divaricatum** Sieb. avis p. 4, rem. p. 5; Raul. cret. p. 840; Bois. fl. or. IV. p. 816; Heldr. chlor. Parn. p. 26; Hal. in ö. b. Z. 1892 p. 375; Bald. viagg. Creta p. 87; Haussk. symb. p. 32. — *T. flavum β.* Willd. sp. III. p. 31; Urv. enum. p. 63. — *T. lucidum* S. et S. pr. I. p. 393, Fl. gr. t. 532; Fraas. fl. class. p. 173; non L. — *T. flavum v. humilior* Ch. et B. exp. p. 162, Fl. pelop. p. 36. — *T. flavum v. purpureum* Benth. lab. p. 682; Weiss in z. b. G. 1869 p. 743. — *T. Sieberi* Celak. in bot. Centralbl. 1883 p. 214; Bois. fl. or. suppl. p. 363. — *T. graecum* Hal. in ö. b. Z. 1896 p. 16, 1897 p. 97; Heldr. fl. Aegina p. 383 *f. minor*, Chlor. Thera p. 18; non Celak. — Exsicc.: Heldr. herb. norm. n. 290 et 868; Rev. pl. cret. n. 261, in Baen. herb. europ. n. 5242; Sint. et Bornm. it turc. n. 1411 (spec. deflorata, forsan ad *T. flavum* spect.); Dörfl. fl. aeg. n. 10.

Basi ramosum; ramis erectis vel adscendentibus, patule vel adpresse hirtis, in racemum saepius laxum interruptum abeuntibus; foliis subcoriaceis, petiolatis, a basi truncata vel cuneata ovatis, crenato-dentatis, utrinque velutino-pubescentibus, subtus pallidioribus, floralibus ovato-lanceolatis lanceolatisve, integris, acutis, calyce subbrevioribus; verticillastris 4—6 floris; calycis campanulati, pedicello eo breviore suffulti, adpresse puberuli vel patule hirsuti, saepissime purpurascentis dentibus lanceolatis, acutis, tubo subbrevioribus; corolla purpurea, pubescente, calyce duplo longiore; staminibus exsertis, corolla longioribus. — Stirps, a praecedente praesertim florum colore discedens, variat indumento e pilis longioribus patulis, uti saepissime specimina cretica (*T. Sieberi* Celak. sensu strict.), vel brevissimis crispulis, uti saepius specimina graeca (*f. vulgaris*) demonstrant, constante; rarius.

β. **graecum** Celak. in bot. Centralbl. 1883 p. 214; Bois. fl. or. suppl. p. 863; pro sp.; Haussk. symb. p. 32. — Inflorescentia tantum

hirta, caules et folia glabra. — Exsicc.: Orph. fl. gr. n. 542 (mixtum cum *f. vulgari*).

In rupestribus regionis inferioris et montanae. Thessalia: in valle Megarema in Olympo (Sint.); mt. Parnassus (Sibth.); Attica: mt. Hymettus, Lycabettus (Orph.), Parnes, Pateras, ad Piraeum (Heldr.), pr. Kamariza, Laurion (Haussk.); promontorium Icrea, insula Aegina, Poros (Chaub.); Elis: pr. Lintzi (Heldr.); Sporadum insula Peristeri (Reis.) et Pelagonisi (Leon.); Euboea (Heldr.); Cycladum insula: Syra (Weiss), St. Istad pr. Kimolos (Urv.), Antimilos (Reis.), Thera (Heldr.), Amorgos (Leon.); Creta: pr. Acroteri (Weiss), Kissamos, Malaxa (Rev.), mt. Kedros pr. Spili, pr. Avdu (Raul.), mt. Volakia (Bald.); — *β*. mt. Lycabettus (Celak.), Hymettus (Orph.), peninsula Methana pr. Vromolimni (Haussk.). — Apr. Jul. ♄

12. **T. microphyllum** Desf. choix pl. p. 18 t. 11; Raul. cret. p. 840; Bois. fl. or. IV. p. 817; Bald. viagg. Creta p. 87. — *T. quadratulum* S. et S. pr. I. p. 391, Fl. gr. VI. p. 25 t. 530, non Schreb. — *T. ramosissimum* Sieb. avis p. 4, rem. p. 5, in Flora I. p. 274, V. p. 639, non Desf. — Exsicc.: Rev. pl. cret. n. 206, in Baen. herb. europ. n. 5241; Bald. it. cret. n. 37; Dörfl. pl. cret. n. 130.

Basi ramosum; ramis tenuibus, flexuosis, prostratis vel adscendentibus, adpresse canis, in racemum brevem, pauciflorum abeuntibus; foliis minimis, petiolatis, cuneato-ovatis, crenato-dentatis, supra viridibus, subtus canis, floralibus subintegris, calyce subbrevioribus; verticillastris bifloris, calycis campanulati, pedicello eo triplo breviore suffulti, adpresse canescentis dentibus lanceolatis, acutis, tubo brevioribus; corolla rosea, pubescente, calyce duplo longiore; staminibus exsertis, corolla subaequilongis. — Ramis flexuosis, diffusis, foliis minutis et verticillastris bifloris a praecedente egregie discedit.

In saxosis regionis inferioris. Creta (Sibth.): pr. Kissamos (Rev.), Epanochorio, Akroteri, promontorium Meleka, pr. Apokorona, Pedhiada, Hierapetra, Voriza, Embaro, mt. Lassiti, pr. Anatoli (Raul.). — Maio, Jun. ♄

Obs. *T. marum* L. sp. p. 564, species maditerranea occidentalis in Ch. et B. fl. pelop. p. 36, fide schedulae mendacis Gittardii erronee pr. Armyros Peloponnesi indicatur.

6. Sectio. *Polium* Benth. lab. p. 684.

a. Folia integra; corolla calyce subduplo longior.

13. **T. montanum** L. sp. p. 565. — *T. supinum* L. sp. p. 566 (f. foliis anguste linearibus, valde revolutis).

Suffruticosum, laxe caespitosum, basi ramosissimum; caulibus adscendentibus, foliosis, cano-tomentellis; foliis coriaceis, oblongo-lanceolatis, lanceolatis linearibusve, in petiolum brevem attenuatis, margine parum revolutis, supra glabrescentibus, nitidis, viridibus, subtus adpresse cano-tomentellis, floralibus minoribus, angustioribus, magis revolutis;

floribus brevissime pedicellatis, in capitula corymbosa, densiflora congestis; calycis glabri dentibus lanceolatis, acuminato-subulatis, tubo duplo brevioribus; corolla lutescente, puberula, calyce subduplo longiore; staminibus exsertis, corolla subaequilongis.

β. **parnassicum** Celak. in bot. Centralbl. 1883 p. 153; Bois. fl. or. suppl. p. 364; Heldr. chlor. Parn. p. 26; Hal. Beitr. fl. Epir. p. 34; Bald. riv. coll. bot. Alb. 1896 p. 88. — *T. montanum* S. et S. pr. I. p. 394, Fl. gr. VI. p. 28 t. 534; Bois. fl. or. IV. p. 818; Form. in Ver. Brünn 1896 p. 73; Haussk. symb. p. 33. — *T. prostratum* Hal. Beitr. fl. Achaia p. 28, non Schur (= *T. pannonicum* Kern in ö. b. Z. 1863 p. 384), quod habitu robustiori, foliis latioribus caulibusque patule hispidis et calycis dentibus ciliatis discedit. — Caules praesertim superne dense et adpresse albo-tomentosi; folia supra adpresse pubescentia, obscure viridia, subtus cano-tomentosa, calyx adpresse pubescens. Variat (*f. lanceoatum*) follis oblongo-lanceolatis lanceolatisve, parum revolutis et (*f. lineare*) anguste linearibus, valde revolutis. Ambae formis intermediis crebris conjunguntur. — Exsicc.: Orph. fl. gr. n. 435; Heldr. herb. norm. n. 393, 1364 et 1464 (*v. modestum*, f. capitulis paucifloris, minoribus); Bald. it. alb. epir IV. n. 140 (*f. lineare*).

γ. **hirsutum** Bois. fl. or. IV. p. 819. — Caules, folia linearia vel lineari-lanceolata et calyces patule hispidi. — Exsicc.: Sint. it. or. a. 1889 n. 1886.

In rupestribus regionis subalpinae et alpinae. Planta typica in ditione non provenit; — *β.* Thessalia: mt. Zygos (Haussk.), mt. Dokimi, Mandra Hodza (Form.) in Pindo; Aetolia: mt. Vardussia (Tunt.); mt. Parnassus (Heldr.); Achaia: mt. Olenos, Panachaicon (Hal.), Chelmos (Orph.); Taygetos (Heldr.); — *f. lineare.* Epirus: ad Vromonero, mt. Vratedon (Bald.), mt. Strungala, Tsumerka (Hal.); Laconia: mt. Malevo (Orph.); — *γ.* mt. Olympus Thessalia et mt. Dirphys Euboeae (Heldr.). — Jun. Aug. ♃

b. Folia crenata (cf. *T. polium v. capitatum*).

α. Capitula laxiflora; corolla calyce subduplo longior.

14. **T. cuneifolium** S. et S. pr. I. p. 395, Fl. gr. VI. p. 30 t. 537; Sieb. avis p. 4, rem. p. 5, in Flora I. p. 272; Raul. cret. p. 841; Bois. fl. or. IV. p. 820. — Exsicc.: Heldr. herb. n. 1671.

Dense tomentoso-lanatum, ramosum; ramis crassis, inferne nudis, procumbentibus, superne adscendentibus, ramulosis, foliosis; foliis crassiusculis, cuneato-ovatis vel rotundis, basi attenuatis, crenatis, floralibus minoribus, summis integris; verticillastris paucifloris, in capitula terminalia approximatis; calycis tubulosi, dense tomentosi dentibus tubo 4plo brevioribus, obtusiusculis; corolla pubescente alba, basi flavescenti; staminibus exsertis, corolla subbrevioribus. — Indumento dense tomentoso insigne

In rupestribus regionis inferioris et montanae, rarissime. Creta: pr. Suia (Raul.), Hagia Rumeli (Heldr.), mt. Sphaciotici (Sibth.). — Maio, Jun. ♃

15. **T. alpestre** S. et S. pr. I. p. 395, Fl. gr. VI. p. 31 t. 538; Raul. cret. p. 841; Bois. fl. or. IV. p. 819; Spreitz. in z. b. G. 1890 p. 297 cum *f. laxior.* — *T. saxatile* Sieb. avis p. 4, rem. p. 5, non Cav. — Exsicc.: Rev. pl. cret. n. 262.

Patule villosulum, canescens, suffruticosum, ramosissimum; ramis vetustis tortuosis, lignosis, procumbentibus, novellis brevibus, simplicibus vel saepius opposite ramulosis; foliis minimis, subsessilibus, obovato vel rotundato-cuneatis, crenatis, floralibus caulinis similibus, summis trilobis; verticillastris paucifloris, in capitula terminalia parva approximatis; calycis tubuloso-campanulati, hirsuti dentibus tubo 2plo brevioribus, acutiusculis; corolla albida, pubescente; staminibus exsertis, corolla subbrevioribus. — Foliis minutis, 4—6 mm. longis insigne.

β. **majus** Bois. fl. or. IV. p. 819. — Rami crassiusculi, firmiores, divaricatim ramulosiores; folia submajora, obovato-oblonga, margine saepe revoluta; capitula majora, 7—12 flora, densiora. — Exsicc.: N. v.

In saxosis regionis montanae et alpinae. Creta: pr. Omalos (Rev.), Askyphos, mt. Hagios Theodoros (Spreitz.) in mt. Sphacioticis (Sibth.), pr. Drakona, Mirabello (Raul.); — *β.* pr. Norisa ad latus meridionale mt. Ida et in arenosis insulae Gaidaronisi (Heldr.). — Jun. Jul. ♄

β. Capitula densiflora; corolla calyce parum longior.

16. **T. polium** L. sp. p. 566; S. et S. pr. I. p. 394, Fl. gr. VI. p. 29 t. 535; Ch. et B. exp. p. 162, Fl. pelop. p. 36; Friedr. Reise p. 264 et 277; Fraas fl. class. p. 173; Clem. sert. p. 79; Raul. cret. p. 841; Weiss in z. b. G. 1869 p. 743; Spreitz. in z. b. G. 1877 p. 715, 1888 p. 666; Bois. fl. or. IV. p. 821; Heldr. fl. cephal. p. 60, chlor. Thera p. 18, chlor. Mykon p. 248; Form. in D. bot. Mon. 1891 p. 31, in Ver. Brünn 1895 p. 37, 1896 p. 72, 1897 p. 54; Hal. Beitr. fl. Epir. p. 34, Beitr. fl. Thessal. p. 16; Bald. in bull. herb. Bois. 1896 p. 206.

Plus minus albo-, cano- vel griseo-tomentosum, basi suffrutescens, ramosum; ramis adscendentibus, apice breviter paniculatis vel saepius corymbosis; foliis sessilibus, oblongis vel lineari-oblongis linearibusve, capitulis densifloris, subglobosis ovatisve, breviter pedunculatis; bracteis lineari-spathulatis, flores non aequantibus; calycis subsessilis, breviter campanulati dentibus tomentosis, brevibus, obtusis; corolla pubescente, calycem parum superante; filamentis exsertis, corolla subbrevioribus. — Stirps maxime polymorpha.

α. **pseudohyssopus** Schreb. unilab. p. 45; Sieb. avis p. 4; pro sp. — *T. teuthrion* Schreb. l. c. p. 36, f. pedunculis abbreviatis. — *T. polium v. vulgare* Benth. lab. p. 685; Bois. fl. or. IV. p. 821; Heldr. fl. Aegina p. 383; Form. in D. bot. Mon. 1898 p. 79. — *T. polium v. lanuginosum* Cel. in bot. Centralbl. 1883 p. 1883 p. 152; Haussk. symb. p. 33. — Folia oblonga plana vel margine non vel parum revoluta, crenata; corolla alba. — Variat (*f. virescens*) indumento tenui usque (*f. bombycinum* Heldr. herb. norm. n. 1363, = *T.*

polium v. gnaphalodes Raul. cret. p. 841, non *T. gnaphalodes* Vahl symb. I. p. 41, quod sec. Willk. et Lange pr. fl. hisp. II. p. 480 speciem propriam sistit) bombycino. — Exsicc.: Heldr. pl. fl. hellen. a. 1890, 1896, 1899, 1900 (*f. virescens*); herb. norm. n. 1363 (*f. bombycinum*).

β. **intermedium** Celak. in bot. Centralbl. 1883 p. 152; Haussk. symb. p. 33. — *T. polium v. angustifolium* Benth. lab. p. 686 p. p.; Marg. et R. fl. Zante p. 76; Weiss in z. b. G. 1869 p. 753. — Folia angusta, oblongo-linearia vel linearia, crenata, magis revoluta; corolla alba. — Variat uti α. indumento plus minus denso et (*f. virens*) tenui. — Exsicc.: Heldr. herb. norm. n. 1362; Heldr. herb. fl. hellen. a. 1899 (*f. virens*).

γ. **Achaemenis** Schreb. unilab. p. 44, Hal. in z. b. G. 1888 p. 764; pro sp. — *T. capitatum* S. et S. pr. I. p. 395, Fl. gr. VI. p. 30 t. 536; Fraas fl. class. p. 174; non L. — *T. polium v. purpurascens* Benth. lab. p. 686. — *T. polium v. roseum* Bois. fl. or. IV. p. 821; Heldr. fl. cephal. p. 60. — Folia oblonga vel lineari-oblonga, plus minus revoluta et crenata; corolla purpurea. — Variat uti α. et β. indumento copiosiori vel (*f. viridula*) parciori et foliis latioribus vel angustioribus. — Exsicc.: Baen. herb. europ. n. 9352; Sint. it. thessal. n. 1038; Heldr. pl. fl. hellen. a. 1899 (*f. viridula*).

δ. **capitatum** L. sp. p. 566 pro sp. — *T. polium v. angustifolium* Benth. lab. p. 686 p. p. — *T. polium angustissimum* Celak. l. c., an *T. angustissimum* Schreb. unilab. p. 48. — Folia linearia vel lanceolato-linearia, revoluta, fere omnia integerrima, breviter tomentella; capitula minora, globosa; corolla minor, alba vel rosea. — Exsicc.: Sint. et Bornm. it. turc. n. 1313.

In collibus siccis, locis glareosis, rupestribus regionis inferioris in subalpinam adscendens, var. α. β. et γ. passim per totam ditionem, var. δ autem ut videtur rarissime, cum tantum e Thessalia: ad monasterium Metoji pr. Litochoron vidi. — Maio, Aug. ♃

II. Tribus. **PRASIEAE** Benth. lab. p. 646.

3. **Prasium** L. gen. n. 302.

1. **P. majus** L. sp. p. 601; S. et S. pr. I. p. 426, Fl. gr. VI. t. 584; Sieb. avis. p. 4; Urv. enum. p. 72; Ch. et B. exp. p. 174, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 76; Friedr. Reise p. 269; Ung. Reise p. 127; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 839; Bois. fl. or. IV. p. 798; Heldr. fl. cephal. p. 59, in ö. b. Z. 1898 p. 184, Fl. Aegina p. 383, chlor. Thera p. 18; Spreitz. in z. b. G. 1887 p. 666; Haussk. symb. p. 30. — Exsicc.: Orph. fl. gr. n. 527.

Fruticosum, glabrum vel parce hispidulum; ramis virgatis; foliis petiolatis, a basi truncata vel cordata ovatis, crenatis vel serratis, floralibus integris; verticillastris bifloris, plus minus approximatis; calycis breviter pedicellati dentibus ovatis, mucronatis; corolla albida vel pallide purpurascente.

β. **creticum** Rchb. fl. germ. exc. p. 334 pro sp.; Haussk. symb. p. 30. — Folia floralia serrata. — Exsicc.: Rev. pl. cret. n. 194; Dörfl. pl. cret. n. 98.

In dumosis, rupestribus regionis inferioris et submontanae. Attica: pr. Athenas, in Pharmacusarum insula Lero (Heldr.); Boeotia (Ung.); Peloponnesus (Sibth.): pr. Corinthum, in peninsula Methana (Haussk.), pr. Psatopyrgos Achaiae (Reis.), ad radices mt. Taygetos (Zahn); insula Aegina (Friedr.); Sporadum insula Jura (Leon.); Cycladum insula Syra, Tenos (Weiss), Melos (Urv.), Naxos, Thera (Heldr.); Creta: pr. Canea, Acroteri, Malaxa, insula Gaidaronisi (Raul.), pr. Males distr. Hierapetra (Leon.); in Strophadum insula majore (Reis.); Zante (Sibth.); Cephalonia: pr. Livatho, Pessada, Lixuri (Heldr.); Leucas: pr. Amaxichi (Spreitz.); Corcyra (Ung.). — Mart. Jun. ♄

3. Tribus. **MONARDEAE** Benth. lab. p. 190.

4. **Salvia** L. gen. n. 39.

Dispositio specierum:

A. Tubus corollae intus pilorum annulo munitus.
  I. Antherarum connectiva crure postico loculum cassum ferentia.
    1. Calyx post anthesin auctus, membranaceo-dilatatus.

1. Sectio. *Hymenosphace* Benth. in Hook. bot. misc. III. p. 373. — Calyx campanulatus, labio superiore integro, repando vel brevissime tridentato; corollae labium superius subrectum.

1. S. calycina S. et S. **2. S. pomifera** L.

2. Calyx post anthesin, vix auctus.

2. Sectio. *Eusphace* Benth. in Hook. bot. misc. III. p. 372. — Calyx campanulutus, labio superiore inaequaliter tridentato; corollae labium superius subrectum.

a. Folia indivisa vel triloba; corollae tubus subrectus.

α. Herbacea; folia ampla, cum petiolo ad 30 cm. longa, basi cordata.

3. S. Eichleriana Heldr.

β. Fruticosae; folia multo minora, basi rotundata vel attenuata.

4. S. officinalis L. **5. S. triloba** L.

b. Folia pinnatisecta; corollae tubus recurvo-adscendens.

6. S. ringens S. et S.

3. Sectio. *Drymosphace* Benth. in Hook. bot. misc. III. p. 273. — Calyx campanulatus vel tubulosus, labio superiore integro vel brevissime tridentato; corollae labium superius falcatum, compressum.

7. S. glutinosa L.

II. Antherarum connectiva crure postico loculo casso carentia.

4. Sectio. *Hemisphace* Benth. in Hook. bot. misc. III. p. 374. -- Calyx tubulosus, post anthesin vix auctus, labio superiore truncato, 3dentato; corollae labium superius erecto-patens.

8. S. **verticillata** L. 9. S. **peloponnesiaca** Bois. et Heldr.

B. Tubus corollae pilorum annulo carens; antherarum connectiva loculo casso carentia, extremitate callosa connexa.

5. Sectio. *Horminum* Benth. in Hook. bot. misc. III. p. 373. — Calyx tubulosus, fructifer auctus, labio superiore truncato, remote bidentato; corollae labium superius rectum.

10. S. **viridis** L. 11. S. **horminum** L.

6. Sectio. *Aethiopis* Benth. in Hook. bot. misc. III. p. 373. — Calyx campanulatus vel tubulosus, fructifer vix auctus, labio superiore tridentato; corollae labium superius falcatum, compressum.

a. Folia floralia membranacea, purpurascentia, calyce longiora, tandem deflexa.

12. S. **sclarea** L.

b. Folia floralia herbacea, virescentia, calyce parum — duplo breviora, erecta.

13. S. **Aethiopis** L. 14. S. **argentea** L.

c. Folia floralia minima, submembranacea, pedicello aequilonga.

15. S. **candidissima** Vahl.

6. Sectio. *Piethiosphace* Benth. in Hook. bot. misc. III. p. 373. — Calyx ovatus, vel campanulatus; fructifer vix auctus, labio superiore concavo-bisulcato, dentibus tribus brevissimis conniventibus; corollae labium superius erectum vel falcato-compressum.

a. Folia ad basin caulis nunquam rosulata, sub anthesi jam emarcida; caulis crebre foliosus.

16. S. **amplexicaulis** Lam.

c. Folia ad basin caulis rosulata, sub anthesi vigentia; caulis paucifolius.

α. Calyx campanulatus.

× Corolla magna, ad 2 cm. longa, pulcherrime azurea vel rosea.

17. S. **Tenorii** Spreng.

×× Corolla mediocris, 10—15 mm. longa, purpureo-violacea vel albida.

18. S. **virgata** Ait. 19. S. **similata** Haussk.

β. Calyx ovatus.

20. S. **verbenaca** L.

1. Sectio. *Hymenosphace* Benth. in Hook. bot. misc. III. p. 373.

1. S. **calycina** S. et S. pr. I. p. 13, Fl. gr. I. p. 13 t. 16; Ch. et B. exp. p. 17, Fl. pelop. p. 2; Fraas fl. class. p. 184; Bois. fl. or. IV. p. 605; Haussk. symb. p. 34; Hal. in ö. b. Z. 1897 p. 97, in z. b. G. 1899 p. 190; Form. in D. bot. Mon. 1898 p. 79; Heldr. prosth. chlor. Thera p. 3. — Exsicc.: Heldr. herb. norm. n. 173 et 1364; Orph. fl. gr. n. 400.

Fruticosa, adpressissime cana; ramis elongatis, simplicibus vel apice anguste paniculatis; foliis petiolatis, rugulosis, planis, minutissime crenulatis, a basi rotundata vel subcordata ovatis vel oblongis; verticillastris 2—6 floris, racemos breves formantibus, inferioribus remotis; bracteis membranaceis, coloratis, deciduis; calycis brevissime pedicellati, late

campanulati, hirti, labiis membranaceis, margine purpureis, superiore integro, repando vel brevissime mucronato, inferiore bilobo, lobis obovato-cuneatis; corolla roseo-violacea, puberula, calyce duplo longiore.

In rupestribus regionis inferioris et montanae. Boeotia (Auch.); Attica: mt. Lycabettus, Parnes, Pentelicon (Heldr.), Hymettus (Sibth.), pr. Sunium (Sprun.); Argolis: in peninsula Methana, pr. Nauplia (Haussk.), Poros (Zuccar.), Kranidi (Chaub.); Sporadum insula Scopelos (Leon.); Cycladum insula: Naxos (Urv.), Melos (Chaub.), Thera (Heldr.). — Mart. Jul. ♄

2. **S. pomifera** L. sp. p. 24; S. et S. pr. I. p. 13, Fl. gr. I. p. 12 t. 15; Sieb. avis p. 2, rem. p. 2, in Flora I. p. 273; Raul. cret. p. 832; Bois. fl. or. IV. p. 604; Heldr. in ö. b. Z. 1880 p. 346; Spreitz. in z. b. G. 1887 p. 666, 1890 p. 297; Bald. viagg. Creta p. 85; Fors. in bull. herb. Bois. V. p. 399. — *S. fragifera* Ettl. salv. n. 5. — Exsicc.: Rev. pl. cret. n. 135.

A praecedente, cui maxime affinis, ramis simplicibus, foliis oblongis vel oblongo-linearibus, saepe undulato-crispis, calycibus majoribus, minus hirtis differre dicitur. — Ad ramorum apicem saepe gallas esculentas, cerasi vel nucis magnitudine, foliis abortivis coronatas, fert; inde nomen „pomifera".

In aridis, rupestribus regionis inferioris et montanae. Creta: pr. Neocorio (Tourn. voy. p. 30 cum icone „S. cretica frutescens pomifera"), Kissamos (Rev.), Malaxa, Rustica, Omalos (Raul.), Madaro (Bald.); Cythaera: in fauce Kalamo (Spreitz.); indicatur quoque a S. et S. l. c. in Graecia, et a Mazz. in ant. ion. II. p. 446 pr. Spartilla et Messongi Corcyrae. — Jun. Jul. ♄

2. Sectio. *Eusphace* Benth. in Hook. bot. misc. III. p. 372.

a. Folia indivisa vel triloba; corollae tubus rectus.

α. Herbacea, folia ampla, cum petiolo ad 30 cm. longa, basi cordata.

3. **S. Eichleriana** Heldr. it. thessal. a. 1882 n. 40, in Sitzungsb. acad. wiss. Berlin 1883 p. 157; Hal. in z. b. G. 1899 p. 191.

Herbacea; caule elato, pubescenti-tomentello, simplici vel apice anguste paniculato; foliis amplis, remotis, oppositis, e basi cordata ovatis, crenato-dentatis, supra viridibus, rugulosis, parce pubescentibus, subtus cinereo-tomentellis, inferioribus et mediis longe petiolatis, superioribus sessilibus, verticillastris 4—10 floris, confertis; floribus brevissime pedicellatis; calyce campanulato viscidulo-hirto, subbilabiato, dentibus triangulari-lanceolatis, acuminatis; corolla coerulescente, puberula, calyce triplo longiore. — Planta orgyalis, foliis cum petiolo ad 30 cm. longis. 15 cm. latis, floribus 5—6 cm. longis; *S. grandiflorae* Ettl. statura humiliori, foliis multo minoribus basi rotundatis, verticillastris remotis, floribus minoribus longius pedicellatis et indumento tenuiori diversae, affinis.

Inter *Pterides* regionis superioris mt. Pelion supra Drakia Thessaliae (Heldr.). — Jul. Aug. ♃

b. Fruticosae; folia multo minora, basi attenuata vel rotundata.

4. S. **officinalis** L. sp. p. 23; S. et S. pr. I. p. 13; Pieri corc. fl. p. 5; Dallap. prosp. p. 8; Fraas fl. class. p. 184; Heldr. Nutzpfl. p. 33, chlor. Thera p. 29; Gelmi in bull. soc. bot. ital. 1889 p. 451. — Icon: Lam. ill. t. 515.

Fruticosa, adpresse pubescenti-incana, ramosissima, ramis erectis, simplicibus vel parce ramulosis, eglandulosis; foliis e basi rotundata vel attenuata ovato- vel oblongo-lanceolatis, crenulatis, rugulosis, inferioribus petiolatis, superioribus sessilibus, verticillastris 3—5 floris, distantibus; floribus pedicellatis; calycis campanulati, adpresse hirti, bilabiati dentibus triangulari-lanceolatis, aristatis, tubo subbrevioribus; corolla violacea puberula, calyce 2—3 plo longiore.

In collibus regionis calidae (Sibth.): Phthiotis, Tzakonia (Fraas); Cycladum insula: Syra (Fraas), Thera (Heldr.); Cephalonia (Dall.); Corcyra (Pieri): mt. Pantocrator (Gelmi); loca nonnulla probabiliter ad sequentem spectant. — Apr. Jul. ♄ N. v.

Obs. *S. cretica* L. sp. p. 23; Sieb. in Flora I. p. 273; Raul. cret. p. 831. — Secundum Bois. fl. or. IV. p. 594 probabiliter forma hortensis submonstrosa speciei praecedentis est.

5. S. **triloba** L. fil. suppl. p. 88; S. et S. pr. I. p. 14, Fl. gr. I. p. 13 t. 17; Sieb. avis p. 2, rem. p. 2; Urv. enum. p. 3; Ch. et B. exp. p. 17, Fl. pelop. p. 1; Marg. et R. fl. Zante p. 17; Friedr. Reise p. 272; Ung. Reise p. 126; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 831; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 595; Heldr. fl. cephal. p. 58, Fl. Aegina p. 283, prosth. chlor. Thera p. 3; Haussk. symb. p. 33; Hal. in ö. b. Z. 1897 p. 97. — *S. baccifera* Ettl. salv. n. 4; ? Urv. enum. p. 3. — Exsicc.: Orph. fl. gr. n. 548 (Chios); Rev. pl. cret. n. 197 (*v. integrifolia*); Baen. herb. europ. n. 9331; Dörfl. pl. cret. n. 80.

Fruticosa, cano-tomentosa, ramosissima, ramis erectis, simplicibus vel ramulosis, tomentellis et superne saepe glandulosis, rarius glabriusculis; foliis petiolatis a basi truncata vel subcordata oblongis vel lanceolatis, crenulatis, rugulosis, saepius ad laminae basin segmentis binis parvis auctis; verticillastris 4—6 floris, plus minus distantibus; floribus pedicellatis; calycis campanulati, glanduloso-hirti, viscosi, subbilabiati dentibus triangularibus, acutis, tubo multo brevioribus; corolla violacea, puberula, calyce 3—4 plo longiore. — Differt a praecedente statura elatiori, indumento copiosiore, foliis saepius trilobis, calyce glanduloso-hirto, dentibus brevibus, non aristatis; corolla violacea, puberula, calyce triplo longiore.

In rupestribus regionis calidae, frequens. Aetolia: mt. Chalkis pr. Krioneri (Hal.); Attica: mt. Parnes pr. Panagia-Kliston, insula Aegina (Heldr.); Peloponnesus (Chaub.): pr. Acrocorinthum, Nauplia (Haussk.)

Troezene (Friedr.); Sporadum insula Skiathos (Leon.); Cycladum insula Cythnos (Tunt.), Syra (Weiss), Melos (Leon.), Thera (Heldr.); Creta pr. Kissamos, Malaxa (Rev.), promontorium Meleka, Aradhena, insula Dia (Raul.), pr. Parsas distr. Viano (Leon.); Zante (Marg.); Cephalonia: pr. Argostoli, mt. Gerolaki, pr. Valeriano, Marcopulo, Valtes, Koroni, Same (Heldr.); Corcyra: pr. San Deca, Barbati, Signes (Spreitz.), Scripero (Baen.), Palaeokastro (Ung.); et certe alibi. — Mart. Maio. ♄

b. Folia pinnatisecta; corollae tubus recurvo-adscendens.

6. **S. ringens** S. et S. pr. I. p. 14, Fl. gr. I. p. 14 t. 18; Ch. et B. exp. p. 17, Fl. pelop. p. 2; Bois. fl. or. IV. p. 600; Heldr. chlor. Parn. p. 25. — Exsicc.: Orph. fl. gr. n. 160; Heldr. herb. norm. n. 380 et 1269; Sint. it. or. a. 1889 n. 1879; Sint. et Bornm. it. turc. n. 1405 b.

Rhizomate indurato, ramoso; caulibus erectis, pruinosis, glabris, superne glanduloso-pilosis, in racemum vel in paniculam parce ramosam abeuntibus; foliis virentibus, rugulosis, inferne praesertim pubescentibus, irregulariter pinnatisectis, segmentis 2—4 jugis, crenatis, lateralibus ovatis ellipticisve, saepe sufpetiolulatis, terminali multo majore, ovato vel oblongo, saepe acutiusculo; verticillastris 2—6 floris, distantibus; floribus pedicellatis; calycis campanulati, glanduloso-pilosi, bilabiati, labio superiore breviter tridentato, inferiore bifido; corolla speciosa, coeruleo-violacea, glanduloso-pilosa, calyce 4 plo longiore. — Species elegans, floribus ad 4 cm. longis.

In petrosis regionis submontanae et abietinae, rare. Thessalia: mt. Olympus (Heldr.); Attica: mt. Parnes, Pateras (Heldr.); mt. Parnassus (Sprun.); Achaia: pr. Megaspilaeon (Sibth.), mt. Kyllene pr. Trikala (Orph.). — Maio, Sept. ♃

3. Sectio. *Drymosphace* Benth. in Hook. bot. misc. III. p. 373.

7. **S. glutinosa** L. sp. p. 26; Form. in D. bot. Mon. 1891 p. 28, in Ver. Brünn 1896 p. 66, 1897 p. 49; Haussk. symb. p. 34. Icon: Rchb. t. 45. — Exsicc.: Orph. fl. gr. n. 1046 (Macedonia); Sint. it. thessal. n. 1115.

Herbacea; molliter pubescens, superne pedicellis, calycibus corollisque glandulosa; caule erecto, simplici vel ramoso; foliis petiolatis, ovato-oblongis, grosse serratis, acuminatis, cordato-hastatis, verticillastris 2—6 floris, racemum elongatum, interruptum formantibus; floribus pedicellatis; calycis campanulati, bilabiati, labio superiore integro vel minute tridentato, inferiore bifido; corolla speciosa, flava.

In silvaticis montanis et subalpinis, rarissime. Thessalia: ad Ochsa Despot (Form.), in oropedio Neuropolis, pr. Sermeniko (Haussk.) in Pindo. — Jul. Sept. ♃

4. Sectio. *Hemisphace* Benth. in Hook. bot. misc. III. p. 374.

8. **S. verticillata** L. sp. p. 26; Benth. lab. p. 311; Mazz. in ant. ion. II. p. 448, Bois. fl. or. IV. p. 634; Haussk. symb. p. 37;

Form. in Ver. Brünn 1897 p. 49. — Huc probabiliter quoque: *S. napifolia* Mazz. l. c.; vix Jacq. hort. vind. II. t. 152, quae verticillastris paucifloris (5—10), floribus fere duplo majoribus, intense violaceis discedit. — Icon: Rchb. t. 54. — Exsicc.: Heldr. it. gr. septentr. a. 1879.

Herbacea, breviter hirta, viridis; caulibus erectis vel adscendentibus, ramosis; foliis crenatis, inferioribus petiolatis, ovatis, basi cordatis, saepius lyratis, unijugis, segmentis lateralibus auriculiformibus, parvis, foliis superioribus a basi sessili, cordato-amplexicauli ovatis; verticillastris distantibus, 20—40 floris, racemum elongatum formantibus; floribus pedicellatis; calycis tubulosi, subbilabiati labio superiore breviter tridentato, inferiore bidentato; corollae purpureo-violaceae, puberulae, calyce duplo longioris tubo breviter exserto. — Floribus parvis, 8—12 mm. longis, in verticillos globosos, multifloros congestis insignis.

In dumosis, silvaticis regionis montanae rarissime. Thessalia: pr. Klinovo in Pindo (Haussk.), pr. Phlambures et mt. Mitriza in mt. Chassia (Form.); Aetolia: pr. Musinitza in mt. Korax (Heldr.); Boeotia (Swainson ex Benth. l. c.); Corcyra: pr. Castrades et Cacochieri (Mazz.). — Jun. Aug. ♃

9. **S. peloponnesiaca** Bois. et Heldr. diagn. VII. p. 47, Fl. or. IV. p. 634; Hal. Beitr. fl. Aetol. p. 9, Fl. Achaia p. 1899 p. 191. — *S. verticillata* S. et S. pr. I. p. 17, quod pl. peloponn.; Ch. et B. exp. p. 19, Fl. pelop. 2. — Exsicc.: Heldr. herb. norm. n. 1270; Bald. it. alb. a. 1892 n. 33.

Praecedenti maxime affinis, ab ea indumento sericeo-canescenti, foliis pinnatipartito-lyratis, segmentis lateralibus 2—3 jugis, sat magnis, terminali cuneato-oblongo, distincta.

In herbidis, vineis, dumosis regionis inferioris et submontanae. Epirus: pr. Vallona (Bald.); Aetolia: pr. Mesolongion (Heldr.), Krioneri (Hal.); Achaia: pr. Psatopyrgos (Heldr.), Patras (Hal.); Elis: pr. Lintzi (Heldr.), Olympia (Reis.); Arcadia: pr. Zatuna (Orph.); Messenia: pr. Pylos, mt. Ithome (Heldr.), mt. Kupe, pr. Methone, Messene, Kalamata (Chaub.). — Maio, Jul. ♃

4. Sectio. *Horminum* Benth. in Hook. bot. misc. III. p. 373.

10. **S. viridis** L. sp. p. 24; Mazz. in ant. ion. II. p. 446; Marg. et R. fl. Zante p. 73; Friedr. Reise p. 269; Ung. Reise p. 126; Raul. cret. p. 832; Bois. fl. or. IV. p. 630; Heldr. fl. cephal. p. 58; Gelmi in bull. soc. bot. ital. 1889 p. 451; Haussk. symb. p. 37. — *S. horminum v. viridis et comosula* Heldr. fl. Aegina p. 384. — Icon: Fl. gr. t. 19. — Exsicc.: Orph. fl. gr. n. 434; Heldr. herb. norm. n. 578 et 1167; Dörfl. pl. cret. n. 82.

Annua, villoso-hispidula, virens, simplex vel a collo pluricaulis; caulibus erectis, simplicibus vel 1—2 ramulos auctis; foliis a basi rotundata vel subcordata ovatis oblongisve, crenulatis, inferioribus petiolatis, superioribus sessilibus, floralibus late cordato-ovatis, persisten-

tibus, basi utrinque bractea subulata auctis, terminalibus diminutis, viridibus vel parce coloratis; verticillastris 4—6 floris, distantibus; calyce tubuloso, sulcato-striato, labio superiore oblique truncato, dentibus binis parvis, remotis, inferiore tridentato, dentibus ternis, lanceolatis, spinosis, fructifero aucto, reflexo, corollo rosea vel violacea, parce pubescente, calyce vix duplo longiore. — Corolla parva, 10—12 mm. longa.

In arvis, collibus arenosis, saxosis regionis inferioris. Attica: pr. Athenas, in colle Lycabetto (Orph.), pr. Heracleon, Piraeum (Heldr.), insula Aegina (Friedr.); Acrocorinthus, Nauplia (Haussk.); Achaia: pr. Patras (Hal.); Laconia: ad radices mt. Taygetos (Zahn); Cycladum insula Cythnos (Tunt.); Creta: pr. Canea, Drapano (Raul.), Hierapetra (Leon.); Zante: mt. Scopo (Marg.); Cephalonia: pr. Argostoli (Ung.); Corcyra: pr. Ipso (Mazz.), San Deca, Signes (Gelmi); et certe alibi. — Mart. Jun. ⊙

11. **S. horminum** L. sp. p. 24; S. et S. pr. I. p. 14, Fl. gr. I. p. 15 t. 20; Pieri corc. fl. p. 6; Ch. et B. exp. p. 18, Fl. pelop. p. 2; Mazz. in ant. ion. II. p. 446; Friedr. Reise p. 275; Fraas fl. class. p. 184; Ung. Reise p. 126; Raul. cret. p. 833; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 631; Gelmi in bull. soc. bot. ital. 1889 p. 451; Form. in D. bot. Monat. 1891 p. 28, in Ver. Brünn 1895 p. 35, 1896 p. 66, 1897 p. 49; Hal. Beitr. fl. Epir. p. 35, in ö. b. Z. 1896 p. 16; Bald. riv. coll. bot. alb. 1896 p. 86; Haussk. symb. p. 37; Heldr. fl. Aegina p. 384. — *S. viridis v. comata* Heldr. fl. cephal. p. 58. — Exsicc.: Rev. pl. cret. n. 136; Sint. et Bornm. it. turc. n. 1406; Sint. it. thessal. n. 34.

Differt a praecedente foliis floralibus terminalibus amplis, intense violaceis, comantibus.

In cultis, arvis, collinis regionis inferioris, passim per totam Graeciam. — Apr. Jul. ⊙

5. Sectio. *Aethiopis* Benth. in Hook. bot. misc. III. p. 373.

a. Folia floralia membranacea, purpurascentia calyce longiora, tandem deflexa.

12. **S. sclarea** L. sp. p. 27; Mazz. in ant. ion. II. p. 448; Bois. fl. or. IV. p. 616; Heldr. fl. cephal. p. 58, chlor. Parn. p. 25; Form. in D. bot. Mon. 1891 p. 18, in Ver. Brünn 1895 p. 35, 1897 p. 49; Hal. Beitr. fl. Epir. p. 34, Beitr. fl. Thessal. p. 16, in ö. b. Z. 1897 p. 325; Bald. riv. coll. bot alb. 1896 p. 86; Haussk. symb. p. 34. — Icon: Fl. gr. t. 25. — Exsicc.: Hal. it gr. secund. a. 1893.

Viscoso-villosa; caule erecto, in paniculam ramosam strictam abeunte; foliis viridibus, cordato-ovatis, irregulariter crenatis, inferioribus petiolatis, superioribus amplexicaulibus, floralibus bracteaeformibus, subito cuspidato-acuminatis, membranaceis, purpurascentibus, calyces superantibus; verticillastris confertis, 4—6 floris; floribus breviter pedicellatis; calyce campanulato, glanduloso, dentibus in setam rigidulam abeuntibus; corolla pallide rosea, parce glandulosa, calyce triplo longiore. — Foliis floralibus magnis, concavis, purpureo-suffusis insignis.

In herbidis regionis inferioris et subalpinae. Epirus: ad monasterium Hagios Paraskevi, in valle Dipotami, mt. Prosgoli (Form.), mt. Micikeli (Bald.), pr. Theodoriana, Pramanta, Mazuki, Chaliki (Hal.), Kastania, Vendista, Velitsena, Koturi, Lepeniza, Kalabaka (Form.), Korona, Bezula, Sermeniko (Haussk.) in Pindo, pr. Kastri, Vlachava, Kucuro in mt. Chassia (Form.); Aetolia: mt. Korax; mt. Parnassus (Heldr.); Peloponnesus: mt. Kyllene (Heldr.), Malevo (Leon.); Cephalonia: pr. Drapano, Peratata (Heldr.); Corcyra: mt. Deka, insula Merlere (Mazz.). — Jun. Jul. ♃.

b. Folia floralia herbacea, virescentia, calyce parum — duplo breviora, erecta.

13. **S. Aethiopis** L. sp. p. 27; S. et S. pr. I. p. 18; Fraas fl. class. p. 185; Bois. fl. or. IV. p. 616. — Icon: Jacq. fl. austr. III. t. 211.

Molliter lanato-tomentosa; pilis glanduliferis destituta; caule erecto, a parte inferiori in paniculam ramosissimam, ample pyramidatam abeunte; foliis canescentibus, fere omnibus ad collum confertis, expansis, breviter petiolatis, cordato-ovatis, irregulariter inciso-dentatis, caulinis paucis ovatis, amplexicaulibus, floralibus bracteaeformibus, subrotundo-ovatis, subito cuspidato-acuminatis, herbaceis, calyce subbrevioribus; verticillastris distantibus, 6—10 floris; floribus breviter pedicellatis; calyce campanulato, albo-lanato, dentibus in setam rigidulam abeuntibus; corolla alba, pubescente, calyce duplo longiore.

Attica: circa Athenas (Sibth.); Corinthus (Fraas); a recentioribus non lecta et forsan cum sequente confusa. — Graeciae civis dubia. — Jun. Jul. ♃ N. v.

14. **S. argentea** L. sp. ed. 2 p. 38; S. et S. pr. I. p. 18, Fl. gr. I. p. 20 t. 27; Ten. adnot. fl. gr. p. 3; Fraas fl. class. p. 185; Raul. cret. p. 833; Bois. fl. or. IV. p. 618; Form. in Ver. Brünn 1895 p. 35, in D. bot. Mon. 1898 p. 79; Haussk. symb. p. 34 *v. thessala* (f. foliis rosularibus elliptico-lanceolatis, calycis dentibus longius spinulosis). — Huc spectare videtur quoque *S. macedonica* Griseb. spic. II. p. 110 pro var. *S. argenteae;* Haussk. symb. p. 34; = *S. alpestris* Haussk. in Nym. consp. suppl. II. 248; Bald. riv. coll. bot. alb. 1896 p. 86. — Exsicc.: Orph. fl. gr. n. 547; Heldr. herb. norm. n. 172.

Caule erecto, ample paniculato, villosulo et saltem superne glanduloso-piloso; foliis canescentibus, fere omnibus ad collum confertis, breviter petiolatis, ovatis vel oblongis, basi cuneatis, rotundatis vel subcordatis, sinuato-lobatis, eroso-dentatis, utrinque sed subtus densius arachnoideo-lanatis, caulinis paucis, ovato-lanceolatis, amplexicaulibus, floralibus bracteaeformibus, subrotundo-ovatis, subito acuminatis vel breviter cuspidatis, herbaceis, calyce duplo brevioribus; verticillastris distantibus, 4—10 floris, summis saepius abortientibus; floribus breviter pedicellatis; calyce campanulato, glanduloso-piloso, dentibus breviter spinosis; corolla alba, parce glandulosa, calyce triplo longiore. — A

praecedente indumento glanduloso, foliis floralibus minoribus et corolla majore specifice discedit.

β. **alpina** Heldr. chlor. Parn. p. 25. — Caulis pumilus, 20—25 cm. altus, parce breviterque ramosus, ramis apice saepius verticillastro fertili terminatis. — Exsicc.: Pichl. pl. gr. a. 1876; Heldr. pl. fl. hellen. a. 1876; Bald. it. alb. epir. IV. n. 43.

In campis, segetibus. Epirus: pr. Ephemiades (Form.); Thessalia: pr. Pharsalus (Haussk.); Attica: pr. Athenas (Orph.), pr. Kallitea, ad Phaleron (Heldr.); Creta (Linné), sed hic a recentioribus non reperta; — β. Epirus: mt. Micikeli (Bald.); Thessalia: ad Gionskala pr. Sermeniko (Haussk.); mt. Parnassus (Sibth.); Euboea: mt. Dirphys (Heldr.). — Apr. Jun. ♃

c. Folia floralia minima, submembranacea, pedicello aequilonga.

15. **S. candidissima** Vahl enum. I. p. 278; Bois. fl. or. IV. p. 621; Bald. riv. coll. bot. alb. 1896 p. 86. — *S. odorata* Willd. en. berol. I. p. 43; Jacq. ecl. II. p. 26 t. 16; sec Bois. l. c. — *S. candidissima v. odorata* Bald. riv. coll. bot. alb. 1895 t. 61. — Exsicc.: Bald. it. alb. epir. III. n. 194, IV. n. 138.

Caule erecto, anguste vel ample paniculato, tomentello, superne minute glanduloso-piloso; foliis ad partem inferiorem caulis congestis, petiolatis, a basi subcordata, rotundata vel attenuata ovatis, oblongis vel oblongo-lanceolatis, crenatis, utrinque albo-pannosis, superioribus diminutis, sessilibus, floralibus bracteaeformibus, minutis, setaceo-acuminatis, pedicello aequilongis; verticillastris remotis, 2—6 floris, summis saepe abortientibus; floribus breviter pedicellatis; calyce campanulato, tomentello, dentibus in setam rigidulam abeuntibus; corolla alba, parce puberula, calyce triplo longiore. — A praecedente distinctissima foliis albo-pannosis, pilis glanduliferis caulis brevissimis, verticillastris paucifloris et foliis floralibus minutis.

In saxosis, alveis, aridis, collibus siccis regionis inferioris et montanae rarissime. Epirus: ad Vromonero distr. Ljaskovik, mt. Smolika distr. Konitza, in alveo fl. Metzovo pr. Han Drysko distr. Janina (Bald.); Aetolia: mt. Tymphrestus (Nicolaidis); mt. Geranion in isthmo Corinthiaco (Stuart Mill). — Jun. Aug. ♃

6. Sectio. *Plethiosphace* Benth. in Hook. bot. misc. III. p. 373.

a. Folia ad basin caulis nunquam rosulata, sub anthesi jam emarcida; caulis crebre foliosus.

16. **S. amplexicaulis** Lam. ill. I. p. 68; Hal. Beitr. fl. Epir. p. 35, Beitr. fl. Thessal. p. 17; Form. in Ver. Brünn 1896 p. 66, 1897 p. 49; Haussk. symb. p. 35. — *S. silvestris* Mazz. in ant. ion. II. p. 446; Bois. fl. or. IV. p. 628 quoad pl. olymp.; Bald. riv. coll. bot. Alb. 1896 p. 86; non L. — *S. villicaulis* Borb. a magy. homok. növ. p. 83 (1886); Form. in D. bot. Mon. 1891 p. 27. — Exsicc.: Bald. it. alb. epir. IV. n. 139; Sint. it. thessal. n. 991.

Caule erecto, crebre foliato, bracteis, calycibus corollisque pilis patentibus eglandulosis villosulo, in paniculam strictam vel amplam abeunte; foliis e basi cordata vel rotundata oblongo-lanceolatis, crenatis, utrinque sed subtus densius hispidulis, infimis breviter petiolatis, mediis amplexicaulibus, superioribus semiamplexicaulibus, floralibus bracteaeformibus, cordato-ovatis vel subrotundis, acutis, saepe coloratis, calyce brevioribus; verticillastris subsexfloris, approximatis; calyce tubuloso-campanulato, breviter pedicellato, dentibus acuminato-mucronatis; corolla violacea, calyce duplo longiore, labio superiore falcato. — Maxime affinis *S. silvestri* L. sp. p. 24, p. p.; Jacq. fl. austr. III. p. 7 et plur. aut. (*S. nemorosae* L. sp. ed. 2 p. 35), indumento minute et brevissime tomentello, foliis mediis semiamplexicaulibus, floralibus longioribus calycem superantibus, verticillis remotioribus diversae.

In herbidis regionis inferioris et montanae. Epirus: mt. Maria et pr. Dobra distr. Ljaskovik (Bald.), pr. Kalarrytes, Mazuki, Pramanta, Theodoriana, Vulgarelion (Hal.); Thessalia: frequens in Pindo tymphaeo et dolopico, mt. Oxia, Chassia, nec non in planitie pr. Trikala, Karditza, Pharsalus, Orman Magula et in Olympo; Corcyra: pr. Viro, cum *S. silvestris* Mazz. probabiliter huc spectat. — Jun. Jul. ♃.

b. Folia ad basin caulis rosulata, sub anthesi vigentia; caulis paucifolius.

α. Calyx campanulatus.

× Corolla magna, ad 2 cm. longa, pulcherrime azurea vel rosea.

17. **S. Tenorii** Spreng. pug. I. p. 2; Roem. et Schult. syst. I. p. 242; ? Mazz. in ant. ion. II. p. 448. — *S. pratensis* ? S. et S. pr. I. p. 15, ? Mazz. in ant. ion. II. p. 446; ? Form. in D. bot. Mon. 1890 p. 28; Bois. fl. or. IV. p. 626, quoad pl. gr.; Heldr. fl. cephal. p. 58; non L. — *S. Barrelieri* Ten. fl. nap. I. p. 9, t. 2; Hal. Beitr. fl. epir. p. 34, Beitr. fl. Achaia p. 28; Form. in Ver. Brünn 1896 p. 66, 1897 p. 49; Heldr. fl. Aegina p. 384; non Ettl. salv. 46, quae = *S. inamoena* Vahl. — Exsicc.: Hal. it. gr. secund. a. 1893.

Caule erecto, pilis brevibus asperulo, in paniculam amplam abeunte; foliis viridibus, supra glabris, subtus parce asperulis, radicalibus petiolatis, e basi cordata vel rotundata ovatis vel oblongis, duplicato-crenatis, interdum lobulatis, caulinis sessilibus, saepe semiamplexicaulibus, floralibus bracteaeformibus, cordato-ovatis rotundatisve, acutis vel acuminatis, calyce triplo vel ultra brevioribus; verticillastris subsexfloris, distantibus; calyce campanulato, breviter pedicellato, glandulis sessilibus lucidis pilisque albis hirto, labio superiore brevissime tridenticulato, inferiore dentibus binis, a basi ovata mucronatis; corolla pilis albis eglandulosis glandulosisque paucis intermixtis et glandulis sessilibus lucidis obsita, calyce triplo longiore, labio superiore falcato. — Species decora, caule ad 80 cm. alto, saepissime ample paniculato, floribundo, foliis radicalibus saepe amplis, ad 30 cm. longis et 12 cm. latis; ab affini *S. pratensi*

L., quacum ab autoribus commutata fuit, caule elatiore ample paniculato, foliis caulinis superioribus amplexicaulibus, indumento asperulo, pilis glandulosis corolla excepta destituto et praesertim florum colore pulcherrime azureo specifice distincta. — Sec. Heldr. fl. Aegina p. 384 occurrit quoque flore roseo.

In herbidis, dumosis regionis inferioris et montanae, rare. Epirus: pr. Kalentini (Hal.); Thessalia: pr. Malakasi, Sermeniko, Neuropolis, Trikala (Form.); Aetolia: pr. Aetolikon (Heldr.); Attica (Orph.); insula Aegina: mt. Oros (Heldr.); Achaia: pr. Iliospulos, Kalavryta (Hal.); Arcadia: pr. Zatuna (Orph.); Messenia: mt. Ithome (Heldr.); Cephalonia: pr. Same, Lixuri (Heldr.); Corcyra: pr. Casopo, insula Vido (Mazz.). — Maio, Jul. ♃

×× Corolla mediocris, 10—15 mm. longa, purpureo-violacea vel albida.

18. **S. virgata** Ait. hort. Kew. I. p. 39; Jacq. hort. vind. I. p. 14 t. 37; Ch. et B. exp. p. 18, Fl. pelop. p. 2; Raul. cret. p. 833; Bois. fl. or. IV. p. 627; Hal. Beitr. fl. Thessal. p. 17, Beitr. fl. Achaia p. 29; Form. in Ver. Brünn 1895 p. 35, 1896 p. 66, 1897 p. 49; Haussk. symb. p. 35. — *S. Sibthorpii* S. et S. pr. I. p. 16, Fl. gr. I. p. 17 t. 22; Urv. enum. p. 3; Marg. et R. fl. Zante p. 73; Mazz. in ant. ion. II. p. 448. — Huc forsan: *S. viscosa* Mazz. l. c. p. 446, vix Jacq. — Exsicc.: Heldr. herb. norm. n. 679; Orph. fl. gr. n. 161; Sint. it. thessal. n. 946; Dörfl. fl. gr. n. 403.

Caule erecto, bracteis calycibusque glanduloso-piloso, simplici vel virgato-ramoso; foliis lobulato-crenatis, utrinque crispule-pubescentibus, radicalibus petiolatis, e basi rotundata vel subcordata ovatis oblongisve, caulinis paucis, diminutis breviter petiolatis et sessilibus, acutis, floralibus bracteaeformibus, ovatis, acuminatis, saepe coloratis, calyce triplo vel ultra brevioribus; verticillastris subsexfloris, distantibus, longe racemosis; calyce campanulato, breviter pedicellato, labio superiore breviter tridentato, inferiore dentibus binis, ovatis, subulato-attenuatis; corollae pubescentis, calyce 2—2½ plo longioris, labio superiore falcato. — Species caulibus elongatis, virgatis, simplicibus vel pauciramosis insignis.

In collibus, campis, herbidis regionis inferioris et montanae. Epirus: pr. Han Balduma, in valle Dipotami (Form.); Thessalia: pr. Malakasi (Sint.), Kastania (Hal.), Korona, Klinovo, Ormanmagula, Karditza, Pharsalus (Haussk.), Kastri, Trikala, Kopraena, Vlachava, Palaeocastro, Ruso, Stylida, Avlaki (Form.); Euboea: pr. Kastaniotissa (Heldr.); Aetolia: pr. Mesolongion (Heldr.); mt. Parnassus (Sibth.); Boeotia (Sprun.); Peloponnesus (Sibth.): pr. Patras, Kalavryta, Planideri (Hal.), Sudena (Leon.) Achaiae, pr. Messene, Methone, Corone Messeniae (Chaub.), mt. Malevo Laconiae (Orph.); insula Syra (Urv.); Creta (Raul.); Zante (Sibth.); Corcyra: mt. Pantaleone, pr. Lavcki, Anaplades (Mazz.). — Maio, Jul. ♃

16 × 18. **S. amplexicaulis × virgata** (*S. adulterina*) Haussk. symb. p. 35. — Differt a *S. amplexicauli* indumento glanduloso-piloso

foliis radicalibus sub anthesi vigentibus, verticillastris remotioribus; — a *S. virgata* caule folioso, magis ramoso, foliis radicalibus brevius petiolatis, in petiolum attenuatis, crenatis nec lobulatis. — Inter parentes supra Klinovo et prope monasterium Korona in Pindo (Haussk.). — N. v.

19. **S. similata** Haussk. symb. p. 36. —

Ex affinitate *S. virgatae*, a qua differt ramis magis elongatis, foliis radicalibus brevius petiolatis, angustioribus longioribusque, oblongo-lanceolatis, acutis, nec obtusis, caulibus ramosissimis, pilis brevissimis, eglandulosis asperulis, verticillastris magis approximatis, calycibus violaceis, minoribus, angustioribus, duplo longius pedicellatis, pilis simplicibus, sparse nec dense glandulosis obsitis, dentibus inferioribus longius productis, floribus minoribus.

In schistosis regionis mediae mt. Pindus supra Klinovo (Haussk.). — Jul. ♃ N. v.

β. Calyx ovatus.

20. **S. verbenaca** L. sp. p. 25; S. et S. pr. I. p. 16; Marg. et R. fl. Zante p. 73; Mazz. in ant. ion. II. p. 446; Gelmi in bull. soc. bot. ital. 1889 p. 451; Heldr. chlor. Thera p. 19. — *S. Sibthorpii* Ch. et B. exp. p. 18, Fl. pelop. p. 2, sec. Bois. fl. or. IV. p. 629. — *S. clandestina* Ch. et B. exp. p. 19, Fl. pl. p. 2. — *S. verbenaca α. serotina* Bois. voy. bot. esp. p. 484, Fl. or. IV. p. 629; Raul. cret. p. 833; Heldr. fl. cephal. p. 58, Fl. Aegina p. 384; Haussk. symb. p. 37. — Icon: Rchb. t. 703. — Exsicc.: Orph. fl. gr. n. 546; Heldr. herb. norm. n. 577; herb. dimorph. n. 58.

Caule erecto vel adscendente, simplici vel ramoso, molliter villoso vel breviter pubescente, superne saepius glanduloso; foliis ovatis vel oblongis, crenatis vel lobulatis, basi cordatis vel rotundatis, glabriusculis, inferioribus petiolatis, superioribus sessilibus, saepe amplexicaulibus, floralibus bracteaeformibus, cordato-orbicularibus, apiculatis, calyce brevioribus; verticillastris 4—6 floris, distantibus; calycis ovati, breviter pedicellati, hirsuti labio superiore late ovato, brevissime denticulato, inferiore dentibus binis, ovato-lanceolatis, cuspidatis; corollae coeruleae rarius albidae, glabriusculae, calyce subaequilongae vel duplo longioris, labio superiore subincurvo. — Species polymorpha.

β. **clandestina** L. sp. ed. 2 p. 36; Marg. et R. fl. Zante p. 73; Friedr. Reise p. 266; Raul. cret. p. 833 *v. multifida*; pro sp. — *S. praecox* Savi fl. pis. I. p. 22; Mazz. in ant. ion. II. p. 448. — *S. multifida* S. et S. pr. I. p. 16, Fl. gr. I. p. 17 t. 23. — *S. verbenaca* Ch. et B. exp. p. 18, Fl. pelop. p. 2. — *S. verbenaca β. vernalis* Bois. voy. bot. esp. p. 484, Fl. or. IV. p. 629; Heldr. fl. Aegina p. 384, chlor. Mykon. p. 248; Haussk. symb. p. 37. — Folia profunde incisa vel pinnatifida. — Sine limites ad typum transit. — Exsicc.: Heldr. herb. dimorph. n. 44; Sint. it. thessal. n. 1512; Dörfl. fl. gr. n. 464.

In herbidis, arvis, collibus regionis inferioris per omnem Graeciam. — Febr. Jun. ♃

Obs. Species ex confusione quadam in ditione indicatae: *S. Forskahlei* L. mant. p. 26, a Mazz. in ant. ion. II. p. 448, pr. Sisermones Corcyrae; *S. nilotica* Vahl. en. I. p. 258, a Pieri corc. fl. p. 5, in Corcyra; *S. pinnata* L. sp. p. 27, a Willd. sp. I. p. 149 in Creta.

5. Rosmarinus L. gen. n. 38.

1. **R. officinalis** L. sp. p. 23; S. et S. pr. I. p. 12, Fl. gr. I. p. 11, t. 14; Dallap. prosp. p. 7; Ch. et B. exp. p. 17, Fl. pelop. p. 2; Mazz. ant. ion. II. p. 446; Marg. et R. fl. Zante p. 73; Friedr. Reise p. 270; Fraas fl. class. p. 183; Ung. Reise p. 126; Heldr. Nutzpfl. p. 33, Fl. cephal. p. 58, Fl. Aegina p. 385, chlor. Thera p. 29; Raul. cret. p. 833; Bois. fl. or. IV. p. 636. — *R. prostratus* Mazz. l. c. — Exsicc.: Baen. herb. europ. n. 9326.

Frutex; ramis crebre foliosis; foliis persistentibus, linearibus, margine revolutis, supra glabris, nitidis, subtus cano-tomentellis; floribus breviter pedicellatis, in racemos breves, axillares dispositis; bracteis parvis, caducis; calyce pulverulento-puberulo; corolla coerulescente. — Planta valde aromatica.

In collibus regionis calidae, sponte ut videtur rare, sed frequenter colitur et tunc plerumque subsponte occurrens. Indicatur: Boeotia (Sibth.): mt. Messapios, Klephtovuni (Fraas); Attica: pr. Athenas, Kephissia, in colle Turcovuni (Heldr.), insula Aegina (Friedr.); Laconia (Chaub.); Cycladum insula Melos (Sibth.), Thera (Heldr.); Creta: pr. Melezes in eparchia Pedhiada (Raul.); Zante (Sibth.): pr. Sculicado (Marg.); Cephalonia (Dall.); Corcyra: pr. Kastrades (Baen.), Garunna, Synarades, Afiona, Palaeokastrizza (Mazz.). — Sept. Jun. ♄

6. Zizyphora L. gen. n. 30.

1. **Z. capitata** L. sp. p. 21; Friedr. Reise p. 276; Fraas fl. class. p. 183; Bois. fl. or. IV. p. 586; Hal. Beitr. fl. Epir. p. 35, Beitr. fl. Achaia p. 29; Bald. riv. coll. bot. Alb. 1895 p. 61, 1896 p. 86; Haussk. symb. p. 37; Heldr. fl. Aegina p. 385. — Icon: Fl. gr. I. p. 10 t. 13. — Exsicc.: Heldr. herb. norm. n. 600.

Caule erecto vel adscendente, simplici vel saepius a basi divaricatim ramoso, tomentello; foliis lanceolatis vel linearibus, adpresse canescentibus, floralibus dilatatis, rhombeo-ovatis, acuminatis, margine longe ciliatis, capitula involucrantibus et superantibus; verticillastris in capitulum terminale congestis; calycibus brevissime pedicellatis, hispidis; corolla rosea.

In collibus saxosis, ad agrorum margines regionis inferioris et montanae. Epirus: pr. Arta, Pramanta (Hal.), mt. Micikeli, Kurenta, Prosgoli (Bald.); Thessalia: pr. Klinovo, Kalabaka, Trikala, Aivali,

Pharsalus (Haussk.); Aetolia (Sprun.); Boeotia: pr. Thebas, Lebadia (Fraas); Attica: pr. Athenas, insula Aegina (Heldr.); Achaia: pr. Kalavryta (Hal.); Argolis: pr. Poros (Friedr.); Laconia: mt. Malevo (Orph.). — Apr. Jul. ⊙

4. Tribus. **STACHYDEAE** Benth. in DC. pr. XII. p. 407.

7. Scutellaria L. gen. n. 734.

1. Sectio. *Lupulinaria* Hamilt. monogr. p. 11. — Flores tetragono-spicati vel racemosi, non secundi.

1. **S. orientalis** L. sp. p. 598; Raul. cret. p. 834, var. *angusta*; Bois. fl. or. IV. p. 682; Hal. in z. b. G. 1888 p. 761; Heldr. chlor. Parn. p. 761. — Icon: Fl. gr. t. 580.

A basi ramosissima; caulibus decumbenti-adscendentibus, pubescentibus; foliis petiolatis, ovatis vel oblongis, crenato-dentatis, supra virentibus, glabris vel pubescentibus, subtus adpresse canis, floralibus imbricatis, viridibus, integris vel utrinque 1—2 dentatis, calyces occultantibus; floribus in axillis solitariis, spicam densam formantibus; corolla pubescente, flava, rarius labio inferiore purpurascente.

β. **pinnatifida** Rchb. ic. pl. crit. I. p. 10; Bois. fl. or. IV. p. 682; Hal. in ö. b. Z. 1897 p. 325; Bald. riv. coll. bot. alb. 1896 p. 86. — Folia profunde pinnatifida. — Exsicc.: Heldr. herb. norm. n. 684; Bald. it. alb. epir. IV. n. 195.

In saxosis regionis subalpinae; in ditione ut videtur β. tantum hucusque observata. Epirus: mt. Smolika (Bald.); Aetolia: mt. Tymphrestus (Sprun.), Korax (Tunt.); mt. Kiona (Hal.), Parnassus (Heldr.); Achaia: mt. Kyllene supra Trikala (Pichl.); Creta (Raul.). — Jun. Aug. ♃

2. **S. alpina** L. sp. p. 599; Bois. fl. or. IV. p. 683; Hal. Beitr. fl. Epir. p. 35. — Icon: W. et K. pl. rar. Hung. t. 137. — Exsicc.: Hal. it. gr. sec. a. 1893.

A basi ramosissima; caulibus decumbenti-adscendentibus, pubescentibus; foliis ovatis vel oblongis, crenatis, utrinque viridibus, pubescentibus, inferioribus breviter petiolatis, superioribus subsessilibus, floralibus imbricatis, purpurascentibus, integris, calyces occultantibus; floribus in axillis solitariis, spicam densam formantibus; corolla pubescente, violacea, labio inferiore albido.

In saxosis regionis alpinae, rarissime. Hucusque tantum in cacumine mt. Peristeri in Epiro (Hal.) et mt. Olenos Elidis (Heldr.). — Jul. Aug. ♃

2. Sectio [illegible] Benth. et Hook. gen. II. p. 1202. — Flores op[illegible]

[illegible]ula petiolata, floralia bracteaeformia.

[illegible]ngior, purpurea.

3. **S. Columnae** All. fl. ped. I. p. 40 t. 84; Ch. et B. exp. p. 173, Fl. pelop. p. 39; Bois. fl. or. IV. p. 687; Hal. Beitr. fl. Epir. p. 35; Bald. riv. coll. bot. Alb. 1895 p. 63; Haussk. symb. p. 37. — Exsicc.: Orph. fl. gr. n. 545; Bald. it, alb. epir. III. n. 82.

Caulibus erectis, simplicibus vel parce ramosis, inferne breviter tomentellis, superne bracteis calycibusque glanduloso-villosis, petiolatis, pubescentibus, a basi truncata vel subcordata ovatis vel ovato-oblongis, grosse crenatis; racemis elongatis, laxis; bracteis subpetiolatis, ovatis, acutis, integris, calyce brevioribus; corolla purpurea, glanduloso-pubescenti; nuculis granulatis, minutissime stellatim-puberulis.

In umbrosis montanis et subalpinis. Epirus: mt. Zalongos distr. Prevesa (Bald.), pr. Kalentini supra Arta (Hal.); Thessalia: mt. Ghavellu, in oropedio Neuropolis (Haussk.), mt. Olympus pr. Hagios Dionysios (Orph.); Aetolia: pr. Karpenisi (Samar.); Euboea: mt. Telethrion (Heldr.), pr. Kurbatzi (Wild.); Peloponnesus: mt. Malevo (Orph.), pr. Scardamyle, Petalidi, Methone, Phigalea, mt. Kupe (Chaub.), sed loca a Ch. et B. indicata, saltem nonnulla probabiliter ad *S. Sibthorpii* pertinent. — Apr. Jul. ♃

β. Corolla calyce 3—4 plo longior.

× Corolla violacea.

4. **S. peregrina** L. sp. p. 599; Clem. sert. p. 78; Bois. fl. or. IV. p. 688; Haussk. symb. p. 37. — *S. adenotricha* Bois. et Heldr; in Heldr. pl. exsicc. a 1852 n. 2749. — *S. peregrina v. adenotricha* Bois et Heldr. diagn. ser. 2 IV. p. 27; Hal. Beitr. fl. Epir. p. 35. Form. in Ver. Brünn 1895 p. 37. — *S. Linnaeana* Caruel in Parl. fl. ital. VI. p. 322; Bald. in nuovo giorn. bot. it. 1894 p. 100. — *S. Columnae v. Sibthorpii* Bald. riv. coll. bot. alb. 1895 p. 63, non Benth. — Exsicc.: Heldr. l. c., herb. norm. n. 84; Orph. fl. gr. n. 430; Bald. it. alb. epir. III. n. 195; Sint. it. thessal. n. 691.

Caulibus erectis vel adscendentibus, paniculatim-ramosis, glabris vel glabriusculis, superne praeter indumentum breve crispulum, breviter glanduloso-pilosis; foliis petiolatis, laete viridibus, glabris, a basi truncata vel subcordata ovatis, grandidentatis; racemis elongatis, laxis; bracteis subpetiolatis, ovatis, acutis, integris, calyce longioribus; corolla coeruleo-violacea, dense pubescente; nuculis punctulatis, pube stellata minutissima canescentibus. — Differt a praecedente caulibus paniculatim ramosis, inferne glabris, superne breviter hirtis, foliis minoribus glabris, bracteis longioribus, corolla breviore, coerulea, dense pubescente et nuculis magis puberulis.

In rupestribus regionis montanae et subalpinae. Epirus: pr. Hagios Paraskevi, Sadovica, mt. Micikeli (Form.), pr. Syraku (Bald.) et Kalarrytes (Hal.) ad radices mt. Peristeri; Thessalia: mt. Baba, pr. Chaliki, mt. Karava, Ghavellu (Haussk.); Aetolia: mt. Korax (Heldr.); Phocis: mt. Parnassus (Heldr.); Boeotia (Sprun.); Attica: mt. Parnes (Orph.); Corcyra (Clem.): pr. Spartilla (Bald.). — Maio, Jul. ♃

5. **S. Sibthorpii** Benth. in DC. pr. XII. p. 419 pro var. *S. Columnae*; Bois. et Reut. diagn. ser. 2 IV. p. 28; Hal. in ö. b. Z. 1892 p. 375; Haussk. symb. p. 37. — *S. peregrina* S. et S. fl. gr. VI. p. 66 t. 582, non L. — *S. peregrina v. Sibthorpii* Bois. et R. l. c., Fl. or. IV. p. 688; Hal. Beitr. fl. Achaia p. 29; Form. in Ver. Brünn 1896 p. 72, 1897 p. 53. — *S. peregrina v. parnassica* Bois. diagn. ser. 2 IV. p. 28, Fl. or. IV. p. 688; Heldr. chlor. Parn. p. 25; Haussk. symb. p. 37; (forma humilior, plerumque simplex). — *S. altissima* Form. in D. bot. Mon. 1891 p. 31, non L. — Exsicc.: Orph. fl. gr. n. 66; Heldr. herb. norm. n. 85; Sint. et Bornm. it. turc. n. 1421; Sint. it. thessal. n. 465.

Caulibus erectis vel adscendentibus, paniculatim ramosis, rarius simplicibus, velutino-hirtis, superne simulque glanduloso-pilosis; foliis petiolatis, velutino-canescentibus, a basi truncata vel subcordata ovatis, grandidentatis; racemis elongatis, laxis; bracteis subpetiolatis, ovatis. integris, calyce longioribus; corolla coeruleo-violacea, dense pubescente; nuculis punctulatis, pube stellata minutissima canescentibus. — Indumento canescenti praesertim foliorum a praecedente meo sensu specifice discedit.

In rupestribus montanis et subalpinis. Thessalia: pr. Kalabaka, Meteora, mt. Agrapha, Chassia, pr. Limogardi, Neraida, Portaria (Form.), Pharsalus, Aivali, Ormanmagula (Hausk.), in fauce Megarema pr. Litochori in Olympo (Sint.); mt. Parnassus, mt. Cithaeron, Pateras et Parnes Atticae, mt. Dirphys Euboeae (Heldr.); Achaia: mt. Kyllene pr. Gura (Hal.) et Trikala (Heldr.); Laconia: mt. Malevo pr. Hagios Joannes (Orph.); insula Naxos (Sart.). — Maio, Jul. ♃

×× Corolla purpurea.

6. **S. rupestris** Bois. et Heldr. diagn. VII. p. 60. — *S. hirta β. brachystegia* Bois. fl. or. IV. p. 690. — *S. peregrina v. Sibthorpii* Hal. in ö. b. Z. 1896 p. 16, 1897 p. 98; non Bois. — Exsicc.: Pichl. pl. gr. a. 1876; Heldr. et Hal. fl. sporad. a. 1896.

Caulibus diffusis vel adscendentibus, ramosis, patule hispidis, superne simulque glanduloso-pilosis; foliis petiolatis, pilosis, canescentibus, a basi truncata vel cordata ovatis, grosse crenatis; racemis laxis; bracteis subpetiolatis, ovatis, acutis, integris, calyce sublongioribus; calyce villoso; corolla purpurea, dense pubescente; nuculis granulatis, pube stellata minutissima canescentibus. — Species certe propria, nec cum *S. hirta*, racemis densifloris et corolla lutea diversa, nec cum *S. Sibthorpii*, indumento brevi et corolla violacea distincta, commutanda. Planta Sporadum certe non *S. Sibthorpii* et omnibus notis huc spectare videtur.

In saxosis. Laconia: in regione alpina mt. Taygetos (Heldr.); Cephalonia: mt. Aenos (Heldr.); Sporadum insula Scopelos et Jura (Leon.). — Maio Jul. in alpinis Aug. ♃

× ochroleuca.

7. **S. Sieberi** Benth. in DC. pr. XII. p. 420; Raul. cret. p. 834; Bois. fl. or. IV. p. 689. — *S. hirta* Sieb. avis. p. 4, non S. et S. — Huc spectare videtur: *S. cretica* Sieb. in Flora I. p. 271, non L., et *S. albida* Weiss in z. b. G. 1869 p. 743, Raul. cret. p. 834. — Exsicc.: Rev. pl. cret. n. 144; Dörfl. pl. cret. n. 83.

Caulibus erectis, 30—50 cm. altis, ramosis, brevissime et adpresse pubescentibus, superne bracteis calycibusque pilis patulis, glandulosis intermixtis, villosis; foliis petiolatis, adpresse canescentibus, a basi truncata vel subcordata ovatis, grosse crenatis; racemis confertifloris; bracteis subpetiolatis, oblongis, integris, acuminatis, calyce longioribus; corolla ochroleuca, labio superiore rubello, pubescente, calyce 4 plo longiore; nuculis granulatis, pube stellata minutissima obsitis.

In rupestribus regionis inferioris et montanae. Creta: pr. Kissamos (Rev.), ad promontorium Meleka, pr. Malaxa, Canea, Aradhena, Voriza (Raul.) ad mt. Ida, in fauce Perivoglia (Sieb.), pr. Anatoli (Heldr.) et Males (Leon.) distr. Hierapetra, pr. Kritza distr. Mirabello (Heldr.). — Apr. Jun. ♃

8. **S. hirta** S. et S. pr. I. p. 425, Fl. gr. VI. p. 66 t. 583; Raul. cret. p. 834; Bois. fl. or. IV. p. 690; Spreitz. in z. b. G. 1890 p. 298. — *S. decumbens* Sieb. avis p. 4, rem. p. 5. — Exsicc.: Rev. pl. cret. a. 1883.

Caulibus erectiusculis vel adscendentibus, 10—25 cm. altis, simplicibus vel ramosis, praeter indumentum adpressum pilis longis patulis, superne glandulosis intermixtis, obsitis; foliis petiolatis, utrinque patule pilosis, a basi truncata vel subcordata ovatis, grosse crenatis, canescentibus; racemis confertifloris; bracteis subpetiolatis, ovatis, acutis, integris, calyce sublongioribus; corolla ochroleuca, labio superiore rubello, pubescente, calyce 3 plo longiore; nuculis granulatis, pube stellata minutissima obsitis. — Praecedenti maxime affinis, sed humilis et caulibus fragilibus, saepe decumbentibus, bracteis non acuminatis, corolla breviore et indumento patulo ut videtur specifice distincta.

In rupestribus regionis subalpina et superioris. Creta (Sibth.): ad Hellinoseli, mt. Hagion Pneuma, Stravopodia (Raul.), Hagios Theodoros, Mavrus Lakus (Spreitz.) in mt. Sphacioticis, mt. Ida et Aphendi Kavutsi (Heldr.). — Maio, Jul. ♃

b. Folia caulina subpetiolata, floralia conformia.

9. **S. galericulata** L. sp. p. 599; Form. in Ver. Brünn 1895 p. 36; Bald. riv. coll. bot. alb. 1896 p. 86. — Icon: Rchb. t. 55. — Exsicc.: Orph. fl. gr. n. 1041 (Macedonia); Heldr. it. gr. septentr. a. 1879.

Parce brevissimeque puberula; rhizomate repente; caulibus erectis, saepissime ramosis; foliis a basi cordata vel truncata oblongo-lanceolatis, remote et obtusissime crenatis, floralibus sursum decrescentibus; floribus axillaribus, oppositis, secundis; corolla coerulea, calyce multoties longiore; nuculis tuberculatis, glabris.

In pratis humidis regionis inferioris et montanae, rarissime. Epirus: ad lacum Janina (Form.); Eurytania: ad Laspi pr. Karpenisi (Heldr.). — Jul. Aug. ♃

8. **Prunella** L. gen. n. 735.

1. **P. vulgaris** L. sp. p. 600; S. et S. pr. I. p. 425; Pieri corc. fl. p. 81; Ch. et B. exp. p. 174, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 273; Ung. Reise p. 127; Raul. cret. p. 834; Bois. fl. or. IV. p. 691; Form. in D. bot. Mon. 1891 p. 31, in Ver. Brünn 1895 p. 37, 1896 p. 72, 1897 p. 53; Hal. Beitr. fl. Epir. p. 35, Beitr. fl. Thessal. p. 17; Haussk. symb. p. 37 *f. parvi- et grandiflora*. — Icon: Fl. dan. t. 910. — Exsicc.: Sint. it. thessal. n. 990; Baen. herb. europ. n. 9315 *f. Herculis* Borb. („f. robusta grandifolia, foliis oblongo-ovatis").

Glabriuscula vel pubescens; rhizomate repente; caulibus adscendentibus vel erectis, simplicibus vel ramosis; foliis petiolatis, ovato-oblongis oblongisve, integris vel dentatis; verticillastris 6 floris, spicam oblongam, densam formantibus; bracteis orbiculatis, ciliatis, abrupte acuminatis; calycis dentibus ciliatis, labii superioris brevissimis, truncatis, mucronatis, inferioris lanceolatis; corolla violacea.

β. **cretica** Bois. fl. or. IV. p. 692. — Pusilla, macra, calycis labium superius subintegrum. — Exsicc.: N. v.

In olivetis, nemorosis, silvaticis regionis inferioris et montanae. Frequens in Pindo epirotico et thessalo; in mt. Oxya, Pelion (Form.); Attica: pr. Athenas, mt. Pentelicon (Haussk.); Peloponnesus: pr. Carithena, mt. Kupe, Taygetos, pr. Kardamyle (Chaub.), insula Poros (Friedr.) et certe alibi; Creta: pr. Platania Enneachoria (Raul.); Zante (Marg.); Corcyra (Pieri): pr. Psarari (Baen.); non indicatur in insulis maris Aegaei; — *β.* in quercetis meridionalibus mt. Ida supra Voriza. — Maio, Aug. ♃

2. **P. laciniata** L. sp. p. 600; Ch. et B. exp. p. 174, Fl. pelop. p. 39; pro var. *P. vulgaris*; L. sp. ed. 2 p. 837; S. et S. pr. I. p. 426; Pieri corc. fl. p. 81; Marg. et R. fl. Zante p. 75; Raul. cret. p. 834; Form. in D. bot. Monat. 1891 p. 31, in Ver. Brünn 1895 p. 37; Hal. Beitr. fl. Epir. p. 35, Beitr. fl. Thessal. p. 17, Beitr. fl. Achaia p. 29; Haussk. symb. p 37, *f. micrantha, macrantha, laciniata et integrifolia*. — *P. alba* Pall. ap. M. a B. fl. taur. cauc. II. p. 67; Bois. fl. or. IV. p. 692; Form. in Ver. Brünn 1896 p. 72, 1897 p. 54. — Icon: Jacq. fl. austr. t. 378. — Exsicc.: Sint. it. thessal. n. 464.

Pubescens; rhizomate repente; caulibus adscendentibus, simplicibus vel ramosis; foliis petiolatis, ovato-oblongis oblongisve, subintegris vel dentatis vel pinnatifidis; verticillastris 6 floris, spicam oblongam, densam formantibus; bracteis orbiculatis, hirtis, abrupte acuminatis; calycis dentibus ciliatis, labii superioris late ovatis, truncatis, abrupte mucronatis, inferioris lanceolatis; corolla alba; rarius carnea.

In nemorosis, silvaticis regionis inferioris et montanae. Frequens in Pindo epirotico et thessalo; mt. Chassia (Form.), Olympus (Heldr.); Attica: mt. Parnes (Heldr.), Pentelicon (Haussk.); Peloponnesus: pr. Patras (Hal.), pr. Carithena, mt. Kupe, pr. Kardamyle, mt. Taygetos (Chaub.), mt. Selitza pr. Kalamata (Zahn), mt. Malevo (Orph.); Creta (Sibth.): pr. Voriza (Raul.); Zante (Marg.); Corcyra (Pieri): pr. Psorari (Baen.). — Maio, Aug. ♃

1 × 2. **P. vulgaris × laciniata.** — *P. intermedia* Link in Annal. Naturgesch. I. (1791) sec. Kern. fl. exs. austro-hung. n. 1420. — *P. hybrida* Knaf in Lotos XIV. p. 84; Haussk. symb. p. 37. — Hybrida, characteribus inter parentes intermedia, foliis pinnatifidis, floribus violaceis; nunc parce pubescens, calycibus ut in *P. vulgaris*, sed foliis pinnatifidis; nunc dense pubescens, calycibus ut in *P. laciniata*, sed floribus violaceis. — Prope Malakasi, Klinovo, mt. Ghavellu et pr. Neuropolis Thessaliae, et in mt. Pentelicon Atticae (Haussk.). — N. v.

9. Sideritis L. gen. n. 712.

1. Sectio. *Empedoclea* Benth. lab. p. 574. — Perennes, basi suffrutescentes; folia floralia flores amplectentia, inferioribus difformia.

a. Folia floralia verticillastris 2—3 plo longiora.

α. Folia basi cordato-amplexicaulia.

1. **S. perfoliata** L. sp. p. 575; Heldr. in delt. syllog. Parn. 1900 p. 4. — *S. pindicola* Heldr. l. c. — Exsicc.: Heldr. it. thessal. n. 115.

Breviter glanduloso-hirta, in partibus junioribus lana detersili albo-hirsuta; caulibus erectis, simplicibus vel superne opposite ramosis; foliis coriaceis, valde reticulatis, oblongis vel oblongo-lanceolatis, acutis, denticulatis vel integris, infimis in petiolum attenuatis, ceteris semi-amplexicaulibus, floralibus late ovato-lanceolatis, acuminato-attenuatis, erecto-patulis, verticillastro 2—3 plo longioribus; verticillastris distinctis, inferioribus distantibus; calycis dentibus lanceolatis, tubo dimidio brevioribus; corollae flavae tubo incluso. — Species foliis caulibusque demum glabrescentibus, pube brevissima glandulosa adspersis, coriaceis, virentibus, caulinis basi non attenuata semiamplexicaulibus, floralibus latissimis, verticillastra fere omnino occultantibus, insignis.

In saxosis dumosis regionis montanae, rarissime. Thessalia: mt. Agrapha in Pindo (Heldr.). — Jun. Jul. ♃

β. Folia basi angustata.

2. **S. theezans** Bois. et Heldr. diagn. VII. p. 58, Fl. or. IV. p. 710; Heldr. Nutzpfl. p. 33, in delt. syll. Parn. 1900 p. 4. — *S. cretica* S. et S. pr. I. p. 400 p. p. quoad pl. lacon.; Ch. et B. exp. p. 163, Fl. pelop. p. 36; non L. — *Phlomis clandestina* Ch. et B. exp. p. 170 t. 20. — *S. syriaca* Ch. et B. fl. pelop. p. 36

t. 21; Fraas fl. class. p. 175 p. p.; non L. — Exsicc.: Orph. fl. gr. n. 14; Heldr. herb. norm. n. 1367 et 1469.

Caulibus dense tomentosis, adscendentibus vel erectis, simplicibus vel superne opposite ramosis; foliis crassiusculis, dense cano-tomentosis, obsolete crenulatis vel integris, inferioribus oblongo-spathulatis obtusis, petiolatis, superioribus late lineari-lanceolatis, sessilibus, floralibus subreticulatis, interdum tenuius tomentosis, a basi subrotunda vel ovata abruptiuscule longe lanceolato-acuminatis, patulis, verticillastro duplo longioribus; verticillastris inferioribus distantibus, ceteris in spicam cylindricam condensatis; calycis adpresse pubescentis dentibus lanceolatis, tubo subbrevioribus; corollae luteae tubo incluso. — Indumento cano-tomentoso, non glanduloso, foliis floralibus minus amplis, abruptiuscule acuminatis a praecedente distincta.

β. **peloponnesiaca** Bois. et Heldr. diagn. ser. 2 IV. p. 32; Heldr. Nutzpfl. p. 34, in delt. syll. Parn. 1900 p. 4; pro sp. — *S. theezans v. cyllenea* Bois. fl. or. IV. p. 711. — Verticillastra magis distantia, folia floralia brevius acuminata, tenuius tomentosa. — Exsicc.: Heldr. pl. fl. hellen. a. 1871 et 1881; Pichl. pl. gr. a. 1876.

In saxosis regionis abietinae et superioris. Laconia: mt. Malevo, Taygetos (Heldr.); — β. Achaia: mt. Kyllene (Heldr.). — Jun. Jul. ♃

3. **S. scardica** Griseb. spic. II. p. 144; Bois. fl. or. IV. p. 170; Form. in Ver. Brünn 1896 p. 72; Heldr. in delt. syll. Parn. 1900 p. 4. — *S. florida* Bois. et Heldr. diagn. ser. 2 IV. p. 31; Heldr. Nutzpfl. p. 34. — Exsicc.: Orph. fl. gr. n. 539; Sint. et Bornm. it. turc. n. 1429 et 1881.

Adpresse albo-lanata; caulibus adscendentibus vel erectis, simplicibus vel superne opposite ramosis; foliis obsolete crenulatis, inferioribus oblongo-lanceolatis, obtusis, in petiolum attenuatis, superioribus lineari-lanceolatis, acutiusculis, sessilibus, floralibus reticulatis, brevissime glanduloso-pubescentibus, a basi subrotunda vel ovata abruptiuscule in cuspidem patentissimam, apice subrecurvam attenuatis, verticillastro subduplo longioribus; verticillastris in spicam crassam, oblongam confertis; calycis hirsuti dentibus lanceolatis, tubo aequilongis; corollae luteae tubo incluso. — Spica densissima, ad 8 cm. longa et 3 cm. lata, foliis floralibus e spica horizontaliter exsertis, acutissimis, demum glabrescentibus et flavidis, corolla magna, calyce subduplo longiore, labio profunde bifido ab omnibus affinibus facile distinguitur.

β. **pelia.** — *S. scardica* Heldr. in Sitzungsb. acad. wiss. Berlin 1883 p. 4; Form. in D. bot. Mon. 1891 p. 30, quoad pl. thessal.; Hal. Beitr. fl. thessal. p. 17. — Verticillastra inferiora distantia, superiora in spicam angustiorem conferta; folia floralia brevius acuminata, densius tomentella. — Transitum ad *S. Roeseri* praebere videtur. — Exsicc.: Heldr. it. thessal. n. 3.

In regione silvatica mt. Olympus Thessaliae ad monasterium Hagios Dionysios (Heldr.) et in mt. Godaman (Form.); — β. in regione media et superiore mt. Pelion (Heldr.). — Jul. Aug. ♃

b. Folia floralia verticillastris non vel parum longiora.

α. Folia floralia viridia, reticulata, brevissime glanduloso-puberula.

4. **S. Roeseri** Bois. et Heldr. diagn. ser. 2 IV. p. 30, Fl. or. IV. p. 710; Heldr. Nutzpfl. p. 34, chlor. Parn. p. 25, in delt. syll. Parn. 1900 p. 5; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Epir. p. 36, in ö. b. Z. 1897 p. 325; Bald. riv. coll. bot. alb. 1895 p. 64; Form. in Ver. Brünn 1896 p. 72, 1897 p. 53; Haussk. symb. p. 45. — *S. syriaca* Fraas fl. class. p. 175 p. p., non L. — *S. attica* Heldr. in delt. syllog. Parn. 1900 p. 5, f. magis incana. — Exsicc.: Heldr. herb. norm. n. 490; Orph. fl. gr. n. 429; Sint. it. thessal. n. 1142; Dörfl. fl. gr. n. 212.

Caulibus adpresse tomentosis, adscendentibus vel erectis, simplicibus vel ramosis; foliis integris vel obsolete crenulatis, infimis cano-tomentosis, elliptico-spathulatis, obtusis, petiolatis, ceteris oblongo-linearibus, acutiusculis, sessilibus, plus minus canis vel virescentibus, floralibus a basi rotundata vel ovata breviter acuminatis, reticulatis, glanduloso-puberulis, virescentibus, calyces non aequantibus; verticillastris distantibus; calycis glandulosi dentibus lanceolatis, tubo subbrevioribus; corollae luteae tubo non exserto. — Species ab italica *S. sicula* Ucr. pl. ad Linn. op. add. n. 16 vix specifice diversa, variat quoad indumentum, foliorum formam et verticillorum magnitudinem.

β. **lanceolata.** — Foliis elongatis, anguste lineari-lanceolatis, floralibus amplis, longius acuminatis. — Exsicc.: Bald. it. alb. epir. III. n. 81.

In saxosis regionis mediae et superioris. Epirus: mt. Peristeri pr. Kalarrytes (Hal.); Thessalia: mt. Baba, Karava, Ghavellu, Gionskala in Pindo (Haussk.), mt. Pelion, mt. Pirgo, Alafovris, Mavrika et Katasara in mt. Othrys (Form.), mt. Oeta (Heldr.); Aetolia: mt. Tymphrestus (Fraas), Arapocephala, Chelidoni, Korax (Heldr.); mt. Kiona (Hal.), Parnassus (Heldr.); Attica: mt. Pateras (Pichl.); — β. Epirus: mt. Murga supra Suli et Sirizana distr. Paramythia (Bald.). — Jun. Aug. ♃

β. Folia floralia dense tomentosa.

5. **S. euboea** Heldr. in delt. syll. Parn. 1900 p. 4. — *S. cretica* S. et S. pr. I. p. 400 p. p. quoad pl. euboeam, non L. — *S. syriaca v. condensata* Bois. et Heldr. diagn. ser. 2 IV. p. 31, non *S. condensata* Bois. et Heldr. diagn. XII. p. 71. — *S. taurica* Heldr. Nutzpfl. p. 34. — *S. cretica v. condensata* Bois. fl. or. IV. p. 708. — Exsicc.: Heldr. herb. norm. n. 793.

Albo-tomentosa; caulibus adscendentibus vel erectis, simplicibus vel parce ramosis; foliis oblongis, obtusis, integris vel crenulatis, inferioribus in petiolum attenuatis, ceteris breviter petiolatis subsessilibusve, floralibus ovato-rotundatis, breviter acuminatis, calyces aequantibus; verticillastris omnibus vel superioribus in spicam cylindricam condensatis; calycis lanati dentibus lanceolatis, tubo subduplo brevioribus;

corollae luteae tubo subincluso. — Sequenti affinis, ab ea indumento verticillorum minus denso, foliis mediis obtusis, subpetiolatis et praesertim verticillis condensatis discedit.

In saxosis regionis superioris mt. Dirphys Euboeae (Heldr.). — Jul. Aug. ♃

6. **S. syriaca** L. sp. p. 574; S. et S. pr. I. p. 400, Fl. gr. VI. p. 41 t. 550; Sieb. avis p. 4, in Flora I. p. 275; Heldr. Nutzpfl. p. 34; Raul. cret. p. 835. — *S. cretica* Bois. fl. or. IV. p. 708; Spreitz. in z. b. G. 1890 p. 298; Bald. viagg. Creta p. 85; Heldr. in delt. syll. Parn. 1900 p. 4; non L. sp. p. 574, quae (*S. candicans* Ait. hort. Kew. II. p. 289) insulae Teneriffae incola et erronee ab autore in Creta indicatur. — *S. Boissieri* Magn. in scrin. fl. sel. IV. p. 85. — Exsicc.: Rev. pl. cret. n. 137, in Magn. fl. sel. n. 944; Bald. it. cret. n. 88, it. cret. alt. n. 242.

Niveo-tomentosa; caulibus simplicibus vel rarius ramosis; foliis oblongis oblongo-lanceolatisve, integris vel crenulatis, infimis breviter petiolatis, obtusis, ceteris sessilibus, acutiusculis, floralibus ovato-rotundatis, breviter acuminatis, calyces aequantibus; verticillastris distantibus; calycis dentibus lanceolatis, tubo subduplo brevioribus; corolla luteae tubo subincluso. — Omnium sectionis densissime pannosa, verticillastris parvis, avellanae minoribus.

In saxosis regionis montanae et subalpinae Cretae: frequens in mt. Sphacioticis et in mt. Ida. — ♃

2. Sectio. *Hesiodia* Benth. lab. 582. — Annuae; folia floralia inferioribus conformia, sed minora.

a. Calycis dentes subaequales, supremus ceteris paulo major.

α. Planta patule villosissima; corolla alba, labiis atro purpureis.

7. **S. lanata** L. sp. ed. 2 p. 804; Raul. cret. p. 835; Bois. fl. or. IV. p. 705; Haussk. symb. p. 45; Heldr. fl. Aegina p. 385, chlor. Thera p. 19. — *S. elegans* Murr. nov. comm. Gott. 1778 p. 92 t. 4; Friedr. Reise p. 272; Fraas fl. class. p. 175. — *S. nigricans* Pers. syn. II. p. 117. — Exsicc.: Heldr. herb. norm. n. 871.

Longe et patule villosa, a collo saepius pluricaulis; caulibus saepius ramosis, erectis vel adscendentibus; foliis ovatis, crenatis, inferioribus petiolatis, ceteris sessilibus, floralibus deflexis, verticillastro longioribus; verticillastris numerosis, distinctis, approximatis; calycis villosi subbilabiati dentibus lanceolatis, spinula terminatis, tubo subbrevioribus, supremo ceteris paulo majore; corollae limbo subexserto.

In maritimis, cultis regionis inferioris. Attica (Fraas): insula Aegina (Heldr.); Argolis: pr. Vromolimni in peninsula Methana (Friedr.); Cycladum insula Keos (Heldr.), Thera (Letourn.); Creta (Sieber). — Apr. Maio. ⊙

β. Planta plus minus lanata; corolla lutea, demum fuscescens.

8. **S. montana** L. sp. p. 575; Bois. fl. or. IV. p. 706; Form. in D. bot. Mon. 1891 p. 30, in Ver. Brünn 1895 p. 36, 1896 p. 72,

1897 p. 53; Bald. riv. coll. bot. alb. 1895 p. 64, 1896 p. 87; Haussk. symb. p. 45. — Icon: Jacq. fl. austr. t. 434. — Exsicc.: Sint. et Bornm. it. turc. n. 1434; Sint. it. thessal. n. 533.

Longe et laxe pilosa, a collo saepe pluricaulis; caulibus simplicibus vel ramosis, erectis vel adscendentibus; foliis oblongo-lanceolatis, integris vel antice dentatis, inferioribus subpetiolatis, ceteris sessilibus, floralibus oblongis, spinuloso-mucronatis, verticillastro sublongioribus, summis saepe flavidis, comatis (*f. comata* Bois. l. c.); verticillastris numerosis, distinctis, approximatis; calycis patule pilosi, subbilabiati dentibus subaequalibus, oblongis, spinuloso-mucronatis, tubo subbrevioribus, demum patentibus, supremo ceteris parum majore; corollae limbo breviter exserto.

In collibus siccis regionis inferioris et montanae. Epirus: pr. Prevesa, Mitikas (Form.), mt. Smolika, Olycika, Micikeli (Bald.); Thessalia: mt. Baba pr. Klinovo (Heldr.), pr. Kastania, Vendista, Kalabaka, Trikala, Pharsalus, Velestinos, Hagios Georgios, Kapurna, Koryza, Volo, Konisko, Neraida, pr. Drepani, Miluna, Patsios et Rapsani in Olympo (Form.), pr. Plaka ad Litochori (Sint.); Corcyra: pr. Kanali, Potamo (Form.). — Jun. Jul. ⊙

9. **S. remota** Urv. enum. p. 66; Bois. fl. or. IV. p. 707; Haussk. symb. p. 45; Form. in D. bot. Mon. 1898 p. 79. — *S. montana* S. et S. pr. I. p. 400, Fl. gr. VI. p. 42 t. 551; et probabiliter Ung. Reise p. 127; non L. — Exsicc.: Orph. fl. gr. n. 428; Heldr. herb. norm. n. 386.

Albo-lanata, a collo saepius pluricaulis; caulibus simplicibus vel ramosis, rigidis; foliis oblongo-spathulatis, integris vel remote denticulatis, inferioribus subpetiolatis, ceteris sessilibus, floralibus caulinis similibus, mucronatis, verticillastro brevioribus vel sublongioribus, summis interdum flavidis, comatis; verticillastris numerosis, saepius distantibus; calycis albo-lanati, subbilabiati dentibus subaequalibus, spinuloso-mucronatis, tubo aequilongis, demum patentibus, supremo ceteris vix majore; corollae limbo breviter exserto. — Praecedenti maxime affinis, ab ea caulibus rigidis, crassioribus, verticillastris saepius remotis, indumento lanato et calycis dentibus elongatis, spina validiore terminatis discedit.

In collibus siccis. Attica: pr. Athenas (Heldr.), Rheitro (Ung.), Kamariza Laurii (Haussk.); Euboea (Sprun.); Acrocorinthus (Haussk.). — Apr. Maio. ⊙

b. Calycis dens supremus ceteris multo latior et subduplo longior.

10. **S. romana** L. sp. p. 575; S. et S. pr. I. p. 401, Fl. gr. VI. p. 43 t. 552; Sieb. avis rem. p. 5; Ch. et B. exp. p. 164, Fl. pelop. p. 36; Friedr. Reise p. 270 et 284; Fraas fl. class. p. 175; Ung. Reise p. 127; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 835; Bois. fl. or. IV. p. 706; Heldr. fl. cephal. p. 58, Fl. Aegina p. 385; Spreitz. in z. b. G. 1887 p. 666; Haussk. symb. p. 45; Form. in Ver. Brünn 1897 p. 53, in D. bot. Mon. 1898 p. 79; Hal. in ö. b. Z. 1897 p. 98. — Exsicc.: Orph. fl. gr. n. 537; Dörfl. pl. cret. n. 36.

Molliter pilosa, simplex vel saepius a collo ramosa; caulibus adscendentibus vel diffusis; foliis ovatis vel oblongis, crenato-dentatis integrisve, inferioribus subpetiolatis, ceteris sessilibus, floralibus mucronatis, verticillastro sublongioribus; verticillastris distantibus; calycis hirsuti, basi valde gibbi, subbilabiati dentibus spinuloso-aristatis, demum patentibus vel recurvis, 4 inferioribus lanceolatis, supremo multo majore, ovato; corollae albae limbo breviter exserto.

In siccis regionis inferioris et montanae. Phthiotis: pr. Stylida, Lamia (Form.); Attica: pr. Athenas (Heldr.), mt. Kerata (Haussk.), ad promontorium Perdicca in insula Aegina (Friedr.); Peloponnesus: pr. Nauplia (Friedr.), Argos, Tripoliza, Sparta, Pallantium, Gortys, Megalopolis, Zarnate, Methone, Arcadia (Chaub.); Sporadum insula Scopelos (Leon.); Cyclades (Sibth.): insula Cythnos (Heldr.), Tenos, Syra (Weiss); Creta: pr. Canea, Acroteri, insula Gaidaronisi (Raul.), Hierapetra (Leon.); Cythaera: pr. Hagios Nicolaus (Spreitz.); Cephalonia: pr. Pessada, Metaxata (Heldr.). — Mart. Maio. ⊙

11. **S. purpurea** Talbot in Benth. lab. p. 742; Marg. et R. fl. Zante p. 75; Raul. cret. p. 835; Bois. fl. or. IV. p. 706; Spreitz. in z. b. G. 1887 p. 666; Gelmi in bull. soc. bot. it. 1889 p. 451; Hal. Beitr. fl. Epir. p. 36, Beitr. fl. Aetol. p. 10, in ö. b. Z. 1896 p. 17; Haussk. symb. p. 45. — *S. romana v. purpurea* Bald. riv. coll. bot. alb. 1896 p. 87. — Icon: Rchb. t. 25. — Exsicc.: Orph. fl. gr. n. 538; Heldr. herb. fl. hellen. n. 90, in Baen. herb. europ. n. 3606, herb. norm. n. 1273; Sint. it. thessal. n. 461; Baen. herb. europ. n. 9340.

Molliter villosa, simplex vel saepius a collo ramosa; caulibus adscendentibus vel diffusis; foliis ovatis vel oblongis, crenato-dentatis integrisve, inferioribus subpetiolatis, ceteris sessilibus, floralibus mucronatis, verticillastro sublongioribus; verticillastris distantibus; calycis villosi, basi gibbi, subbilabiati dentibus spinuloso-aristatis, demum patentibus, 4 inferioribus lanceolatis, supremo multo majore, ovato; corollae purpureae limbo exserto. — Praecedenti affinis, ab ea indumento longiore, calyce basi minus gibboso, corolla purpurea, majore, magis exserto discedit.

In siccis, olivetis regionis inferioris. Epirus: pr. Kalentini supra Arta (Hal.), pr. Kastriza distr. Janina (Bald.); Thessalia: pr. Vutades (Sint.), Klinovo, Korona, Ormanmagula, Aivali, Pharsalus (Haussk.), Palaeokastro (Heldr.), ad lacum Karla (Reis.), pr. Volo (Orph.); Acarnania: pr. Kravassaras, Agrinion (Hal.); Aetolia: pr. Aetolikon (Heldr.), ad radices mt. Taphiassos (Hal.); Achaia: pr. Corinthum, Psatopyrgos (Heldr.), Patras (Hal.); Elis: pr. Chlemutzi, Lechaena (Heldr.); Messenia: pr. Diabolitzi (Zahn); Creta (Raul.); Zante (Marg.); Leucas: pr. Amaxichi, monasterium Hagios Georgios (Spreitz.); Corcyra (Talbot): pr. Signes (Gelmi), ad lacum Corissia (Bickn.). — Maio, Jul. ⊙

Obs. *S. tragoriganum* Lag. nov. gen. et sp. p. 18; Raul. cret. p. 835. — In Benth. lab. p. 581 „probabilissime ex schedulae commutatione erronee ex Creta indicatur“ (Bois. fl. or. IV. p. 714).

10. Marrubium L. gen. n. 721.

a. Calycis dentes deni.

1. **M. vulgare** L. sp. p. 853; S. et S. pr. I. p. 412; Dallap. prosp. p. 83; Ch. et B. exp. p. 169, Fl. pelop. p. 38; Friedr. Reise p. 264 et 267; Fraas fl. class. p. 180; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 835; Bois. fl. or. IV. p. 703; Heldr. fl. cephal. p. 58, chlor. Thera p. 19; Hal. Beitr. fl. Epir. p. 36; Form. in Ver. Brünn 1895 p. 36, 1896 p. 71, 1897 p. 53; Haussk. symb. p. 43. — Icon: Fl. dan. t. 1036. — *M. canescens* Sieb. avis rem. p. 5. — Exsicc.: Sint. it. thessal. n. 562; Baen. herb. europ. n. 9287.

Caulibus erectis vel adscendentibus, albo-lanatis, simplicibus vel ramosis; foliis petiolatis, ovato-orbiculatis vel ovatis, irregulariter crenatis, plus minus tomentellis, supra virescentibus, subtus canis, floralibus subsessilibus, deflexis, oblongo-cuneatis, verticillastro-longioribus; verticillastris globosis, distantibus; bracteis subulatis, uncinatis; calycis villosi dentibus subulatis, patentibus, uncinatis; corolla parva, alba, puberula.

β. **apulum** Ten. fl. nap. V. p. 16; Link. in Linn. IX. p. 576; pro sp.; Heldr. fl. aeg. p. 385, chlor. Mykon. p. 249. — *M. vulgare v. lanatum* Benth. lab. p. 591; Bald. viagg. Creta p. 85. — ? *M. candidissimum* Form. in Ver. Brünn 1895 p. 36, 1896 p. 71; vix L. sp. p. 583, quod a nemini antea vel postea nec in Epiro nec in Thessalia observatum fuit; neque in Creta, ubi in Linnaeo ipso cum ? indicatur, crescit. — Folia minora, suborbicularia, utrinque molliter et niveo-lanato-tomentosa. — Formis intermediis, in ditione vulgatioribus, ad typum transit. — Exsicc.: Heldr. herb. norm. n. 1170a.

In ruderatis, ad vias regionis inferioris et montanae per totam Graeciam. — Maio, Sept. ♃

1 × 2. **M. vulgare × peregrinum.** — *M. remotum* Kit. in Schult. öst. fl. ed. 2 II. p. 161. — *M. pannonicum* Rchb. fl. germ. p. 325. — *M. peregrinum × vulgare* Reichardt in z. b. G. 1861 p. 342; Haussk. symb. p. 44. — *M. peregrinum β. platyphyllum* Haussk. symb. p. 43 (f. dentibus calycinis, semper quinis, rectis). — Hybrida, nunc ad unum, nunc ad alterum parentium magis vergens; a *M. vulgare* caule magis ramoso, indumento densiori, foliis angustioribus, floribus paucioribus verticillastrorum, calycis dentibus 5—10, plerisque rectis; a *M. peregrino* caule minus ramoso, indumento minus denso, foliis latioribus, floribus verticillastrorum numerosioribus, calycis dentibus 5—10, saepius patentibus, discedens. — Thessalia: ad monasterium Korona, pr. Trikala, Karditza, Pharsalus (Haussk.); Euboea: pr. Kastaniotissa (Heldr.). — Exsicc.: Heldr. herb. norm. n. 1171.

b. Calycis dentes quini.

α. Caules patentim ramosissimi.

2. **M. peregrinum** L. sp. p. 582; S. et S. pr. I. p. 411; Ch. et B. fl. pelop. p. 38; Raul. cret. p. 835; Bois. fl. or. IV. p. 694;

Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 8; Hal. in z. b. G. 1888 p. 761, in ö. b. Z. 1890 p. 40, 1897 p. 325, Beitr. fl. Thessal. p. 17; Form. in D. bot. Mon. 1891 p. 30, in Ver. Brünn 1895 p. 36, 1896 p. 71, 1897 p. 53; Bald. in nuovo giorn. bot. ital. 1894 p. 100, riv. coll. bot. alb. 1896 p. 87; Haussk. symb. p. 43. — *M. creticum* Mill. dict. n. 3; S. et S. pr. I. p. 412; Fraas fl. class. p. 180. — Icon: Jacq. fl. austr. II. t. 160. — Exsicc.: Heldr. herb. norm. n. 60 et 1170 b; Orph. fl. gr. n. 521; Sint. et Bornm. it. turc. n. 1424 et 1885; Sint. it. thessal. n. 1110; Dörfl. fl. gr. n. 204.

Adpresse cano-tomentosum; caulibus erectis, patentim ramosissimis; foliis breviter petiolatis, oblongo-cuneatis, crenato-dentatis, floralibus diminutis, verticillastro longioribus; verticillastris 6—10 floris, ad ramorum apicem congestis; bracteis calyce multo brevioribus; calycis adpresse tomentosi dentibus lanceolatis, erectis, tubo triplo brevioribus; corolla parva, alba, puberula.

In campis, ruderatis, olivetis regionis inferioris et montanae. Frequens in Epiro et Thessalia tota, a Pindo per planitiem usque ad mare Aegaeum, nec non in Phthiotide; Locris: pr. Granitza (Tunt.): Doris: pr. Amphissa, Itea (Hal.); ad radices Parnassi pr. Rachova (Heldr.); Boeotia: pr. Orchomenos (Sibth.); Attica borealis (Fraas); Euboea: pr. Kastaniotissa (Heldr.), Chalkis (Fraas); Arcadia: ad lacum Pheneon (Orph:); Creta (Sibth.). — Jun. Sept. ♃

β. Caules simplices. Species inter se valde affines et notis minus gravioribus tantum distinctae.

× Calyces dentes tubo manifeste breviores, corollam non aequantes.

3. **M. thessalum** Bois. et Heldr. diagn. ser. 2 IV. p. 50, Fl. or. IV. p. 699; Form. in Ver. Brünn 1896 p. 71. — Exsicc.: Orph. fl. gr. n. 520.

Caulibus erectis vel adscendentibus, lanato-tomentosis; foliis petiolatis, ovatis oblongisve, basi cuneatis, crenatis, supra sericeo-lanatis, subtus cano-tomentosis, floralibus subsessilibus, oblongo-cuneatis, verticillastro longioribus; verticillastris multifloris, globosis, distantibus; bracteis subulatis, lanato-villosis, tubo calycino subbrevioribus; calyce breviter tomentoso, dentibus subulatis, rectis, villosis, tubo duplo brevioribus; corolla alba, tomentella, calycis dentes superante. — Indumento foliorum pannoso et verticillastris lana longa, bracteas et calyces recondenti, obsitis, ab affinibus facile distinguenda.

In regione subalpina et alpina mt. Olympus et Ossa Thessaliae (Heldr.). — Jul. Aug. ♃

4. **M. cyllenum** Bois. et Heldr. diagn. ser. 2 IV. p. 51. — *M. velutinum v. cylleneum* Bois. fl. or. IV. p. 702. — Exsicc.: Heldr. herb. norm. n. 967; Orph. fl. gr. n. 299; Dörfl. fl. gr. n. 423.

Caulibus erectis vel adscendentibus; adpresse-tomentosis; foliis petiolatis, orbiculatis vel ovato-orbiculatis, basi rotundatis crenatis, supra adpresse sericeis, subtus cano-tomentosis, floralibus subsessilibus, sub-

orbiculatis vel ovato-oblongis, verticillastro longioribus; verticillastris multifloris, globosis, distantibus; bracteis subulatis, breviter tomentosis, tubo calycino subbrevioribus; calyce breviter tomentoso, dentibus subulatis, patulis, breviter tomentosis, tubo triplo brevioribus; corolla flavida, tomentella, calycis dentes superante. — Indumento flavido, minus copioso, foliis rotundatis, verticillastris non lanato-villosis et calycis dentibus brevioribus, patulis a praecedente; calycis dentibus tubo triplo brevioribus a sequente discedit.

In regione silvatica et alpina mt. Chelmos et Kyllene Achaiae (Heldr.). — Jun. Aug. ♃

×× Calycis dentes tubo subaequilongi vel aequilongi, corollam superantes.

5. **M. velutinum** S. et S. pr. I. p. 412, Fl. gr. VI. p. 49 t. 561; Bois. fl. or. IV. p. 702; Hal. in z. b. G. 1888 p. 761; Heldr. chlor. Parn. p. 25; Bald. riv. coll. bot. alb. 1895 p. 63. — Exsicc.: Heldr. herb. norm. n. 364; Orph. fl. gr. n. 431; Bald. it. alb. epir. III. n. 193; Dörfl. fl. gr. n. 242.

Caulibus erectis vel adscendentibus, adpresse tomentosis; foliis petiolatis, orbiculatis vel ovato-orbiculatis, basi rotundatis, crenatis, supra adpresse-sericeis, subtus cano-tomentosis, floralibus subsessilibus, suborbiculatis vel ovato-oblongis, verticillastro longioribus; verticillastris multifloris, globosis, distantibus; bracteis subulatis, villosis vel breviter tomentosis, tubo calycino subbrevioribus; calyce breviter tomentoso, dentibus subulatis, patulis, saepe villosis, tubo subbrevioribus vel aequilongis; corolla flavida, tomentella, calycis dentes subaequante.

β. **Frivaldszkyanum** Bois. diagn. XII. p. 74; Hal. Beitr. fl. Epir. p. 36; Form. in Ver. Brünn 1896 p. 71, 1897 p. 53, pr. sp. — *M. Haussknechtii* Uechtr. in Haussk. symb. p. 44. — Folia adpresse breviterque tomentosa, superiora saepius acute dentata; verticillastra breviter tomentosa; calycis dentes tubo aequilongi. — Intermediis ad typum transit. — Planta nostra cum rumelica (Deg. pl. rum. or. exs. n. 123) optime congruit. — Exsicc.: Sint. it. thessal. n. 686.

In rupestribus regionis mediae et superioris. In mt. Parnasso (Sibth.), Kiona (Hal.), Velugo (Sprun.), Oeta (Heldr.), mt. Strungula et Micikeli (Bald.); specimina e mt. Velugo, Oeta et ex Epiro transitum ad var. *β.* praebent; — *β.* Epirus: mt. Tsumerka pr. Vulgarelion et Theodoriana, mt. Peristeri (Hal.); Thessalia: mt. Baba, Zygos, Karava, Ghavellu (Haussk.), pr. Vlachi, Kalivia, Numaria, Karamanol, Beluja, mt. Oxya, Dokimi in Pindo, mt. Pirgo, Alafovris et Mavrika in mt. Othrys, pr. Lokatamburia et Godaman in Olympo (Form.). — Jun. Aug. ♃

11. Melittis L. gen. n. 731.

1. **M. melissophyllum** L. sp. p. 597; S. et S. pr. I. p. 424; Ch. et B. exp. p. 173, Fl. pelop. p. 39; Ung. Reise p. 126; Bois. fl. or. IV. p. 754; Hal. Beitr. fl. Achaia p. 29; Bald. riv. coll. bot. alb.

1895 p. 66; Haussk. symb. p. 38. — *M. grandiflora* Sm. fl. brit. II. p. 644. — *M. albida* Guss. pr. II. p. 140. — *M. cordifolia* Bois. et Heldr. exs. — Icon: Jacq. fl. austr. t. 26. — Exsicc.: Orph. fl. gr. n. 1165.

Pubescens vel hirsuta; caule erecto, simplici vel rarius ramoso; foliis petiolatis, ovatis oblongisve, crenatis, basi truncatis vel cordatis; verticillastris ad axillas superiores 2—6 floris, subsecundis; pedicellis calyce brevioribus; corolla pollicari et ultra, albo-rosea, purpureo-maculata, vel omnino alba.

In silvaticis regionis montanae et subalpinae. Epirus: mt. Olycika (Bald.); Thessalia: mt. Ghavellu, in oropedio Neuropolis (Haussk.) in Pindo, mt. Olympus (Auch.); Euboea (Ung.): mt. Telethrion (Heldr.); Aetolia: mt. Arapocephala (Heldr.); Achaia: mt. Chelmos pr. Planideri et Syvista (Hal.), mt. Kyllene (Orph.); in valle fl. Neda, pr. Phigalea, Androvista (Chaub.) et Anavryti (Heldr.) in mt. Taygetos, insula Poros (Wiedem.). — Jun. Aug. ♃

12. Moluccella L. gen. n. 296.

1. **M. spinosa** L. sp. p. 587; S. et S. pr. I. p. 415, Fl. gr. VI. p. 54 t. 567; Ch. et B. exp. p. 170, Fl. pelop. p. 38; Raul. cret. p. 837.

Glabra; caule erecto, procero, simplici vel inferne parce ramoso; foliis petiolatis, ovatis, grosse vel incise mucronato-serratis, floralibus verticillastra subsuperantibus; verticillastris multifloris, distantibus; bracteis tenuiter subulato-spinosis, deflexis, calycis tubo brevioribus; calycis tubo obconico, papillari-glanduloso, labio superiore triangulari, spinescente, inferiore 7—8 dentato, dentibus latis, spinosis; corolla albida, calyce vix longiore, extus villosa.

In mt. Parnasso (Sibth.); in collibus aridis pr. Navarin Messeniae (Chaub.); Creta: pr. Annato (Sieb. ex Raul.); a recentioribus ut videtur non lecta. — Jun. Jul. ⊙ N. v.

13. Phlomis L. gen. n. 723.

a. Frutices.

α. Folia ovato-oblonga vel oblonga, obtusa vel acutiuscula, ad 10 cm. longa.

1. **P. fruticosa** L. sp. p. 584; S. et S. pr. I. p. 414, Fl. gr. VI. p. 51 t. 563; Pieri corc. fl. p. 78; Sieb. in Flora I. p. 270; Urv. enum. p. 70; Ch. et B. exp. p. 38, Fl. pelop. p; 170; Marg. et R. fl. Zante p. 75; Friedr. Reise p. 273; Fraas fl. class. p. 181; Clem. sert. p. 77, Weiss in z. b. G. 1869 p. 743; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 784; Heldr. fl. cephal. p. 59. Fl. Aegina p. 385; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Epir. p. 35, Beitr. fl. Aetol. p. 2, Beitr. fl. Achaia p. 30; Gelmi in bull. soc. bot. it. 1889 p. 451; Form. in D. bot. Mon. 1891 p. 30, in Ver.

Brünn 1895 p. 36, 1896 p. 71, 1897 p. 53; Bald. viagg. Creta p. 87; Haussk. symb. p. 43; Fors. in bull. herb. Bois. III. p. 88. — Exsicc.: Rev. pl. cret. n. 143; Sint. it. thessal. n. 38; Bald. it. cret. alt. n. 348.

Ramis novellis adpresse cano-tomentosis; foliis ovato-oblongis oblongisve, basi truncatis vel cuneatis, cano-tomentosis, subtus magis canis, inferioribus breviter petiolatis, superioribus subsessilibus; verticillastris binis, remotis, vel unico terminali, multifloris, foliis floralibus binis sessilibus, eis longioribus suffultis; bracteis ovato-ellipticis vel late lanceolatis, acutis, tomentosis, calyce brevioribus; calyce tomentoso pilisque elongatis plus minus hispido, dentibus brevissimis, abrupte in mucronem subulatum, patulum vel rubrecurvum abeuntibus; corolla lutea, tomentosa, calyce duplo longiore; nuculis glabris vel apice hirtis.

In siccis apricis regionis inferioris in subalpinam usque adscendens, saepe dumeta extensa formans, per omnem Graeciam. — Febr. Jul. ♄

2. **P. viscosa** Poir. enc. V. p. 271 (1804). — *P. ferruginea* Ten. fl. nap. I. pr. p. 35 (1811), II. p. 36 t. 57; Sieb. avis. p. 4, rem. p. 5; Bois. fl. or. IV. p. 387. — *P. virens* DC. cat. monsp. p. 132 (1813). — *P. cretica* Presl del. prag. p. 84; Weiss in z. b. G. 1869 p. 743. — *P. ferruginea β. cretica* Benth. lab. p. 627; Raul. cret. p. 838. — *P. polymorpha* Clos in bull. soc. bot. Fr. XV. p. 6. — Exsicc.: Orph. fl. gr. n. 523; Rev. pl. cret. n. 143 et 202.

Ramis novellis laxe floccoso-tomentosis et superne pilis glanduliferis plus minus obsitis; foliis supra velutinis, virentibus, subtus floccoso-tomentosis, canis, inferioribus longiuscule petiolatis, oblongo-lanceolatis oblongisve, basi truncatis vel attenuatis, superioribus breviter petiolatis, cordato-ovatis; verticillastris 2—3, remotis, vel unico terminali, multifloris, foliis floralibus binis, sessilibus, eis longioribus suffultis; bracteis lanceolatis vel subulatis, acutis, villosis, calyce subaequilongis; calyce tomentoso, pilisque elongatis hispido, dentibus brevissimis, abrupte in mucronem subulatum, patulum abeuntibus; corolla lutea, tomentosa, calyce duplo longiore; nuculis glabris. — Praecedenti similis, ab ea indumento floccoso, sicco cito ferrugineo, foliis longius petiolatis, superioribus e basi profunde cordata subtriangularibus, bracteis angustis, longioribus, longe villosis, calycis dentibus longius mucronatis specifice distincta.

In regione inferiori et montana, rare. Laconia: mt. Malevo pr. Hagios Joannes (Orph.); Creta: pr. Kissamos, Canea (Rev.), Suda, Acroteri (Weiss), Malaxa, Sphakia, Retymo (Raul.). — Mart. Jun. ♄

b. Folia ovato-subrotunda vel elliptica, obtusissima, ad 25 mm. longa.

3. **P. lanata** Willd. en. hort. berol. suppl. p. 41; Raul. cret. p. 838; Bois. fl. or. IV. p. 785. — *P. parvifolia* Presl del. prag. p. 86. — *P. microphylla* Sieb. avis p. 4, in Flora 1822 I. p. 244, Reise II. p. 319 t. 8. — Exsicc.: Heldr. pl. cret. n. 1425; Bald. it. cret. alt. n. 43; Dörfl. pl. cret. n. 34.

Ramis novellis floccoso-tomentosis; foliis ovato-subrotundis vel ellipticis, basi rotundatis vel attenuatis, breviter petiolatis, supra velutinis, virentibus, subtus albo-lanatis; verticillastris binis, remotis, vel unico, terminali, 6—8 floris, foliis floralibus binis, sessilibus, eis subbrevioribus suffultis; bracteis oblongo-lanceolatis, acutis, tomentosis, calyce subduplo brevioribus; calyce tomentoso, dentibus brevissimis, truncatis, obtusis vel brevissime mucronatis; corolla lutea, tomentosa, calyce duplo longiore; nuculis apice hirtis. — Species egregia, praecedentibus omnibus partibus minor.

β. **biflora.** — Indumentum tenuius, rami graciles, parce decalvati, verticillastri saepissime 2, rarius 3—4 flori. — Exsicc.: Bald. it. cret. alt. n. 349.

In aridis, dumetis regionis inferioris Cretae: inter Hagia Rumeli et Lutro (Raul.), pr. Retymo (Tourn. voy. p. 15), Spilia distr. Candia (Bald.), Melidoni, Aludha pr. Spinalunga (Heldr.), pr. Parsas distr. Viano (Leon.); — β. in dumetis Nida sub mt. Psiloriti versus distr. Malevisi (Bald.). — Apr. Jun. ♄

b. Herbae.

α. Caulis simplex vel parce ramosus; folia ampla, ovata, basi profunde cordata; verticillastri magni, multiflori.

4. **P. samia** L. sp. p. 585; S. et S. pr. I. p. 414, Fl. gr. VI. p. 52 t. 564; Bois. fl. or. IV. p. 790; Heldr. in Sitzungsb. acad. Wiss. Berl. 1883 p. 4, chlor. Parn. p. 25; Hal. in ö. b. Z. 1890 p. 40, Beitr. fl. Thessal. p. 17, Beitr. fl. Achaia p. 30; Form. in Ver. Brünn 1896 p. 72, 1897 p. 53; Haussk. symb. p. 43. — *P. samia v. graeca et achaia* Bornm. in ö. b. Z. 1892 p. 115. — Exsicc.: Orph. fl. gr. n. 522; Sint. et Bornm. it. turc. n. 1425 et 1883; Sint. it. thessal. n. 1510; Heldr. herb. norm. n. 1573.

Caule erecto, glanduloso-pubescente, simplici vel opposite ramoso; foliis ovatis, acutiusculis, crenatis, basi profunde cordatis, supra viridibus, tomentellis, subtus cano-tomentosis, radicalibus longe, caulinis breviter petiolatis, floralibus subsessilibus, verticillastra superantibus; verticillastris 2—7, maximis, multifloris, distantibus vel superioribus approximatis; bracteis subulatis, pungentibus, calyces aequantibus, eisque patule glanduloso-viscidis; calycis dentibus lanceolatis, pungentibus, erectis, tubo subtriplo brevioribus; corolla purpurea, tomentosa, calyce duplo longiore, nuculis glabris. — Caulis crassus, 30—100 cm. altus, folia radicalia cum petiolo ad 35 cm. longa, calyx 2 cm. longus, corolla ampla.

In silvaticis regionis montanae et subalpinae. Thessalia: pr. Kastania (Hal.), Kalabaka (Sint.), mt. Oxya, pr. Vlachava, in valle Asproklisia, pr. Longi, Jerakuri, Konisko, Kucuro, mt. Mitrica et Phlambures in mt. Chassia, pr. Selicani et Spilia in mt. Ossa, pr. Tafilvris, Lokatamburia (Form.) et Hagios Dionysios (Sint.) in Olympo, mt. Pelion (Heldr.); Euboea (Sibth.); mt. Parnassus (Heldr.); Achaia: pr. Megaspilaeon, Planideri ad radices mt. Chelmos (Hal.), mt. Kyllene (Pichl.); Laconia: mt. Malevo (Orph.), Taygetos (Heldr.). — Jun. Aug. ♃

5. **P. lunarifolia** S. et S. pr. I. p. 414; Ch. et B. exp. p. 170, Fl. pelop. p. 38.

Foliis cordatis, crenatis, subtus tomentosis; bracteis ovato-lanceolatis, fasciculato-ciliatis, mucronatis. — Habitus praecedentis, sed bracteae latae, dense ciliatae ac setosae, setis fasciculatis; calycis dentes horizontaliter patentes, spinosi, rigidi; corolla flava.

In variis Peloponnesi locis (Sibth.): mt. Diaforti (Chaub.). — Species autoribus hodiernis ignota („exemplar in herb. Sibthorpiano pessimum" Benth. lab. p. 629), denuo inquirenda. — N. v.

β. Caulis patentim ramosissimus; folia oblongo-lanceolata, basi attenuata; verticillastri parvi, 2—8 flori.

6. **P. pungens** Willd. sp. III. p. 121; Heldr. chlor. Parn. p. 26; Form. in D. bot. Monat. 1891 p. 30, in Ver. Brünn 1896 p. 72, 1897 p. 53; Haussk. symb. p. 43. — *P. herba venti* S. et S. pr. I. p. 415, quoad pl. graecam; Fl. gr. VI. p. 51 t. 564; Ch. et B. exp. p. 170, Fl. pelop. p. 38; non L. — *P. herba venti β. tomentosa* et probabiliter quoque *α. typica* Bois. fl. or. IV. p. 791. — Exsicc.: Orph. fl. gr. n. 65; Heldr. herb. norm. n. 77.

Caule erecto, fere jam a basi divaricatim ramosissimo, adpresse-tomentoso, inferne simulque plus minus patule piloso; foliis breviter petiolatis, a medio minute serratis vel omnino integris, supra lucidis, viridibus, sparsim hirtis, subtus tenuiter cano-tomentellis; verticillastris 2—5, remotis, foliis floralibus binis longe superatis; bracteis subulatis, subpungentibus, calyces subsuperantibus, eisque minute tomentellis et pilis patulis, basi tuberculatis sparse hirsutis; calycis dentibus subulatis, subpungentibus, erecto-patulis, inaequalibus, calyce 2—3 plo brevioribus; corolla purpurea, tomentella, calyce duplo longiore; nuculis glabris. — Affinis *P. herba venti* L. distinguitur, caule robustiore, ramisque patule pilosis, foliis latioribus ovato-oblongis, crenatis, saepius utrinque viridibus, verticillastris plurifloris, pilosissimis, corollis majoribus.

β. **hispidula**. — Caulis ramisque, praeter indumentum tomentosum, ad apicem usque pilis patulis longis hispidula, folia subtus magis cano-tomentosa. — Exsicc.: Hal. it. gr. a. 1888.

In sterilibus siccis regionis inferioris et subalpinae. Thessalia: pr. Pharsalus (Heldr.), Orman Magula (Haussk.), Aivali, Persufli, Velestinos, Larissa, Limogardi et Metalion pr. Lamia (Form.); Boeotia (Heldr.); Attica (Sibth.); Laconia: mt. Malevo pr. Sarantapsychron (Orph.), Kalamata (Bois.); Arcadia: pr. Francovritzi, Pallantium (Chaub.); — β. pr. Livadi in mt. Parnasso (Heldr.). — Jun. Jul. ♃

14. Lamium L. gen. n. 716.

1. Sectio. *Galeobdolon* Benth. lab. p. 515. — Corollae tubus intus annulato-pilosus; antherae glabrae.

1. **L. galeobdolon** L. sp. p. 580 (*Galeopsis*); Crantz stirp. austr. ed. 2 p. 262. — *Galeobdolon luteum* Huds. fl. angl. ed 2 I. p. 258. — Icon: Fl. dan. t. 1272.

Parce pubescens; caulibus erectis vel adscendentibus, simplicibus: foliis petiolatis, ovatis vel ovato-lanceolatis, acute dentatis, floralibus verticillastra superantibus; verticillastris distantibus, plurifloris; bracteis lanceolatis, calyce brevioribus; calycis dentibus lanceolatis, subulato-spinulosis, tubo brevioribus; corolla lutea, puberula.

In nemorosis Pindi dolopici pr. Sermeniko (Haussk.). — Apr. Jun. ♃ N. v.

2. Sectio. *Lamiotypus* Dum. fl. belg. p. 45. — Corollae tubus intus annulato-pilosus; antherae hirsutae.

a. Corollae tubus calyce brevior.

2. **L. moschatum** Mill. dict. n. 4; Raul. cret. p. 837; Bois. fl. or. IV. p. 765; Haussk. symb. p. 38. — *L. calycinum* Urv. enum. p. 67; Ch. et B. exp. p. 341, Fl. pelop. p. 37. — *L. ochroleucum* Link in Linnaea 1834 p. 574. — *L. maculatum* var. *β*. Ch. et B. fl. pelop. p. 37. — *L. album* Friedr. Reise p. 266, non L. — Exsicc.: Orph. fl. gr. n. 441; Heldr. herb. norm. n. 469, herb. fl. hellen. n. 91, in Baen. herb. europ. n. 3601; Dörfl. fl. gr. n. 380.

Caulibus crassiusculis, glabris, fistulosis, saepe ramosis; foliis petiolatis, ovatis, saepe cordatis, viridibus, pilosulis, crenatis, floralibus subsessilibus, verticillastra superantibus; verticillastris distinctis, plurifloris; bracteis lanceolatis, ciliatis, tubo calycino brevioribus; calyce glabriusculo, fructifero aucto, dentibus triangulari-lanceolatis, tubo longioribus, demum stellatim patentibus; corolla alba, lobis lateralibus triangularibus, galeae basi valde attenuata oblonga, incurva, villosa. — Species tubo corolliño brevissimo insignis; corolla in toto circa 2 cm. longa.

In ruderatis, saxosis regionis inferioris. Phocis: pr. Delphi (Leonis); Attica: pr. Athenas (Sprun.), in Acropoli (Heldr.); Argolis: pr. Nauplia (Link.), Mykene (Daenzer), Epidaurus (Zuccar.); Creta: pr. Almyros (Heldr.). — Mart. Maio. ⊙

b. Corollae tubus calyce longior.

α. Perennia.

3. **L. album** L. sp. p. 579; Pieri corc. fl. p. 77; Ung. Reise p. 126. — Icon: Fl. dan. t. 594.

Plus minus pubescens, caulibus e basi prostrata, radicante adscendentibus vel erectis, simplicibus vel ramosis; foliis petiolatis, ovato-cordatis, acuminatis, inaequaliter serratis, floralibus breviter petiolatis verticillastra superantibus; verticillastris distinctis, plurifloris; bracteis parvis; calycis dentibus lanceolatis, longe subulato-acuminatis, tubo longioribus, demum patulis; corolla alba, tubo recurvo-porrecto, annulo pilorum obliquo, lobis lateralibus in 2—3 dentes, altero saepe subulato abeuntibus, galea e basi subattenuata, oblonga, subincurva, villosa. — Differt a praecedente radice perenni, caulibus saepissime patule pilosis, calyce fructifero vix aucto, dentibus angustioribus et corolla e tubo calyce longiore.

Indicatur in Corcyra (Pieri): ad sepes (Ung.). — Maio, Jul. ♃ N. v.

4. **L. maculatum** L. sp. p. 579 pro var. *L. albi*; L. sp. ed. 2 p. 809; S. et S. pr. I. p. 405, Fl. gr. VI. p. 46 t. 556; Ch. et B. fl. pelop. p. 37; Bois. fl. or. IV. p. 763. — Exsicc.: Orph. fl. gr. n. 543.

Differt a praecedente corolla purpurea, puberula vel tamen minus villosa, annulo pilorum transverso, nec obliquo munita. — Forma albiflora in Graecia hucusque ut videtur non reperta fuit.

In umbrosis regionis montanae. Argolis (Sibth.); Laconia: mt. Malevo pr. Hagios Petros, rarissime (Orph.). — Apr. Maio. ♃

β. Annuum.

5. **L. purpureum** L. sp. p. 579; Ch. et B. exp. p. 166, Fl. pelop. p. 37; Bois. fl. or. IV. p. 766; Haussk. symb. p. 39.

Plus minus pubescens; caulibus e basi saepe prostrata et radicante adscendentibus vel erectis, simplicibus vel ramosis, inferne et superne foliatis, medio nudis; foliis petiolatis, ovato-cordatis, crenatis, inferioribus obtusis, floralibus acutis, verticillastra superantibus; verticillastris in capitulum confertis, plurifloris, inferiore interdum subdistante; bracteis subulatis, brevibus; calycis dentibus lanceolato-subulatis, tubo longioribus; corolla purpurea, rarissime alba, tubo recto, lobis lateralibus subbidentatis, galea ovata, obtusa, pubescente.

In ruderatis, cultis ut videtur rare. Indicatur tantum in Thessalia: ad monasterium Meteora (Haussk.); in Peloponneso et Archipelago (Chaub.). — Mart. Nov. ⊙ N. v.

3. Sectio. *Lamiopsis* Dum. fl. belg. p. 45. — Corollae tubus annulo pilorum carens; antherae hirsutae.

a. Perennia.

6. **L. garganicum** L. sp. p. 578 (*Orvala*); L. sp. ed. 2 p. 808; S. et S. pr. I. p. 504; Ch. et B. exp. p. 165, Fl pelop. p. 36; Friedr. Reise p. 286; Raul. cret. p. 837; Haussk. symb. p. 38.

Plus minus pubescens; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis cordato-ovatis, crenatis vel dentatis, inferioribus longe petiolatis, saepe obtusis, superioribus brevius petiolatis, saepius acutis vel acuminatis, floralibus saepius brevissime petiolatis, verticillastro brevioribus vel eum superantibus; verticillastris plurifloris, inferioribus distantibus; bracteis lanceolatis, calyce brevioribus; calycis dentibus triangulari-lanceolatis, dimidio tubo sublongioribus; corolla rosea, purpureo-striata, tubo recto, lobis lateralibus obsolete denticulatis, galea elongata, integra, denticulata emarginata vel bifida, pubescente. — Stirps quoad indumentum, foliorum formam et marginatudinem, galeae marginem valde variabilis et ab auctoribus in species permultas divisa, quae tamen sine limites una in alteram transeunt, quamobrem eas ut varietates *L. garganici* sensu amplo enumerare malui.

α. **typicum.** — Molliter villosulum; galea integra vel leviter emarginata. — Exsicc.: Hal. it. gr. sec. a. 1893.

β. **nivale** Bois. et Heldr. diagn. VII. p. 54; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Achaia p. 29. — *L. striatum v. cylleneum*

Bois. et Heldr. diagn. ser. 2 IV. p. 45. — *L. cylleneum* Bois. et Heldr. fl. or. IV. p. 757. — *L. striatum v. minus* Bois. fl. or. IV. p. 757; Heldr. chlor. Parn. p. 25. — *L. pictum* Hal. in z. b. G. 1888 p. 761; Beitr. fl. Epir. p. 35; Form. in Ver. Brünn 1896 p. 71; non Bois. et Heldr. — Pubescens, caules saepe breviores, flores subminores, galea profunde emarginata. Transitum praebet ad var. *striatum*. — Exsicc.: Orph. fl. gr. n. 1166; Dörfl. fl. gr. n. 388.

γ. **longiflorum** Ten. fl. nap. I. prodr. p. 34, Fl. nap. V. p. 10 t. 152; Gelmi in bull. soc. bot. ital. 1889 p. 451; pro sp. — Glabriusculum; galea emarginata, lobis saepe bidenticulatis. — Exsicc.: Sint. it. thessal. n. 36, 304 et 682; Baen. herb. europ. n. 9277.

δ. **glabratum** Griseb. spic. II. p. 133; Haussk. symb. p. 38. — *L. bithynicum* Benth. in DC. pr. XII. p. 505. — *L. garganicum v. nivale* Bald. riv. coll. bot. alb. 1895 p. 65. — Glabriusculum; corolla integra. — Exsicc.: Heldr. it. thessal. n. 41.

ε. **striatum** S. et S. pr. I. p. 405, Fl. gr. VI. p. 46 t. 557; Ch. et B. fl. pelop. p. 37; Ung. Reise p. 126; Raul. cret. p. 837; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. p. 757; Heldr. fl. cephal. p. 59; Boissieu in bull. soc. bot. Fr. 1896 p. 287; Haussk. symb. p. 39; pro sp.; Bald. riv. coll. bot. alb. 1895 p. 65. — *Dracocephalum lamiifolium* Desf. choix p. 33 t. 15. — *L. grandiflorum* Sieb. avis rem. p. 5, non Pourr. act. tolos. III. p. 322, quod ad *L. maculatum* pertinet. — *L. rugosum* Ch. et B. exp. p. 165, non Ait. hort. Kew. II. p. 296, quod etiam ad *L. maculatum* pertinet. — *L. Columnae* Friedr. Reise p. 280, non Ten. syll. p. 285, quod sec. Benth. lab. p. 514 ad *L. tomentosum* Willd., sec. Bert. fl. ital. VI. p. 113 ad *L. maculatum*, sec. Car. in Parl. fl. ital. VI. p. 208 ad *L. longiflorum* spectat. — Galea bifida. — Variat *f. hymettium* Heldr. herb. norm. n. 468 (f. robustior, pubescentia majore et colore florum saturatiori) et *f. glabrescens* Heldr. herb. norm. n. 1271 (glabriuscula, calyce breviore, fauce angustiore). — Exsicc.: Heldr. l. c.; Orph. fl. gr. n. 97; Dörfl. fl. gr. n. 363.

In rupestribus regionis montanae et alpinae omnium montium; non indicatur in insulis Archipelagi. — Mart., in alpinis ad Aug. ♃

7. **L. pictum** Bois. et Heldr. diagn. ser. 2 IV. p. 44; Heldr. chlor. Parn. p. 25. — *L. striatum v. pictum* Bois. fl. or. IV. p. 758. — Exsicc.: Heldr. herb. norm. n. 727 et 1467; Orph. fl. gr. n. 440.

Glabriusculum vel parce puberulum; caulibus filiformibus, diffusis, simplicibus; foliis parvis, reniformibus vel ovato-orbiculatis, obtusis, crenatis, inferioribus longe, superioribus brevius petiolatis, floralibus calycem aequantibus vel superantibus; verticillastris 2—4 floris, paucis vel unico terminali; bracteis minutis; calycis dentibus triangularibus, tubo dimidio brevioribus; corolla rosea, purpureo-striata, hirtula, tubo recto, lobis lateralibus angulatis, galea elongata, biloba. — Caespites densiusculos formans, caulibus brevibus filiformibus, foliorum forma et

verticillis paucifloris a praecedente distincta; formas intermedias hucusque non vidi.

In rupestribus regionis alpinae mt. Korax, Parnassus et Taygetos (Heldr.). — Jul. Aug. ♃

b. Annua.

α. Corollae galea bifida.

8. **L. bifidum** Cyr. pl. rar. neap. I. p. 22; Ch. et B. exp. p. 165, Fl. pelop. p. 37; Marg. et R. fl. Zante p. 75; Friedr. Reise p. 280; Raul. cret. p. 837; Bois. fl. or. IV. p. 760; Haussk. symb. p. 39. — Icon: Ten. fl. nap. t. 153. — Exsicc.: Heldr. fl. thessal. a. 1883.

Parce pubescens; caulibus adscendentibus, simplicibus vel basi ramosis, medio saepe nudis; foliis inferioribus longe petiolatis, cordato-ovatis, crenatis vel subincisis, floralibus breviter petiolatis, ovato-triangularibus, inciso-dentatis; verticillastris plurifloris, inferioribus distantibus; bracteis linearibus, calyce brevioribus; calycis hirti dentibus lanceolato-subulatis, tubo brevioribus, demum patentibus; corolla alba, pubescente, tubo recto, calyce 3—4 plo longiore, galea bifida, lobis obovatis; acheniis granulatis.

In arvis, ruderatis regionis inferioris. Thessalia: pr. Chaliki in Pindo tymphaeo (Haussk.), pr. Mares in mt. Pelion (Heldr.); Euboea (Wiedem.); Attica (Bois.); Arcadia: pr. Carithena (Friedr.); Messenia: pr. Pylos, Methone (Chaub.), Kalamata (Zahn); Creta: ad ripas fl. Kladiso pr. Garipa (Heldr.); Zante: pr. Litakia (Marg.); Corcyra: pr. urbem (Baen.). — Mart. Maio. ⊙

β. Corollae galea integra.

9. **L. hybridum** Vill. hist. pl. Dauph. I. p. 291. — *L. incisum* Willd. sp. III. p. 89; Friedr. Reise p. 279; Boissieu in bull. soc. bot. Fr. 1896 p. 287. — Icon: Rchb. t. 3. — Exsicc.: Sint. it. thessal. n. 1516.

Parce pubescens; caulibus adscendentibus, simplicibus vel basi ramosis, medio saepe nudis; foliis inferioribus longe petiolatis, subcordato-ovatis, crenatis vel subincisis, floralibus breviter petiolatis, ovato-triangularibus, inciso-dentatis; verticillastris plurifloris, inferioribus distantibus; bracteis linearibus, calyce brevioribus; calycis hirti dentibus lanceolato-subulatis, tubo brevioribus, demum patentibus; corolla purpurascenti, pubescente, tubo recto, calyce breviore, galea integra, acheniis laevibus. — Habitu praecedentis, a qua corolla purpurea, tubo calyce breviore et acheniis laevibus distinguitur.

In herbidis, arvis, ruderatis regionis inferioris. Thessalia: in valle Negerli pr. Chaliki in Pindo tymphaeo (Sint.), pr. Kalabaka (Boissieu); Arcadia: mt. Trikorfo pr. Sinano (Friedr.). — Apr. Jun. ⊙

10. **L. amplexicaule** L. sp. p. 579; S. et S. pr. I. p. 406; Pieri corc. fl. p. 77; Ch. et B. exp. p. 166, Fl. pelop. p. 37; Marg. et R. fl. Zante p. 75; Friedr. Reise p. 266; Clem. sert. p. 76; Ung. Reise

p. 126: Raul. cret. p. 837; Weiss in z. b. G. 1869 p. 743; Bois. fl. or. IV. p. 760: Heldr. fl. cephal. p. 59. Fl. Aegina p. 385. chlor. Thera p. 19. chlor. Mykon p. 249 *f. microphylla*: Haussk. symb. p. 39. — Icon: Fl. dan. t. 752. — Exsicc.: Sint. it. thessal. n. 37.

Parce pubescens: caulibus adscendentibus, simplicibus vel basi ramosis, medio saepe nudis: foliis inferioribus petiolatis, subcordato-ovatis, crenatis, floralibus sessilibus, reniformibus, amplexicaulibus: verticillastris plurifloris, distantibus: bracteis nullis: calycis villosi dentibus lanceolato-subulatis, tubo subbrevioribus, conniventibus: corolla purpurea, rarissime alba, pubescente, tubo recto, calyce triplo longiore, galea integra.

β. **clandestinum** Rchb. ic. pl. crit. t. 708. — *L. mesogaeum* Heldr. fl. hellen. a. 1875. — Corolla abbreviata, calyce subinclusa, abortiente.

In cultis, arvis, vinetis regionis inferioris et montanae per totam Graeciam. — Febr. Oct. ⊖

15. Betonica L. gen. n. 718.

a. Corollae tubus intus nudus.

α. Corolla purpurea vel alba: antherae flavidae.

1. **B. officinalis** L. sp. p. 573: S. et S. pr. I. p. 407; Dallap. prosp. p. 83; Ch. et B. fl. pelop. p. 38; Bois. fl. or. IV. p. 752. — *Stachys betonica* Benth. lab. p. 532. — Icon: Fl. dan. t. 726.

Pubescens vel glabrata; caulibus erectis vel adscendentibus, simplicibus: foliis inferioribus a basi cordata ovatis vel ovato-oblongis, crenatis, inferioribus longe petiolatis, caulinis remotis, sessilibus, floralibus inferioribus lanceolatis, serratis, summis linearibus, integris: spica oblongo-cylindrica, basi interrupta: bracteis ovato-lanceolatis, mucronatis: calycis dentibus lanceolatis, spinulosis, tubo vix brevioribus; corollae pubescentis, purpureae rarius albae tubo calyce longiore.

In dumosis regionis montanae. Laconia (Sibth.); Cephalonia Dallap.). — Jun. Aug. ♃ N. v.

2. **B. Hausskneechtii** Uechtr. in Haussk. symb. p. 39. — *B. officinalis* Form. D. bot. Mon. 1891 p. 30, quoad pl. thessal., non L. quae in Pindo sec. Haussk. l. c. non obvia. — Exsicc.: Sint. it. thessal. n. 1112.

Praecedenti valde affinis, ab ea differt statura robustiore, bracteis ad margines longius densiusque ciliatis, calycis tubo fauce excepto glabro, sinubus acutis, nec rotundatis, corolla submajore, constanter niveo, tubo crassiore, galea longiore, integerrima, nec crenulata, labio inferiore latiore, subtruncato.

In quercinis oropedii Neuropolis in Pindo dolopico (Haussk.) et in mt. Zygos Pindi tymphaei (Form.). — Jun. Aug. ♃

β. Corolla ochroleuca, roseo suffusa; antherae purpureae.

3. **B. scardica** Griseb. Reise Rumel. II. p. 189 (*Stachys*); Griseb. spic. II. p. 138; Hal. Beitr. fl. Thessal. p. 17; Bald. riv. coll. bot. alb. 1896 p. 87; Haussk. symb. p. 39. — *B. graeca* Bois. et Sprun. diagn. V. p. 27, Fl. or. IV. p. 751; Form. in Ver. Brünn 1896 p. 70 *v. tymphaea*, 1897 p. 52 *v. petiolata*; ambae sic dictae varietates typo nullo modo discrepant. — Exsicc.: Heldr. herb. norm. n. 870; Sint. it. thessal. n. 565 b; Bald. it. alb. n. 88.

Hirsuta; caulibus erectis vel adscendentibus, simplicibus; foliis e basi subcordata oblongo-lanceolatis, crenato-dentatis, subtus canescentibus, inferioribus breviter petiolatis, superioribus sessilibus, floralibus inferioribus saepe deflexis; spica ob verticillastra inferiora distincta laxe cylindrica; bracteis oblongo-lanceolatis, acuminatis; calycis dentibus lanceolato-subulatis, spinulosis tubo duplo brevioribus; corollae pubescentis tubo calyce vix longiore.

In quericetis, dumosis regionis montanae et subalpinae. Epirus: mt. Cika in mt. Acroceraunicis, mt. Smolika (Bald.), mt. Peristeri (Form.); Thessalia: mt. Zygos, pr. Malakasi (Haussk.), mt. Oxya (Hal.), ad monasterium Korona, in oropedio Neuropolis (Heldr.), mt. Beluja, Itamos in mt. Agrapha, mt. Said Pascha, Dokimi, pr. Vlachava, Hagios Paraskevi, mt. Kerasia Sina, Cuka et mt. Mitrica in mt. Chassia (Form.); Eurytania: mt. Tymphrestus (Sprun.). — Maio, Aug. ♃

b. Corollae tubus intus annulatus.

4. **B. Jacquini** Gr. et Godr. fl. fr. II. p. 694; Bois. fl. or. IV. p. 749; Heldr. chlor. Parn. p. 25; Hal. in ö. b. Z. 1896 p. 16. — *B. alopecurus* L. sp. p. 573 p. p.; Jacq. fl. austr. I. p. 50 t. 78; S. et S. pr. I. p. 407. — Exsicc.: Orph. fl. gr. n. 311; Dörfl. fl. gr. n. 183.

Hirsuta; caulibus erectis vel adscendentibus, simplicibus; foliis cordato-ovatis, crenatis vel dentatis, inferioribus longe petiolatis, superioribus remotis, subsessilibus, floralibus parvis, ovatis ovato-lanceolatisve; spica ob verticillastra inferiora distincta laxe cylindrica; bracteis lanceolatis, mucronatis; calycis dentibus lanceolato-subulatis, spinulosis, tubo duplo brevioribus; corollae flavae, pubescentis, tubo calyce vix longiore. — Planta alpium Europae occidentalis — *B. alopecurus* L. pro maxima parte — differt parum a nostra spica ovata, densa, crassiori et foliis floralibus infimis amplis, late ovatis, verticillastro multo longioribus.

In rupestribus regionis abietinae et alpinae. Thessalia: mt. Olympus mt. Oeta (Heldr.); mt. Kiona (Reiser), Parnassus (Sibth.); Achaia: mt. Chelmos et Kyllene (Orph.). — Jun. Aug. ♃

16. Stachys L. gen. n. 719.

Dispositio specierum:

1. Sectio. *Eriostachys* Bois. fl. or. IV. p. 715. — Perennes; verticillastra multiflora, bracteis dimidium calycem aequantibus vel superantibus; calycis dentes rigidi vel pungentes; corollae tubus inclusus.

a. Corollae labium superius integrum vel retusum.

α. Corolla ochroleuca.

1. S. orientalis L.

β. Corolla purpurascens, rarissime alba.

× Folia saltem radicalia oblonga vel lineari-oblonga, basi attenuata.

2. S. cretica L. **3. S. italica** Mill. **4. S. cassia** Bois.

×× Folia radicalia ovata vel ovato-oblonga, basi cordata, vel late rotundata.

○ Plantae albo-pannosae, sericeae; calyx ore rectus.

5. S. Tournefortii Poir. **6. S. sericophylla** Hal.

○○ Plantae pubescentes vel plus minus tomentosae; calyx ore subobliquus.

. Calyx eglandulosus.

7. S. germanica L.

.. Calyx praeter indumentum villosum ad dentes pilis glanduliferis parce obsitum.

8. S. Reinerti Heldr.

b. Corollae labium superius profunde bifidum.

9. S. acutifolia Ch. et B.

2. Sectio. *Stachyotypus* Bois. fl. or. IV. p. 715. — Perennes; verticillastra saepius pauciflora, bracteis (exc. *S. decumbens*) brevissimis vel nullis.

a. Rhizoma repens; corolla purpurea, rarissime alba.

10. S. silvatica L. **11. S. palustris** L.

b. Rhizoma non repens.

α. Herbaceae.

× Rhizoma caules floriferos tantum edens.

○ Verticillastra 4—10 flora.

. Caulis praeter pilos patentes, pilis brevissimis glandulosis obsitus

12. S. viridis Bois. et Heldr. **13. S. Freynii** Haussk.

.. Caulis eglandulosus.

; Caules adscendentes, patule hirsuti; folia molliter tomentella, plus minus canescentia.

14. S. messeniaca Bois. **15. S. canescens** Ch. et B.

;; Caules erecti, rigiduli breviter hirti; folia parce hirta, viridia.

, Corolla ochrolenca; antherae labium superius subaequantes.

16. S. Parolinii Vis. **17. S. recta** L.

,, Corolla purpurea; antherae inclusae.

18. S. iberica M. a. B.

○○ Verticillastra 2 flora.

19. S. virgata Ch. et B. **20. S. tenuifolia** Pall.

×× Rhizoma caules floriferos et rosulas steriles edens.

21. S. maritima L. **22. S. pubescens** Ten.

β. Basi fruticulosae.

× Bracteae calyce subaequilongae.

23. S. decumbens Pers.

×× Bracteae minutae vel nullae.

○ Rami inermes.

. Folia elliptica vel oblonga, nunquam dense pannosa.

, Flores subsessiles vel breviter pedicellati.

, Caules patule hirsuti; verticillastra saltem superiora in spicam ovatam vel oblongam congesta.

24. S. argolica Bois. **25. S. Swainsoni** Benth. **26. S. Spruneri** Bois.

,, Caules virgati, saltem in parte superiori glabri; verticillastra remotissima.

27. S. tetragona Bois. et Heldr.

;; Flores crassiuscule pedicellati; pedicelli tubo calycis subaequilongi.

28. S. mucronata Sieb.

.. Folia ovato-subrotunda, dense pannosa.

; Calyx eglandulosus.

29. S. candida Ch. et B. **30. S. chrysantha** Bois. et Heldr.

; Calyx glandulosus.

31. S. Spreitzenhoferi Heldr.

○○ Rami spinescentes.

32. S. spinosa L.

3. Sectio. *Olisia* Bois. fl. or. IV. p. 716. — Species annuae.

a. Corolla albido-ochroleuca.

33. S. annua L. **34. S. spinulosa** S. et S.

b. Corolla purpurea.

35. S. serbica Panc. **36. S. arvensis** L.

1. Sectio. *Eriostachys* Bois. fl. or. IV. p. 715. — Perennes; verticillastra multiflora, bracteis dimidium calycem aequantibus vel superantibus; calycis dentes rigidi vel pungentes; corollae tubus inclusus.

a. Corollae labium superius integrum vel retusum.

α. Corolla ochroleuca.

1. **S. orientalis** L. sp. p. 582?; Vahl. symb. II. p. 64; S. et S. pr. I. p. 409; Fl. gr. VI. p. 48 t. 560; Bald. in bull. herb. Bois. 1896 p. 205.

Molliter villosa, virens; caulibus erectis, simplicibus vel subsimplicibus; foliis oblongo-lanceolatis, crenulatis, basi subattenuatis vel rotundatis, rarius subcordatis, inferioribus longe, mediis breviter petiolatis, floralibus sessilibus, ovato-lanceolatis; verticillastris 6—10 floris, inferioribus remotis; bracteis elliptico-rhombeis, acutis, calyces subaequantibus vel superantibus; calycis villosi, recti, dentibus lanceolato-subulatis, dimidium tubum aequantibus; corollae villosae labio superiore integro vel retuso. — Caulis elatus, foliorum radicalium lamina ad 13 cm. longa, petiolo aequilongo suffulta, indumentum e pilis elongatis sed non densis constans, ideo folia viridia; hac nota et corolla ochroleuca ab affinibus distinctissima.

In agro Messeniaco (Sibth.); in saxosis vinearum pr. Eukluvi insulae Leucas (Bald.). — Jun. Aug. ♃ N. v.

β. Corolla purpurascens, rarissime alba.

× Folia saltem radicalia oblonga vel lineari-oblonga, basi attenuata.

2. **S. cretica** L. sp. p. 581; S. et S. pr. I. p. 408, Fl. gr. VI. p. 47 t. 558; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 836; Bois. fl. or. IV. p. 719 p. p.; Hal. in ö. b. Z. 1897 p. 98; Haussk. symb. p. 10, cum *v. condensata*, f. caulibus pumilis, verticillastris approximatis. — *S. salviaefolia* Friedr. Reise p. 284, non Ten. — *S. germanica v. Boissieri* Briq. lab. alp. marit. p. 222. — Exsicc.: Heldr. herb. norm. n. 288, 1168 *f. canescens*, 1169 *f. virens*; Orph. fl. gr. n. 77; Rev. pl. cret. n. 138, in Baen. herb. europ. n. 4892; Dörfl. fl. aeg. n. 28, pl. cret. n. 127.

Adpresse tomentosa, canescens; caulibus erectis vel adscendentibus, simplicibus vel subsimplicibus; foliis oblongis vel oblongo-linearibus, crenulatis, basi attenuatis, inferioribus petiolatis, mediis subsessilibus, floralibus sessilibus, ovato-oblongis; verticillastris multifloris, remotis; bracteis lanceolatis, calyces subaequantibus; calycis lanati, ore subobliqui dentibus triangulari-lanceolatis, inter lanam suboccultatis, spinula lutescenti vel purpurascenti terminatis, fructiferis erectis, tubo triplo brevioribus; corollae roseae labio superiore lanato, integro vel retuso.

In lapidosis regionis inferioris et montanae, ut videtur in parte magis orientali ditionis. Boeotia (Sprun.); Attica: pr. Athenas, in colle Lycabettus, mt. Pentelicon, Hymettus (Heldr.), pr. Kamariza, Laurion, Sunium (Haussk.); Argolis: pr. Nauplia (Sart.), mt. Palamidi pr. Nauplia, Tolon (Haussk.); Sporadum insula Scopelos (Leon.); Euboea: mt. Telethrion, Dirphys (Heldr.), in Petalium insula Tragonisi (Holzm.); Cyclades (Sibth.): insula Naxos (Leon.); Creta: pr. Kissamos (Reverch.), Canea (Weiss), Acroteri, promontorium Meleka, pr. Malaxa, Hagios Deka, Omalos (Raul.). — Maio, Jul. ♃

3. **S. italica** Mill. dict. n. 3; Marg. et R. fl. Zante p. 75; Raul. cret. p. 835. — *S. cretica* Pieri corc. fl. p. 78; ? Ch. et B. exp. p. 168, Fl. pelop. p. 37; Bois. fl. or. IV. p. 719 p. p.; Heldr. fl. cephal. p. 59; Hal. Beitr. fl. Achaia p. 30; non L. — *S. salviaefolia* Ten. fl. nap. II. p. 23 t. 53; Link. in Linnaea IX. p. 576. — *S. cretica* Hal. Beitr. fl. Achaia p. 30. — *S. Heldreichii* Form. in Ver. Brünn 1896 p. 70, non Bois. — Exsicc.: Dörfl. fl. gr. n. 279.

Adpresse tomentosa, canescens vel pallide virens; caulibus erectis vel adscendentibus, simplicibus; foliis oblongis vel oblongo-linearibus, crenulatis, basi attenuatis, inferioribus petiolaiis, mediis subsessilibus, floralibus sessilibus, ovato-oblongis; verticillastris multifloris, superioribus saepius contiguis; bracteis lanceolatis, calyce brevioribus; calycis lanati ore subobliqui, dentibus triangulari-lanceolatis, spinula saepius purpurascenti terminatis, fructiferis erecto-patulis, tubo subtriplo brevioribus, inter lanam non occultatis; corollae roseae labio superiore lanato, integro vel retuso. — Praecedenti maxime affinis et ab ea verticillastris minus remotis, superioribus saepissime approximatis, calycis dentibus submajoribus, brevius spinulosis, demum erecto-patulis, inter lanum non occultatis vix specifice diversa et ut videtur intermediis cum ea conjuncta. A *S. germanica* foliis radicalibus basi non cordatis differt.

β. **albiflora** Hal. Beitr. fl. Achaia p. 30 pro var. *S. creticae.* — Corolla alba. — Exsicc.: Hal. it. gr. sec. a. 1893.

In lapidosis regionis inferioris et montanae, ut videtur in parte magis occidentali ditionis. Thessalia: pr. Pharsalus (Heldr.), Velestinos (Form.); Aetolia: mt. Korax pr. Musinitza (Leon.); Doris: pr. Amphissa (Hal.); Elis: pr. Manolas, Chlemutzi (Heldr.); Achaia: pr. Patras (Link.), Megaspilaeon cum var. β. (Hal.); Messenia: pr. Messene, mt. Diaforti (Chaub.); Laconia: pr. Sparta, mt. Taygetos (Chaub.); Creta: pr. Guves, Avdu, inter Pyrgos et Karakas (Heldr.); Zante (Marg.); Cephalonia: pr. Argostoli, Pessada, Erisso, Lixuri (Heldr.); Corcyra (Pieri). — Maio, Jul. ♃

4. **S. cassia** Bois. diagn. XII. p. 76, Fl. or. IV. p. 721. — *S. italica v. chondrostachys* Heldr. in Bois. diagn. ser. 2 IV. p. 37. — Exsicc.: Orph. fl. gr. n. 535 (mt. Korfiati); Heldr. it. gr. septentr. a. 1879, it. thessal. IV. a. 1885.

Differt a praecedente indumento saepius parciore et praesertim calycis dentibus longe subulato-spinosis, tubo subbreviorrbus. — Specimina graeca a macedonicis indumento densiore et foliis caulinis vix subcordatis parum discedunt.

In lapidosis, collibus siccis pr. Tyrnovo Thessaliae et in mt. Korax Aetoliae (Heldr.). — Jun. Jul.

×× Folia radicalia ovata vel ovato-oblonga, basi cordata vel late rotundata.

⊙ Plantae albo-pannosae, sericeae; calyx ore rectus.

5. **S. Tournefortii** Poir. dict. V. p. 227; Raul. cret. p. 837; Bois. fl. or. IV. p. 718. — *S. cretica* Sieb. avis p. 4, rem. p. 5, non L. — *S. Sieberi* C. Koch in Linnaea XXI. p. 685. — Icon: Tourn. voy. I. p. 29. — Exsicc.: Rev. pl. cret. n. 140.

Dense pannoso-lanata; caulibus elatis, simplicibus vel parce ramosis; foliis inferioribus petiolatis, ovatis, obtusis, crenulatis, basi manifeste cordatis, floralibus sessilibus, oblongis vel oblongo-lanceolatis, superioribus verticillastro brevioribus; verticillastris multifloris, distantibus; bracteis lanceolatis, calyce brevioribus; calycis albo-lanati dentibus rectis, lanceolatis, spinula purpurascenti terminatis, tubo triplo brevioribus; corollae purpureae labio superiore lanato, integro. — Indumento albopannoso sericeo et calyce ore recto, nec obliquo, ideo dentibus fere aequilongis a *S. germanica* et affinibus distincta.

In locis aridis regionis inferioris et montanae. Creta: pr. Kissamos (Rev.), Canea (Tourn.), Malaxa, Omalo (Bois.), pr. Gortyna distr. Messara (Heldr.). — Maio, Jul. ♃

6. **S. sericophylla.** — *S. germanica v. penicillata* Bald. in bull. herb. Bois. IV. p. 646, non (Heldr. et Sart.) Bois. — Exsicc.: Bald. it. alb. alt. 1894 n. 102.

Molliter sericeo-lanata, nitidula; caulibus humilibus, adscendentibus; foliis inferioribus breviter petiolatis, ovatis, acutis, crenulatis, basi rotun-

datis vel subcordatis, floralibus sessilibus, ovato-lanceolatis, verticillastro longioribus, verticillastris plurifloris, distantibus; bracteis lineari-subulatis, calyce brevioribus; calycis albo-lanati dentibus rectis, lanceolato-subulatis, spinula elongata purpurascenti terminatis, tubo subtriplo brevioribus; corollae purpureae labio superiore lanato, integro. — Stirps elegans, caulibus humilibus, 15—30 cm. altis, indumento sericeo-nitente, foliis inferioribus brevibus, cum petiolo circa 4 cm. longis, basi lata rotundatis vel subcordatis, floralibus verticillastra superantibus, verticillastris paucis (2—4), bracteis subulatis, calycis dentibus longioribus a praecedente specifice distincta.

In saxosis alpinis mt. Nimercka l. d. Pljepista distr. Premeti (Bald.) Albaniae meridionalis, jam extra ditionem nostram; in montibus Epiri inquirenda. — Jul. Aug.

○○ Plantae pubescentes vel plus minus tomentosae; calyx ore subobliquus.

. Calyx eglandulosus.

7. **S. germanica** L. sp. p. 581; S. et S. pr. I. p. 408; Ch. et B. fl. pelop. p. 37; Fraas fl. class. p. 180; Bois. fl. or. IV. p. 720; Form. in D. bot. Mon. 1891 p. 30. — Icon: Jacq. fl. austr. t. 319. — Exsicc.: Heldr. herb. norm. n. 1468.

Plus minus tomentosa, cana vel canescens; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis inferioribus petiolatis, ovatis, crenulatis, basi cordatis, floralibus sessilibus, oblongis vel lanceolatis, inferioribus verticillastrum superantibus; verticillastris multifloris, superioribus approximatis; bracteis lanceolatis, calyce brevioribus; calycis villosi, ore subobliqui dentibus triangularibus, acutis, spinula brevi terminatis, erectis, tubo triplo brevioribus; corollae roseae labio superiore lanato, integro vel retuso.

β. **penicillata** Heldr. et Sart. in Bois. diagn. ser. 2 IV. p. 37; Heldr. in Sitzungsb. acad. Wiss. Berlin 1883 p. 4, chlor. Parn. p. 25; Hal. Beitr. fl. Epir. p. 35, Beitr. fl. thessal. p. 17; Haussk. symb. p. 40; pro sp.; Bois. fl. or. p. 720. — *S. lanata* S. et S. pr. I. p. 408?; Ch. et B. exp. p. 169, Fl. pelop. p. 37; non Jacq. ic. rar. I. p. 11, quae indumento densissime pannoso sericeo et foliis radicalibus in petiolum attenuatis differt. — *S. dasyanthes Raf. v. alpina* Heldr. herb. norm. n. 91. — Corollae labium superius longe albo-penicillatum. — Intermediis ad typum transit. — Exsicc.: Heldr. herb. norm. n. 91, it. thessal. n. 39; Orph. fl. gr. n. 439; Dörfl. fl. gr. n. 411.

γ. **Heldreichii** Bois. fl. or. IV. p. 719; Heldr. chlor. Parn. p. 25; Hal. Beitr. fl. Epir. p. 35; Haussk. symb. p. 40; pro sp. — *S. germanica v. stricta* Form. in Ver. Brünn 1896 p. 70, 1897 p. 52. — Elatior, ramosa vel paniculata, foliis majoribus, profundius crenatis, floralibus saepius dentatis, verticillastra numerosa, spicam elongatam formantia. — Exsicc.: Orph. fl. gr. n. 1045; Sint. et Bornm. it. turc. n. 1428; Sint. it. thessal. n. 1111.

In collibus siccis, silvaticis regionis inferioris in subalpinam adscendens. Epirus: mt. Tsumerka (Hal.); Thessalia: mt. Zygos, pr. Malakasi, Chaliki, mt. Baba, pr. Klinovo, Kalabaka, Korona, mt. Ghavellu (Haussk.), mt. Chassia, pr. Karditza, Velestinos, Hadzi Obasi, Larissa, Selicany, Ambelakia, Volo (Form.), mt. Pelion, Ossa, Olympus, Oeta (Heldr.); Aetolia: mt. Tymphrestus (Fraas); mt. Kiona (Hal.), Parnassus (Heldr.); Euboea: mt. Dirphys (Bois.); Achaia (Sibth.): mt. Olenos, Chelmos, Kyllene (Heldr.); Laconia: mt. Taygetos (Heldr.); Corcyra (Form.); — γ. rarior: Epirus: pr. Kalentini (Hal.); Thessalia: pr. Sermenikon (Sint.), mt. Ghavellu, Chassia, Velestinos, Hagios Georgios, Trikala, Sophades, Karditza, Pharsalus (Haussk.), pr. Godaman (Form.) et Catherine (Heldr.) in Olympo; Euboea: pr. Oreos (Orph.); Boeotia: pr. Lebadea, mt. Parnassus (Heldr.). — Loca nonnulla forsan ad *S. italicam* spectant, quacum defectu foliorum radicalium facillime confunditur. — Maio, Aug. ♃

.. Calyx praeter indumentum villosum ad dentes pilis glanduliferis parce obsitum.

8. **S. Reinerti** Heldr. herb. norm. n. 743 (1857); Murb. Beitr. fl. Südbosn. p. 62; Hal. Beitr. fl. Epir. p. 35. — *S. alpina v. discolor* Bois. fl. or. IV. p. 719. — *S. tymphaea* Haussk. in Mitth. thür. bot. Ver. V. II. p. 70, symb. p. 40 cum. f. f. *lacmonica et petiolata.* — *S. alpina v. Reinerti* Bald. riv. coll. bot. alb. 1895 p. 64. — *S. alpina* Form. in D. bot. Monat. 1891 p. 30; forsan etiam S. et S. pr. I. p. 408. — *S. alpina v. turcica* Form. in Ver. Brünn 1896 p. 69, 1897 p. 52. — Exsicc.: Heldr. l. c., Orph. fl. gr. n. 1044 mt. Peristeri Macedoniae; Sint. it. thessal. n. 950.

Molliter tomentella, virens; caulibus adscendentibus vel erectis, simplicibus vel subsimplicibus; foliis ovatis vel ovato-oblongis, acutis vel obtusis, crenulatis, inferioribus petiolatis, basi cordatis, floralibus ovato-lanceolatis, sessilibus; verticillastris multifloris, distinctis, inferioribus remotis; bracteis lanceolatis, calyces subaequantibus; calycis villosi, ore subobliqui dentibus triangulari-lanceolatis, spinula terminatis, parce glanduloso-pilosis, tubo triplo brevioribus; corollae roseae labio superiore lanato, integro. — Habitu *S. alpinae* L., indumento tenuiore, inflorescentia glandulosa, calycis dentibus latioribus diverso, accedens.

β. **abbreviata.** — *S. penicillata* Hal. in z. b. G. 1888 p. 761, non Heldr. et Sart. — Caulibus nanis, 8—15 cm. altis; foliis parvis, brevissime petiolatis, cum petiolo 10—20 mm. longis, ovato-rotundatis vel ovatis, plerisque obtusissimis; verticillastris 2—4; corollae labio superiore longe albo-penicillato. — Exsicc.: Hal. it. gr. a. 1888; Reis. fl. gr. a. 1894.

In herbidis saxosis regionis subalpinae et alpinae. Epirus: mt. Tsumerka (Bald.), Peristeri (Hal.); Thessalia: mt. Zygos, pr. Malakasi (Haussk.), mt. Ghavellu (Heldr.), Ochsa Despot, Dokimi, Gisel Tepe, Oxya, Karava, pr. Phlambures, mt. Cuka et Mitrica in mt. Chassia, pr. Suilismen, Godaman et Rapsani in Olympo, mt. Pelion,

mt. Alafovris, Mavrika et Katasara in mt. Othrys (Form.), mt. Oeta (Heldr.); Eurytania: mt. Tymphrestus (Samar.); Laconia (Sibth.), si *S. alpina* S. et S. huc pertineat. — *β.* in cacumine mt. Kiona (Hal.). — Jun. Aug. ♃

b. Corollae labium superius profunde bifidum.

9. **S. acutifolia** Ch. et B. exp. p. 168; Link in Linnaea IX. p. 575. — *S. lusitanica* Ch. et B. fl. pelop. p. 37; non Brot. — *S. graeca* Bois. et Heldr. diagn. XII. p. 77, Fl. or. IV. p. 723; Hal. Beitr. fl. Achaia p. 30; Haussk. symb. p. 40. — Exsicc.: Heldr. fl. gr. n. 688; Orph. fl. gr. n. 1043; Sint. it. thessal. n. 466.

Caulibus erectis vel adscendentibus, simplicibus vel ramosis, patule hispidis; foliis ovatis vel ovato-oblongis, obtusis vel acutis, basi rotundatis vel subcordatis, crenatis, adpresse hirsutis, virentibus, inferioribus petiolatis, floralibus sessilibus; verticillastris multifloris, distantibus; bracteis lanceolato-subulatis, calyce subbrevioribus; calycis hirsuti, recti dentibus triangulari-lanceolatis, spinula terminatis, saepius parce glanduloso-pilosis, tubo triplo brevioribus; corollae roseae limbo amplo, labio superiore lanato, bifido. — Ab omnibus antecedentibus corolla majore, ad 15 mm. longa, labio inferiore amplo, ad 8 mm. lato, superiore profunde bifido distincta.

In rupestribus, herbidis, olivetis regionis inferioris et subalpinae. Thessalia: pr. Vutades in Pindo tymphaeo (Sint.), pr. Aivali, Orman Magula (Haussk.); Aetolia: pr. Angelocastron (Heldr.); Boeotia: pr. Lebadea (Orph.); Attica: mt. Cithaeron, Pateras, Parnes (Heldr.); Euboea: mt. Dirphys (Sprun.); Elis: mt. Olenos (Hal.); Achaia: pr. Patras (Link.); Argolis: pr. Nauplia (Berger), Tolon, in peninsula Vromolimni (Haussk.), insula Poros (Wiedem.); Laconia: mt. Malevo (Sprun.); Messenia (Chaub.).

2. Sectio. *Stachyotypus* Bois. fl. or. IV. p. 715. — Perennes; verticillastra saepius pauciflora, bracteis (excl. *S. decumbens*) brevissimis vel nullis.

a. Rhizoma repens; corolla purpurea, rarissime alba.

10. **S. silvatica** L. sp. p. 580; S. et S. pr. I. p. 407; Ch. et B. fl. pelop. p. 37; Bois. fl. or. IV. p. 726; Haussk. symb. p. 41. — Icon: Fl. dan. t. 1102.

Herbacea, hirsuta, viridis; caulibus erectis, simplicibus vel ramosis, superne glanduloso-pilosis; foliis petiolatis, cordato-ovatis, acuminatis, crenatis vel serratis, floralibus lanceolatis, subsessilibus; verticillastris distinctis, 2—6 floris; calycis glandulosi dentibus triangulari-lanceolatis, tubo subbrevioribus; corollae glanduloso-puberulae tubo exserto.

β. **pallida** Haussk. symb. p. 41. — Corolla pallide-rosea vel alba. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

In umbrosis, silvaticis regionis montanae, rarissime. Thessalia: pr. Pezula (Heldr.) et Sermeniko in Pindo dolopico, cum. *β.* (Haussk.); in agro Laconico (Sibth.). — Jun. Jul. ♃

11. **S. palustris** L. sp. p. 580; Bois. fl. or. IV. p. 726; Form. in Ver. Brünn 1895 p. 36; Haussk. symb. p. 41. — Icon: Fl. dan. t. 1103. — Exsicc.: Sint. et Bornm. it. turc. n. 1430.

Herbacea, viridis; caulibus erectis, simplicibus vel ramosis, retrorsum setulosis, eglandulosis; foliis oblongis vel oblongo-lanceolatis, acutis, crenatis, pubescentibus, inferioribus breviter petiolatis, ceteris subsessilibus sessilibusve; verticillastris 6—8 floris, superioribus approximatis; calycis hirsuti dentibus lanceolato-subulatis, tubo vix brevioribus; corollae puberulae tubo incluso.

Ad ripas, in humidis, rarissime. Epirus: pr. Safikbi (Form.); Thessalia: pr. Trikala, Karditza (Haussk.); Boeotia: pr. Lebadea (Orph.). — Jul. Aug. ♃

b. Rhizoma non repens.

α. Herbaceae.

× Rhizoma caules floriferos tantum edens.

⊙ Verticillastra 4—10 flora.

. Caulis praeter pilos patentes, pilis brevissimis glandulosis obsitus.

12. **S. viridis** Bois. et Heldr. diagn. ser. 2 IV. p. 39, Fl. or. IV. p. 727; Haussk. symb. p. 42. — Exsicc.: Sint. et Bornm. it. turc. 1431, 1431b, 1432.

Laete viridis; caulibus erectis, ramosis, pilis brevissimis glandulosis aliisque elongatis patulis sparsis hirsutis; foliis sessilibus, oblongo-lanceolatis, acutis, crenulatis, basi attenuatis, brevissime glanduloso-pubescentibus; verticillastris numerosis, 6—8 floris, in spicam elongatam congestis, inferioribus distantibus; calycis breviter glandulosi, pilisque patulis sparsissimis obsiti, dentibus lanceolatis, longe aristatis, tubo aequilongis; corollae carneae tubo incluso, labio superiore puberulo. — Caules elati, ad 60 cm. alti.

In rupestribus regionis inferioris et montanae, rare. Thessalia: pr. Malakasi, Gionscala, Kalabaka in Pindo (Haussk.), ad monasterium Metoji (Heldr.), in fauce Megarema et ad Plaka pr. Litochori (Sint.) in Olympo. — Jun. Jul. ♃

13. **S. Freynii** Haussk. in Mitth. thür. bot. Ver. 1886 p. 70, symb. p. 42. — *S. dolopica* Form. in D. bot. Mon. 1897 p. 75, in Ver. Brünn 1897 p. 52. — Exsicc.: Heldr. it. thessal. a. 1885.

Obscure viridis; caulibus erectis, simplicibus vel ramosis, pilis brevissimis glandulosis aliisque elongatis patulis hirsutis; foliis ovato-oblongis vel ovato-lanceolatis, acutis, crenatis, basi attenuatis, brevissime glanduloso-pubescentibus pilisque longioribus obsitis, inferioribus petiolatis, superioribus sessilibus; verticillastris numerosis, 6—15 floris, in spicam elongatam congestis, inferioribus distantibus; calycis breviter glandulosi, pilisque patulis crebris obsiti, dentibus lanceolatis, longe aristatis, tubo aequilongis; corollae sordidae ochroleucae tubo incluso, labio superiore piloso. — Praecedenti maxime affinis, ab ea colore

obscure viridi, foliis densius pubescentibus, calyce pilis elongatis crebris subvilloso, corolla majore, ochroleuca specifice distincta; affinis quoque *S. menthaefoliae* Vis. dalmaticae et nuper a Baldacci in Albania australi detectae, quae indumento minus hirsuto et praesertim calycis dentibus latioribus, breviter mucronatis, nec in aristam elongatis differt.

In subalpinis Pindi dolopici ad Gionskala supra Sermeniko (Haussk.) et in mt. Ghavellu (Form.(. — Jul. Aug. ♃

.. Caulis eglandulosus.

; Caules adscendentes, patule hirsuti; folia molliter tomentella, plus minus canescentia.

14. **S. messeniaca** Bois. diagn. XII. p. 81, Fl. or. IV. p. 728. — *S. Tournefortii* Ch. et B. exp. p. 167, Fl. pelop. p. 37, non Poir., sec. Bois. l. c. — Exsicc.: Heldr. pl. fl. hellen. a. 1862.

Caulibus adscendentibus, simplicibus vel parce ramosis, pilis patulis mollibus dense hirsutis; foliis ovato-oblongis, obtusis, obtuse-crenatis, basi cordatis, utrinque sed subtus densius tomentellis, inferioribus petiolatis, superioribus sessilibus; verticillastris 3—6 floris, in spicam brevem approximatis; calycis glandulosi pilisque longioribus patulis hirsuti dentibus triangulari-lanceolatis, breviter mucronatis, rectis, tubo triplo brevioribus; corollae ochroleucae, purpureo-maculatae tubo incluso, labio superiore glanduloso-puberulo. — Differt a duabus antecedentibus statura minori, foliis minoribus, obtusis, basi cordatis et calycis dentibus breviter mucronatis, nec longe aristatis, a *S. menthaefolia* Vis. indumento densiore, foliis cordatis, foliis fere duplo minoribus.

In rupestribus regionis inferioris et montanae. Messenia: pr. Pylos usque ad mt. Diaforti, pr. Phigalea, Androvista ad radices mt. Taygetos (Chaub.); indicatur quoque in mt. Dia insulae Naxos (Chaub.), sed probabiliter ex confusione. — Apr. Jun. ♃

15. **S. canescens** Ch. et B. exp. p. 167, Fl. pelop. p. 37; Bois. fl. or. IV. p. 728. — Exsicc.: Heldr. herb. norm. n. 1365 sub *S. messeniaca*.

Differt a praecedente, cui valde affinis, foliis oblongis vel oblongo-lanceolatis, angustioribus, acutis vel acutiusculis, basi rotundatis vel attenuatis, magis canescentibus et praesertim calycis dentibus lanceolato-subulatis, erecto-patulis, tubo aequilongis.

In rupestribus regionis inferioris et submontanae. Laconia: mt. Selitza, pr. Kalamata (Zahn.), Chimova, Vitilos, Kisternes, ad promontorium Taenaron, pr. Monembasia (Chaub.). — Apr. Jun. ♃

;; Caules erecti, rigiduli, breviter hirti; folia parce hirta, viridia.

, Corolla ochroleuca; antherae labium superius subaequantes.

16. **S. Parolinii** Vis. in mem. ist. venet. I. p. 10 t. 4; Bois. fl. or. IV. p. 730; Heldr. fl. cephal. p. 59; Hal. Beitr. fl. Aetol. p. 2, Beitr. fl. Achaia p. 30. — Exsicc.: Hal. it. gr. a. 1893.

Caulibus retrorsum hispidulis, inferne ramulosis; foliis oblongis vel lanceolatis, acutis, serratis, basi attennatis, inferioribus breviter petiolatis, superioribus sessilibus; verticillastris 4—6 floris, in spicam brevem approximatis, inferioribus distantibus, summis capitulum ovatum formantibus; calycis glanduloso-pilosi, pilisque longioribus obsiti dentibus lanceolatis, aristatis, erecto-patulis, tubo subbrevioribus; corollae tubo subexserto, labio superiore piloso. — Sequenti affinis, ab ea spica brevi, verticillastris summis in capitulum ovatum conglobatis et calycis dentibus longe aristatis discedit.

In rupestribus, dumosis regionis inferioris et montanae. Aetolia: ad radices mt. Taphiassos pr. Antirrhion (Hal.), pr. Naupactos (Parol.); Achaia: in fauce Diakophto (Heldr.), pr. Megaspilaeon (Hal.); Cephalonia: pr. Argostoli, Omala, Valeriano, mt. Atro (Heldr.). — Maio, Jun. ♃

17. **S. recta** L. mant. p. 82; S. et S. pr. I. p. 410; Ch. et B. fl. pelop. p. 38; Raul. cret. p. 836; Bois. fl. or. IV. p. 729; Bald. riv. coll. bot. alb. 1895 p. 65. — Icon: Jacq. fl. austr. t. 359.

Caulibus breviter hispidis, simplicibus vel ramosis; foliis oblongis vel lanceolatis, acutis, crenatis, basi attenuatis, inferioribus petiolatis, superioribus sessilibus; verticillastris 6—10 floris, in spicam elongatam, interruptam dispositis; calycis breviter hirti dentibus triangularibus, breviter mucronatis, erectis, tubo triplo brevioribus; corollae tubo incluso, labio superiore pubescente.

Indicatur in agro Laconico (Sibth.), Creta (Benth. lab. p. 557) et forma quaedam alpina collectori ipso dubia in mt. Tsumerka in Epiro (Bald.). — Species pro ditione nostra valde dubia. — Maio, Jul. ♃ N. v.

„ Corolla purpurea; antherae subinclusae.

18. **S. iberica** M. a. B. fl. taur. cauc. II. p. 51; Bois. fl. or. IV. p. 731. — Exsicc.: Ozanon in Magn. fl. sel. n. 1269, e semine thessalo culta.

Caulibus sparse hirsutis, simplicibus vel ramosis; foliis oblongis vel lanceolatis, acutis, serratis, basi attenuatis, inferioribus subpetiolatis, superioribus sessilibus; verticillastris 6—10 floris, distinctis; calycis hirsuti dentibus triangulari-lanceolatis, purpureo-aristatis, erecto-patulis, tubo subaequilongis; corollae hirsutae tubo subincluso.

Thessalia: mt. Oeta supra Mustapha Bey (Sprun.). — Jun. Jul. ♃

○○ Verticillastra 2 flora.

19. **S. virgata** Ch. et B. exp. p. 166 t. 17; Fl. pelop. p. 37 t. 18; Bois. fl. or. IV. p. 731. — *S. Zuccarinii* Benth. lab. p. 731.

Adpresse sericeo-lanata; caulibus virgatis, ramosis, sparsim foliatis; foliis integris, linearibus, inferioribus obtusis, in petiolum attenuatis, superioribus acutis; verticillastris remotis; floribus subsessilibus; calycis dense sericei dentibus lanceolatis, mucronatis, tubo 4plo brevioribus; corollae albae, sericeae tubo calyce breviore.

In rupestribus regionis inferioris. Acrocorinthus (Zuccar.); Argolis: pr. Kranidi, Troezena (Chaub.). — Jun. Jul. ♃ N. v.

20. **S. tenuifolia** Pall. cat. esp. veg. spont. Taur. in nov. act. acad. Petrop. X. (1797) p. 313 (nomen solum). — *S. angustifolia* M. a. B. fl. taur. cauc. II. p. 51; Form. in Ver. Brünn 1896 p. 70, 1897 p. 52; Haussk. symb. p. 43. — *S. abrotanifolia* Bois. et Heldr. pl. exs. a. 1851. — Exsicc.: Orph. fl. gr. n. 536 (mt. Korfiati Macedoniae); Sint. it. thessal. n. 1037.

Glabra; caulibus virgatis, ramosis; foliis sessilibus, inferioribus in lacinias angustissime lineares pinnatifidis partitisve, superioribus remotis; angustissime linearibus, integris vel lacinula auctis; verticillastris remotis; floribus pedicellatis; calycis glabri vel parce hirto-scabridi dentibus lanceolatis, mucronatis, tubo duplo brevioribus; corollae ochroleucae, roseo suffusae, pubescentis tubo breviter exserto.

In rupinis regionis inferioris et montanae. Thessalia: pr. monasterium Hagios Stephanos supra Kalabaka (Haussk.), mt. Cuka et Hagios Elias in mt. Chassia, pr. Patsios in Olympo (Form.). — Jun. Aug. ♃.

×× Rhizoma caules floriferos et rosulas steriles edens.

21. **S. maritima** L. mant. p. 82; S. et S. pr. I. p. 409; Ch. et B. fl. pelop. p. 37; Bois. fl. or. IV. p. 745. — Icon: Jacq. hort. vind. t. 70.

Breviter tomentella, canescenti-virens; caulibus floriferis infra rosulas steriles adscendentibus, subsimplicibus; foliis oblongis, obtusis, crenulatis, basi rotundatis vel attenuatis, inferioribus petiolatis, superioribus sessilibus; verticillastris 4—6 floris, in spicam brevem approximatis; calycis tomentoso-villosi dentibus lanceolatis, breviter mucronatis, tubo triplo brevioribus; corollae ochroleucae, pubescentis tubo incluso.

In arenosis maritimis regionis occidentalis Peloponnesi (Chaub.) et in rupibus maritimis insulae Hydrae (Sibth.). — Apr. Jul. ♃ N. v.

22. **S. pubescens** Ten. syll. p. 289, Fl. nap. t. 239; Link in Linnaea IX. p. 575; Marg. et R. fl. Zante p. 75. — Huc forsan: *S. arenaria* Friedr. Reise p. 278, vix Vahl. e Nauplia.

Differt a praecedente, cui maxime affinis, caulibus elongatis, decumbentibus, foliis minus rugosis et calycis dentibus lanceolatis, aristatis, tubo duplo brevioribus.

Indicatur in insula Zacyntho et pr. Nauplia Argolidis (Link); a recentioribus non reperta. — Apr. Jul. ♃ N. v.

β. Basi fruticulosae.

× Bracteae calyce subaequilongae.

23. **S. decumbens** Pers. syn. II. p. 123. — *S. mollissima* Willd. en. hort. berol. p. 60 t. 60; Aschers. in bot. Zeit. 1872 p. 629; Bois. fl. or. IV. p. 727; Spreitz. in z. b. G. 1887 p. 686; Gelmi in bull. soc. bot. it. 1889 p. 451. — *S. hirta* Pieri corc. fl. p. 78, non L. — Exsicc.: Spreitz. it. ion. II. n. 139; Baen. herb. europ. n. 9349.

Molliter villosa, canescens; caulibus adscendentibus vel erectis, simplicibus vel ramosis; foliis petiolatis, cordato-ovatis, acutis vel obtusiusculis, crenatis, floralibus sessilibus; verticillastris 5—multifloris, in spicam brevem, ovatam vel oblongam congestis; bracteis lineari-lanceolatis; calycis villosi dentibus late lanceolatis, aristatis, calyce subaequilongis; corollae ochroleuca, pubescentis tubo subexserto. — Species basi manifeste suffruticosa, ramis vetustis lignosissimis, duris, ab omnibus aliis hujus sectionis bracteis evolutis discedens et hac nota ad *Eriostachydes* accedens, habitu toto, sequenti valde simili, tamen huic inserenda.

Ad muros, in rupestribus regionis inferioris et montanae Corcyrae: pr. Potamo, Peleka (Bickn.), Benizze, Gasturi (Baen.), mt. Deca, pr. Palaeokastrizza, Lacones (Spreitz.), Scripero, San Pantaleone, Spartilla (Gelmi). — Maio, Jun. ♃

×× Bracteae minutae vel nullae.

○ Rami inermes.

. Folia elliptica vel oblonga, nunquam dense pannosa.

; Flores subsessiles vel breviter pedicellati.

, Caules patule hirsuti; verticillastra saltem superiora in spicam ovatam vel oblongam congesta.

24. **S. argolica** Bois. diagn. XII. p. 83. — *S. Swainsoni v. argolica* Bois. fl. or. IV. p. 735; Haussk. symb. p. 43. — *S. Swainsoni* Haussk. l. c., non Benth. — Huc forsan: *S. circinnata* Friedr. Reise p. 274, non L'Hér. — Exsicc.: Orph. fl. gr. n. 1167; Heldr. herb. norm. n. 1067.

Patule hirsuta, viridis; caulibus adscendentibus, simplicibus vel ramosis; foliis petiolatis, rugosis, ellipticis vel oblongis, obtusis, crenatis, basi rotundatis vel cuneatis, floralibus sessilibus, verticillastra superantibus, verticillastris 6—10 floris in spicam brevem, ovatam vel oblongam condensatis; bracteis minutis; calycis hirsuti, glandulosi dentibus triangularibus, breviter mucronatis, tubo paulo brevioribus; corollae pallide carneae, pubescentis tubo subexserto. — Caules caespites e fissuris rupium oriundos formantes, basi prostrati, 10—30 cm. longi.

In rupibus regionis inferioris, rarissime. Argolis: pr. Nauplia (Sprun.), Tolon, Mykene (Haussk.), pr. Vromolimni, insula Hydra (Heldr.). — Apr. Maio. ♃

25. **S. Swainsoni** Benth. lab. p. 535; Vis. in mem. ist. ven. I. p. 9 t. 3; Bois. fl. or. IV. p. 735 cum var. *β. scyronica*; Heldr. chlor. Parn. p. 25. — *S. scyronica* Bois. diagn. V. p. 31, XII. p. 83; Heldr. fl. cephal. p. 59 et Spreitz. in z. b. G. 1887 p. 666 pro var. *S. Swainsoni*. — Probabiliter huc: *S. candida* Ung. p. 127, non Ch. et B., sec. Heldr. fl cephal. p. 39. — Exsicc.: Heldr. herb. norm. n. 90; Orph. fl. gr. n. 436 et 438.

Molliter sericeo-cinerea; caulibus prostratis vel adscendentibus; foliis parvis, rugulosis, oblongis vel ellipticis, obtusiusculis, denticulatis, in-

ferioribus in petiolum brevissimum attenuatis; floralibus sessilibus, calycem aequantibus; verticillastris 2—4 floris, in spicam brevem, oblongam, inferne interdum laxiusculam condensatis; bracteis minutis; calycis hirsuti, glandulosi dentibus triangularibus, breviter mucronatis, tubo paulo brevioribus; corollae purpurascentis, pubescentis tubo subexserto. — Differt a praecedente statura minori, indumento densiori cinereo, foliis parvis, verticillis paucifloris et floribus minoribus.

In rupium fissuris. Megara: ad Scyronides petras — nunc Kakiskala —, pr. Lutraki in isthmo Corinthiaco (Heldr.), ad radices mt. Parnassus pr. Rachova (Swains.); Cephalonia: pr. Karakas, Asso, Poros, Steno, Kipuria, Tafius (Heldr.); Leucas: ad promontorium Zuana (Spreitz.); Ithaca (Webb.): ad castellum Ulysses (Vis.). — Apr. Jul. ♃.

26. **S. Spruneri** Bois. diagn. XII. p. 82, Fl. or. IV. p. 736; Ung. Reise p. 127; Haussk. symb. p. 43. — Exsicc.: Orph. fl. gr. n. 437; Heldr. herb. norm. n. 389 et 1272.

Sparsim et patule hispida, viridis; caulibus rigidis, simplicibus vel parce ramosis; foliis rigidis, valde rugosis, argute serratis, infimis oblongo-ellipticis, obtusiusculis, in petiolum attenuatis, ceteris sessilibus linearilanceolatis, acutis, floralibus basi dilatatis, acuminatis, subintegris, flores superantibus; verticillastris 4—6 floris, in spicam ovatam congestis, inferioribus saepe remotis; bracteis nullis; calycis hirsuti, eglandulosi, dentibus lanceolatis, mucronatis, tubo dimidio brevioribus; corollae albidae tubo incluso, labio superiore piloso. — A duabus praecedentibus, caulibus et foliis rigidis, indumento rariore obductis, foliis anguste serratis, floralibus dilatatis acuminatis specifice differt.

In rupestribus regionis submontanae in abietinam adscendens, rare. Attica: mt. Parnes (Sprun.), Kerata, Pateras, Cithaeron (Heldr.). — Maio, Jun. ♃.

„ Caules virgati, saltem in parte superiori glabri; verticillastra remotissima.

27. **S. tetragona** Bois. et Heldr. in Bois. fl. or. IV. p. 736; Hal. in ö. b. Z. 1896 p. 16, 1897 p. 98. — Icon: Rouy. pl. europ. rar. t. 270. — Exsicc.: Heldr. et Hal. fl. sporad. a. 1896.

Viridis; caulibus virgatis, flexuosis, simplicibus vel ramosis, glabris vel inferne plus minus setulosis; foliis ad caulis basin congestis, oblongo-spathulatis, obtusis, dentatis, adpressiuscule hirtis, in petiolum attenuatis, superioribus sessilibus, oblongo-linearibus, integris, floralibus minimis, superioribus calyce multo brevioribus; verticillastris 2—6 floris, remotissimis; calycis parce setosi, dentibus triangularibus, breviter mucronatis, tubo triplo brevioribus; corollae flavidae, purpureo-maculatae tubo incluso, labio superiore hispido. — Caules rigidi, flexuosi, ad 30 cm. alti, acute quadranguli, ad basin dense foliosi, ceterum subnudi.

In rupestribus regionis inferioris et montanae, rarissime. Euboea: pr. Steni in mt. Dirphys (Heldr); Sporadum insula Jura (Reis.). — Maio, Jun. ♃.

;; Flores crassiuscule pedicellati; pedicelli tubo calycis subaequilongi.

28. **S. mucronata** Sieb. avis p. 4, rem. p. 5; Raul. cret. p. 837; Bois. fl. or. IV. p. 737. — Exsicc.: Heldr. pl. cret. n. 1458.

Molliter lanata; caulibus rigidis, ramosis; foliis oblongis, obtusis, crenatis, in petiolum attenuatis, floralibus sessilibus, superioribus calyce brevioribus; verticillastris 2—4 floris, approximatis; calycis coriacei, sulcato-costati, hirsuti glandulisque sessilibus obsiti dentibus pungentibus, tubo subaequilongis; corollae carneae, villosae tubo incluso. — Caules 5—15 cm. alti; floribus manifeste pedicellatis, calycibus elevato-costatis, dentibus pungentibus insignis.

In saxosis regionis inferioris Cretae (Sieb.): ad promontorium Sidero (Heldr.), inter Palaeokastro et Zakro (Raul.). — Apr. Maio. ♃

.. Folia ovato-subrotunda, dense pannosa.

; Calyx eglandulosus.

29. **S. candida** Ch. et B. exp. p. 167 t. 19, Fl. pelop. p. 37 t. 20; Bois. fl. or. IV. p. 742. — Exsicc.: Pichl. pl. gr. a. 1876.

Niveo-pannosa, sericeo-nitens; caulibus prostratis vel adscendentibus, subsimplicibus; foliis breviter petiolatis, ovato-subrotundis, obscure crenulatis; verticillastris 4—6 floris, in spicam oblongam approximatis; bracteis subnullis; calycis lanato-tomentosi dentibus lanceolatis, mucronatis, tubo subaequilongis; corolla alba, pubescente, fauce purpureo-maculata. — Caules 10—20 cm. longi, fragiles.

In rupium fissuris regionis submontanae. Arcadia: mt. Diaforti, pr. Thuria Messeniae, pr. Mistra, mt. Taygetos et pr. Zarnate Laconiae (Chaub.). — Maio, Jul. ♃

30. **S. chrysantha** Bois. et Heldr. diagn. VII. p. 56, Fl. or. IV. p. 743. — *S. candida β. chrysantha* Benth. in DC. pr. XII. p. 475. — Exsicc.: Orph. fl. gr. n. 76; Heldr. herb. norm. n. 1366.

Praecedenti maxime affinis, ab ea indumento absque nitore et corolla lutea vix specifice differre videtur.

In rupium fissuris regionis submontanae. Laconia: mt. Malevo pr. Ajanni (Orph.) et Kastanitza (Heldr.). — Maio, Jul. ♃

;; Calyx glandulosus.

31. **S. Spreitzenhoferi** Heldr. in ö. b. Z. 1880 p. 344; Spreitz. in z. b. G. 1887 p. 666; Bois. fl. or. suppl. p. 362; Fors. in bull. Bois. V. p. 399. — Exsicc.: Spreitz. it. ion. IV. a. 1880.

Cano-pannosa, opaca; caulibus prostratis vel adscendentibus, simplicibus; foliis breviter petiolatis, ovato-subrotundis, crenulatis; verticillastris 4—6 floris, in spicam oblongam approximatis; bracteis lineari-setaceis, calyce brevioribus; calycis cano-tomentosi et glandulosi dentibus triangularibus, muticis, tubo triplo brevioribus; corolla alba, pubescente, fauce roseo-picta.

In rupium fissuris insulae Cythaera ad castrum Kapsali et in faucibus Kalamo ad antrum Mylopotamo (Spreitz.). — Maio, Jul. ♃

○○ Rami spinescentes.

32. **S. spinosa** L. sp. p. 581; S. et S. pr. I. p. 409; Sieb. avis p. 4. rem. p. 5; Raul. cret. p. 837; Bois. fl. or. IV. p. 736; Bald. viagg. Creta p. 85. — Icon: Fl. gr. t. 559. — Exsicc.: Rev. pl. cret. n. 139; Bald. it. cret. n. 35; Dörfl. fl. aeg. n. 43.

Dumuloso-erinacea, adpresse sericeo-lanata; caulibus humilibus, rigidis, ramosis, ramis apice spinescentibus; foliis oblongo-linearibus, integris, inferioribus in petiolum attenuatis, ceteris sessilibus, floralibus calyce brevioribus; verticillastris 1—4 floris; floribus sessilibus; calycis sericei dentibus triangulari-lanceolatis, spinosis, tubo triplo brevioribus; corollae albidae, pilosae tubo incluso.

In collibus aridis regionis inferioris et montanae. Creta: pr. Hagia Triada (Bald.), Lakus (Rev.), Suia, Hagios Paulos, Embaro, Rhizokastron, Hodhegetria (Raul.); Cycladum insula Anaphi (Leon.). — Maio, Jul. ♃

Obs. *S. glutinosa* L. sp. p. 581, species corsica, erronee a S. et S. pr. I. p. 409 in Creta indicatur.

3. Sectio. *Olisia* Bois. fl. or. IV. p. 716. — Species annuae.

a. Corolla albido-ochroleuca.

33. **S. annua** L. sp. p. 573 (*Betonica*); L. sp. ed. 2 p. 813; Bois. fl. or. IV. p. 745; Heldr. chlor. Parn. p. 25; Hal. Beitr. fl. Epir. p. 35; Bald. riv. coll. bot. alb. 1896 p. 87. — *S. annua* v. *parnassica* Heldr. et Sart. in Heldr. herb. norm. n. 287.

Glabriuscula, superne pubescens et saepius plus minus glandulosa: caulibus erectis vel adscendentibus, simplicibus vel divaricato-ramosis; foliis breviter petiolatis, ovatis oblongisve, crenatis, basi rotundatis vel cuneatis, floralibus lanceolatis, sessilibus; verticillastris 3—6 floris, summis approximatis; bracteis minutis; calycis tubuloso-campanulati, hirsuti glandulisque sessilibus adspersi, dentibus lanceolato-subulatis, longe cuspidatis, tubo duplo brevioribus; corollae albido-ochroleucae, pubescentis tubo exserto.

β. **adenocalyx** C. Koch in Linnaea XXI. p. 691 pro sp.; Haussk. symb. p. 43. — *S. thessala* Sint. it. thessal. n. 684. — Caules superne, folia floralia et calyces dense glanduloso-pilosi. — Intermediis ad typum transit. — Exsicc.: Sint. l. c.; Bald. it. alb. epir. IV. n. 42.

In saxosis, dumosis regionis montanae, rare. Epirus: mt. Micikeli (Bald.), Peristeri, Tsumerka (Hal.), pr. Chaliki (Haussk.); mt. Parnassus (Heldr.); in ditione formae plus minusve glandulosae tantum occurrere videntur. — Maio, Aug. ⊙

34. **S. spinulosa** S. et S. pr. I. p. 410; Sieb. avis p. 4, rem. p. 5, Marg. et R. fl. Zante p. 75; Friedr. Reise p. 275; Ung. Reise p. 177; Raul. cret. p. 826; Spreitz. in z. b. G. 1877 p. 716; Bois. fl. or. IV. p. 746; Heldr. fl. cephal. p. 59; Gelmi in bull. soc. bot. ital. 1889 p. 451; Hal. in ö. b. Z. 1896 p. 16; Haussk. symb. p. 42. — *S. betonicaefolia* Desf. choix p. 22 t. 14; Ch. et B. exp. p. 166, Fl. pelop. p. 37. — Exsicc.: Heldr. herb. norm. n. 966 et 966b.

Hispidula, superne glanduloso-pilosa; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis cordato-ovatis, crenatis, parce hirtis, inferioribus longe petiolatis, floralibus oblongis, sessilibus; verticillastris 4—6 floris, distinctis; bracteis subnullis; calycis late campanulati, glanduloso-pilosi dentibus late ovatis, breviter mucronatis, tubo subtriplo brevioribus; corollae albido-ochroleucae, glandulosae tubo subexserto. — Foliis inferioribus longe petiolatis, cordatis, floribus duplo majoribus a praecedente facillime distinguitur.

In collibus apricis, arvis derelictis regionis inferioris, rarior. Aetolia: pr. Aetolikon (Reis.); Attica: pr. Athenas, Porto Rhaphti (Heldr.); Achaia: pr. Psatopyrgos (Heldr.); insula Poros (Friedr.); Messenia: pr. Methone, Pylos (Chaub.); Creta (Sibth.): pr. Almyros, Lutro (Raul.); Zante (Marg.); Cephalonia: pr. Aphrato (Heldr.); Ithaca (Spreitz.); Corcyra: pr. Potamo (Spreitz.). — Apr. Jun. ⊙

b. Corolla purpurea.

35. **S. serbica** Panc. fl. princ. Serb. p. 564; Haussk. symb. p. 43.

Patule hirsuta; caulibus erectis, simplicibus vel ramosis; foliis petiolatis, ovatis, crenatis, basi rotundatis vel subcordatis, floralibus subsessilibus, varticillastra superantibus; verticillastris 4—8 floris, distantibus; bracteis lanceolatis, calyce aequilongis; calycis hirsuti, demum aucti dentibus lanceolatis, cuspidatis, tubo aequilongis; corollae puberulae limbo calycem vix excedente. — Caulis 10—30 cm. altus.

In herbidis oropedii Neuropolis in Pindo dolopico (Haussk.). — Maio, Jul. ⊙ N. v.

36. **S. arvensis** L. sp. p. 578 (*Glecoma*); L. sp. ed. 2 p. 814; Ch. et B. fl. pelop. p. 37; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 836; Bois. fl. or. IV. p. 747; Gelmi in bull. soc. bot. ital. 1889 p. 451; Hal. in ö. b. Z. 1897 p. 325. — Icon: Fl. dan. t. 587. — Exsicc.: Heldr. herb. norm. n. 1572.

Patule hirsuta; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis petiolatis, ovato-cordatis, crenatis, floralibus sessilibus, oblongis, calyce sublongioribus; verticillastris 4—6 floris, distantibus; bracteis nullis; calycis hirsuti dentibus lanceolatis, mucronatis, tubo subaequilongis; corollae puberulae limbo calycem vix excedente.

In locis aridis, ad vias regionis inferioris, rare. Messenia: pr. Messene, Kalamata (Chaub.), Petalidi (Zahn); Creta: pr. Canea (Weiss); Corcyra (Gelmi). — Apr. Jul. ⊙

Obs. *S. palaestina* L. sp. ed. 2 p. 1674, species asiatica, a S. et S. pr. I. p. 409 certe ex confusione quadam „per totam Graeciam copiose“ indicatur. — *S. lamioides* Sieb. in Flora I. p. 273, solum nomen: indicatur in Creta, mihi ignota.

17. Ballota L. gen. n. 720.

a. Calycis limbus 5 dentatus.

1. **B. nigra** L. sp. p. 582; S. et S. pr. I. p. 411; Ch. et B. exp. p. 169; Fraas fl. class. p. 180; Clem. sert. p. 77; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 838; Bois. fl. or. IV. p. 775; Heldr. fl. cephal. p. 59; Form. in D. bot. Mon. 1891 p. 30, in Ver. Brünn 1895 p. 36, 1896 p. 71, 1897 p. 53; Hal. Beitr. fl. Epir. p. 35; Haussk, symb. p. 43. — *B. alba* L. fl. suec. p. 206; Ch. et B. fl. pelop. p. 38; Friedr. Reise p. 274; f. floribus albis. — *B. foetida* Lam. fl. fr. II. p. 381. — *B. rupestris* Form. in D. bot. Mon. 1891 p. 30, in Ver. Brünn 1896 p. 71, non Vis. — Icon: Rchb. t. 17. — Exsicc.: Sint. et Bornm. it. turc. n. 1423; Dörfl. fl. gr. n. 202.

Plus minus hirsuta; caulibus erectis vel adscendentibus, ramosis vel simplicibus; foliis petiolatis, ovatis, grosse dentatis, floralibus verticillastra superantibus; verticillastris multifloris, distantibus; calycis dentibus triangulari lanceolatis, brevibus, in spinulam subpungentem attenuatis; corollae roseae, rarius albae tubo subexserto.

In ruderatis regionis inferioris et montanae passim per totam Graeciam. — Jun. Aug. ♃

b. Calycis limbus 10 dentatus.

2. **B. hirsuta** Willd. sp. III. p. 113; Sieb. in Flora V. II. p. 639; (*Marrubium*); Nym. syll. p. 98; Hal. Beitr. fl. Epir. p. 35; non Benth. lab. p. 595 et in DC. pr. XII. p. 518, quae ad *Marrubium hispanicum* L. sp. p. 583 spectat. — *Marrubium rupestre* Biv. manip. II. p. 1 t. 1. — *B. hispanica* Benth. lab. p. 597, quoad pl. ital. et dalm. — *B. rupestris* Vis. fl. dalm. II. p. 216; Form. in Ver. Brünn 1895 p. 36. — *B italica* Benth. in DC. pr. XII. p. 519; Raul. cret. p. 838. — *B. saxatilis* Guss. fl. sic. syn. II. p. 82, non Sieb. in Benth. lab. p. 596. — Exsicc.: Hal. it. gr. secund. a. 1893.

Hirsuta; caulibus erectis, simplicibus vel ramosis; foliis petiolatis, ovato-cordatis, crenato-dentatis, subtus canescentibus, floralibus verticillastra superantibus vel aequantibus; verticillastris multifloris, distantibus; calycis dentibus lanceolatis, brevibus, abrupte aristatis; corollae carneae tubo exserto.

In dumosis regionis montanae, rarissime. Epirus: pr. Govrica. mt. Micikeli (Form.), infra Kalarrytes (Hal.); ?Creta (Sieb.). — Jul. Aug. ♃

c. Calycis limbus late membranaceo-expansus, crenatus.

3. **B. acetabulosa** L. sp. p. 584; Urv. enum. p. 69; Ch. et B. exp. p. 169, Fl. pelop. p. 38; Link in Linn. IX. p. 576; (*Marrubium*); Benth. lab. p. 595; Clem. sert. p. 77; Raul. cret. p. 838; Bois. fl. or. IV. p. 772; Hal. in z. b. G. 1888 p. 761; Hal. Beitr. fl. Aetol. p. 9, Beitr. fl. Thessal. p. 17, Beitr. fl. Achaia p. 30, in ö. b. Z. 1896 p. 16; Form. in D. bot. Mon. 1891 p. 30, 1898 p. 79, in Ver. Brünn

1896 p. 71, 1897 p. 53; Haussk. symb. p. 43; Heldr. fl. Aegina p. 385, chlor. Thera p. 19, chlor. Mykon. p. 249. — *Marrubium pseudodictamnus* S. et S. pr. I. p. 413, Fl. gr. VI. p. 50, t. 562; Friedr. Reise p. 274; Fraas fl. class. p. 181; non L. — *Beringeria acetabulosa* Neck. elem. bot. p. 312; Weiss in z. b. G. 1869 p. 743; Spreitz. in z. b. G. 1887 p. 666. — Exsicc.: Orph. fl. gr. n. 526; Heldr. herb. norm. n. 215, herb. dimorph. n. 42; Sint. it. thessal. n. 1513.

Caulibus erectis vel adscendentibus, simplicibus vel ramosis, patule villosulis; foliis breviter petiolatis, ovatis vel orbiculatis, basi cordatis, grosse crenatis, junioribus utrinque dense albo-lanatis, demum tenuiter canescentibus, floralibus verticillastra aequantibus vel superantibus; verticillastris parum distantibus vel contiguis; calycis tomentelli limbo expanso, membranaceo, tubo multo longiore, irregulariter multicrenato, crenis muticis vel brevissime mucronatis; corollae purpurascentis tubo breviter exserto, labio superiore hirsuto.

In ruderatis, saxosis regionis inferioris et montanae. Thessalia: frequens a Meteora et Kalabaka per planitiem usque ad mare Aegaeum, in Phthiotide et Attica; dein in Aetolia: ad radices mt. Chalkis ad sinum Patranum; pr. Segditza ad radices mt. Kiona (Hal.); Peloponnesus: pr. Nauplia (Link), Poros (Friedr.), in valle fl. Voreikos pr. Megaspilaeon, pr. Kalavryta, Planideri Achaiae (Hal.), mt. Diaforti, pr. Gortys, Francovritzi, Vitiolos, Sparta, Scutari, Monembasia (Chaub.); Sporadum insula Jura (Reiser); Petalium insula Tragonisi (Holzm.); insula Aegina (Heldr.); Cycladum insula: Syra (Weiss), Tenos, Naxos (Chaub.), Rhenea (Tunt.), Melos, Thera (Urv.); Cythaera: mt. Hagios Georgios (Spreitz.); Creta: pr. Aradhena, Anopolis (Raul.). — Apr. Jun. ♃

4. **B. pseudodictamnus** L. sp. p. 583; Sieb. avis p. 4, rem. p. 5, in Flora I. p. 273; (*Marrubium*); Benth. lab. p. 594; Raul. cret. p. 838; Bois. fl. or. IV. p. 772; Spreitz. in ö. b. Z. 1880 p. 346, in z. b. G. 1887 p. 666, 1890 p. 298; Bald. viagg. Creta p. 86. — *Marrubium acetabulosum* S. et S. pr. I. p. 413; non L. — *Beringeria pseudodictamnus* Neck. elem. bot. p. 312; Weiss in z. b. G. 1869 p. 743. — Exsicc.: Rev. pl. cret. n. 142, in Baen. herb. europ. n. 4887, in Magn. fl. sel. n. 941; Bald. it. cret. n. 36.

Praecedenti maxime affinis, ab ea differt foliis multo minoribus, ad 2 cm. tantum (nec ad 6 cm.) longis, etiam adultis dense albo-lanatis, verticillastris distantibus; calycis dense lanati limbo tubo parum longiore, 5 lobo, lobulis accessoriis interdum quinis minimis aucto et corollae labio superiore dense villoso.

In ruderatis, locis aridis regionis inferioris. Creta: pr. Kissamos (Rev.), Canea, Akroteri, Malaxa, Sphakia (Raul.), Herakleion (Heldr.), Kalyves, Vamos, Anopolis (Spreitz.); Anticythera et Cythera (Spreitz.). — Jun. Jul. ♃

18. Leonurus L. gen. n. 722.

1. **L. cardiaca** L. sp. p. 584; Ch. et B. fl. pelop. p. 38; Bois. fl. or. IV. p. 753; Form. in D. bot. Mon. 1891 p. 30, in Ver. Brünn 1895 p. 36, 1896 p. 71, 1897 p. 53; Haussk. symb. p. 43. — Icon: Fl. dan. t. 727. — Exsicc.: Sint. et Bornm. it. turc. n. 1403.

Adpresse pubescens vel glabriusculus; caule elato, ramoso; foliis petiolatis, ovatis oblongisve, inferioribus palmatifidis, floralibus trifidis, verticillastro multo longioribus; verticillastris distantibus; calycis glabri dentibus tubo subbrevioribus; corollae roseae, hirsutae tubo intus oblique annulato, labio inferiore trifido, lobo medio integro.

In collibus, ruderatis regionis inferioris et montanae. Epirus: pr. Han Kauberga, Govrica (Form.); Thessalia: mt. Zygos, pr. Chaliki, Klinovo, Korona, Kalabaka, Karditza (Haussk.), Velitsena, Koturi, Said Pascha, Bisula, Mavreli, Vlachava, Jerakari, pr. Portaria et Plessidi in mt. Pelion (Form.), pr. Litochori in Olympo (Sint.); Eurytania: pr. Karpenisi (Samarit.); Achaia: pr. Patras (Daenzer). — Jun. Aug. ♃

2. **L. marrubiastrum** L. sp. p. 584; Bois. fl. or. IV. p. 754. — *Chaiturus marrubiastrum* Rchb. fl. germ. p. 317, icon. XXVIII. t. 32. —

Adpresse pubescens; caule elato, ramoso; foliis petiolatis, ovato-lanceolatis lanceolatisve, subinciso-dentatis, floralibus verticillastro multo longioribus; verticillastris distantibus; calycis pubescentis dentibus tubo 2—3 plo brevioribus; corollae pallide roseae, pubescentis tubo exannulato, labio inferiore trifido, lobo medio subintegro.

In dumosis humidiusculis pr. Katerina in Olympo thessalo (Heldr.). — Jul. Aug. ♃ N. v.

5. Tribus. NEPETEAE Benth. lab. p. 462.

19. Glechoma L. gen. n. 714.

1. **G. hederacea** L. sp. p. 578; S. et S. pr. I. p. 404; Ch. et B. fl. pelop. p. 36; Dallap. prosp. p. 82. — Icon: Fl. dan. t. 789.

Glabra vel pubescens; caulibus repentibus, ramis adscendentibus; foliis petiolatis, reniformi-orbiculatis, crenatis; verticillastris 2—3 floris, axillaribus; calycis dentibus triangulari-lanceolatis, tubo 3—4 plo brevioribus; corollae violaceae tubo calyce subtriplo longiore.

In umbrosis Laconiae (Sibth.) et in Cephalonia (Dall.). — Apr. Jun. ♃ N. v.

20. Nepeta L. gen. n. 710.

a. Nuculae laeves.

1. **N. cataria** L. sp. p. 570; Pieri corc. fl. p. 76; Bois. fl. or. IV. p. 643; Heldr. chlor. Parn. p. 25; Form. in D. bot. Mon. 1891 p. 29, in Ver. Brünn 1895 p. 36, 1896 p. 69, 1897 p. 51; Hal. in

ö. b. Z. 1897 p. 325; Haussk. symb. p. 45. — Icon: Fl. dan. t. 580. — Exsicc.: Orph. fl. gr. n. 551; Sint. et Bornm. it. turc. n. 1422.

Pubescenti-canescens; caulibus erectis, ramosis; foliis petiolatis, cordato-ovatis, grosse dentatis, subtus magis canescentibus; cymis breviter pedunculatis, in racemos oblongos, subinterruptos, interdum ramosos congestis; bracteis subulatis, calycis tubo brevioribus; calycis pubescentis, ovati, ore obliqui dentibus lanceolatis, spinuloso-mucronatis, tubo subbrevioribus, superioribus sublongioribus; corollae hirtae, albidae, saepe rubello-punctatae tubo incluso.

In dumosis regionis praesertim montanae et subalpinae. Epirus: pr. Ephemiades, mt. Micikeli (Form.); Thessalia: pr. Sermeniko, mt. Karava, pr. Kastreiki (Haussk.), Kalabaka (Heldr.), Koturi, Kastania et Kataphygi in Pindo, pr. Velestinos, pr. Spilia in mt. Ossa, pr. Miluna (Form.) et Hagios Dionysios (Heldr.) in Olympo; Aetolia: mt. Korax pr. Vustinitza (Tuntas); mt. Parnassus pr. Delphi (Orph.); Achaia: mt. Kyllene (Heldr.); Corcyra (Pieri). — Jun. Aug. ♃

b. Nuculae tuberculatae.

α. Bracteae late ovatae, acutae.

2. **N. scordotis** L. sp. ed. 2 p. 798; Sieb. avis rem. p. 5; Raul. cret. p. 833; Weiss in z. b. G. 1869 p. 743. — *N. tomentosa* Sieb. avis p. 4. — *N. incana* Sieb. avis p. 4, rem. p. 5, sec. Raul. l. c. — Exsicc.: Rev. pl. cret. n. 141.

Tomentoso-lanata; caulibus adscendentibus, elatis, simplicibus vel inferne ramosis; foliis petiolatis, cordato-ovatis, grosse crenatis, subtus magis canescentibus, floralibus sessilibus; verticillastris multifloris, densis, inferioribus distinctis, superioribus approximatis; bracteis calycem aequantibus; calycis villoso-lanati dentibus subaequalibus, lanceolato-subulatis, tubo aequilongis; corollae puberulae, coerulescentis tubo calyce sesquilongiore. — Stirps pulchra, habitu *Stachydis circinnatae.*

In incultis, siccis regionis inferioris Cretae: pr. Kissamos (Rev.), Canea, promontorium Meleka, Perama (Raul). — Mart. Maio. ♃

β. Bracteae oblongo-lanceolatae, lineares vel subulatae.

× Radix fasciculato-fusiformis.

3. **N. melissaefolia** Lam. dict. I. p. 703; Desf. choix pl. p. 19 t. 12; Urv. enum. p. 65; Raul. cret. p. 633; Bois. fl. or. IV. p. 659. — Exsicc.: Heldr. pl. cret. n. 1460.

Pubescens; caulibus adscendentibus, simplicibus; foliis petiolatis, cordato-ovatis, crenatis, floralibus oblongis, sessilibus; verticillastris plurifloris, distinctis; bracteis subulatis, calyce multo brevioribus; calycis breviter hirti, ore obliqui dentibus lanceolatis, acutis, tubo 4plo brevioribus, superioribus submajoribus; corollae puberulae, coerulescentis tubo breviter exserto, labio inferiore coeruleo, rubro-punctato. — Ab omnibus radice fibris fusiformibus, fasciculatis distinctissima.

In dumosis regionis inferioris. Creta: inter Chersonesos et Malea (Raul.), pr. Sakro distr. Sitia (Heldr.); Cycladum insula Melos (Urv.). — Apr. Maio. ♃

×× Radix non fasciculato-fusiformis.

○ Folia pleraque sessilia; verticillastra laxiuscula, laxe racemosa; calycis dentes acuti; flores saepissime violacei.

4. **N. nuda** L. sp. p. 570; S. et S. pr. I. p. 398, Fl. gr. VI. t. 547; Benth. lab. p. 486; Bois. fl. or. IV. p. 663, suppl. p. 362; Freyn. in Flora 1884 p. 682; Hal. in z. b. G. 1888 p. 761; Heldr. chlor. Parn. p. 25. — *N. violacea* Vill. fl. delph. II. p. 367; Form. in D. bot. Mon. 1891 p. 29; Hal. Beitr. fl. Epir. p. 36, in ö. b. Z. 1897 p. 325; non L. — Exsicc.: Orph. fl. gr. n. 432; Heldr. herb. norm. n. 261; Dörfl. fl. gr. n. 211.

Minute pubescens vel glabriuscula; caulibus erectis, stricte paniculatis, ramis subaequilongis; foliis sessilibus vel inferioribus breviter petiolatis, oblongis vel oblongo-lanceolatis, crenatis, basi cordatis; verticillastris secus paniculae ramos densiuscule racemosis, breviter pedunculatis inflorescentiam plus minus confertam formantibus; bracteis subulatis, calyce brevioribus; calycis pubescentis, fructiferi ovati dentibus subaequalibus, lanceolatis, acutis, tubo subbrevioribus; corollae violaceae, pubescentis tubo subexserto. — Bracteae, calycis dentes et corollae plantae graecae fere semper violaceae; flores nunc androdynami, majores, nunc gynodynami minores.

β. **epirota.** — Verticillastra longiuscule pedunculata, multiflora, racemum simplicem, densiusculum, angustum, elongatum formantia; flores parvi; ulterius observanda. — Exsicc.: Hal. it. gr. sec. a. 1893.

In herbidis saxosis regionis subalpinae et alpinae mt. Tymphrestus, Oeta (Heldr.), Korax (Tunt.), Kiona (Hal.), Parnassus (Sibth.), Ossa (Form.); — *β.* mt. Tsumerka pr. Vulgarelion in Epiro (Hal.). — Jun. Aug. ♃

5. **N. pannonica** L. sp. p. 570; Jacq. en. pl. vind. p. 106, Fl. austr. II. t. 129; Haussk. symb. p. 45; Form. in Ver. Brünn 1897 p. 51. — Exsicc.: Sint. it. thessal. n. 1144.

Differt (an specifice) a praecedente ramis paniculae inferioribus patulis, longioribus, inflorescentiam laxam pyramidatam formantibus, verticillastris distantibus.

In dumosis, silvaticis subalpinis. Thessalia: in oropedio Neuropolis, pr. monasterium Korona (Haussk.), mt. Agrapha, Karava, Ghavellu, pr. Kalivia, Sermenikon in Pindo, mt. Pirgo, Alafovris, Mavrika et Katasara in mt. Othrys (Form.), mt. Ossa (Heldr.); Doris: mt. Kiona pr. Segditza (Hal.). — Jun. Aug. ♃

○○ Folia petiolata; verticillastra densa, superiora in spicam oblongam congesta; calycis dentes aristato-mucronati; flores albi. Species valde affines.

. Calycis dentes tubo aequilongi.

6. **N. Sibthorpii** Benth. lab. p. 474; Bois. fl. or. IV. p. 652. — *N. italica* S. et S. pr. I. p. 399, Fl. gr. VI. t. 548; Link in

Linnaea IX. p. 575; Ch. et B. fl. pelop. p. 36; Friedr. Reise p. 277; non L. sp. p. 571, quae indumento tenuiore, verticillastris distinctis et calyce minus profunde fisso specifice differre videtur. — *N. argolica* Ch. et B. exp. p. 163 t. 16, Fl. pelop. p. 36 t. 17; Link l. c. — Exsicc.: Heldr. herb. norm. n. 555, herb. fl. hellen. n. 76, in Baen. herb. europ. n. 3219; Orph. fl. gr. n. 210 et 210 bis; Dörfl. fl. gr. n. 422.

Caulibus erectis vel adscendentibus, adpressissime cinereo-velutinis, simplicibus vel parce ramosis; foliis adpresse cano-tomentosis, breviter petiolatis, ovato-oblongis, cordatis, crenatis; verticillastris in spicam oblongam condensatis, inferioribus saepius distantibus; bracteis oblongo-lanceolatis, subulato-aristatis, late scarioso-marginatis, calycem aequantibus; calycis adpresse tomentelli dentibus subaequalibus, aristatis, tubo aequilongis; corollae albae, pubescentis tubo calycem non excedente.

In dumosis saxosis regionis inferioris, in subalpinam adscendens. Attica: pr. Athenas (Sibth.), mt. Parnes, Pateras (Heldr.); isthmus Corinthiacus (Chaub.); Argolis: pr. Nauplia (Link), Argos, in peninsula Methana (Chaub.), pr. Poros (Friedr.); Achaia: mt. Kyllene (Orph.), pr. Patras (Link). — Maio, Jul. ♃

7. **N. parnassica** Heldr. et Sart. in Bois. diagn. ser. 2. IV. p. 22, Fl. or. IV. p. 654; Heldr. chlor. Parn. p. 25. — *N. Sibthorpii* Form. in Ver. Brünn 1896 p. 69; Haussk. symb. p. 45; non Benth. — Huc probabiliter: *N. Spruneri* Form. in Ver. Brünn 1896 p. 69, 1897 p. 51, vix Bois. — Exsicc.: Heldr. herb. norm. n. 66, it. thessal. IV. p. 1885; Orph. fl. gr. n. 433; Dörfl. fl. gr. n. 198.

Caulibus erectis vel adscendentibus, pilis patulo-crispulis, viscidulis villosulis, simplicibus vel ramosis; foliis villoso-sublanatis, viscidulis, virentibus, breviter petiolatis, ovato-oblongis, truncato-subcordatis, crenatis; verticillastris in spicam oblongam condensatis, inferioribus saepius distantibus; bracteis lanceolatis. subulato-aristatis, anguste scarioso-marginatis, calycem subaequantibus; calycis breviter patuleque hirti dentibus subaequalibus, aristatis, tubo aequilongis; corollae albae, pubescentis tubo calycem non excedente. — *N. Sibthorpii* maxime affinis, ab ea indumento patulo, viscidulo, foliis virentibus, bracteis anguste marginatis discedit.

β. **Orphanidea** Bois. diagn. ser. 2 IV. p. 22 pro sp.; Bois. fl. or. IV. p. 654. — *N. laconica* Orph. fl. gr. n. 84. — Indumentum subbrevius, vix viscidulum, folia canescentia.

In rupestribus regionis mediae et superioris. Thessalia: mt. Baba pr. Klinovo in Pindo tymphaeo (Heldr.), pr. Miluna, Patsios, Godaman et Diodendron in Olympo, pr. Longici in mt. Othrys (Form.), mt. Ossa et in mt. Parnassus (Heldr.); — *β.* mt. Malevo supra Kastanitza (Orph.). — Jun. Aug. ♃

8. **N. Spruneri** Bois. diagn. ser. 2 IV. p. 23, Fl. or. IV. p. 654; Form. in D. bot. Monat. 1890 p. 29, in Ver. Brünn 1895 p. 36; Hal. Beitr. fl. Epir. p. 36, in ö. b. Z. 1897 p. 325; Haussk. symb.

p. 45. — *N. tymphrestea* Heldr. et Sart. in Heldr. herb. norm. n. 663. — *N. italica* v. *Spruneri* Bald riv. coll. bot. alb. 1895 p. 63. — Exsicc.: Heldr. l. c.; Bald. it. alb. epir. III. n. 80; Sint. it. thessal. n. 683.

Caulibus erectis vel adscendentibns, adpressissime cinereo-velutinis, stricte ramosis; foliis adpresse tomentellis, virentibus, breviter petiolatis, ovato-oblongis, cordatis, crenatis; verticillastris in spicam oblongam condensatis, inferioribus distantibus; bracteis lineari-lanceolatis, breviter aristatis, angustissime marginatis; calycis adpresse tomentelli dentibus subaequalibus, breviter aristatis, tubo subaequilongis; corollae albae, pubescentis tubo calycem vix excedente. — Differt a *N. Sibthorpii*, cui indumento affinis, verticillastris magis dissitis et praesertim bracteis lineari-lanceolatis, angustissime marginatis; a *N. parnassica* indumento adpresso, non viscidulo, calyce minus profunde fisso.

In saxosis subalpinis. Epirus: mt. Micikeli (Form.), pr. Dodona (Bald.), pr. Vulgarelion in mt. Tsumerka, pr. Kalarrytes in mt. Peristeri (Hal.); Thessalia: pr. Chaliki, Lepeniza (Form.), mt. Oeta (Heldr.); Aetolia: mt. Tymphrestus (Samar.), Korax (Tunt.). — Jun. Aug. ♃

.. Calycis dentes tubo subduplo breviores.

9. **N. dirphya** Heldr. herb. norm. n. 777. — *N. Sibthorpii* β. *dirphya* Bois. fl. or. IV. p. 652.

Caulibus erectis vel adscendentibus, tomentoso-lanatis, simplicibus vel ramulosis; foliis adpresse cano-tomentosis, breviter petiolatis, ovato-oblongis, cordatis, crenatis; verticillastris in spicam oblongam condensatis, inferioribus saepe distantibus; bracteis lineari-subulatis, angustissime marginatis, calycis tubum aequantibus vel eo brevioribus; calycis tomentelli dentibus subaequalibus, breviter aristatis, tubum dimidium aequantibus; corollae albae, purpureo-punctatae, pubescentis tubo calycem non excedente. — A *N. Sibthorpii* caulibus tomentoso-lanatis, nec adpressissime cinero-velutinis, a *N. parnassica* indumento breviore, vix viscidulo, foliis manifeste canis, ab ambabus bracteis brevioribus subulatis, angustissime marginatis et calycis dentibus tubo subduplo brevioribus, nec eum aequantibus, a *N. Spruneri* indumento tomentoso-lanato, foliis canis, verticillastris approximatis et calyce minus profunde fisso distinguitur.

In rupestribus cacuminis mt. Dirphys Euboeae (Heldr.). — Jul. Aug. ♃

10. **N. camphorata** Bois. et Heldr. diagn. VII. p. 49, Fl. or. IV. p. 653. — Exsicc.: Heldr. herb. a. 1876; Pichl. pl. gr. a. 1876.

Caulibus erectis vel adscendentibus, pilis patulo-crispulis, viscidulis villosulis, inferne ramosis; foliis villoso-sublanatis, viscidulis, virentibus, breviter petiolatis, ovatis, cordatis, crenatis; verticillastris remotis, longe racemosis; bracteis subulatis, calyce duplo brevioribus; calycis patule hirti dentibus subaequalibus, aristatis, tubum dimidium aequantibus; corollae albae, purpureo-punctatae, pubescentis tubo calycem vix exce-

dente. — Ab omnibus antecedentibus gracilitate, foliis parvis, verticillastris omnibus remotis, parvis, avellanae minoribus distincta.

In alveo torrentis Pentaplon in regione abietina mt. Taygetos (Heldr.). — Jun. Jul. ♃

11. **N. Heldreichii.** — *N. camphorata* Heldr. herb. norm. n. 1470, non Bois. et Heldr. l. c.

Caulibus erectis vel adscendentibus, adpressissime cinereo-velutinis, inferne ramosis; foliis minutis, adpresse canis, breviter petiolatis, ovatis, cordatis, crenulatis; verticillastris remotis, longe racemosis; bracteis subulatis, calyce 2—3 plo brevioribus; calycis viridis, tenuissime adpresse tomentelli dentibus subaequalibus, aristatis, tubum dimidium aequantibus; corollae albae, purpureo-punctatae, pubescentis tubo calycem vix excedente. — Stirps gracilis, foliis circiter 5—10 mm. longis, adultis interdum majoribus. A praecedente, cui habitu maxime affinis, indumento adpressissimo caulis, non viscidulo, foliis minoribus, canis, calycibus viridibus, sub lente tantum tenuissime tomentellis, glandulis sessilibus pellucidis punctatis, statim diagnoscitur, id est ab ea fere eodem modo differt, uti *N. Spruneri* a *N. parnassica*.

In regione abietina mt. Taygetos Laconiae (Zahn). — Jun. Jul. ♃

6. Tribus. SATUREIEAE Benth. lab. p. 324.

21. Melissa L. gen. n. 728.

1. **M. officinalis** L. sp. p. 592; S. et S. pr. I. p. 423; Pieri corc. fl. p. 80; Dallap. prosp. p. 84; Ch. et B. exp. p. 173, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 275; Heldr. Nutzpfl. p. 33, Fl. cephal. p. 58; Bois. fl. or. IV. p. 584; Form. in D. bot. Mon. 1891 p. 29, in Ver. Brünn 1895 p. 36, 1896 p. 68, 1897 p. 51; Hal. Beitr. fl. Epir. p. 36; Haussk. symb. p. 45. — *M. cordifolia* Pers. syn. II. p. 132. — *M. altissima* S. et S. pr. I. p. 423, Fl. gr. VI. p. 62 t. 579; Sieb. avis p. 4, rem. p. 5; Ch. et B. l. c.; Fraas fl. class. p. 182; Heldr. Nutzpfl. p. 33; Raul. cret. p. 831. — Exsicc.: Orph. fl. gr. n. 534; Rev. pl. cret. n. 133; Sint. et Bornm. it. turc. n. 1427; Sint. it. thessal. n. 1034.

Hirsuta vel villosa; caulibus elatis, ramosis; foliis ovatis, serrato-dentatis, basi saepe cordatis, floralibus sensim diminutis; cymis axillaribus subsessilibus, 3—5 floris; calyce campanulato, bilabiato, labio superiore breviter tridentato, inferiore bipartito; corolla alba, calyce subduplo longiore. — Variat indumento, magnitudine foliorum, forma eorum baseos nunc truncato, nunc cordato, longitudine corollarum.

In dumosis regionis inferioris et montanae passim per totam Graeciam. — Jun. Aug. ♃.

22. Calamintha Tourn. inst. t. 92.

1. Sectio. *Clinopodium* L. gen. n. 725. — Flores in verticillastra globosa congesti, bracteis setaceis numerosis involucrati; calycis tubus non gibbus.

1. **C. clinopodium** Benth. in DC. pr. XII. p. 233; Bois. fl. or. IV. p. 579; Heldr. fl. cephal. p. 58; Form. in Ver. Brünn 1896 p. 68, 1897 p. 51, in D. bot. Mon. 1898 p. 79. — *Clinopodium vulgare* L. sp. p. 587; S. et S. pr. I. p. 416; Ch. et B. exp. p. 171, Fl. pelop. p. 38; Raul. cret. p. 831; Hal. Beitr. fl. Epir. p. 36, Beitr. fl. Thessal. p. 17; Form. in Ver. Brünn 1895 p. 36. — *C. plumosum* Sieb. in Flora 1822 p. 242. — *C. Plumieri* Fraas fl. class. p. 181. Icon: Fl. dan. t. 930. — Exsicc.: Heldr. it. thessal. n. 60.

Patule pubescens vel villosa; caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis breviter petiolatis, ovatis vel ovato-oblongis, subcrenatis; verticillastris multifloris, distantibus, summo capitulum terminale formante; bracteis calyces aequantibus, longe ciliatis; calycis breviter pedicellati, incurvi, breviter hirti ciliisque longis obsiti, fauce nudi dentibus lanceolatis, labii inferioris multo profundius fissis; corolla purpurea, rarissime alba, calyce duplo longiore.

In silvaticis dumosisque regionis montanae. Frequens in Epiro et Thessalia, in montibus Atticae et Peloponnesi; Euboea: mt. Telethrion (Heldr.); Creta (Sibth.): pr. Enneachoria (Raul.); Cephalonia: pr. Spartia, Omala, mt. Gerolaki, Akoli (Heldr.); Corcyra (Form.). — Jun. Aug. ♃

2. Sectio. *Eucalamintha* Gr. et Godr. fl. fr. II. p. 662. — Flores in cymas 3—multifloras dispositi, ad apicem caulis vel ramorum racemum plus minus laxum formantes; bracteae parvae; calycis tubus vix gibbus.

a. Corolla ampla, ad 3 cm. longa.

2. **C. grandiflora** L. sp. p. 592; S. et S. pr. I. p. 423; Ch. et B. exp. p. 173, Fl. pelop. p. 39; (*Melissa*); Moench meth. p. 408; Bois fl. or. IV. p. 576; Hal. Beitr. fl. Thessal. p. 17; Form. in Ver. Brünn 1896 p. 69, 1897 p. 51; Haussk. symb. p. 46. — Icon: Rchb. t. 75. — Exsicc.: Orph. fl. gr. n. 293; Heldr. herb. norm. n. 21.

Pubescens vel glabrescens; rhizomate repente; caulibus erectis vel adscendentibus, simplicibus vel parce ramosis; foliis petiolatis, ovatis vel oblongis, grosse serratis, basi rotundatis vel cuneatis; cymis pedunculatis, 3—7 floris, laxis; bracteis lanceolatis; calycis sparsim hirti, fauce subnudi dentibus lanceolatis, superioribus adscendentibus, inferioribus sublongioribus, porrectis; corolla rosea, calyce 2—3 plo longiore; nuculis atris. — Foliis magnis, laete viridibus, profunde serratis et corolla ampla eximia.

In silvis regionis subalpinae. Thessalia: mt. Oxya pr. Chaliki (Hal.), mt. Zygos, Karava, Ghavellu (Haussk.), pr. Mandra Kalivia et Munari in mt. Agrapha, pr. Hepdominta Aderfia in mt. Oxya, mt. Mitrica et pr. Phlambures in mt. Chassia (Form.), mt. Olympus (Orph.).

mt. Oeta (Heldr.); Euboea: mt. Dirphys (Leon.); mt. Parnassus (Guicc.); Achaia: mt. Kyllene (Orph.); Laconia (Sibth.); mt. Taygetos (Heldr.). — Jun. Aug. ♃

b. Corolla parva, ad 15 mm. longa.

α. Folia dentato-serrata.

× Calycis dentes valde inaequales, inferiores multo longiores tubo aequilongi.

3. **C. officinalis** Moench meth. p. 409; Bald. riv. coll. bot. alb. 1896 p. 85; Haussk. symb. p. 46; Form. in Ver. Brünn 1897 p. 51. — *Melissa calamintha* L. sp. p. 593; Friedr. Reise p. 264. — *Thymus calamintha* Sm. fl. brit. II. p. 641. — Icon: Rchb. t. 75. — Exsicc.: Sint. it. thessal. n. 1215.

Rhizomate subrepente; caulibus erectis vel adscendentibus, simplicibus vel ramosis, hirtis; foliis majusculis, petiolatis, ovatis, dentato-serratis, utrinque pubescentibus; cymis axillaribus, paucifloris, longiuscule pedunculatis, inferioribus folio florali aequilongis; pedicellis calyce aequilongis vel longioribus; bracteis parvis, subulatis; calycis hirti, fauce subnudi dentibus superioribus lanceolatis, recurvis, inferioribus multo longioribus, lineari-subulatis, rectis, tubo aequilongis; corolla rosea, calyce 2—3 plo longiore; nuculis subglobosis.

In dumosis montanis, rare. Epirus: mt. Micikeli (Bald.); Thessalia: mt. Baba, Ghavellu (Haussk.), pr. Sermeniko et Bisula in mt. Agrapha, pr. Phlambures et mt. Mitrica in mt. Chassia (Form.); Achaia: pr. Patras (Friedr.). — Jul. Aug. ♃

×× Calycis dentes minus inaequales, inferiores parum longiores, tubo breviores.

4. **C. thessala** Haussk. symb. p. 46. — Exsicc.: Sint. it. thessal. n. 1032.

Rhizomate subrepente; caulibus erectis vel adscendentibus, simplicibus vel ramosis, sparsim pilosis; foliis majusculis, petiolatis, ovatis, dentato-serratis, laete viridibus, supra glabris, subtus ad nervos parce pilosis; cymis axillaribus, paucifloris, longiuscule pedunculatis, inferioribus folio florali aequilongis; pedicellis calyce brevioribus; bracteis parvis, subulatis; calycis parce hirtuli, fauce subnudi dentibus lanceolato-subulatis, superioribus subrecurvis, inferioribus longioribus, rectis, tubo brevioribus; corolla rosea, calyce duplo longiore; nuculis ovoideis. — Differt a praecedente indumento parcissimo, foliis laete viridibus et calycis dentibus minus inaequalibus.

In rupestribus ad Kastreiki (Sint.) et ad monasterium Meteora pr. Kalabaka Thessaliae (Haussk.). — Jul. Aug. ♃

5. **C. nepetoides** Jord. obs. IV. p. 16 t. 2. — *C. officinalis* Hal. in ö. b. Z. 1890 p. 40, non Moench. — Exsicc.: Orph. fl. gr. n. 531 (*f. cana*); Sint. it. or. a. 1889 n. 1878 (*f. virescens*).

Rhizomate repente; caulibus erectis vel adscendentibus, simplicibus vel ramosis, molliter hirtis; foliis mediocribus, petiolatis, ovatis, dentatis,

pubescenti-canescentibus; cymis axillaribus, saepius plurifloris, longe pedunculatis, folia floralia superantibus; pedicellis calyce aequilongis; bracteis minutis; calycis hirti, fauce villis exsertis vestiti dentibus superioribus lanceolatis, subrecurvis, inferioribus parum longioribus, lineari-subulatis, rectis, tubo brevioribus; corolla rosea, calyce duplo longiore; nuculis ovoideis. — A praecedentibus cymis longe pedunculatis, folio florali multo longioribus discedit.

In dumosis, silvaticis. Thessalia: pr. Litochoron (Orph.) et Hagios Dionysios (Sint.) in Olympo; Eurytania: mt. Chelidoni (Heldr.). — Jul. Aug. ♃.

β. Folia crenata.

6. **C. nepeta** L. sp. p. 593; Ch. et B. exp. p. 173, Fl. pelop. p. 39; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 264; (*Melissa*); Savi fl. pis. II. p. 63; Raul. cret. p. 830; Bois. fl. or. IV. p. 577; Heldr. fl. cephal. p. 58; Form. in Ver. Brünn 1895 p. 36, 1896 p. 69, 1897 p. 51; Haussk. symb. p. 46. — *Thymus nepeta* Sm. fl. brit. II. p. 642; S. et S. pr. I. p. 422. — *Melissa calamintha* Dall. prosp. p. 85, sec. Heldr. l. c., non L. — *Thymus calamintha* Fraas fl. class. p. 177, non Sm. — *C. Spruneri* Bois. diagn. XII. p. 53. — *C. incana* Clem. sert. p. 75 p. p., non Bois. et Heldr. — *C. adscendens* Form. in D. bot. Mon. 1891 p. 29, non Jord. — Icon: Rchb. t. 76. — Exsicc.: Heldr. herb. norm. n. 222 et 872, herb. fl. hellen. n. 113, in Magn. fl. sel. 3830 et 3831; Sint. it. thessal. n. 1252.

Molliter pubescenti-canescens; rhizomate obliquo, lignoso; caulibus adscendentibus, simplicibus vel ramosis; foliis parvis, petiolatis, ovato-rhombeis, crenatis; cymis axillaribus, paucifloris, breviter pedunculatis, folio florali longioribus; pedicellis calyce subbrevioribus; bracteis minutis; calycis pubescentis, fauce villis exsertis vestiti dentibus superioribus lanceolatis, subrecurvis, inferioribus parum longioribus, lineari-subulatis, rectis, tubo brevioribus; corolla rosea, calyce subduplo longiore; nuculis ovoideis. — A praecedente rhizomate lignoso, foliis minoribus, crenatis, cymis paucifloris, breviter pedunculatis, pedicellis brevioribus et floribus minoribus discedit.

In saxosis, locis sterilibus regionis calidae, passim per totam Graeciam. — Jul. Aug. ♃.

γ. Folia parva, subintegerrima.

7. **C. incana** S. et S. pr. I. p. 421, Fl. gr. VI. p. 62 t. 577; Ch. et B. fl. pelop. p. 38; Fraas fl. class. p. 178; (*Thymus*); Bois. et Heldr. in DC. pr. XII. p. 226, diagn. XII. p. 52, Fl. or. IV. p. 578; Clem. sert. p. 75 p. p.; Haussk. symb. p. 46; Heldr. fl. Aegina p. 225. — *Thymus lanuginosus* Ch. et B. exp. p. 172, non fl. gr. n. 258; Heldr. herb. norm. 419 et n. n. 7274, it. thessal. n. 61.

ana; rhizomate obliquo, lign so; caulibu
parvis, petiolatis, orbiculat s, subinteg

rimis; cymis axillaribus, 2—6 floris, brevissime pedunculatis, folio florali longioribus; pedicellis calyce longioribus; bracteis minimis; calycis fauce villosi labio superiore lato, brevissime tridentato, inferioris aequilongi dentibus lanceolatis; corolla rosea, calyce duplo longiore; nuculis ovato-globosis. — Transitum ad sectionem sequentem praebet.

β. **calvescens** Heldr. herb. norm. n. 873 b. — Tomento praesertim in parte superiori plantae rarescenti; corolla majore, calyce triplo longiore.

In campis incultis, aggeribus regionis inferioris et montanae. Attica: ubique pr. Athenas, pr. Stadion in colle Turcovuni, mt. Hymettus, insula Aegina (Heldr.); Argolis: pr. Troezene, insula Tenos Cycladum (Chaub.). — Sept. Nov. ♃

Obs. Secundum Bois. fl. or. IV. p. 578 circa Athenas formae inter *C. nepetam* et *C. incanam* hybridae (*C. Boissieri* Briqu. lab. alp. marit. p. 432 sub *Satureia*) interdum occurrunt.

8. **C. cretica** L. sp. p. 593 (*Melissa*); Benth. in DC. pr. XII. p. 227; Raul. cret. p. 830; Bois. fl. or. IV. p. 578; Bald. viagg. Creta p 84. — *Thymus hirtus* Sieb. avis p. 4, rem. p. 5, non Willd. nec alior. — Huc probabiliter: *Micromeria marifolia* Raul. cret. p. 830, non Benth. — Exsicc.: Rev. pl. cret. n. 131; Bald. it. cret. n. 86, it. cret. alt. n. 347.

Tota dense cano-villosa; rhizomate obliquo, lignoso; caulibus decumbentibus, ramosis; foliis parvis, petiolatis, ovatis, subintegris; cymis axillaribus, 2—6 floris, breviter pedunculatis, folio florali sublongioribus; pedicellis calyce subbrevioribus; bracteis minimis; calycis fauce villosi dentibus labii superioris lanceolatis, inferioribus parum longioribus lanceolato-subulatis, tubo brevioribus; corolla albida, calyce duplo longiore; nuculis globosis. — Indumento e pilis patulis flexuosis constante egregia.

In lapidosis regionis subalpinae. Cretae: pr. Askyphos, Omalos, mt. Volakia (Raul.), ad Hellinoseli (Bald.), mt. Lakus (Rev.). — Jun. Aug. ♃

3. Sectio. *Acinos* Moench. meth. p. 407. — Flores axillares, singillatim pedicellati; bracteae minutae vel nullae, calycis tubus supra basin gibbus.

a. Perennes vel perennantes.

α. Dentes calycis labii superioris eis inferioris duplo latiores, fructiferi patuli.

9. **C. alpina** L. sp. p. 591; S. et S. pr. I. p. 420; Pieri corc. fl. p. 80 (*Thymus*); Lam. fl. fr. II. p. 394; Bois. fl. or. IV. p. 581; Heldr. fl. cephal. p. 58, cblor. Parn. p. 25; Hal. Beitr. fl. Epir. p. 36, Beitr. fl. Thessal. p. 17, Beitr. fl. Achaia p. 30; Bald. riv. coll. bot. alb. 1896 p. 85; Form. in Ver. Brünn 1897 p. 51; Haussk. symb. p. 47. — *Acinos alpinus* Moench. meth. p. 407; Raul. cret.

p. 828. — *Thymus acinos* Ch. et B. exp. p. 172, Fl. pelop. p. 38 p. p. — Icon: Jacq. fl. austr. I. t. 97. — Exsicc.: Orph. fl. gr. n. 532.

Caespitosa; caulibus e basi diffusa, radicanti adscendentibus, reflexo-puberulis, basi lignosis, flexuosis; foliis petiolatis, ovatis vel ellipticis, acutis, a medio dentatis vel subintegerrimis, glabris vel glabriusculis, floralibus calyces aequantibus; floribus 2—5 ad axillas superiores breviter pedunculatis; calycis patule hirti, fauce villosi, basi gibbi labiis tubo duplo brevioribus; corollae violaceae, calyce $2^1/_2$ plo longioris tubo superne valde ampliato.

β. **nebrodensis** Kern. et Strobl. in ö. b. Z. 1897 p. 171, Fl. nebrod. p. 398, pro sp. — *C. alpina v. meridionalis* Haussk. symb. p. 47. — *C. suaveolens* Hal. Beitr. fl. Achaia p. 30, non Bois. — Folia ad nervos vel omnino puberula; calyx breviter vel longe hirsutus. — Intermediis ad typum transit. — Exsicc.: Orph. fl. gr. n. 283.

γ. **granatensis** Bois. et Reut. pug. p. 94 pro sp. — *C. aetnensis* Strobl. in ö. b. Z. 1874 p. 29. — *C. meridionalis* Nym. consp. p. 589. — Minor, folia parva, circa 5 mm. longa, glabriuscula vel puberula; flores fere dimidio minores; calyx pilis subaduncis hirtus. — Intermediis ad typum transit. — Exsicc.: Dörfl. fl. gr. n. 331.

δ. **pentelica** Haussk. symb. p. 48. — Folia magis conferta, utrinque pube brevi denso tomentella, incana, floralia calyces superantia; verticillastra conferta. — Exsicc.: Leon. pl. gr. a. 1897.

ε. **major** Haussk. symb. p. 47, pro var. *C. patavinae.* — Caules firmiores; folia majora, elliptica, acutissima, glabra vel margine ciliata, floralia flores superantia; calyx submajor, longe denseque hirsutus, fauce valde villosus. — Propter calycis labii superioris dentes patentes meo sensu ad formas *C. alpinae*, nec *C. patavinae* pertinet. — Exsicc.: Sint. it. thessal. n. 198.

In lapidosis regionis subalpinae et alpinae. Epirus: mt. Vratedon distr. Zagorion (Bald.), mt. Tsumerka, Peristeri (Hal.); Thessalia: mt. Oxya pr. Chaliki (Hal.), Zygos (Haussk.) et Karava in Pindo, mt. Pirgo et Mavrika in mt. Othrys (Form.), mt. Olympus; mt. Korax Aetoliae, mt. Parnassus (Heldr.); Peloponnesus: pr. Megaspilaeon (Hal.), mt. Chelmos, Kyllene, Malevo (Orph.), Diaforti (Chaub.); Creta: mt. Hellinoseli (Heldr.) in mt. Sphacioticis (Sibth.), mt. Ida (Raul.); Cephalonia: mt. Aenos (Heldr.); Corcyra (Pieri); — *γ.* mt. Parnes (Heldr.) et Pentelicon (Haussk.) Atticae; — *δ.* pr. Klinovo in Pindo tymphaeo)Sint.). — Maio, Aug. ♃

β. Dentes calycis labii superioris eis labii inferioris subsimiles, fructiferi porrecti.

10. **C. suaveolens** S. et S. pr. I. p. 420 (*Thymus*); Bois. fl. or. IV. p. 582; Hal. in z. b. G. 1888 p. 761; Heldr. chlor. Parn. p. 25; Form. in D. bot. Mon. 1891 p. 29, in Ver. Brünn 1896 p. 69, 1897 p. 51; Haussk. symb. p. 47 *v. canescens*; Bald. riv. coll. bot. alb.

1896 p. 36. — Probabiliter: *T. acinos* Ch. et B. exp. p. 172, Fl. pelop. p. 38, p. p. — Exsicc.: Heldr. herb. norm. n. 221; Orph. fl. gr. n. 550.

Perennis, molliter et patule canescens; caulibus e rhizomate indurato adscendentibus; foliis petiolatis, ellipticis vel elliptico-lanceolatis, acutis, parce denticulatis, floralibus calycem parum superantibus; verticillastris 5—9 floris, spicam longam, laxiusculam formantibus; calycis patule hirsuti, fauce villosi, basi gibbi dentibus subaequilongis, a basi sublatiore subulatis, rectis, tubo triplo brevioribus; corollae roseae calyce duplo longioris tubo exserto.

β. **viridis** Heldr. et Haussk. symb. p. 47. — Caulibus brevissime, interdum ad angulos tantum pilosiusculis; foliis glabris, viridibus, elliptico-lanceolatis, acute denticulatis; spica abbreviata; calyce breviter et adpressiuscule hirto. — Exsicc.: Sint. it. thessal. n. 463 et 1033; Bald. it. alb. epir. IV. n. 85.

γ. **meteorica** Haussk. symb. p. 47. — Caulibus glabris vel ad angulos tantum sparse pilosiusculis; foliis glaberrimis, viridibus, anguste lanceolatis, acutissimis, acute denticulatis; spica abbreviata; calyce breviter et adpressiuscule hirto. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

In rupestribus regionis montanae et subalpinae. Thessalia: mt. Zygos, Said Pascha, Karava, Oxya, Chassia, pr. Kalabaka, Trikala, Velestinos (Form.), Tyrnovo, Pharsalus, Aivali (Haussk.), mt. Olympus (Auch.), Ossa, Pelion (Form.), Oeta (Heldr.); mt. Kiona pr. Segditza (Hal.), mt. Parnassus, Pateras (Heldr.); Peloponnesus: mt. Kyllene, Taygetos (Heldr.); — *β.* Epirus: mt. Olycika, Micikeli (Haussk.); Thessalia: pr. Chaliki, Kalabaka (Haussk.); — *γ.* pr. Malakasi, Meteora (Haussk.). — Maio, Jul. ♃

11. **C. patavina** Jacq. obs. IV. p. 7 t. 87 (*Thymus*); Host. fl. austr. II. p. 133; Haussk. symb. p. 46. — *C. suaveolens* Hal. Beitr. fl. Thessal. p. 36, non Bois. — Exsicc.: Sint. it. thessal. n. 678.

Perennans, pubescens; caulibus e basi diffusa, radicanti adscendentibus; foliis petiolatis, ellipticis, acutis, parce denticulatis, floralibus calycem superantibus; verticillastris 3—6 floris, spicam interruptam formantibus; calycis patule hirti, fauce villosi, basi gibbi dentibus subaequilongis, labii superioris lanceolatis, subinclinatis, inferioris subulatis, porrectis, tubo subtriplo brevioribus; corollae roseae, calyce duplo longioris tubo exserto. — An a praecedente specifice diversa; Briquet in lab. alp. marit. p. 448—454 omnes tres species sectionis *Acinos* hucusque enumeratas pro subspecies vel varietates unius et ejusdem speciei (*C. alpinae*) habet.

In rupestribus regionis montanae et subalpinae. Epirus: pr. Kalarrytes ad radices mt. Peristeri (Hal.); Thessalia: pr. Chaliki, mt. Baba, Zygos, Gionskala, Ghavellu, Karava in Pindo, mt. Pelion (Haussk.). — Jun. Aug. ⊙

b. Annuae.

12. **C. acinos** L. sp. p. 591; S. et S pr. I. p. 420; Fraas fl. class. p. 177; (*Thymus*); Clairv. in Gaud. fl. helv. IV. p. 84; Bois. fl. or. IV. p. 583; Form. in Ver. Brünn 1897 p. 51. — Icon: Fl. dan. t. 814.

Pubescens vel villosula; caulibus adscendentibus, simplicibus vel ramosis; foliis petiolatis, ovatis vel ellipticis, integris vel pauciserratis, acutis, floralibus flores aequantibus; verticillastris 2—6 floris; calycis basi gibbi dentibus tubo duplo brevioribus, fructiferis porrectis; corolla violacea, tubo exserto.

Indicatur pr. Lamia Phthiotidis (Form.), pr. Corinthum (Fraas) et in agro Laconico (Sibth.), sed probabiliter ex confusione cum sequente. — Maio, Jul. ♃ N. v.

13. **C. exigua** S. et S. pr. I. p. 421, Fl. gr. VI. p. 61 t. 575 (*Thymus*). — *Thymus graveolens* M. B. fl. taur. cauc. II. p. 60. — *Acinos graveolens* Link. en. hort. berol. II. p. 117; Raul. cret. p. 8 v. *integrifolia*. — *C. graveolens* Benth. in DC. pr. XII. p. 231; Bois. fl. or. IV. p. 583; Heldr. chlor. Parn. p. 25, Fl. Aegina p. 385; Haussk. symb. p. 48; Form. in D. bot. Mon. 1898 p. 79. — Exsicc.: Heldr. herb. norm. n. 619 et 1274; Dörfl. pl. cret. n. 22.

Differt a praecedente foliis elliptico-subrotundis, abrupte acuminatis, floralibus flores superantibus, calycis dentibus dimidio tubo sublongioribus, corollae tubo incluso.

In collibus siccis, arvis derelictis regionis inferioris et montanae. Thessalia: ad monasterium Korona, pr. Kalabaka (Haussk.); Attica: pr. Kephissia ad radices mt. Pentelicon, mt. Hymettus (Heldr.), Kerata (Haussk.); mt. Parnassus pr. Rachova (Heldr.); Laconia: mt. Malevo (Sprun.); insula Aegina (Heldr.); Cyclades (Oliv.); Creta: mt. Aphendi Kavutsi (Raul.), pr. Anatoli distr. Hierapetra (Leon.). — Apr. Jun. ⊙

23. Micromeria Benth. in bot. reg. XV. n. 1282.

a. Flores sessiles.

1. **M. juliana** L. sp. p. 567; S. et S. pr. I. p. 396, Fl. gr. VI. p. 32 t. 540; Sieb. avis p. 4, in Flora 1818 p. 270; Ch. et B. exp. p. 162, Fl. pelop. p. 36; Link in Linnaea IX. p. 574; Fraas fl. class. p. 174: (*Satureia*); Benth. lab. p. 378; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 265; Clem. sert. p. 74; Weiss in z. b. G. 1869 p. 744; Raul. cret. p. 829; Bois. fl. or. IV. p. 569; Heldr. fl. cephal. p. 57, Fl. Aegina p. 385, prosth. chlor. Thera p. 3; Form. in D. bot. Mon. 1891 p. 29, in Ver. Brünn 1895 p. 35, 1896 p. 68, 1897 p. 50; Bald. in nuovo giorn. bot. 1894 p. 99; Haussk. symb. p. 48. — *Hyssopus officinalis* Dallap. prosp. p. 80, sec. Heldr. l. c., non L. — Exsicc.: Orph. fl. gr. n. 530; Heldr. herb. norm. n. 153 et 968, herb. dimorph. n. 43; Rev. pl. cret. n. 260; Sint. it. or. a. 1889 n. 1882, it. thessal. n. 1042; Sint. et Bornm. it. turc. n. 1420.

Pubescens, caulibus e rhizomate suffrutescenti numerosis, erectis vel adscendentibus, simplicibus vel ramosis; foliis sessilibus, obtusiusculis,

inferioribus ovato-oblongis, superioribus linearibus; cymulis breviter pedunculatis, densifloris, racemum interruptum formantibus; floribus singulis sessilibus; bracteis calycem subaequantibus; calycis fauce nudi dentibus subulatis, strictis, tubo subbrevioribus; corollae purpureae tubo incluso; nuculis mucronatis.

In rupestribus regionis inferioris et subalpinae totius Graeciae. — Maio, Jul. ♃

2. **M. myrtifolia** Bois. et Hoh. diagn. V. p. 19. — *M. juliana β. myrtifolia* Bois. fl. or. IV. p. 570; Haussk. symb. p. 48. — *M. Reinholdi* Heldr. sched. sec. Bois. l. c.

Differt a praecedente foliis inferioribus ovatis, breviter petiolatis, superioribus oblongo-lanceolatis, floribus dimidio minoribus et nuculis obtusiusculis. Cf. Murb. Beitr. fl. Südbosn. p. 57.

In rupestribus regionis inferioris, pr. Aivali Thessaliae, pr. Chalkis Euboeae, in peninsula Methana (Haussk.) et in insula Poros Argolidis (Heldr.). — Maio, Jul. ♃ N. v.

b. Flores pedicellati.

α. Nuculae mucronatae.

3. **M. cremnophila** Bois. et Heldr. fl. or. IV. p. 570; Heldr. chlor. Parn. p. 25; Form. in D. bot. Mon. 1891 p. 29, in Ver. Brünn 1896 p. 68, 1897 p. 50 *v. thessala*, quae autem nullo modo a typo discedit; Hal. Beitr. fl. Epir. p. 36, Beitr. fl. Thessal. p. 17; Haussk. symb. p. 48. — Exsicc.: Sint. it. thessal. n. 1040.

Minutissime pubescens; caulibus e rhizomate suffrutescenti numerosis, tenuibus, adscendentibus, simplicibus vel parce ramosis; foliis subsessilibus, obtusiusculis, inferioribus ellipticis, superioribus linearibus; cymulis longiuscule pedunculatis, 2—5 floris, laxis, racemum interruptum formantibus; floribus singulis pedicellatis; bracteis pedicello sublongioribus; calycis fauce sparsim villosi dentibus lanceolatis, rectis, tubo quadruplo brevioribus; corollae purpureae tubo incluso. — Colore laete viridi, caulibus filiformibus, cymulis longiuscule pedunculatis paucifloris, laxis, calycis dentibus brevibus et nuculis mucronatis insignis.

In rupestribus regionis inferioris et subalpinae. Epirus: ad radices mt. Peristeri pr. Kalarrytes (Hal.); Thessalia: mt. Baba, pr. Klinovo, Krania, Kotura, Chaliki, Malakasi, Meteora, Kalabaka (Haussk.), Kastania, Kastri, Vlachava (Form.) in Pindo; mt. Oeta Phthiotidis, mt. Parnassus pr. Gurna (Heldr.). — Jul. Aug. ♃

1 × 3. **M. juliana × cremnophila.** — *M. meteorica* Haussk. symb. p. 48. — Exsicc.: Heldr. it. thessal. IV. a. 1885. — Differt a *M. juliana* caulibus tenuioribus, cymulis paucifloris, floribus breviter pedicellatis, bracteis brevioribus; a *M. cremnophila* caulibus firmioribus, foliis latioribus, cymulis densiusculis, floribus brevius pedicellatis, calycis dentibus tubo subbrevioribus et indumento copiosiori. — Inter parentes pr. Kalabaka (Heldr.).

β. Nuculae obtusiusculae.

× Planta plus minus hirsuta, superne tantum interdum hispida.

○ Caules firmi, virgati.

4. **M. nervosa** Desf. fl. atl. II. p. 9 t. 121; S. et S. pr. I. p. 396; Sieb. in Flora I. p. 270; Urv. enum. p. 65; (*Satureia*); Benth. lab. p. 376; Marg. et R. fl. Zante p. 74; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 829; Bois. fl. or. IV. p. 569; Heldr. fl. cephal. p. 57, prosth. chlor. Thera p. 3; Boissieu in bull. soc. bot. Fr. 1896 p. 287; Hal. in ö. b. Z. 1897 p. 98, 1899 p. 25; Haussk. symb. p. 48. — *S. filiformis* Sieb. avis p. 4, rem. p. 5; Link. in Linn. IX. p. 574; non (Ait.) Benth. — *Thymus mastichina* S. et S. pr. I. p. 422; Ch. et B. fl. pelop. p. 38, non L. — *S. graeca* Ch. et B. fl. pelop. p. 36 p. p., non Benth. — *S. plumosa* Hampe in Flora 1842 p. 127. — *M. plumosa* Heldr. herb. norm. n. 259, Fl. Aegina p. 385. — Exsicc.: Heldr. l. c.; Orph. fl. gr. n. 1040; Rev. pl. cret. n. 130; Dörfl. pl. cret. n. 99.

Caulibus e basi suffrutescenti numerosis, pubescentibus, erectis vel adscendentibus, simplicibus vel parce ramosis; foliis brevissime petiolatis, ovatis, acutis, pubescentibus, cymulis breviter pedunculatis, 2—10 floris, laxiusculis, racemum inferne interruptum, superne densum formantibus; floribus singulis pedicellatis; bracteis pedicello longioribus; calycis patule longeque villosi, fauce hispiduli dentibus setaceis, erecto-patulis, tubo subbrevioribus; corollae purpureae tubo incluso. — Caules usque 50 cm. alti; cymulis valde villosis insignis.

In collibus apricis, saxosis regionis inferioris et submontanae. Boeotia (Sprun.); Attica: pr. Athenas, mt. Hymettus (Sibth.), Phaleron, insula Aegina (Heldr.), Laurion (Haussk.); Achaia: pr. Vostizza (Chaub.); Argolis: pr. Nauplia (Link), insula Hydra (Heldr.); Sporadum insula Jura (Leonis); Cycladum insula Keos (Boissieu), Syra, Tenos (Weiss), Melos (Urv.), Thera (Heldr.); Creta: pr. Kissamos (Rev.), Canea, Akroteri, Rethymo (Raul.), in fauce Serisso (Sieb.), pr. Males distr. Hierapetra (Leon.); Strophadum insula major (Reiser); Zante (Sibth.); Cephalonia: pr. Argostoli (Heldr.). — Mart. Jun. ♃

5. **M. graeca** L. sp. p. 567; S. et S. pr. I. p. 397, Fl. gr. VI. p. 34 t. 542; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 162, Fl. pelop. p. 36 p. p.; Link in Linnaea IX. p. 574; (*Satureia*); Benth. lab. p. 373; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 270 et 273; Clem. sert. p. 74; Raul. cret. p. 829; Bois. fl. or. IV. p. 571; Heldr. fl. cephal. p. 58, Fl. Aegina p. 385, chlor. Thera p. 19; Gelmi in bull. soc. bot. ital. 1889 p. 451; Haussk. symb. p. 48. — Exsicc.: Heldr. herb. norm. n. 152.

Caulibus e rhizomate suffrutescenti numerosis, foliisque adpresse pubescentibus, erectis vel adscendentibus, simplicibus vel parce ramosis; foliis sessilibus, inferioribus ovatis vel ovato-oblongis, acutis, superioribus lanceolatis vel linearibus; cymulis pedunculatis, 2—7 floris, laxiusculis, racemum interruptum formantibus; floribus singulis pedicellatis; bracteis

pedicello longioribus; calycis adpresse pubescentis, fauce villosi dentibus setaceis, erecto-patulis, tubo subduplo brevioribus; corollae purpureae tubo breviter exserto. — Differt a praecedente foliis sessilibus, superioribus lineari-lanceolatis, calyce adpresse pubescente, nec patule villoso, dentibus brevioribus.

β. **villicaulis** Borb. ap. Form. in Ver. Brünn 1895 p. 35. — Huc forsan: *S. canescens* Link in Linn. IX. p. 574, vix Guss., = *M. canescens* Marg. et R. fl. Zante p. 74; Raul. cret. p. 829; vix Benth. — Caulibus, foliis calycibusque patentim pilosis; cymulis 1—4 floris. — Exsicc.: Form. pl. corc. a. 1894.

In collibus apricis, saxosis regionis inferioris et submontanae, Attica: mt. Parnes, Pentelicon (Heldr.), insula Aegina (Friedr.); Euboea (Sprun.); Peloponnesus (Sibth.): peninsula Methana (Haussk.), insula Poros (Friedr.) Argolidis, pr. Methone, Corone, Navarin, Pylos usque Messene, pr. Androvista, in Maina (Chaub.); Cyclades (Bois.): insula Thera (Heldr.); Creta (Sibth.); Zante (Marg.); Cephalonia; pr. Argostoli (Heldr.); Corcyra (Clem.); — *β.* Corcyra: pr. Manducchio (Spreitz.), Analypsis, Kanali (Form.); Zante (Link); Creta: pr. Canea (Raul.), si *M. canescens* aut. supra cit. huc spectet. — Apr. Jul. ♃

○○ Caules filiformes, saepius decumbentes.

6. **M. microphylla** Urv. enum. p. 71 (*Thymus*); Benth. lab. p. 377; Raul. cret. p. 829; Bois. fl. or. IV. p. 572. — *M. sphaciotica* Bois. et Heldr. diagn. XII. p. 48.

Pubescens; caulibus e rhizomate suffrutescenti numerosis, tenuibus, simplicibus vel parce ramosis, decumbentibus vel adscendentibus; foliis parvis, subsessilibus, ovatis vel oblongis, acutiusculis; cymulis pedunculatis, 1—3 floris, racemum laxum formantibus; floribus singulis pedicellatis; bracteis pedicello saepius brevioribus; calycis ovato-cylindrici, patule hispiduli, fauce villosi dentibus lanceolatis, erecto-patulis, tubo 2—3 plo brevioribus; corollae purpureae tubo brevissime exserto. — Differt ab omnibus foliis minutis, 5 mm. longis, 2—3 mm. latis et calycibus maximum 3 mm. longis, vix 1 mm. latis.

In rupestribus regionis inferioris. Cretae: ad Hagios Rumeli et Aradhena (Raul.). — Apr. Jul. ♃ N. v.

×× Planta tota dense patentim hispida, grisea.

7. **M. hispida** Bois. et Heldr. diagn. XII. p. 48, Fl. or. IV. p. 572; Raul. cret. p. 829. — Exsicc.: Heldr. herb. n. 1422.

Caulibus e rhizomate suffrutescenti numerosis, diffusis, simplicibus vel parce ramosis; foliis parvis, subsessilibus, ovatis vel ellipticis, acutis; cymulis longiuscule pedunculatis, 1—3 floris, racemum laxum formantibus; floribus singulis pedicellatis; bracteis pedicello brevioribus; calycis cylindrici, hispidi, fauce hirti dentibus lanceolato-subulatis, erecto-patulis tubum dimidium aequantibus; corollae purpureae tubo subincluso. — Folia et inflorescentia pauciflora praecedentis, a qua abunde differt indumento totius plantae multo copiosiori, calyce cylindrico, 4 mm. longo.

In rupestribus submontosis Cretae: pr. Aradhena distr. Sphakia (Raul.), pr. Kaenuriokorio distr. Mirabello (Heldr.). — Apr. Maio. ♃

24. Satureia L. gen. n. 707.

a. Verticillastra dense globosa, multiflora, remota, bracteis longis suffulta.

1. **S. thymbra** L. sp. p. 567; S. et S. pr. I. p. 396, Fl. gr. VI. p. 33 t. 541; Pieri corc. fl. p. 75; Sieb. avis rem. p. 5, in Flora I. p. 270; Urv. enum. p. 65; Link in Linnaea IX. p. 574; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 277 et 286; Fraas fl. class. p. 174; Clem. sert. p. 75; Heldr. Nutzpfl. p. 33, chlor. Thera p. 19; Ung. Reise p. 126; Raul. cret. p. 829; Weiss in z. b. G. 1869 p. 743; Bois. fl. or. IV. p. 567; Bald. viagg. Creta p. 84; Haussk. symb. p. 49; Form. in D. bot. Mon. 1898 p. 79. — *Thymus tragoriganum* L. mant. p. 84; S. et S. pr. I. p. 421; Sieb. avis p. 4, in Flora I. p. 270; Ch. et B. exp. p. 172, Fl. pelop. p. 38. — *Thymbra hirsuta* Pers. syn. II. p. 114. — Exsicc.: Orph. fl. gr. n. 395; Heldr. herb. norm. n. 175; Rev. pl. cret. n. 129.

Suffruticosa, scabrido-canescens; caulibus erectis, ramosis; foliis obovato-cuneatis, acutis, subsessilibus; verticillastris multifloris, globosis, valde distantibus, bracteis numerosis, oblongis, acuminatis suffultis; calycis patule albo-hirsuti dentibus subaequalibus, lanceolato-subulatis, tubo subbrevioribus; corollae roseae tubo incluso.

In collibus siccis regionis inferioris et submontanae. Boeotia (Sibth.), pr. Klephto Vuno (Ung.); Attica: pr. Athenas, mt. Parnes, Pentelicon, Hymettus (Orph.), insula Salamis, Lero Pharmacusarum (Heldr.), pr. Laurion (Haussk.); Euboea: mt. Dirphys (Heldr.); Argolis: pr. Nauplia (Haussk.), insula Poros (Friedr.); Laconia: mt. Taygetos (Zahn); pr. Messene (Chaub.); Cyclades (Sibth.): insula Syra (Weiss), Keos, Thera (Heldr.); Creta: pr. Kissamos (Rev.), Canea (Weiss), Akroteri, Suda, Rethymo (Raul.), in fauce Serisso (Sieb.); Zante: pr. Lagana, mt. Scopo (Marg.); Corcyra (Pieri). — Maio, Jul. ♃

b. Verticillastra pauciflora, saepius approximata.

α. Rami inermes.

2. **S. cuneifolia** Ten. fl. nap. pr. p. 33, Fl. nap. p. 3 t. 155; Bois. fl. or. IV. p. 564; Heldr. fl. cephal. p. 57; Form. in Ver. Brünn 1895 p. 35. — *S. fasciculata* Tausch in syll. soc. Ratisb. II. p. 248, non Raf. — *S. virgata* Vis. in Benth. lab. p. 353, Fl. dalm. t. 18. — Exsicc.: Heldr. fl. cephal. a. 1867.

Suffruticosa, grisea; caulibus erectis vel adscendentibus, virgatis, pubescentibus, ramosis, in spicas longas, tenues, interruptas abeuntibus; foliis sessilibus, lineari-lanceolatis, acutis vel inferioribus oblongo-spathulatis, hirtis, punctatis, floralibus saepius abbreviatis; cymulis subtrifloris, subsessilibus; calycis scabrido-hirsuti dentibus aequalibus, lanceolatis, subaequilongis; corollae albo-roseae tubo subincluso. —

A *S. montana* differt imprimis inflorescentia; calycis dentes etiam parum breviores et flores minores.

In saxosis montosis. Epirus: mt. Micikeli (Form.); Cephalonia: mt. Aenos, Rhudi et Phalaris (Heldr.). — Aug. Sept. ♃

3. **S. montana** L. sp. p. 568; Bald. riv. coll. bot. alb. 1895 p. 61, 1896 p. 85. — Icon: Rchb. t. 72. — Exsicc.: N. v.

Suffruticosa, viridis; caulibus erectis vel adscendentibus, in racemos spicaeformes densiusculos abeuntibus; foliis sessilibus, lineari-lanceolatis, acutis, pellucide-punctatis, glabris vel ciliatis, floralibus flores aequantibus vel superantibus; cymulis 3—5 floris, pedunculatis; calycis hirtuli dentibus subaequalibus, lanceolatis, acutis, tubo subaequilongis; corollae albae tubo subincluso.

β. **pisidica** Wettst. in Sitzungsb. acad. Wiss. Wien 1889 p. 374; Hal. in ö. b. Z. 1890 p. 40; pro sp. — *S. olympica* Hal. l. c. — *S. cuneifolia β. Wiedemanniana* Bois. fl. or. IV. p. 564, saltem pro p. — Griseo-viridis; caulibus et foliis magis hirtis; cymulis sessilibus subsessilibusve. — Exsicc.: Sint. it. or. a. 1889 n. 1876; Sint. et Bornm. it. turc. n. 1435.

γ. **hellenica** Heldr. ap. Hal. in z. b. G. 1899 p. 192 pro sp. — *S. parnassica* Bois. fl. or. IV. p. 563, quoad pl. euboeam. — Caulibus abbreviatis, caespitosis; foliis densius hirtis; racemis brevissimis; cymulis sessilibus; calycis dentibus minus acutis, tubo brevioribus; corolla albido-rosea. — Foliis anguste lineari-lanceolatis a sequente discedit. — Exsicc.: Heldr. pl. fl. hellen. a. 1895.

In saxosis montosis. Epirus: ad Paleochori Syraku distr. Janina, pr. Cepelovon distr. Zagorion (Bald.); — *β.* Thessalia: pr. Hagios Dionysios et Litochoron in Olympo (Sint.); — *γ.* mt. Dirphys Euboeae (Heldr.). — Aug. Sept. ♃

Obs. *S. pygmaea* Sieb. in Koch syn. p. 559, a praecedente caulibus tetragonis, saepius glabris, calycis purpurascentis dentibus inferioribus longioribus, corollae purpureae lobo intermedio labii inferioris latiore obcordato discedens, indicatur a Form. in D. bot. Mon. 1891 p. 28, probabiliter ex confusione, pr. Chaliki, Velitsena et Vendista Thessaliae.

4. **S. parnassica** Heldr. et Sart. pl. exs. a. 1846 n. 1871, herb. norm. n. 80, pro var. *S. spinosae*; Heldr. et Sart. in Bois. fl. or. IV. p. 563; Heldr. chlor. Parn. p. 25; Hal. in ö. b. Z. 1897 p. 325, in . b. G. 1899 p. 191. — *S. approximata* Friv. in Flora 1836 p. 438, non Biv. stirp. in sic. man. IV. p. 13.

Suffruticosa, griseo-viridis; caulibus brevibus, caespitosis, procumbentibus, hispidis, ramosis, in racemos breves, densiusculos, paucifloros abeuntibus; foliis sessilibus, hispidulis, pellucido-punctatis, inferioribus obovatis, obtusis, superioribus spathulatis, acutis, floralibus flores aequantibus vel superantibus; cymulis 1—3 floris, subsessilibus; calycis hispidi dentibus aequalibus, lanceolatis, tubo subbrevioribus; corollae

albidae tubo incluso. — Differt a *S. montana* rhizomate crasso, tortuoso, caulibus nanis, 3—5 cm. longis, caespitosis, procumbentibus, foliis parvis, 5—7 mm. latis, 3—4 mm. latis, hispidis, inferioribus obtusis, verticillastris paucis.

β. **stygia** Orph. fl. gr. exs. n. 622 pro var. *S. spinosae.* — Foliis canis, dense hispido-tomentellis, fere omnibus obovatis, obtusis.

γ. **macrophylla** Hal. in z. b. G. 1899 p. 191. — Caulibus ad 10 cm. longis, foliis virentibus, minus hispidis, majoribus, ad 25 mm. longis et 7 mm. latis; cymulis numerosioribus, breviter pedunculatis. — Exsicc.: Heldr. pl. fl. hellen. a. 1896; Sint. it. thessal. n. 1141b.

In saxosis regionis subalpinae et alpinae mt. Parnassus et Taygetos (Heldr.); — *β.* mt. Chelmos (Orph.); — *γ.* Thessalia: mt. Kokino Lithari pr. Sermeniko (Sint.); Aetolia: mt. Korax (Tunt.). — Jul. Aug. ♃

β. Rami spinescentes.

5. **S. spinosa** L. sp. ed. 2. p. 795; S. et S. pr. I. p. 398, Fl. gr. VI. p. 36 t. 545; Sieb. avis p. 4, rem. p. 5; Raul. cret. p. 828; Bois. fl. or. IV. p. 563; Bald. viagg. Creta p. 83. — Exsicc.: Bald. it. cret. n. 134, it. cret. alt n. 188.

Suffruticosa, ramosissima; ramis crassis, lignosis, intricatis, spinescentibus, hornotinis brevissimis, tomentellis; foliis minimis, sessilibus, oblongo-lanceolatis, acutis, sparsim hirtis, pellucide-punctatis; floribus paucis, axillaribus, solitariis, subsessilibus; calycis scabriusculi dentibus aequalibus, lanceolatis, tubo subbrevioribus; corollae albidae tubo incluso. — Dumulus erinaceus, palmaris.

In saxosis regionis subalpinae et alpinae. Creta: mt. Sphakia, Ida et Lassiti. — Jul. Aug. ♃

25. Origanum L. gen. n. 726.

1. Sectio. *Amaracus* Moench meth. suppl. p. 137. — Folia floralia ampla; calyx bilabiatus, labium superius elongatum, integrum, inferius multo brevius, dentibus angustis vel obsoletis.

1. **O. dictamnus** L. sp. p. 589; S. et S. pr. I. p. 416; Sieb. avis p. 4, rem. p. 5; Fraas fl. class. p. 181; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 547; Spreitz. in z. b. G. 1890 p. 299; Bald. viagg. Creta p. 83. — *O. pseudodictamnus* Sieb. in Flora l. p. 273. — *Amaracus dictamnus* Benth. lab. p. 333; Raul. cret. p. 826. — Icon: Curt. bot. mag. t. 298. — Exsicc.: Spreitz. it. cret. a. 1882.

Rhizomate suffrutescenti; caulibus numerosis, adscendentibus, lanatis, simplicibus vel parce paniculato-ramosis, foliosis, superne subnudis; foliis suborbiculatis, utrinque lanatis, inferioribus breviter petiolatis, ceteris sessilibus; spicis terminalibus, solitariis, ovatis vel oblongis; bracteis coloratis, calyces superantibus; calycis labio inferiore subretuso; corollae roseae, calyce duplo longioris tubo non gibbo.

Ad rupes umbrosas faucium regionis inferioris et montanae. Creta: ad Akroteri pr. Canea (Weiss), promontorium Meleka (Tourn. voy. p. 12), pr. Katholiko, promontorium Spada, pr. Sina, Aradhena, Laki, Asprokremnos, in fauce Nipros, mt. Ida, Kophino, Lassiti supra Males (Raul.), Asomatos-Kavutsi (Bald.); Argolis: in montibus pr. Poros (Sart.). — Maio, Jul. ♃

2. **O. Tournefortii** Sibth. in Ait. Kew. II. p. 311; S. et S. pr. I. p. 416, Fl. gr. VI. p. 56 t 569; Bois. fl. or. IV. p. 547; Heldr. in ö. b. Z. 1898 p. 184. — *O. dictamnus* Ch. et B. fl. pelop. p. 38, non L. — *Amaracus Tournefortii* Benth. lab. p. 333. — Exsicc.: Orph. fl. gr. n. 1168.

Rhizomate crasso, suffrutescenti; caulibus numerosis, fragillimis, adscendentibus, lanatis, subsimplicibus, tota longitudine foliosis; foliis suborbiculatis vel ellipticis, lanatis vel ciliatis, glaucis; spicis terminalibus, solitariis vel 2—3 nis, oblongis; bracteis purpureis, calyces superantibus; calycis labio inferiore bidentato; corollae roseae, calyce triplo longioris tubo medium versus gibbo. — Differt a praecedente indumento tenuiore, caulibus omnino foliatis, foliis coriaceis, glaucis, corollae tubo longiore, gibbo.

In rupibus insulae Amorgos, prope divae virginis coenobium (Tourn.). — Apr. Aug. ♃

2. Sectio. *Anatolicon* Benth. in DC. pr. XII. p. 192. — Folia floralia ampla, colorata; calyx bilabiatus, labium superius distincte tridentatum, inferius bidentatum, dentibus angustioribus vel abbreviatis.

a. Caulis glaber; folia subcordata.

3. **O. scabrum** Bois. et Heldr. diagn. VII. p. 48, Fl. or. IV. p. 549. — *Amaracus scaber* Briqu. lab. in Engl. et Prantl. nat. Pflanzenfam. IV. p. 306. — Exsicc.: Orph. fl. gr. n. 42; Heldr. herb. norm. n. 1471.

Rhizomate repente; caulibus erectis, foliosis, apice breviter racemosis; foliis sessilibus, coriaceis, glaucis, punctatis, cordato-ovatis, acutiusculis, glabris, margine scabris; spicis ovatis, nutantibus; bracteis coloratis, ovatis, glabris, calyces superantibus; calycis glabriusculi, punctati labio superiore ad tertiam partem obtuse tridentato, inferiore triplo breviore, in dentes lanceolatos bipartito; corollae roseae, tubo calyce duplo longiore. — Species pulcherrima.

In rupestribus regionis silvaticae mt. Malevo pr. Vromopigadon (Orph.) et mt. Taygetos (Heldr.). — Jul. Aug. ♃

4. **O. pulchrum** Bois. et Heldr. diagn. ser. 2. IV. p. 11, Fl. or. IV. p. 549. — *O. sipyleum* S. et S. pr. I. p. 417 quoad pl. euboeam; Fraas fl. class. p. 182; non L. — *Amaracus pulcher* Briqu. lab. in Engl. et Prantl nat. Pflanzenfam. IV. p. 306. — Exsicc.: Heldr. herb. norm. n. 784 et 969.

Differt a praecedente, cui maxime affine, foliis margine laeviusculis, spicis longius pedunculatis, bracteis obtusiusculis, calycis labio superiore ad medium tridentato, inferiore vix breviore.

In lapidosis regionis superioris mt. Dirphys et Kandyli (Heldr.) Euboeae (Sibth.). — Jul. Aug. ♃

b. Caulis breviter hirtulus; folia basi rotundata.

5. **O. lirium** Heldr. herb. norm. n. 753 b; Hal. in z. b. G. 1899 p. 192. — *O. hybridum* Heldr. herb. norm. n. 753, non Mill.

Rhizomate repente; caulibus erectis, foliosis, in paniculam plus minus compositam abeuntibus; foliis sessilibus, coriaceis, glaucis, punctatis, ovatis, basi rotundatis; spicis oblongis, nutantibus; bracteis coloratis, ovatis, hirtulis, calyces superantibus; calycis glabriusculi, punctati labio superiore breviter obtusiuscule tridentato, inferiore vix breviore, breviter bidentato; corollae roseae tubo calyce duplo longiore. — Spicae eis praecedentium multo minores, saepe confertae. — „Cum cl. Boissier olim pro stirpe inter *O. pulchrum* et *O. hirtum* hybrida habui, sed nunc potius species propria mihi videtur" (Heldr. l. c.).

Euboea: in regione abietina mt. Dirphys l. d. Elatakia, circa fontem Liri, nec non in mt. Xerovuni (Heldr.). — Jul. Aug. ♃

3. Sectio. *Euoriganum* Vog. in Limnaea 1841 p. 79. — Folia floralia mediocria, interdum colorata; calyx quinquedentatus, dentibus subaequilongis.

6. **O. viride** Bois. fl. or. IV. p. 551; Haussk. symb. p. 49; Form. in Ver. Brünn 1897 p. 49: pro var. *O. vulgaris*; Hal. Beitr. fl. Thessal. p. 17. — *O. viridulum* Heldr. in ö. b. Z. 1898 p. 184, an Mart. Don. fl. Tarn. p. 551. — Huc probabiliter: *O. vulgare* S. et S. pr. I. p. 418; Pieri corc. fl. p. 79; Ch. et B. exp. p. 171, Fl. pelop. p. 38; Fraas fl. class. p. 181 v. *album*; Heldr. chlor. Parn. p. 25; Form. in D. bot. Mon. 1891 p. 28, in Ver. Brünn 1895 p. 35, 1896 p. 66, 1897 p. 49; vix L. sp. p. 590, quod indumento copiosiori, floribus dimidio majoribus, saepissime purpureis, in paniculam saepius corymbiformem dispositis, bracteis saepe coloratis discedit et in Graecia vix occurrit. — Exsicc.: Hal. it. gr. secund. a. 1893.

Rhizomate obliquo; caulibus erectis, foliatis, foliisque pilosulis; foliis petiolatis, ovatis vel oblongis, punctatis, subtus glaucescentibus; spicis densifloris, ovatis vel oblongis, glomeratis, racemoso-paniculatis; bracteis viridibus, ovatis, acutis, glabris, non punctatis, calycem superantibus; calycis glabri vel pilosuli, vix punctati dentibus ovatis, acutis; corollae albae, rarissime roseae tubo breviter exserto.

In dumosis regionis praesertim montanae. Frequens ut videtur in Epiro in Pindo thessalo, dein pr. Pharsalum (Haussk.), pr. Selicany et Spilia in mt. Ossa (Form.), mt. Olympus (Heldr.), indicatur quoque in mt. Parnasso (Heldr.); Euboea (Fraas), pr. Gortys, Marathonisi et [illegible] (Chaub.), Cephalonia (Dall.), Corcyra (Pieri), [illegible] loca nonnulla probabiliter ad speciem

sequentem spectant, quacum ut videtur saepe confunditur. — Jun. Aug. ♃

7. **O. heracleoticum** L. sp. p. 589; S. et S. pr. I. p. 418; Benth. lab. p. 336; Marg. et R. fl. Zante p. 73; Ch. et B. fl. pelop. p. 38; Friedr. Reise p. 275; Fraas fl. class. p. 181. — *O. smyrnaeum* S. et S. pr. I. p. 417, Fl. gr. VI. p. 57 t. 571; ? Link in Linnaea IX. p. 576; Raul. cret. p. 827; non L. sp. p. 589, quod sec. Benth. lab. p. 339 = *O. onites* L. — *O. hirtum* Link en hort. berol. II. p. 114; Vog. in Linnaea 1841 p. 86; Clem. sert. p. 74; Raul. cret. p. 827; Bois. fl. or. IV. p. 552; Heldr. fl. cephal. p. 57, in Sitzungsb. acad. Wiss. Berl. 1883 p. 4, chlor. Parn. p. 25; Hal. Beitr. fl. Epir. p. 36; Haussk. symb. p. 49. — *O. parviflorum* Urv. enum. p. 71; Ch. et B. exp. p. 171. — *O. neglectum* Vog. l. c. p. 81; Raul. cret. p. 827. — ? *O. vulgare v. latebracteatum* Form. in Ver. Brünn 1895 p. 35; vix Beck fl. Südbosn. p. 142, quod bracteis latissimis intense purpureis gaudet et in Graecia vix provenit.

Rhizomate obliquo; caulibus erectis, foliatis, foliisque plus minusve hirtis; foliis ovatis vel oblongis, punctatis, subtus pallidioribus; spicis densifloris, ovatis vel oblongis, glomeratis, racemoso-paniculatis; bracteis viridibus, glabris vel hirsutis, glanduloso-punctatis, calyce sublongioribus; calycis glabri vel hirti, glanduloso-punctati dentibus lanceolatis acutis; corollae albae, rarius roseae tubo subincluso. — A praecedente praesertim indumento et bracteis calycibusque glanduloso-punctatis vix specifice discedit. Variat:

α. **albiflorum** Haussk. symb. p. 49, pro var. *O. hirti.* — Spicae breves; bracteae virides, glabrae, corolla alba. — Exsicc.: Heldr. it. thessal. n. 24; Sint. it. thessal. n. 1515; Dörfl. fl. gr. n. 200.

β. **rubriflorum** Haussk. l. c. pro var. *O. hirti.* — Spicae breves; bracteae supernae purpurascentes, glabrae; corolla purpurascens. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

γ. **trichocalycinum** Haussk. symb. p. 50, pro var. *O. hirti.* — Spicae breves; bracteae et calyces dense patule hirsuti; indumentum caulium et foliorum copiosius. — Exsicc.: Orph. fl. gr. n. 528; Heldr. herb. norm. n. 782; Rev. pl. cret. n. 134; Dörfl. fl. aeg. n. 133.

δ. **creticum** L. sp. p. 589?; S. et S. pr. I. p. 417; Sieb. avis p. 4, rem. p. 5; Urv. enum. p. 70; Fraas fl. class. p. 181; pro sp. — *O. vulgare v. prismaticum* Raul. cret. p. 826, non Gaud. fl. helv. IV. p. 78. — *O. hirtum v. prismaticum* Haussk. symb. p. 49. — *O. vulgare v. elongatum* Form. in ö. b. Z. 1890 p. 92, in Ver. Brünn 1895 p. 35, 1896 p. 67, 1897 p. 49. — Spicae fructiferae elongatae, prismaticae; bracteae virides nunc glabrae nunc hirsutae. — Exsicc.: Sint. et Bornm. it. turc. n. 1404b (*f. glabra*); Heldr. pl. fl. hellen. a. 1897 (*f. hirsuta*).

In dumosis regionis inferioris et montanae passim per totam Graeciam. — Jun. Aug. ♃

4. Sectio. *Majorana* Moench meth. p. 406. — Folia floralia minima, non colorata, dense imbricata; calyx bilabiatus, labium superius integrum vel rarius denticulatum, inferius minimum, bidentatum truncatum vel nullum.

a. Folia mediocria, viridi-canescentia; corolla alba.

8. **O. onites** L. sp. p. 590; S. et S. pr. I. p. 418, Fl. gr. VI. p. 58 t. 572; Bois. fl. or. IV. p. 553; Bald. viagg. Creta p. 83; Hal. in ö. b. Z. 1896 p 17, 1897 p. 98, 1898 p. 184; Haussk. symb. p. 50; Heldr. chlor. Thera p. 19. — *O smyrnaeum* L. sp. p. 589, sec. Benth. lab. p. 339; Sieb. avis p. 4, rem. p. 5; Ch. et B. exp. p. 171, Fl. pelop. p. 38. — *Majorana onites* Benth. lab. p. 339; Raul. cret. p. 828. — *Schizocalyx smyrnaeus* Scheele in Flora 1843 p. 575. — Exsicc.: Orph. fl. gr. n. 143 et 1112; Dörfl. fl. aeg. n. 25.

Basi suffruticosum; caulibus erectis, foliatis, hirsutis; foliis ovatis, tomentellis, punctatis, subserratis, infimis brevissime petiolatis, ceteris sessilibus, acutis; spicis ovatis, compactis, in corymbum planiusculum congestis; bracteis ovato-orbiculatis, acutiusculis, hirtis, punctatis, calyce sublongioribus; calyce ovato-spathulato, antice fisso, unilabiato, obsoletissime tridenticulato; corollae albae tubo incluso.

In dumosis, collibus siccis regionis inferioris. Attica: mt. Pateras (Pichl.); Acrocorinthus, pr. Vromolimni (Heldr.) et Nauplia (Orph.) Argolidis, ad fl. Eurotas versus Scala, pr. Marathonisi, Scutari, Kardamyle, Vitilos, Monembasia (Chaub.); Sporadum insula Jura (Reis.); Cycladum insula: Tenos (Chaub.), Cythnos (Orph.), Jos, Naxos, Thera (Heldr.); Creta: pr. Melidoni (Bald.), Suda, Aludha pr. Spinalonga, Zakro (Raul.). — Apr. Jul. ♃

9. **O. dubium** Bois. fl. or. IV. p. 553.

Basi suffruticosum; caulibus erectis, foliatis, glabriusculis vel tomentellis; foliis petiolatis, adpresse canescentibus, ovatis, obtusis, integris; spicis ovatis, glomeratis, paniculatis; bracteis tomentellis, ovato-spathulatis, obtusis, calyce sublongioribus; calyce ovato-spathulato, antice fisso, unilabiato, obtuso; corollae albae tubo incluso. — Species mihi ignota, a praecedente inflorescentia paniculata et foliis integris differre dicitur.

In Cycladum insula Naxos (Lenormand). — ♃

b. Folia minima, ad 5 mm. longa, albo-tomentosa; corolla purpurea.

10. **O. maru** L. sp. ed. 2 p. 825; S. et S. pr. I. p. 419, Fl. gr. VI. p. 59 t. 573; Sieb. avis p. 4, rem. p. 5, in Flora I. p. 275. — *O. microphyllum* Sieb. in Benth. lab. p. 338; Bois. fl. or. IV. p. 552; Spreitz. in z. b. G. 1890 p. 298; Bald. viagg. Creta p. 83. — *Majorana microphylla* Benth. l. c.; Raul. cret. p. 827. — Exsicc.: Rev. pl. cret. n. 128; Bald. it. cret. n. 85.

Basi suffruticosum; caulibus erectis, remote foliatis, rubellis, glabriusculis; foliis minimis, ovatis vel oblongis, integris, breviter petiolatis;

spicis subglobosis, solitariis vel ternis, albo-lanatis; bracteis spathulato-ovatis, obtusis, calyce longioribus; calyce glabro, glanduloso-punctato, turbinato, oblique truncato, edentulo, fauce barbato; corollae purpureae tubo vix exserto. — Species eximia, habitu *O. majoranae* L., capitulis piso minoribus. — Linnaeus sub suo *O. maru* certe nostram plantam habuit, uti ex descriptione sua accurata atque plantae patriae patet; *Maru creticum* Alp. exot. t. 288 a Linnaeo ipso pro sua specie citatum insuper nostram bene exhibet.

In saxosis regionis montanae Cretae: mt. Lakus (Rev.), Omalos (Raul.), pr. Askyphos (Spreitz.) in mt. Sphacioticis (Sibth.). — Jun. Jul. ♃

Obs. *O. majorana* L. sp. p. 590; Dallap. prosp. p. 84; Heldr. Nutzpfl. p. 32. — Colitur ad usum culinarem in hortis. — *O. crassifolium* Benth. lab. p. 339; Raul. cret. p. 827; (*Majorana*); a Bentham ex confusione cum *O. maru* L. in Creta indicatur.

26. Thymbra L. gen. n. 708.

1. **T. spicata** L. sp. p. 569; S. et S. pr. I. p. 398, Fl. gr. VI. p. 37 t. 546; Benth. lab. p. 413; Ch. et B. fl. pelop. p. 36; Friedr. Reise p. 263; Fraas fl. class. p. 176; Raul. cret. p. 831; Bois. fl. or. IV. p. 561. — ? *Satureia spicata* Sieb. in Flora I. p. 270. — Exsicc.: Orph. fl. gr. n. 529.

Fruticulus dumulosus; ramis erectis, novellis bifariam puberulis; foliis sessilibus, linearibus, punctatis, glabris; spicis oblongo-cylindricis; bracteis lanceolatis, ciliatis, corollae tubum aequantibus; calycis ciliati labio superiore dentibus breviter triangularibus, inferiore aequilongo dentibus lineari-subulatis; corollae roseae tubo exserto; staminibus vix exsertis.

In collibus apricis regionis inferioris, rare. Boeotia: pr. Lebadea (Orph.); Attica: pr. Marathon (Fraas), Oropos (Heldr.); Achaia (Sibth.); Cycladum insula Melos (Beaupré); Creta (Sibth.): Corcyra: pr. urbem (Friedr.). — Jun. Aug. ♄

2. **T. capitata** L. sp. p. 568; S. et S. pr. I. p. 397, Fl. gr. VI. p. 36 t. 544; Sieb. avis p. 4, rem. p. 5; Urv. enum. p. 65; Link in Linnaea 1834 p. 574; Fraas fl. class. p. 174; (*Satureia*); Griseb. spic. II. p. 127; Clem. sert. p. 75; Heldr. Nutzpfl. p. 33; Raul. cret. p. 830. — *Thymus capitatus* Hoffm. et Link fl. port. I. p. 123; Marg. et R. fl. Zante p. 74; Friedr. Reise p. 274; Weiss in z. b. G. 1869 p. 744; Bois. fl. or. IV. p. 560; Heldr. fl. cephal. p. 57, Fl. Aegina p. 385, chlor. Thera p. 19, chlor. Mykon p. 249; Spreitz. in z. b. G. 1887 p. 666; Form. in D. bot. Mon. 1891 p. 28, 1898 p. 79, in Ver. Brünn 1895 p. 35, 1896 p. 67, 1897 p. 50; Hal. Beitr. fl. Epir. p. 38; Bald. viagg. Creta p. 82; Haussk. symb. p. 50. — *T. creticus* Brot. lusit. I. p. 174; Ch. et B. exp. p. 171, Fl. pelop. p. 38. — *T. vulgaris* Pieri corc. fl. p. 79; Dallap. prosp. p. 79.

non L. — Exsicc.: Orph. fl. gr. n. 525; Rev. pl. cret. n. 132; Bald. it. cret. n. 135; Heldr. herb. norm. n. 1369.

Fruticulus dumulosus; ramis erectis, novellis undique tomentellis; foliis sessilibus, brevissimis, oblongo-linearibus, punctatis, basi ciliatis; spicis ovatis; bracteis ovatis vel oblongis, ciliatis, calycem aequantibus; calycis ciliati labio superiore dentibus brevissimis, inferiore sublongiore dentibus subulatis; corollae roseae, rarissime albae tubo breviter exserto; staminibus longe exsertis.

β. **albo-spinosa** Bald. viagg. Creta p. 82 pro var. *Thymi capitati.* — Erinacea, ramis induratis, apice spinosis; spicis depauperatis; floribus albis. — Exsicc.: Bald. it. cret. n. 164.

In aridis, campis, collibus saxosis regionis calidae copiosissime per totam Graeciam; — *β.* in alpestribus mt. Ida Cretae (Bald.). — Maio, Sept. ♃

27. Thymus L. gen. n. 727.

Dispositio specierum:

1. Sectio. *Euthymus* Borb. in math. term. Közl. XXIV. p. 48. — Calycis non gibbi labium superius ovatum, saepissime ad tertiam vel mediam partem in dentes lanceolatos vel triangulares, aequales divisum.

a. Folia margine revoluta.

α. Folia floralia caulinis subconformia.

1. T. Boissieri Hal. **2. T. ciliatopubescens** Hal.

β. Folia floralia caulinis manifeste latiora.

× Folia praeter marginem parce ciliatum glabra.

3. T. tymphresteus Hal.

×× Folia undique hirta.

○ Calyx pellucide punctatus.

4. T. euboeus Hal.

○○ Calyx impunctatus.

5. T. leucotrichus Hal. **6. T. parnassicus** Hal.

b. Folia margine non revoluta.

β. Folia pube minutissima velutina.

7. T. leucadicus (Rchb.).

β. Folia glabra vel hirsuta, sed nunquam velutina.

× Folia linearia, rigida; calycis dentes labii superioris lanceolato-acuminati

8. T. atticus Celak. **9. T. striatus** Vahl.

×× Folia latiora, lineari-elliptica, oblonga vel ovata, vix rigida; calycis dentes labii superioris triangulari-lanceolati.

○ Rami teretiusculi, folia parva vel medioria.

. Rami floriferi plus minus elongati; capitula non involucrata.

10. T. Sibthorpii Benth. **11. T. serpyllum** L. **12. T. lanuginosus** Mill.

.. Rami floriferi brevissimi; capitula foliis summis involucrata.

13. T. ocheus Heldr. et Sart.

○○ Rami tetrogoni; falia majuscula, ad 15 mm. longa et 8 mm. lata

14. T. montanus W. et K.

2. Sectio. *Piperella* Willk. et Lange pr. fl. hisp. II. p. 404. — Calycis subgibbi labium superius late ovatum, ad quartam partem in dentes breviter et late triangulares, acutos, subaequales, laterales nempe medio subminores, divisum.

15. T. teucrioides Bois. et Spr.

1. Sectio. *Euthymus* Borb. in math. termész. Közl. XXIV. p. 48. — Calycis non gibbi labium superius ovatum saepissime ad tertiam vel mediam partem in dentes lanceolatos vel triangulares, aequales divisum.

a. Folia margine revoluta.

α. Folia floralia caulinis subconformia.

1. **T. Boissieri** Hal. Beitr. fl. Epir. p. 37; Form. in Ver. Brünn 1896 p. 67. — *T. hirsutus* Bois. fl. or. IV. p. 557 p. p.; non M. a B. — *T. hirsutus v. humillimus* Bald. riv. coll. bot. alb. 1895 p. 61; non *T. humillimus* Celak. in Flora 1884 p. 534, qui = *T. cherlerioides* Vis. ill. alc. piante Grec. c. As. min. 1842 p. 8 (*T. hirsutus* Sint. it. trojan. 1883 n. 1176) et a nostro differt ramis brevissimis, caespites densissimos formantibus, capitulis e caespite vix exsertis, foliis minoribus, velutino-puberulis et calycis minus ciliati dentibus labii superioris brevioribus. — Exsicc.: Hal. it. gr. sec. a. 1893; Bald. it. alb. epir. III. n. 184.

Rhizomate suffruticoso; caulibus numerosissimis, caespites pulvinatos formantibus, ramis brevibus, dense foliosis, circumcirca patule pilosis; foliis anguste linearibus, obtusis, viridibus, impunctatis, margine revoluto longe albo-ciliatis, ceterum glaberrimis, nervis secundariis obsoletis instructis, floralibus sublatioribus, rubentibus, penninerviis, calyces parum superantibus; floribus pedicellatis, in capitula subglobosa vel ovata congestis; calycis bilabiati, rubentis, patule hirsuti, impunctati labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis; corollae parvae, purpureae tubo calycis dentes vix superante. — Species elegans, *T. hirsuto* M. a. B. foliis punctatis, undique hirsutis, canescentibus, calycis labii superioris dentibus latioribus brevioribusque et corolla pellucide punctata egregie diverso, affinis.

In herbidis saxosis regionis superioris. Epirus: mt. Tsumerka (Bald.), frequentissime in cacumine mt. Peristeri (Hal.); Thessalia: mt. Olympus et Ossa (Heldr.). — Jul. Aug. ♃

2. **T. ciliato-pubescens** Hal. Beitr. fl. Epir. p. 37 pro var. *T. Boissieri*. — *T. hirsutus* Hal. in z. b. G. 1888 p. 761, non M. a B. — Exsicc.: Hal. it. gr. a. 1888.

Rhizomate suffruticoso; caulibus numerosis, caespites densos formantibus, ramis brevibus, dense foliosis, circumcirca patule pilosis; foliis linearibus, obtusis, viridibus, parce punctatis, undique pube minutissima velutinis, margine revoluto longe albo-ciliatis, nervis secundariis obsoletis instructis, floralibus sublatioribus, vix coloratis, penninerviis, calyces parum superantibus; floribus pedicellatis, in capitula ovata vel sub-

globosa congestis; calycis bilabiati, rubentis, patule hirsuti, impunctati labio superiore fere ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis; corollae parvae, roseae tubo calycis dentes vix superante. — Differt a praecedente rhizomate crassiore, foliis latioribus, dense velutinis, parce punctatis, capitulis submajoribus et calycis labii superioris dentibus basi latioribus. — *T. hirsutus* M. a. B. ex speciminibus tauricis (Callier it. taur. sec. a. 1896 n. 228) discedit a nostra foliis rubenti punctatis, patule hirsutis, calycis labii superioris dentibus latioribus brevioribusque et corolla punctata.

In rupestribus regionis alpinae. Doris: mt. Kiona (Hal.). — Jul. Aug. ♃

β. Folia floralia caulinis manifeste latiora.

× Folia praeter marginem parce ciliatum glabra.

3. **T. tymphrestus.** — *T. hirsutus* Bois. fl. or. IV. p. 557 p. p., non M. a B. — Exsicc.: Heldr. it. graec. septentr. a. 1879.

Rhizomate suffruticoso, crasso; caulibus numerosis, lignosis, caespitosis, ramis brevibus, dense foliosis, floriferis circumcirca breviter puberulis, sterilibus glabriusculis; foliis linearibus, obtusis, viridibus, parce punctatis, margine revoluto parce ciliatis, ceterum glabris, nervis secundariis obsoletis vel parum prominentibus instructis, floralibus ovatis vel oblongis, non coloratis, subtus elevatim penninerviis, calyces superantibus; floribus pedicellatis, in capitula subglobosa congestis; calycis bilabiati, rubentis, patule hirsuti, impunctati labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus lanceolato-subulatis; corollae parvae, roseae tubo calycis dentes vix superante. — A *T. Boissieri*, quocum indumento convenit, rhizomate crasso, valde lignoso, foliis rigidis, punctatis, floralibus difformibus discedit.

In regione alpina mt. Tymphrestus Eurytaniae (Heldr.). — Jul. Aug. ♃

×× Folia undique hirta.

○ Calyx pellucide punctatus.

4. **T. euboeus.** — Exsicc.: Leonis pl. gr. a. 1901.

Rhizomate suffruticoso; caulibus numerosis, decumbentibus, ramis praesertim sterilibus elongatis, foliosis, circumcirca hirsutis; foliis linearilanceolatis, acutiusculis, obscure viridibus, rubenti-punctatis, undique dense velutino-pubescentibus, margine subrevolutis, basin versus albociliatis, nervis secundariis obsoletis instructis, floralibus ovato-oblongis, non coloratis, subtus penninerviis, calyce aequilongis; floribus pedicellatis, in capitula magna, ovata vel oblonga, inferne interdum interrupta congestis; calycis bilabiati, vix colorati, hirsuti, pellucide punctati labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori brevioris dentibus subulatis; corollae parvae, intense purpureae tubo calycis dentibus breviore. — Species colore obscure viridi, capitulis magnis, 15 mm. et ultra diametro latis, calycibus pellucide punctatis, corollis parvis, intense purpureis, ab affinibus egregie distincta.

Euboea: in cacumine mt. Dirphys (Leonis). — Jul. Aug. ♃

○○ Calyx impunctatus.

5. **T. leucotrichus.** — *T. hirsutus* Bois. fl. or. IV. p. 557 p. p. — *T. parnassicus* Hal. in z. b. G. 1899 p. 193, non Beitr. fl. Epir. p. 38. — Huc forsan *T. vulgaris* S. et S. pr. I. p. 419, non L. — Exsicc.: Orph. herb. a. 1854; Heldr. pl. fl. hellen. a. 1897.

Rhizomate suffruticoso; caulibus numerosis, decumbentibus, ramis foliosis, circumcirca patule hirsutis; foliis lineari-lanceolatis, obtusiusculis, impunctatis, undique pilis brevibus longisque dense obsitis, canescentibus, margine subrevolutis, nervis secundariis obsoletis instructis, floralibus ovato-oblongis oblongisve, non coloratis, subtus penninerviis, calyce longioribus; floribus pedicellatis, in capitula ovata vel globosa congestis; calycis bilabiati, subrubentis, albo-villosi, impunctati labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis; corollae roseae tubo calycis dentes subsuperante. — *T. revoluto* Celak. in Flora 1883 p. 171, mihi e descriptione tantum noto, ut videtur similis, qui autem foliis floralibus rotundatis, abrupte acuminatis calycibusque rubenti punctatis discedit. — *T. hirsutus* M. a. B. foliis floralibus conformibus, calycis labii superioris dentibus brevioribus et floribus minoribus a nostro differt.

β. **Degenii.** — *T. revolutus v. creticus* Deg. in Bald. viagg. Creta p. 82, non *T. creticus* Brot., nec DC. — Huc probabiliter: *T. serpyllum* S. et S. pr. I. p. 418 quoad pl. cret.; Raul. cret. p. 828; non L. — *T. striatus* Raul. l. c., non Vahl. — *T. hirsutus* Spreitz. in z. b. G. 1890 p. 299, non M. a. B. — Rhizomate tortuoso, crasso, valde lignoso, caulibus abbreviatis, ramis floriferis brevissimis, foliis minoribus, minus copiose hirsutis. — Forma macra ut videtur speciei. — Exsicc.: Bald. it. cret. n. 137.

In rupestribus regionis alpinae. Peloponnesus: mt. Kyllene, Chelmos (Orph.), Taygetos (Heldr.); — *β.* Creta: mt. Mavrus Lakus (Spreitz.) et Hagion Pneuma (Bald.) in mt. Sphacioticis. — Jun. Jul. ♃

6. **T. parnassicus** Hal. Beitr. fl. Epir. p. 38. — *T. hirsutus* Bois. fl. or. IV. p. 557 p. p., non M. B. — Exsicc.: Hal. it. gr. a. 1888.

Rhizomate suffruticoso; caulibus elongatis, decumbentibus, ramis foliosis, brevibus, undique retrorsum puberulis; foliis linearibus, obtusis, parce punctatis, undique pube minutissima velutino-canescentibus margineque revoluto albo-ciliatis, nervis secundariis obsoletis instructis, floralibus ovatis vel oblongis, non coloratis, subtus penninerviis, calyce aequilongis; floribus pedicellatis, in capitula globosa congestis; calycis bilabiati, vix colorati, adpresse hirti, impunctati labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis, omnibus longe pectinato-ciliatis; corollae roseae tubo calycis dentes superante. — Ab affinibus praesertim indumento brevissime velutino discedit.

In rupestribus regionis alpinae mt. Parnassus, rarissime (Hal.); forsan quoque in mt. Pateras Atticae, cum *T. hirsutus* a Boissier ibi indicatur. — Jul. Aug. ♃

b. Folia margine non revoluta.

α. Folia pube minutissima velutina.

7. **T. leucadicus** Rchb. ic. fl. germ. et helv. XVIII. p. 36; Bald. in bull. herb. Bois. IV. p. 205; pro var. *T. striati*. — *T. striatus* Bois. fl. or. IV. p. 557 p. p.; Heldr. fl. cephal. p. 57; non Vahl. — Exsicc.: Heldr. pl. cephal. a. 1867; Bald. it. alb. epir. III. n. 79.

Rhizomate suffruticoso; caulibus decumbentibus, ramis foliosis, undique brevissime tomentellis; foliis lineari-lanceolatis, acutissimis, punctatis, undique pube minutissima velutino-canescentibus, margine ciliatis, nervis secundariis parum prominentibus vel obsoletis instructis, floralibus ovatis, acutis, valide penninerviis, calyce longioribus; floribus pedicellatis, in capitula magna, ovata congestis; calycis bilabiati, non colorati, velutini, pellucide punctati labio superiore vix ad quartam partem in dentes breviter lanceolatos fisso, labii inferioris, superiori subaequilongi dentibus subulatis, omnibus pectinato-ciliatis; corollae roseae tubo calycis dentes superante. — Species foliis acutissimis, velutinis, non revolutis, capitulis magnis, calycis labio superiore brevissime dentato, corolla majuscula distinctissima.

In rupestribus regionis inferioris et montanae. Cephalonia: mt. Aenos, Phalaris (Heldr.); Leucas (Mazz.): pr. Karya (Bald.). — Jun. Aug. ♃

β. Folia glabra vel hirsuta, sed nunquam velutina.

× Folia linearia, rigida; calycis dentes labii superioris lanceolato-acuminati.

8. **T. atticus** Celak. in Flora 1882 p. 564, 1883 p. 165; Bois. fl. or. suppl. p. 360; Hal. Beitr. fl. Achaia p. 30; Form. in Ver. Brünn 1896 p. 67, in D. bot. Mon. 1898 p. 79; Haussk. symb. p. 50. — *T. zygis* S. et S. pr. I. p. 420, Fl. gr. VI. p. 60 t. 574; Ch. et B. fl. pelop. p. 38; Fraas fl. class. p. 178; Clem. sert. p. 75; non L. sp. p. 591, qui Hispaniae civis et foliis tomentellis, verticillastris distinctis, spicam longam interruptam formantibus, foliis floralibus caulinis conformibus, floribus multo minoribus diversissimus est. — *T. striatus* Bois. fl. or. IV. p. 557 p. p., non Vahl. — Exsicc.: Orph. fl. gr. n. 199; Heldr. herb. norm. n. 291 et 874.

Rhizomate suffruticoso; caulibus procumbentibus, ramis floriferis strictis, undique hirtis; foliis viridibus, linearibus, obtusis, basin versus longe spathulato-angustatis, punctatis, ciliatis, ceterum glaberrimis, subtus nervo mediano valido secundariisque parum prominulis instructis, floralibus ovato-lanceolatis, puberulis, pectinato-ciliatis, subtus valide pluri-nerviis; floribus breviter pedicellatis, in capitula globosa vel ovata congestis; calycis bilabiati, pallidi, breviter hirti, pellucide punctati labio

superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis, omnibus pectinato-ciliatis; corollae albae tubo calycis dentes non superante. — Species foliis rigidis, linearibus, glabris et floribus albis egregia.

In saxosis regionis montanae, ut videtur rare. Thessalia: pr. Miluna in Olympo (Form.); Attica: mt. Hymettus, Pentelicon (Heldr.); Achaia: pr. Megaspilaeon (Hal.). — Maio, Jun. ♃

9. **T. striatus** Vahl. symb. p. 78; Bald. riv. coll. bot. alb. 1895 p. 61, 1896 p. 85; Form. in Ver. Brünn 1896 p. 67. — *T. Chaubardi* Hal. Beitr. fl. Epir. p. 36, non Bois. et Heldr. — *T. paronychioides* Haussk. symb. p. 50 ex speciminibus in Sint. it. thessal. n. 462 distributis; non Celak. in Flora 1882 p. 564, qui foliis sat latis, supra longe hirsutis, bracteis maximis, calycis labio superiore ad quartam partem tantum fisso, dente medio lateralibus majore a planta thessala valde discedit. — Huc forsan spectat: *T. serpyllum v. angustifolius* Bois. fl. or. IV. p. 556 quoad pl. e mt. Dirphys Euboeae. — Exsicc.: Hal. it. gr. sec. a. 1893; Sint. it. thessal. n. 462 et 1519.

Rhizomate suffruticoso; caulibus procumbentibus, saepe elongatis, ramis floriferis erectis vel adscendentibus, undique hirtis; foliis viridibus, linearibus, obtusiusculis, basin versus breviter angustatis, punctatis, ciliatis, ceterum glaberrimis, subtus nervo validiori secundariisque parum prominulis instructis, floralibus ovato-lanceolatis, glabriusculis, ciliatis subtus plurinerviis; floribus breviter pedicellatis, in capitula globosa vel ovata congestis; calycis bilabiati, saepius rubentis, hirtuli, parce punctati labio superiore ad tertiam partem in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis, omnibus breviter pectinato-ciliatis; corollae roseae tubo calycis dentes non superante. — Praecedenti valde affinis, ab eo caulibus floriferis tenuioribus, foliis minus rigidis, basin versus non longe attenuatis, floralibus subangustioribus, calycis dentibus labii superioris latioribus, brevius acuminatis et corolla rosea discedit.

In herbidis regionis montanae et subalpinae. Epirus: mt. Maria distr. Ljaskovik, pr. Cepelovon, mt. Olycika, Strungula (Raul.), Tsumerka, Peristeri (Hal.); Thessalia: mt. Zygos (Haussk.), Wutades pr. Sermeniko (Sint.), Chaliki in Pindo, mt. Pelion (Form.). — Jun. Jul. ♃

×× Folia latiora, lineari-elliptica, oblonga vel ovata, vix rigida; calycis dentes labii superioris triangulari-lanceolati.

○ Rami teretiusculi; folia parva vel mediocria.

. Rami floriferi plus minus elongati; capitula involucrata.

10. **T. Sibthorpii** Benth. lab. p. 345; Haussk. symb. p. 50 (cum f. *spicata, capitata, grandiflora et parviflora*). — *T. lanceolatus* S. et S. pr. I. p. 419, non Desf. — *T. glabratus* Ch. et B. exp. p. 38, Fl. pelop. p. 172; Fraas fl. class. p. 178; non Hoffm. et Link. —

T. serpyllum Clem. sert. p. 75, non L. — *T. Chaubardi* Bois. et Heldr. pl. exs. a. 1851 n. 269; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Achaia p. 30, in ö. b. Z. 1897 p. 98 et 325; Haussk. symb. p. 50. — *T. angustifolius v. Chaubardi* Bois. et Heldr. diagn. ser. 2 IV. p. 6. — *T. serpyllum v. Chaubardi* Bois. fl. or. IV. p. 555; Heldr. chlor. Parn. p. 25. — Huc probabiliter: *T. pindicolus* Form. in Ver. Brünn 1896 p. 67. — Exsicc.: Heldr. herb. norm. n. 498 et 1368; Orph. fl. gr. n. 396; Sint. it. thessal. n. 303.

Rhizomate suffruticoso; caulibus procumbentibus, elongatis, ramis floriferis erectis vel adscendentibus, undique pubescentibus; foliis viridibus, ellipticis vel lineari-ellipticis, obtusis, punctatis, glabris, basin versus breviter ciliatis, subtus elevatim nervulosis, floralibus oblongo-ellipticis; floribus breviter pedicellatis, in capitula ovata vel oblonga, basi saepe laxiuscula congestis; calycis bilabiati, plus minus patule pilosi, punctati labio superiore ad tertiam partem in dentes triangulari-lanceolatos fisso, labii inferioris, superiori aequilongi dentibus lanceolato-subulatis, breviter ciliatis; corollae roseae tubo calycis dentes non superante. — *T. serpyllo* maxime affinis, sed omnibus partibus robustior, rami undique puberuli, folia elongata, floralia majora, capitula majora et formis extremis ad illum valde accedit, formae aliae angustifoliae contra transitum ad *T. striatum* praebere videntur.

In herbidis regionis montanae et subalpinae. Epirus: mt. Peristeri (Form.); Thessalia: mt. Zygos, Ghavellu, Karava, in oropedio Neuropolis, ad monasterium Korona, pr. Malakasi, Kalabaka, Pharsalus, Aivali (Haussk.), mt. Olympus, Pelion (Heldr.); Sporadum insula Scopelos (Leon.); Euboea: mt. Hagios Elias (Fraas); Aetolia: mt. Korax (Tunt.); mt. Kiona (Hal.), Parnassus, Cithaeron, Pateras (Heldr.), Parnes, Hymettus (Orph.); Peloponnesus (Chaub.): mt. Chelmos (Hal.), Taygetos (Heldr.); Cyclades (Chaub.). — Apr. Jul. ♃

11. **T. serpyllum** L. sp. p. 590; Fraas fl. class. p. 177; Bois. fl. or. IV. p. 555 *v. genuinus*; Bald. riv. coll. bot. alb. 1896 p. 85. — *T. longicaulis* Hal. Beitr. fl. Epir. p. 36, an Presl. — Exsicc.: Bald. it. alb. epir. IV. n. 41.

Rhizomate suffruticoso; caulibus prostratis, elongatis, radicantibus, ramis floriferis erectis vel adscendentibus, bifariam puberulis; foliis viridibus, ellipticis vel oblongis, obtusis, punctatis, glabris vel glabriusculis, basin versus parce ciliatis, subtus elevatim nervulosis, floralibus subconformibus; floribus breviter pedicellatis, in capitula globosa congestis; calycis bilabiati, pilosi, punctati, saepe colorati labio superiore ad tertiam partem in dentes triangulari-lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis, breviter ciliatis; corollae roseae tubo calycis dentes non superante. — Specimina omnia a me visa ramis bifariam pilosis a typo suecico recedunt, ceterum cum illo conveniunt.

In herbidis regionis subalpinae et alpinae rare. Epirus: mt. Olycika (Bald.), Tsumerka, Peristeri (Hal.); Thessalia: mt. Ghavellu (Heldr.); Euboea: mt. Dirphys (Bois.). — Jun. Jul. ♃

12. **T. lanuginosus** Mill. dict. n. 8. — Huc spectare videtur, quatenus ex frustulo incompleto dijudicare queo: *T. dolopicus* Form. in D. bot. Mon. 1897 p. 75, in Ver. Brünn 1897 p. 50. — Exsicc.: Orph. fl. gr. n. 524 (mt. Korfiati Macedoniae).

Rhizomate suffruticoso; caulibus adscendentibus, ramis elongatis, undique lanuginosis; foliis ellipticis vel oblongis, obtusiusculis, punctatis utrinque pilosis, canescentibus, subtus elevatim nervulosis, floralibus conformibus; floribus petiolatis, in verticillastra multiflora, distincta congestis, racemum saepius elongatum, inferne ramulosum formantibus; calycis bilabiati, villosi labio superiore ad tertiam partem in dentes triangulari-lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis; corollae roseae tubo incluso.

β. **Marschallianus** Willd. sp. III. p. 141; Form. in D. bot. Mon. 1891 p. 28; pro sp. — *T. serpyllum v. Marschallianus* Bois. fl. or. IV. p. 555. — Folia lineari-oblonga, acutiuscula, utrinque glabra, basin versus ciliata. — Exsicc.: ? Sint. et Bornm. it. turc. n. 1408 (specimina incompleta, omnino deflorata).

In collibus herbosis. Thessalia: mt. Ghavellu in Pindo (Form.), si re vera *T. dolopicus* huc spectat; — *β.* pr. Litochori (Sint.), pr. Plessidi et Portaria in mt. Pelion (Form.). — Jun. Aug. ♃

Obs. *T. heterotrichus* Griseb. spic. II. p. 116. — Species ab autoribus varie interpretata (cf. Benth. in DC. pr. XII. p. 199; Bois. fl. or. IV. p. 556; Borb. in math. term. Közl. XXIV. p. 61; Vel. fl. bulg. p. 469), mihi ignota; indicatur a Form. in Ver. Brünn 1896 p. 67, 1897 p. 50, in Thessalia: pr. Kalabaka, Pharsalus, pr. Phlamburo in mt. Oxya, pr. Metochi, Varlani, Vlachava, mt. Cuka, Phlambures et Mitrica in mt. Chassia, pr. Patsios, Tafilvris, Godaman, Nezeros in Olympo, pr. Longici, Pirgo, Alofovris, Mavrika, Katasara, Jurti in mt. Othrys, pr. Neraida.

.. Rami floriferi brevissimi; capitula foliis summis involucrata.

13. **T. ocheus** Heldr. et Sart. in Bois. diagn. ser. 2 IV. p. 6. — *T. serpyllum v. ocheus* Bois. fl. or. IV. p. 555. — Exsicc.: Sart. herb. a. 1847; Sint. et Bornm. it. turc. n. 851 b (Athos).

Rhizomate suffruticoso; caulibus prostratis, radicantibus, ramis floriferis brevissimis, undique patule hirsutis; foliis ovato-oblongis, obtusis, punctatis, setoso-villosis, cinerascentibus, subtus elevatim nervulosis, floralibus conformibus, imis capitulum involucrantibus; floribus breviter pedicellatis in capitula globosa congestis; calycis bilabiati, setuloso-villosi, punctati, rubentis labio superiore ad medium in dentes triangulari-lanceolatos fisso, labii inferioris, superiori aequilongi, dentibus subulatis; corolla roseae tubo calycis dentes superante. — Species pubescentia setulosa omnium partium, caulibus prostratis, capitulis involucratis insignis.

In summis mt. Hagios Elias — Ocha veterum — pr. Karisto

Euboeae Sart.; provenit quoque in cacumine mt. Athos peninsulae Hagion Oros. — Jun. Jul. ♃

○○ Rami tetragoni; folia majuscula, ad 15 mm. longa et 8 mm lata.

14. **T. montanus** W. et K. in Willd. sp. III. p. 143. — *T. serpyllum* r. *latifolius* Bois. fl. or. IV. p. 555. — Huc forsan: *T. pulegioides* S. et S. pr. I. p. 419. — Exsicc.: Sint. it. thessal. n. 948.

Rhizomate suffruticoso; caulibus prostratis, radicantibus, ramis elongatis, floriferis adscendentibus vel erectis, tetragonis, bifariam vel ad angulos reflexo-puberulis; foliis ovato-ellipticis oblongisve, acutiusculis, punctatis, glabris, viridibus, ad petiolum parce ciliatis, subtus elevatim nervulosis, floralibus conformibus; floribus pedicellatis, in capitula ovata, inferne saepe interrupta congestis; calycis bilabiati, glabriusculi, punctati, viridis labio superiore ad medium in dentes lanceolatos fisso, labii inferioris, superiori aequilongi dentibus subulatis, omnibus ciliatis; corollae parvae, roseae tubo calycis dentes non superante. — Foliis magnis, glabris viridibus insignis.

In subalpinis rarissime, hucusque tantum in mt. Sina pr. Malakasi in Pindo tymphaeo (Sint.). — Maio, Jul. ♃

2. Sectio. *Piperella* Willk. et Lange pr. fl. hisp. II. p. 404. — Calycis subgibbi labium superius late ovatum, ad quartam partem in dentes breviter et late triangulares, acutos, subaequales, laterales nempe medio subminores, divisum.

15. **T. teucrioides** Bois. et Spr. diagn. n. V. p. 15, Fl. or. IV. p. 554; Hal. in z. b. G. 1888 p. 761, Beitr. fl. Epir. p. 36, Beitr. fl. Thessal. p. 17, Heldr. chlor. Parn. p. 25; Bald. riv. coll. bot. alb. 1895 p. 61, 1896 p. 84; Form. in Ver. Brünn 1896 p. 68 subsp. *tymphaeus*; Haussk. symb. p. 50. — *T. graveolens* S. et S. pr. I. p. 421, Fl. gr. VI. p. 61 t. 576; Fraas fl. class. p. 178; non M. a. B. — Exsicc.: Heldr. herb. norm. n. 594; Bald. it. alb. epir. III. n. 192; Sint. it. thessal. n. 563 b.

Rhizomate suffruticoso; caulibus prostratis, ramis elongatis, adscendentibus, undique puberulis; foliis minutis, petiolatis, ovato-rhombeis, obtusiusculis, punctatis, glabriusculis, viridibus, margine revolutis, subtus nervo mediano valido nervulisque lateralibus obsoletis instructis, floralibus conformibus; floribus pedicellatis, in axillis superioribus 2—6, verticillastra remota formantibus; calycis bilabiati, basi subgibbi, glabri vel hirtuli, colorati, punctati labio superiore ad quartam partem in dentes late triangulares diviso, labii inferioris longioribus, lanceolati-subulatis, ciliatis; corollae magnae lilacinae tubo calycis dentibus longiore. — Facies *Calaminthae alpinae*.

In herbidis lapidosis regionis alpinae. Epirus: mt. Smolika, Vratedon (Bald.), Peristeri, Strungula, Tsumerka (Hal.); Thessalia: mt. Tringia (Hartl.), Zygos, pr. Malakasi, Uranaeos, Tsungeri, mt. Karava, Ghavellu (Haussk.), Dokimi, Said Pascha (Form.), in Pindo, mt. Oeta

(Sprun.); Aetolia: mt. Tymphrestus, Korax (Heldr.); mt. Kiona (Hal.); mt. Parnassus (Sibth.); Peloponnesus: mt. Chelmos (Heldr.). — Jul. Aug. ♃

Obs. *T. Billardieri* Bois. diagn. ser. 2 IV. p. 8, Fl. or. IV. p. 560 = *T. villosus* S. et S. pr. I. p. 422, Fl. gr. VI. p. 62 t. 578, non L. — Indicatur a Sibthorp in Archipelagi insulis, sed probabiliter erronee.

7. Tribus. **OCIMOIDEAE** Benth. lab. p. 1.

28. **Lavandula** L. gen. n. 711.

1. **L. stoechas** L. sp. p. 573; S. et S. pr. I. p. 399, Fl. gr. VI. t. 549; Sieb. avis p. 4, rem. p. 5, in Flora I. p. 270; Ch. et B. exp. p. 163, Fl. pelop. p. 36; Marg. et R. fl. Zante p. 72; Friedr. Reise p. 270; Fraas fl. class. p. 175; Weiss in z. b. G. 1869 p. 743; Raul. cret. p. 825; Bois. fl. or. IV. p. 540; Gelmi in bull. soc. bot. ital. 1889 p. 451; Bald. viagg. Creta p. 81; Hal. in ö. b. Z. 1897 p. 98; Haussk. symb. p. 37; Heldr. fl. Aegina p. 385. — Exsicc.: Orph. fl. gr. n. 273; Heldr. herb. fl. hellen. n. 60, in Baen. herb. europ. n. 3113; Baen. herb. europ. n. 9280; Dörfl. fl. aeg. n. 107.

Fruticulus erectus, ramis pubescentibus; foliis linearibus oblongo-linearibusve, pube minutissima incanis, margine revolutis, integris, axillis inferioribus fasciculigeris; spicis oblongis, densis, breviter pedunculatis; bracteis rhombeo-cordatis, acuminatis, tomentoso-pubescentibus, violascentibus, calyce sublongioribus, summis sterilibus, membranaceis, comosis, ellipticis vel oblongis, saepe denticulatis, violaceis; calyce ovato, incano; corolla atro-purpurea.

In collibus, dumosis regionis inferioris et submontanae. Attica: mt. Hymettus (Orph.); insula Aegina (Friedr.); Corinthia: pr. Kenada in isthmo (Fraas); Argolis: mt. Chelone in peninsula Methana, pr. Poros, insula Hydra (Heldr.); Laconia: pr. Sparta, Gythion (Heldr.); Sporadum insula Scopelos (Leon.); Cyclades: insula Tenos, Kimolos (Weiss), Melos, Naxos (Leon.); Creta: pr. Canea, Suda (Raul.), Gonia (Bald.), in fauce Serisso (Sieb.); Zante (Marg.); Corcyra: pr. Alippi (Gelmi), Pelleka (Baen.). — Mart. Jul. ♄

2. **L. spica** L. sp. p. 572; S. et S. pr. I. p. 399; Dallap. prosp. p. 81; Ch. et B. exp. p. 163, Fl. pelop. p. 36; Heldr. Nutzpfl. p. 32, Fl. cephal. p. 57. — Icon: Rchb. t. 26.

Fruticulus erectus, ramis pubescentibus; foliis oblongo-linearibus linearibusve, pube minuta incanis, margine revolutis, integris, adultis virentibus, subplanis; spicis cylindricis, interruptis, longe pedunculatis; bracteis ovato-rhombeis, acuminatis, membranaceis, pubescentibus, albicantibus, calyce brevioribus, summis conformibus; calyce tubuloso, amethystino-tomentello; corolla coerulea.

In locis petrosis regionis inferioris. In agro Laconico (Sibth.); Argolis, Cyclades (Chaub.); saepe colitur. — Jun. Aug. ♄ N. v.

Obs. *L. dentata* L. sp. p. 572; S. et S. pr. I. p. 399; Marg. et R. fl. Zante p. 72. — Indicatur in Zacyntho, at vix spontanea. — *Ocimum basilicum* L. sp. p. 597; Dallap. prosp. p. 85; Heldr. Nutzpfl. p. 32, Fl. cephal. p. 57, chlor. Thera p. 29; Form. in D. bot. Mon. 1891 p. 28. — Frequenter colitur in hortis.

8. Tribus. **MENTHOIDEAE** Benth. lab. p. 152.

29. Lycopus L. gen. n. 33.

1. **L. europaeus** L. sp. p. 21; S. et S. pr. I. p. 12; Mazz. in ant. ion. II. p. 444; Friedr. Reise p. 278 et 284; Bois. fl. or. IV. p. 545; Heldr. fl. cephal. p. 57; Form. in D. bot. Mon. 1891 p. 27. in Ver. Brünn 1897 p. 49; Haussk. symb. p. 51. — *L. exaltatus* Fraas fl. class. p. 179, non L. — Icon: Fl. dan. t. 1081.

Caule erecto, simplici vel ramoso, glabro vel leviter pubescente; foliis oblongis vel lanceolatis, grosse vel inciso-dentatis, glabris vel puberulis, in petiolum brevem attenuatis; floribus fasciculatis, in verticillastra axillaria congestis; corolla parva, alba; filamentis sterilibus minutissimis vel nullis.

β. **pubescens** Benth. lab. p. 186. — *L. mollis* Kern. in ö. b. Z. 1866 p. 371; Hal. in ö. b. Z. 1892 p. 400. — Caulis lanuginosus; folia ovata vel oblonga, utrinque canescentia. — Exsicc.: Sint. et Bornm. it. turc. n. 1426; Heldr. pl. fl. hellen. a. 1886 et 1895.

In paludosis, ad ripas, in humidis regionis inferioris. Thessalia: pr. Malakasi, Karditza (Form.), Trikala, Órman Magula (Haussk.), pr. Litochori ad Olympum (Sint.); Attica: in valle Cephissi pr. Athenas (Heldr.); Argolis: ad ripas Lernae, pr. Nauplia (Friedr.); Elis (Sibth.); Cephalonia: pr. Akoli, Livadi (Heldr.); Corcyra: pr. Kanale, Botumia (Mazz.). — Jul. Oct. ♃

2. **L. exaltatus** L. fil. suppl. p. 87; Mazz. in ant. ion. II. p. 444; Hal. in ö. b. Z. 1890 p. 41, 1896 p. 17; Form. in Ver. Brünn 1896 p. 66; Haussk. symb. p. 51. — *L. albus* Mazz. l. c. p. 446. — Icon: Fl. gr. I. p. 9 t. 12. — Exsicc.: Sint. it. or. 1889 n. 1887.

Caule erecto, ramoso, pubescente; foliis oblongis lanceolatisve, pinnatifidis partitisve, puberulis, in petiolum brevem attenuatis; floribus fasciculatis, in verticillastra axillaria congestis; corolla parva, alba; filamentis sterilibus acute capitatis.

In paludosis regionis inferioris. Thessalia: pr. Trikala, Karditza (Haussk.), Phanari (Heldr.), Palaeokastro (Reis.), Katerina (Sint.); Corcyra: ad lacum Corizia (Mazz.). — Jul. Oct. ♃

1 × 2. **L. europaeus × exaltatus** (*L. intermedius*) Haussk. symb. p. 51. — A *L. exaltato* foliis minus incisis, lobis brevioribus, latioribus, plus minus obtusis, nec longe acutatis diversa; notae aliae

in verticillastris, calycibus et filamentis intermediae dicuntur. — Inter parentes pr. Karditza Thessaliae (Haussk.). — N. v.

30. Mentha L. gen. n. 713.

1. Sectio. *Menthastrum* Coss. et Germ. fl. env. Paris p. 387. — Calycis faux nudus, dentes subaequales.

1. Subsectio. *Spicatae* L. sp. ed. 2 p. 804. — Verticillastra spicam conicam vel cylindricam, densam vel interruptam formantia; calyx campanulatus, obsolete 10 nervius; corolla intus glabra.

a. Folia ovato-rotundata, obtusa, crenata.

1. **M. rotundifolia** L. sp. p. 576 pro var. *M. spicatae*; L. sp. ed. 2 p. 805; S. et S. pr. I. p. 402; Ch. et B. exp. p. 164, Fl. pelop. p. 36; Friedr. Reise p. 264 et 275; Raul. cret. p. 825; Bois. fl. or. IV. p. 543; Haussk. symb. p. 50. — *M. rugosa* Lam. fl. fr. II. p. 420. — Icon: Rchb. t. 81. — Exsicc.: Heldr. herb. norm. p. 970.

Caulibus erectis, molliter pubescentibus vel lanato-tomentosis; foliis sessilibus, ovato-rotundatis, basi saepe subcordatis, crenatis, supra pubescentibus, subtus cinereo-vel incano-tomentosis; verticillastris numerosis, in spicam cylindrico-conicam densam dispositis; bracteis lanceolatis; calyce ovato-globoso, hirto, dentibus breviter lanceolatis; corolla alba rarius rosea. — Foliorum forma et indumento e pilis simplicibus ramosisque intermixto egregia.

In paludosis, ad ripas, fossas regionis inferioris, rare. Thessalia: pr. arcem urbis Volo (Haussk.); Boeotia (Sprun.); Peloponnesus (Sibth.): pr. Patras Achaiae (Friedr.), ad fontem Lernae pr. Myli (Heldr.) et pr. Poros (Friedr.) in Argolide (Chaub.); Creta (Sibth.): pr. Epanochorio (Raul.). — Jun. Oct. ♃

b. Folia ovato-lanceolata vel oblonga, acuta vel acuminata, acute serrata.

2. **M. longifolia** L. sp. p. 576 pro var. *M. spicatae*; Huds. fl. angl. p. 221. — *M. silvestris* L. sp. ed. 2 p. 804; S. et S. pr. I. p. 402; Pieri corc. fl. p. 76; Ch. et B. fl. pelop. p. 36; Friedr. Reise p. 275; Bois. fl. or. IV. p. 543; Heldr. chlor. Parn. p. 25, chlor. Thera p. 3; Bald. riv. coll. bot. alb. 1895 p. 60; Form. in Ver. Brünn 1896 p. 66, f. *stenotricha* Borb. in term. füz. IX. p. 24 (f. spica angusta, inferne interrupta, calyce saepe violaceo-suffuso).

Caulibus erectis vel adscendentibus, tomentellis; foliis sessilibus, ovato-lanceolatis oblongisve, acutis, acute serratis, subtus vel utrinque tomentosis; verticillastris numerosis, in spicam conicam, densam vel interruptam, conico-cylindricam dispositis; bracteis lineari-setaceis; calyce campanulato, dense hirto, dentibus subulatis; corolla lilacina. — Speciei quoad indumentum, foliorum magnitudinem, formam et serraturam, spicas breves, densas, crassiusculas vel elongatas, interruptas, angustas maxime polymorphae, habitu saepe diversissimae, sed intermediis innumeris con-

junctae. varietates sequentes, a menthologo quodam perito ulterius observandas, e Graecia mihi innotuerunt.

α. **typica**. — Folia elongato-lanceolata, 6—12 cm. longa, supra viridia, glabriuscula, subtus cano-pubescentia; spica crassiuscula, densa. — Exsicca: N. v.

β. **candicans** Crantz stirp. IV. p. 330 pro sp. — *M. silvestris* v. *candicans* Haussk. symb. p. 50. — Folia lanceolata, 5—7 cm. longa, supra pubescentia, viridia, subtus adpresse albo-tomentosa, spica crassiuscula, densa. — Exsicc.: Heldr. it. thessal. IV. a. 1883; Dörfl. fl. gr. n. 201.

γ. **mollissima** Borkh. fl. Wetter. II. p. 348; Form. in Ver. Brünn 1896 p. 65; pro sp.: Benth. lab. p. 171. — *M. incana* Sm. in Rees Cyclop. XXIII. sub Mentha n. 5. — *M. undulata* Willd. en. berol. VI. p. 608; Friedr. Reise p. 273; f. foliis undulatis. — Folia oblongo-lanceolata, 5—8 cm. longa, supra dense incano-pubescentia, subtus adpresse albo-tomentosa; spica crassiuscula, densa. — Variat (*M. Rocheliana* Borb. et Br. in z. b. G. 1890 p. 384; Form. in Ver. Brünn 1897 p. 49) spica elongata ad 9 cm. longa. — Exsicc.: Sint. it. thessal. n. 1295.

δ. **Wierzbickiana** Opiz in Flora 1824 p. 525; Form. in D. bot. Mon. 1891 p. 27. in Ver. Brünn 1895 p. 34, 1896 p. 65, 1897 p. 49. — Folia elliptico-lanceolata, 20—35 mm. longa, supra dense incano-pubescentia, subtus albo-tomentosa; spica elongata, 6—12 cm. longa, verticillastris 3—4 inferioribus, saepissime distantibus. — Exsicc.: Orph. fl. gr. n. 549; Sint. it. thessal. n. 987 et 1113.

ε. **minutiflora** Borb. in bot. Centralbl. 1886 p. 239; Form. in Ver. Brünn 1896 p. 65, 1897 p. 49. — Folia oblongo-lanceolata, 2—5 cm. longa, supra viridia, pubescentia, subtus cano-tomentosa; spica angustissima, virgata, verticillastris minutis distantibus, violaceis vel (*M. viridescens* Borb. in Békésvárm. fl. p. 74, Form. in Ver. Brünn 1896 p. 65, 1897 p. 49) viridibus. — Exsicc.: Sint. it. or. n. 1877; Sint. et Bornm. it. turc. n. 1417, 1418 et 1419.

ζ. **Sieberi** C. Koch in Linn. 1848 p. 649; Hal. Beitr. fl. Epir. p. 38, in z. b. G. 1899 p. 193; Form. in Ver. Brünn 1895 p. 34, 1896 p. 66. — *M. canescens* Sieb. avis p. 4, non Roth. — *M. tomentosa* Urv. enum. p. 67; Ch. et B. exp. p. 164, Fl. pelop. p. 36; Friedr. Reise p. 264; Fraas fl. class. p. 176; Clem. sert. p. 73; Raul. cret. p. 825; Heldr. fl. cephal. p. 57. — *M. silvestris* v. *stenostachya* Bois. fl. or. IV. p. 543; Bald. viagg. Creta p. 81; Haussk. symb. p. 51; Heldr. in ö. b. Z. 1898 p. 184. — Huc forsan: *M. niliaca* Link. in Linn. IX. p. 576; vix Jacq. hort. vind. p. 46 t. 3, quae hybrida inter *M. rotundifoliam et M. longifoliam*. — Folia ovato-lanceolata vel lanceolata, 2—5 cm. longa, reflexa, utrinque albo-tomentosa; spica angusta, elongata, verticillastris parvis, demum distantibus.

— Variat (*M. cretica* Portenschl. ap. Braun in z. b. G. 1890 p. 42; Form. in Ver. Brünn 1895 p. 34, 1896 p. 66, 1897 p. 49) foliis elongato-lanceolatis, margine breviter, saepe subundulato-serratis, spica breviore, verticillastris minus distantibus. — Huc spectat quoque: *M. ilissia* Heldr. herb. norm. n. 982, quae foliis lanceolatis, minus tomentosis et verticillastris minutis, valde distantibus transitum ad var. ε. manifeste praebet. — Exsicc.: Heldr. herb. norm. n. 64 et 971, 1574 v. *viridula* (f. foliis oblongis, tomento minus denso-obsitis, supra viridulis); Rev. pl. cret. n. 125 et 126, in Baen. herb. europ. n. 4886 et 5051, in Magn. fl. sel. n. 643; Bald. it. cret. n. 84.

Ad ripas, fossas, in paludosis, arvis derelictis regionis inferioris et montanae, formae microphyllae magis tomentosae in calidioribus. — Majo. Sept. ♃

Obs. Formam inter *M. rotundifoliam et M. Sieberi* intermediam ut videtur hybridam, caule pyramidatim-ramoso, foliis late ovatis, acutis, serratis, tomentoso-lanatis, spicis brevibus, verticillastris paucifloris, laxiusculis, calycis dentibus breviter lanceolatis, legit Reverchon pr. Platania Cretae (*M. Sieberi v. rotundifolia* Rev. pl. cret. a. 1883 sine n.). *M. Reverchoni* Hal. in herb. — Alia *M. pubescens* Willd. enum. berol. p. 608 (*mollissima* × *aquatica*) indicatur a Form. in D. bot. mon. 1891 p. 27 pr. Velitsena in Pindo.

3. **M. viridis** L. sp. p. 576 pro var. *M. spicatae*; L. sp. ed. 2 p. 804; Ch. et B. exp. p. 165, Fl. pelop. p. 36; Form. in Ver. Brünn 1895 p. 35 v. *graciliflora* Borb. in ö. b. Z. 1884 p. 169 (f. spica elongata, angusta, interrupta, floribus minoribus); Haussk. symb. p. 51. — Exsicc.: Heldr. it. thessal. IV. a. 1885.

Caulibus erectis, glabris vel sparse pilosulis; foliis sessilibus, lanceolatis vel oblongo-lanceolatis, acutis, acute serratis, glabris vel subtus ad nervos pilosulis; verticillastris numerosis, in spicam conico-cylindricam, inferne interruptam dispositis; bracteis lineari-setaceis; calyce campanulato, glabriusculo, dentibus lineari-subulatis; corolla lilacina. — Glabritie insignis.

In humidis, nemorosis. Corcyra: pr. Kanali (Form.): Epirus: pr. Janina (Form.); Thessalia: ad monasterium Korona in Pindo dolopico (Heldr.); Argolis et Cycladum insula Tenos (Chaub.). — Jun. Sept. ♃

Obs. *M. piperita* L. sp. p. 576; Fraas fl. class. p. 176; Heldr. Nutzpfl. p. 32. — A praecedente foliis petiolatis praesertim diversa, colitur hin inde in hortis.

2. Subsectio. *Capitatae* L. sp. ed. 2 p. 805. — Verticillastra in capitula terminalia congesta; calyx tubulosus, elevatim 13 nervius; corolla intus pilosa.

4. **M. aquatica** L. sp. p. 576; Fraas fl. class. p. 177; Clem. sert. p. 73; Bois. fl. or. IV. p. 544; Haussk. symb. p. 51.

Caulibus erectis, retrorsum hirsutis; foliis petiolatis, ovatis vel oblongis, acutis, serratis, glabratis vel pubescentibus; verticillastris 2—3

terminalibus, remotiusculis vel in capitulum terminale congestis; bracteis lanceolatis; calyce tubuloso, dentibus a basi triangulari subulatis; corolla bilacina.

α. **typica.** — Folia longiuscule petiolata, crenato-serrata, basi subcordata, utrinque sparsim pubescentia. — Exsicc.: Heldr. herb. a. 1886.

β. **hirsuta** L. mant. p. 81; Ch. et B. fl. pelop. p. 36; Hal. in ö. b. Z. 1892 p. 400; pro sp.; Benth. lab. p. 177; Friedr. Reise p. 264 et 274. — Huc forsan: *M. litoralis* Borb. ap. Form. in Ver. Brünn 1896 p. 66. — Folia breviter petiolata, acute serrata, basi subattenuata vel rotundata, utrinque adpresse pubescentia. — Exsicc.: Sint. et Bornm. it. turc. n. 1416.

γ. **crispa** L. sp. ed. 2 p. 805; S. et S. pr. I. p. 403; pro sp.; Benth. lab. p. 177. — Folia breviter petiolata, crispato-plicata, lacerodentata, pubescentia. — Exsicc : N. v.

In humidis regionis inferioris. Thessalia: pr. Krania in Pindo (Form.), pr. Trikala, Palaeokastro, Karditza (Haussk.), Litochori ad Olympum (Sint.); Boeotia: ad Cephissum (Fraas); Attica: pr. Athenas (Clem.), ad Cephissum pr. Podoniphti (Heldr.); Elis (Heldr.); Achaia: pr. Patras (Daenzer); Argolis: ad ripas Erasini pr. Argos (Heldr.), pr. Poros (Friedr.); — γ. in agro Laconico (Sibth.). — Jul. Sept. ♃

2. Sectio. *Pulegium* Lam. fl. fr. III. p. 537. — Calycis faux annulo pilorum clausa, dentes subbilabiati, inferioribus binis angustioribus.

5. **M. pulegium** L. sp. p. 577; S. et S. pr. I. p. 404; Dallap. prosp. p. 82; Ch. et B. exp. p. 165, Fl. pelop. p. 36; Marg. et R. fl. Zante p. 73; Friedr. Reise p. 264; Clem. sert. p. 73; Bois. fl. or. IV. p. 545; Heldr. fl. cephal. p. 57; Form. in D. bot. Mon. 1891 p. 27, in Ver. Brünn 1895 p. 35, 1896 p. 66; Haussk. symb. p. 51.

Caulibus erectis vel adscendentibus, simplicibus vel ramosis; foliis breviter petiolatis, ovatis vel oblongis, acutis vel obtusis, minute et obscure denticulatis, floralibus conformibus; verticillastris numerosis, globosis, remotis; calyce tubuloso-infundibuliformi, dentibus lanceolato-subulatis; corolla rosea.

α. **typica.** — Caulis et folia glabra vel parce puberula, viridia; calyx puberulus. — Variat (*M. erinoides* Heldr. chlor. Mykon p. 249) caulibus prostratis, foliis minutis suborbiculatis, verticillastris paucifloris. — Exsicc.: Heldr. pl. fl. hellen. a. 1887.

β. **pulegioides** Sieb. avis rem. p. 5; pro sp. — *M. pulegium v. subtomentella* Braun in z. b. G. 1890 p. 144; Hal. Beitr. fl. Epir. p. 38. — *M. pulegium v. villicaulis* Borb. ap. Form. in D. bot. Mon. 1891 p. 27, in Ver. Brünn 1896 p. 66. — Caulis dense adpresse vel patule hirsutus, folia patule hirsuta, canescentia; calyx hirsutus. — Variat (*M. cephalonia* Braun in z. b. G. 1890 p. 144, Form. in Ver. Brünn 1895 p. 35) foliis minoribus late ovatis vel subrotundis, verti-

cillastris minoribus. — Exsicc.: Rev. pl. cret. n. 127, in Magn. fl. sel. n. 649; Sint. et Bornm. it. turc. n. 1415; Sint. it. thessal. n. 1249.

γ. **tomentella** Hoffm. et Link fl. port. I. p. 73; Link in Linnaea IX. p. 576; Fraas fl. class. p. 177; Hal. in z. b. G. 1888 p. 761; Bald. in nuov. giorn. bot. ital. 1894 p. 99; Heldr. chlor. Mykon p. 249; pro sp.; Marg. et R. fl. Zante p. 73; Raul. cret. p. 825; Spreitz. in z. b. G. 1897 p. 666. — *M. gibraltarica* Willd. en. berol. p. 611. — *M. pulegium v. tomentosa* Vis. fl. dalm. II. p. 185; Haussk. symb. p. 51. — *M. pulegium v. villosa* Benth. in DC. pr. XII. p. 175; Weiss in z. b. G. 1869 p. 744. — Caulis, folia et verticillastra dense albo-villosa. — Exsicc.: Bald. it. bot. penins. balk. a. 1889.

Ad ripas, in arvis humidis, ruderatis regionis inferioris et montanae totius Graeciae, β. et γ. typo multo vulgatiores. — Jul. Oct. ♃

Index

ordinum, generum, specierum, varietatum et synonymorum.

A.

pag.

Absinthium petrosum Baumg. . . 74
Acanthaceae Juss. . . . 460
Acanthus L. 460
Caroli Alexandri Haussk. 460
v. minor Haussk. 460
v. tenuissimus Haussk. . . 460
longifolius Host . . . 460
mollis Dall. 461
spinosissimus Pers. . . . 461
spinosus L. 460
v. thessalus Form. . . . 461
spinulosus Friedr. 461
spinulosus Host 461
Acarna cancellata All. 99
gummifera Willd. 94
Achillea L. 40
abrotanoides Vis. . . . 44
absinthoides Hal. . . . 48
aegyptiaca L. 50
v. taygetea B. et H. . . 50
ageratifolia Bois. . . . 41
v. aizoon Bois. 41
v. euageratifolia Heim. . . 41
ageratum L. 49
aizoon Griseb. 41
ambrosiaca Bois. . . . 43
Barbeyana Heldr. et Heim. 43
Boissieri Haussk. 43
Boissieriana Heldr. 43
capitata Willd. 42
chrysocoma Friv. . . . 50
Clavennae L. 42
v. capitata Heim. 42
v. depauperata Hal. . . . 42
v. integrifolia Hal. . . . 42
v. pindicola Haussk. . . . 42

pag.

Clavennae × Fraasii Hal. 42
clypeolata S. et S. . . 49
coarctata Poir. 49
compacta Will. 49
compacta × odorata Haussk. 50
cretica L. 51
crithmifolia W. et K. . 47
dolopica Fr. et Sint. . . . 47
fililoba Freyn. 47
flabelliformis S. et S. . . . 48
Fraasii Schultz 44
glomerata M. B. 49
grandifolia Friv. . . . 45
grandifolia × umbellata Heim. 43
holosericea S. et S. . . 48
Kerneri Hal. 42
ligustica All. 45
magna L. 49
magna S. et S. 45
major Heim. 43
micrantha M. B. 51
millefolium L. 47
v. setacea Koch. 47
Millii Heldr. 43
moschata Wulf. 44
v. olympica Heim. . . . 44
Neumayeri Heldr. 44
nobilis L. 46
v. Neilreichii (Kern.) . . . 46
odorata L. 46
v. lacmonica Haussk. . . 46
v. virescens Fenzl. . . . 46
odorata × coarctata Hal. 50
olympica Heim. 44
pallescens DC. 45
pannonica Form. 47
pubescens S. et S. 51
punctata Ten. 46

pag.
santolina S. S. 51
sericea Janka 49
setacea W. et K. . . . 47
silvatica Ten. 45
v. filifolia Bois. 47
v. subvelutina DC. . . . 45
taygetea B. et H. 50
tomentosa Ch. et B. . . . 50
tomentosa Fraas 48
tomentosa L. 51
Tournefortii DC. 51
tymphaea Haussk. 50
umbellata S. et S. . . 42
v. major Bois. 43
v. monocephala Heldr. . . 43
v. pauciloba Heldr. . . . 43
velutina Desf. 49
Achyrophorus cretensis Bois. . . 174
serioloides DC. . 174
Acinos alpinus Moench. 543
graveolens Link 546
v. integrifolia Raul. . . . 546
Aegialophila B. et H. 167
cretica B. et H. . . . 167
pumila (L.) 167
Ajuga L. 466
chamaepitys Schreb. . . 467
chia Schreb. 468
v intermedia B. et O. . . 468
Halácsyana Zahlbr. . . . 468
iva Schreb. 467
v. pseudoiva Benth. . . . 467
orientalis L. 466
v. aenesia Heldr. 466
Piskoi Deg. et Bald. . . 468
pseudoiva Rob. et Cast. . . 467
reptans L 466
Alectorolophus Hall. 441
alpinus Fraas 441
glandulosus Haussk. . . . 441
minor Stern. 441
pindicus Stern. 441
pubescens Stern. . . . 441
Sintenisii Stern. . . . 442
Alkanna Tausch 342
boeotica DC. 344
v versicolor Bois. . . . 344
calliensis Heldr. . . . 342
graeca Baen. 345
graeca Bois. et Spr. . . 344
v. hispidior Bois. 344
v. versicolor Bois. . . . 344
graeca Hal. 343

pag.
graeca Ung. 344
Matthioli Tausch. 346
methanaea Haussk. . . 343
orientalis Bois. 342
v. hellenica Bois. 342
pindicola Haussk. . . . 343
v conferta Haussk. . . . 343
pulmonaria Form. 343
pulmonaria Hal. 343
Sartoriana B. et H. . . 345
scardica Form. 343
Sieberi DC. 446
Stribrnyi Vel. 345
v pelia Hal. 345
tinctoria Tausch. . . . 346
v. incana Weiss. 346
v. Lehmani Tin. 346
Amaracus dictamnus Benth. . . 552
pulcher Briqu. . . . 553
scaber Briqu. 553
Tournefortii Benth. . . 553
Amberboa lancifolia DC. 165
Ambrosia L. 246
maritima L. 246
Ambrosiaceae Cass. . 246
Anacyclus Pers. . . . 65
Candollei Nym. 65
clavatus Pers. 65
creticus L 59
orientalis L. 56
pectinatus Ch. et B. . . . 58
pyrethrum DC. 65
radiatus Lois 65
tomentosus DC. 65
valentinus L. 65
Anarrhinum bellidifolium Dsf. . . 420
Anchusa L. 321
Adami Mazz. 322
aegyptiaca DC. 327
aggregata Lehm. . . . 328
angustifolia L. 323
angustifolia S. et S. . . . 324
arvensis M. B. 329
aspera Bois. 326
bulbosa Mazz. 331
caespitosa Lam. . . . 328
Cesatiana Fenzl. 331
Gmelini Led. 324
hybrida Ten. 324
v. setigera Hal. 325
italica Retz 326
leiosperma Fauch et Ch. . . 321

pag.
linearifolia Urv. 324
lutea Friedr. 344
macrocalyx Haussk. . . . 323
micrantha R. et Sch. . . . 328
obliqua Vis. 324
ochroleuca M. B. . . . 324
v canescens Bois. . . . 326
officinalis Bois. 323
officinalis L. 323
v macrocalyx (Haussk.) . 323
orientalis L. 342
ovata Mazz. 329
paniculata Ait. 326
parnassica B. et O. . . 323
parviflora S. et S. 328
Sartorii Heldr. 326
sempervirens L. . . . 322
Spruneri Bois. 328
stylosa M. B. 328
v. major DC. 328
thessala B. et Spr. . . . 327
tinctoria L. 346
undulata L. 325
v. maritima Heldr. . . . 326
variegata Lehm. . . . 329
ventricosa S. et S. . . . 321
Andryala L. 245
dentata S. et S. . . . 245
integrifolia Urv. 245
lanata S. et S. 239
Anthemis L. 51
absinthifolia B. et Spr. . . 57
ageratifolia S. et S. . . . 41
aizoides B. et O. 41
aizoon Grsb. 41
altissima L. 54
arvensis L. 59
v. incrassata Bois. . . . 60
v. microcephala Heldr. . . 60
auriculata Bois. 63
austriaca Form. 60
Brachmanni B. et H. . 55
brachycentros Gay . . 54
brachyglossa C. Koch . . 57
carpathica Haussk. 56
chia L. 63
v. conica Bald. 63
chrysocephala Heldr. . . . 56
clavata Desf. 65
complanata Hal. 58
cota Koch 54
cota L. 54
cota S. et S. 64

pag.
cotula L. 64
cretica Nym. 59
cronia B. et H. 56
discoidea Willd. 53
fuscata Ch. et B. 63
Gerardiana Jord. 55
graveolens Bois. 58
Guicciardii Heldr. et Sart. 62
incana Bois. 56
incrassata Bois. 60
maritima S et S. 53
melampodina Ung. 63
meteorica Haussk. . . 58
misella Heldr. et Sart. . . 60
mixta L. 64
montana L. 55
v. angustisecta Hal. . . . 56
v. breviloba Hal. 56
v. cronia Bois. 56
v. gracilior Heldr. 56
v. incana Bois. 56
v. Linnaeana G. et G. . . 55
v. Linnaeana Hal. 57
v. macedonica Grisb. . . 55
v. olenaea Hal. 56
v. pentelica Bois. 56
v. saxatilis Heldr. . . . 55
v. Spruneri (B. et H.) . . 56
v. tenuiloba Bois. 56
v. thracica Grisb. 55
Muenteriana Hal. 62
Muenteriana Heldr. 61
multicaulis B. et H. . . . 60
Neilreichii Ortm. 60
nobilis L. 64
panachaica Hal. 54
parnesia B. et H. . . . 60
pectinata B. et R. . . 58
pedunculata Ch. et B. . . 65
peregrina L. 62
v. Guicciardii Bois. . . . 62
v. heracleotica B. et H. . . 62
v. platyloba Haussk. . . . 62
pindicola Heldr. . . . 57
pontica Willd. 55
pseudocota Vis. 54
psorosperma Ten. 64
pyrethrum Fraas 53
pyrethrum L. 65
ruthenica M. B. . . . 60
saxatilis DC. 55
Spruneri B. et H. 57
taygetea B. et H. 58
tenuiloba DC. 56
tinctoria L. 52

pag.
v. discoidea Vahl 53
v. pallida DC. 53
v. parnassica Bois. . . . 53
v. Triumfetti L. 53
tomentosa Hal. 62
tomentosa L. 61
Triumfetti Bald. 53
Urvillei Nym. 59
valentina L 65
Visianii Weiss. 63
Antirrhinum L. . . . 418
aegyptiacum S. et S. . . . 416
altissimum Sieb. 420
arvense L. 413
asarina L. 420
bellidifolium L. 420
calycinum Lam 419
chalepense L. 410
cirrhosum L. 416
cymbalaria L. 446
dalmaticum L. 408
elatine S. et S. 414
genistifolium L. 408
graecum Ch. et B. 415
lanigerum Ch. et B. . . . 414
linaria L. 412
litorale Bernh. 418
majus L. 419
v. angustifolium Chav. . . 419
micranthum Cav. 413
minus L. 418
orontium L. 418
v. grandiflorum Chav. . . 418
parviflorum Jacq. 413
parviflorum Willd. 413
Pelisserianum L. 412
pilosum Ch. et B. 417
purpureum L. 411
reflexum L 411
siculum Ucr. 419
simplex Willd. 413
spurium L. 414
strictum S. et S. 410
triphyllum L. 410
Ammanthus B. et H. . 71
filicaulis B. et H. . . . 71
maritimus B. et H. . . 71
Apargia aspera W. et K. . . . 186
cichoracea Ten. 185
fasciculata Biv. 185
hastilis Host 186
hirta Sm. 184
hispida Host 185
hyoseroides Sieb. 188
saxatilis Ten. 187
strigosa M. B. 186

pag.
tuberosa Willd. 189
Apocynaceae R. Br. . . 293
Arbutus L. 283
andrachne L. 283
andrachne × unedo Haussk. 284
andrachnoides Link. . . . 284
hybrida Ker. 284
integrifolia Lam. 282
intermedia Heldr. 284
nothocomaros Heldr. . . . 284
Sieberi Klotsch 283
unedo L. 283
v. turbinata Pers. . . . 283
unedo × andrachne Bois. 284
Arctium lappa L. 101
majus Bernh. 101
minus Bernh. 102
Arnebia cephalotes DC. 341
cornuta Fisch. et M. . . 336
Arnica cordata Wulf. 76
scorpioides S. et S. . . 76
Arnopogon Dalechampii W. . . 191
picroides W. 191
Artemisia L. 72
abrotanum L. 74
absinthium Dall. 72
absinthium L. 72
v montana Form. . . . 72
arborescens L. 72
Baumgartenii Bess. . . . 74
campestris L. 74
camphorata Vill. . . . 73
eriantha Ten. 74
petrosa (Baumg.) . . . 74
pontica L. 75
scoparia W. et K. . . 74
spicata S. et S. 74
v. eriantha DC. 74
Villarsii G. et G. 74
vulgaris L. 72
Asclepiadaceae R. Br. 290
Asclepias canescens W. 290
Dioscoridis Fraas . . . 291
fruticosa L. 293
nigra L. 291
vincetoxicum Dall. . . 292
vincetoxicum S. et S. . 291
Asperugo L. 361
aegyptiaca L. 327
procumbens L. 361
Aster L. 16
alpinus Bois. 16
alpinus L. 16

pag.
amellus L. 16
cylleneus B. et O. . . . 16
tripolium L. 16
Asteriscus Moench . . 17
aquaticus Less. 17
v. discoideus Weiss . . . 17
v. nanus Bois. . . . 17
citriodorus Heldr et Hal. 18
maritimus Less. . . . 17
Asterothrix asperrima Cass. . . 186
Athanasia maritima L. 39
Atracty is L. 99
cancellata L. 99
gummifera L. 94
Atropa L. 368
belladonna L. 368
mandragora L. 366
mandragora S. et S. . . . 367

B.

Ballota L. 532
acetabulosa Benth. . . 532
alba L. 532
foetida Lam. 532
hirsuta Nym. 532
hispanica Benth. 532
italica Benth. 532
nigra L. 532
pseudodictamnus Benth. . . 533
rupestris Form. 532
rupestris Vis. 532
saxatilis Guss. 532
Barbellina sericea Cass. 100
Bartsia latifolia S. et S. 436
trixago L. 440
versicolor Pers. 440
viscosa L. 437
Batatas edulis Chois. 302
litoralis Chois. 302
Barkhausia foetida DC. 227
hispida Link . . . 228
purpurea Ung. . . . 226
rhoeadifolia M. B. . 227
rubra Moench . . . 226
scariosa Willd. . . . 229
setosa DC. . . . 228
vesicaria Spreng. . . 228
zacynthia M. et R. . 228
Bellardia All. 440
trixago All. 440
Bellis L. 12
amoena Heldr. 12
annua L. 13

pag.
v. dentata Clem. 13
v. minuta DC. 13
dentata DC. 13
hybrida Ten. 12
longifolia Bois. 12
longifolia B. et H. . . 13
v. glabrata Raul. 13
perennis L. 12
v. microcephala Bois. . . 12
perennis × silvestris Haussk. 12
silvestris Cyr. 13
v. glabrescens Heldr. . . 13
Bellium L. 14
bellidioides S. et S. . . . 14
minutum L. 14
Beringeria acetabulosa Neck. . . 533
pseudodictamnus Neck. 533
Betonica L. 514
alopecurus L. 515
annua L. 530
graeca B. et Spr. 515
v. petiolata Form. . . . 515
v. tymphaea Form. . . . 515
Haussknechtii Uechtr. . 514
Jacquini G. et G. 515
officinalis Form. 514
officinalis L. 514
scardica Grisb. 515
Bidens L. 39
cernua L. 39
bipartita L. 39
Borraginaceae Juss. . . 313
Borrago L. 320
cretica Willd. 320
officinalis L. 320
Brotera corymbosa W. 92
Buphthalmum aquaticum L. . . 17
maritimum L. 17
spinosum L. 18

C.

Cacalia verbascifolia S. et S. . . 80
Calamintha Tourn. . . 540
acinos (L.) 546
adscendens Form. 542
alpina Lam. 543
v. major (Haussk.) . . . 544
v. meridionalis Haussk. . 544
v. nebrodensis Kern. et Str. 544
v. pentelica Haussk. . . . 544
Boissieri Briqu. 543
clinopodium Benth. . . 540
cretica Benth. 543

pag.
exigua (S. et S.) . . . 546
grandiflora Moench. . . 540
graveolens Benth. 546
incana Bois. 542
v calvescens Heldr. . . . 543
incana Clem. 542
meridionalis Nym. 544
nebrodensis Kern. et Str. . 544
nepeta Savi 542
nepetoides Jord. . . . 541
officinalis Hal. 541
officinalis Moench . . . 541
patavina Host. 545
v. major Haussk. 544
Spruneri Bois. 542
suaveolens Bois. . . . 544
v. canescens Haussk. . . 544
v. meteorica Haussk. . . 545
v. viridis Heldr. et Haussk. 545
suaveolens Hal. 545
thessala Haussk. . . . 541
Calcitrapa galactites Lam. . . . 126
Calendula L. 86
aegyptiaca Desf. . . . 87
arvensis L. 86
v. bicolor DC. 87
v. pygmaea Clem. . . . 87
v. rugosa Vis. 87
bicolor Raf. 87
micrantha Tin 87
officinalis Pieri 87
parviflora Friedr. 87
sicula Friedr. 87
Calystegia sepium R. Br. . . . 302
silvatica Chois. . . . 302
silvestris R. et Sch. . 302
soldanella R. Br. . . 303
Campanula L. 249
abietina Grisb. et Sch. . 267
aizoon B. et Spr. . . . 269
anchusiflora S. et S. . 254
Andrewsii DC. 255
asperuloides Orph. 278
athoa B. et H. 261
attica B. et H. 266
bicaulis Lap. 273
bononiensis L. 262
calaminthifolia Lam. . 258
v Olivieri (DC.) 259
capitata Sims. 257
carpatha Hal. 252
Celsii DC. 254

pag.
cephallenica Feer. . . . 264
cichoracea S. et S. . . . 257
corymbosa Desf. . . . 251
dichotoma L. 259
drabifolia S. et S. . . 266
eriantha Hampe 254
erinus L. 266
expansa Friv. 271
v. sphaerothrix Bois. . . 271
expansa Hal. 272
falcata R. et Sch. 275
flagellaris Hal. 260
foliosa Form. 259
foliosa Ten. 260
Frivaldszkyi Steud. . . . 271
garganica Bois. 264
garganica Ten. 264
glomerata Bald. 261
glomerata L. 259
v. stenosiphon Bois. . . . 260
v. Vlachavae B. et O. . . 259
graminifolia L. 279
Halácsyana Bald. 264
Hawkinsiana Haussk. et Heldr. 264
heterophylla L. 258
hybrida L. 275
Jacquini DC. 278
incurva Auch. 256
laciniata Andr. 255
laciniata L. 253
Lambertiana DC. 270
lanuginosa W. 255
Leutweinii Heldr. 256
lilifolia L. 273
limonifolia L. 277
lingulata W et K. . . . 257
v cichoracea Grsb. . . . 257
lyrata Lam. 256
lyrata Link. 254
mollis L. 273
muscosa Mazz. 273
nutans Sieb. 248
Olivieri DC. 259
oreadum B. et H. . . . 257
parnassica Bald. 261
parnassica B. et Spr. . . . 279
patula L. 270
v. chassia Form. 270
v. pauciflora Roch. . . . 267
patula S. et S. 268
pauciflora Desf. 273
pauciflora Lam. 267
pelviformis Lam. . . . 251

pag.
v. micrantha DC. 252
pentagonia L. 273
persicifolia L. 267
phrygia Jaub. et Sp. . . . 272
Pichleri Vis. 276
pruinosa Mazz. 273
radicosa Ch. et B. . . 265
ramosissima S. et S. . 272
rapunculoides L. . . . 262
rapunculus L. 270
v Lambertiana Haussk. . 270
Reiseri Hal. 253
v. Leonis Hal. 254
Rochelii Schur 267
rotundifolia L. 263
rumeliana Vatke 278
rupestris S. et S. . . . 255
v. calycina Heldr. . . . 256
v. gracilis Heldr. 256
v. virescens Hal. 256
rupicola B. et Spr. . . 258
Sartorii B. et S. . . . 264
saxatilis L. 252
v. Simonellii Bald. . . . 253
scutellata Grisb. . . . 265
Sibthorpiana Hal. . . . 268
v filicaulis Hal. 269
sparsa Friv. 271
spathulata Friedr. 268
spathulata S. et S. . . . 268
speciosa Pourr. 273
speculum L. 274
sphaerothrix Grsb. . . 271
v. oxya Hal. 272
Spruneri Raul. 268
v. alpina Bois. 269
Spruneriana Hampe . . 268
v hirsuta Heldr. 268
stenosiphon B. et H. . 260
v. majoriflora Hal. . . . 260
stricta S. et S. 256
tomentosa Bois. 255
tomentosa Hal. 255
tomentosa Vent. 254
v. brachyantha Bois. . . . 255
v. bracteosa Heldr. . . . 255
v. calycina Heldr. . . . 256
v. diffusa Heldr. 254
trachelium L. 262
v. balcanica Form. . . . 261
v. orientalis Bois. . . . 261
v. parviflora Form. . . . 261
trachelium Mazz. 261

pag.
trichocalycina Ten. . . . 276
tubulosa Heldr. 252
tubulosa Lam. 252
tymphaea Haussk. . . . 260
uniflora L. 273
urticaefolia Mazz. 261
versicolor Andr. . . . 262
v. thessala Bois. 262
v. tomentella Hal. . . . 262
virgata Lodd. 277
Welandii Heuff. 271
Campanulaceae Juss. . 247
Capsicum annuum L. 366
grossum W. 366
longum DC. 366
Cardopatium Juss. . . 92
Boryi Spach 92
corynbosum Pers. . . 92
orientale Spach 92
Vrionis Heldr. 92
Carduncellus Ad. . . 170
coeruleus DC. 170
v. dentatus DC. 170
v. tingitanus (L.) 170
v. incisus DC. 170
Carduus L. 102
acanthoides L. 104
v. thessalus B. et H. . . 104
v. tymphaeus Form. . . . 105
acarna L. 119
acicularis Bert. 107
afer Jacq. 117
albidus M. B. 106
arabicus Friedr. 106
argentatus L. 107
argyroa Biv 107
armatus B. et H. . . . 104
v. cronius B. et H. . . . 104
armatus Hal. 105
benedictus Camer. 171
candicans W. et K. . . 105
Casabonae Ch et B. . . . 117
collinus Form. 105
creticus Lam. 116
crispus L. 105
cynaroides Lam. 116
ferox Ch. et B. 109
fruticosus Dsf. 127
galactites Ch. et B. . . . 126
glycacanthus S. et S. . . . 171
Halácsyi Heldr. 107
hamulosus Ehrh. . . . 105
hamulosus × pindicolus Haussk. 105

pag.
Hippolyti Ch. et B. . . . 116
intercedens Haussk. . . . 105
italicus Savi 112
lanceolatus L. 112
leiophyllus Petr. 102
v. vestitus Hal. 102
leucographus L. 125
macrocephalus Hal. . . . 103
marianus L. 124
marmoratus B. et H. . 107
v. Halácsyi (Heldr.) . . . 107
v. intermedius Hal. . . . 107
mollis L. 127
neglectus Ten. 107
nutans Bois. 103
nutans Hal. 103
nutans L. 102
v. brachycentros Haussk. . 103
v. sporadum Hal. 103
v. taygeteus Heldr. . . . 103
pindicolus Haussk. 102
polyanthus Ch. et B. . . . 116
pycnocephalus L. . . . 106
v. albidus Bois. 106
rivularis Jacq. 118
seminudus Ch. et B. . . . 125
stellatus L. 118
syriacus L. 119
taygeteus B. et H. . . 103
v. parnassicus Hal. . . . 103
tenuiflorus S. et S. . . . 106
v. albidus Friedr. 106
v. pycnocephalus Friedr. . 106
Carlina L. 94
acanthifolia All. 94
acanthophylla Haussk. . 96
acaulis Dall. 94
corymbosa L. 97
v. gracilis Form. 98
v. graeca Bois. 97
v. involucrata Bois. . . . 98
v. Rothii Bois. 98
corymbosa S. et S. . . . 97
curetum Heldr. 98
dolopica Form. 96
elegans Heldr. 97
frigida B. et H. . . . 96
graeca Heldr. et Sart. . 97
v. actinobola Heldr. . . . 98
v. platyrrhachis Hal. . . 98
v. stenorrhachis Hal. . . 98
gummifera Less. . . . 94
involucrata Poir. 98
lanata L. 99
rigida Form. 95
v. humilis Form. 95
v. othryana Form. . . . 95
v. pallida Form. 95
v. sordida Form. 95
Rothii Heldr et Sart. . 98
semiamplexicaulis Form. . 96
simplex W. et K. 95
v. thessala Form. 95
utzka Hacq. 94
vulgaris Bois. 96
Carthamus L. 167
ambiguus Heldr. . . . 169
Boissieri Hal. 168
coeruleus L. 170
corymbosus L. 92
creticus L. 168
cycladum Heldr. 168
dentatus Bois. 169
dentatus Urv. 169
dentatus Vahl. 169
glaucus Bois. 168
v. syriacus Bois. 168
lanatus L. 167
v. creticus (L.) 168
v. graecus Heldr. 168
leucocaulos S. et S. . . 168
ruber Bald. 168
ruber Link. 169
Sartorii Heldr. 160
tinctorius L. 179
tingitanus L. 170
Caryolopha sempervirens F. et M. 322
Catananche Vaill. . . 172
graeca Ch. et B. 173
graeca L. 172
lutea L. 172
Celsia L. 395
acaulis B. et Ch. . . . 398
arcturus Murr. 395
Boissieri Heldr. et Sart. 397
cretica L. 395
cyllenea B. et H. . . . 397
Daenzeri Heldr. et Sart. 396
v. lyrata B. et H. . . . 397
orientalis L. 398
speciosa Fenzl 396
sublanata Jacq. 396
tomentosa Zucc. 393
Centaurea L. 129
achaia B. et H. 153
v. corinthiaca Bois. . . . 153

pag.
v. ferox B. et H. 153
v. submutica Bois. . . . 153
Adami Willd. 161
aegyptiaca S. et S. . . . 163
affinis Haussk. 145
affinis Friv. 146
v. denudata Hal. 146
v. lacerata Haussk. . . . 145
v. pallidior Bois. 146
v. peloponnesiaca Hal. . . 146
affinis × Grisebachii Haussk. 147
alba L. 136
v. deusta DC. 136
v. princeps Bois. 135
albanica Hal. 134
albida Heldr. 140
albiflora C. Koch 140
amara L. 134
amplifolia B. et H. . . 133
apula Lam. 162
argentea L. 141
armoracifolia S. et S. . . 165
asperula Hal. 144
atropurpurea Oliv. 152
attica Nym. 144
v. pateraea Hal. 144
axillaris Ch. et B. 139
v. angustifolia Form. . . 139
v. cana Bois. 139
Baldaccii Deg. 140
benedicta L. 171
Biebersteinii Form. . . . 147
Boissieri Walp. 144
brevispina Form. 148
brevispina Haussk. . . 148
v fusconigra Haussk. . . 148
brevispina × pelia Haussk. . 148
brunnea Hal. 135
cadmea Bois. 134
calcitrapa L. 162
v. longispina Asch. . . . 163
calcitrapoides DC. 163
cana S. et S. 139
v. albiflora Raul. 140
v. pindicola Bald. . . . 139
v. pindicola Grisb. . . . 140
corduncella DC. 158
Ceccariniana B. et H. . . 155
v. brachycentra Bois. . . 155
Ceccariniana Haussk. . . . 155
centauroides S. et S. . . . 159
Charrelii Hal. et Dörfl. . . 160
chrysolepis Vis. 160
ciliata Frid. 142

pag.
cineraria Sieb. 141
collina S. et S. 159
confusa Hal. 147
cordonis B. et O. 155
corinthiaca B. et H. . . . 153
coronopifolia S. et S. . . . 165
crupina L. 166
crupina S. et S. 166
cyanoides Berg. 139
cyanus L. 138
dalmatica Fraas. 165
depressa M. B. 138
deusta Bald. 134
deusta Ten. 136
v. brunnea Hal. 135
v. epirotica Bald. 134
diffusa Lam. 150
dissecta Bois. 146
v. alpina Heldr. 145
v. microcephala Hal. . . 146
drakiensis Fr. et Sint. . 149
ebenoides Heldr. . . . 152
elegans All. 126
epirota Hal. 139
eryngioides Ch. et B. . . . 154
v. subarachnoidea B. H. . 154
eryngioides Raul. 154
eryngioides S. et S. . . . 153
eumorpha Sieb. 165
exscapa Urv. 159
fruticosa L. 100
galactites L. 126
graeca B. et Spr. 144
v. Grisebachii Bald. . . . 147
v. macedonica Bois. . . . 147
v. ochroleuca Heldr. et Sart. 144
graeca Griseb. 155
v Ceccariniana (B. et H.) . 155
Grisebachii Nym. 147
Guicciardii Bois. . . . 156
v. lineariloba Hal. et Dörfl. 156
v. minutispina Hal. . . . 155
Guicciardii Form. 155
Halácsyi Dörfl. 157
Heldreichii Hal. . . . 133
hellenica B. et Spr. . . . 158
Holzmanniana Heldr. . . . 163
hyalolepis Bois. 163
iberica Frev 163
v Holzmanniana Bois. . . 163
idaea B. et H. 161
incompleta Hal. 137
jacea Form. 137
v. montana Form. . . . 137

pag.
Kalambakensis Fr. et Sint. 143
lacerata Haussk. . . . 145
laconica Bois. 154
lactiflora Hal. 150
lancifolia Sieb. 165
latisquama DC. 159
laureotica Heldr. 144
lingulata Lag. 139
lyrata Sieb. 162
macedonica Bois. 160
macedonica Grisb. . . 147
v. pallidior Hal. 146
v. pallidissima Hal. . . . 147
macedonica Hal. 142
macedonica Heldr. 148
melitensis L. 162
minoa Heldr. 157
mixta DC. 157
v caulescens Hal. . . . 158
montana S. et S. 139
v albida DC. 140
musarum B. et O. . . 133
myconia B. et Sart. . . 158
nana Sieb. 158
napifolia L. 164
nervosa Willd. 138
Niederi Heldr. 140
nigrescens Willd. 137
Oliveriana DC. 152
v. amorgina B. et O. . . 152
orbelica Vel. 140
Orphanidea Hal. 148
Orphanidea Heldr. . . 148
v thessala Haussk. . . . 149
ossaea Hal. 145
ovina Form. 150
pallescens v hyalolepis Bois. 163
pallida Friv. 142
pallida × tymphaea Haussk. 142
paniculata Ch. et B. . . . 146
paniculata S. et S. 144
v. macedonica Grsb. . . . 147
v. sublanata DC. 142
Parlatoris Heldr. 145
v Boissieri H. et S. . . . 144
Parlatoris Haussk. 146
parnonia Hal. 160
parviflora S. et S. 150
paucijuga Hal. 146
v. melanolepis Hal. . . . 146
pelia DC. 149
v. refracta Haussk. . . . 150
pentelica Haussk. 143
pindicola (Grsb.) . . . 140
plumosa (Lam.) 138
princeps B. et H. . . . 135
princeps Haussk. 136
psilacantha B. et H. . . 154
ptarmicoides Hal. . . . 134
pumila S. et S. 167
pumila Urv. 158
ragusina S. et S. 141
raphanina Ch. et B. . . . 157
raphanina S. et S. . . 158
redempta Heldr. . . . 154
romana Ch. et B. 164
rupestris v. minor Bois. . . 160
salmantica L. 165
salonitana Vis. 159
v. macracantha Bois. . . 160
v. subinermis Bois. . . . 160
saxatilis C. Koch 128
saxicola v. Fraasiana Schultz 157
v. Spruneriana Schultz . . 156
scopulorum Raul. 154
seusana Chaix 139
Sibthorpii Hal. 153
sicula S. et S. 161
solstitialis L. 161
v. Adami Bois. 161
v. idaea Bald. 162
v. insularis Heldr. . . . 161
v. pycnoclada Heldr. . . 161
sonchifolia L. 164
squarrosa S. et S. 149
sphaerocephala L. . . 164
spinosa L 151
v. glabrata Heldr. . . . 152
v. tomentosa Hal. . . . 151
Spruneri B. et H. 156
Spruneriana (Schultz) . 156
subarachnoidea (B. et H.) 154
subciliaris B. et H. . . 136
sublanata Bois. 142
thessalonica Hal. 160
transiens Hal. 143
Tuntasia Heldr. 159
tymphaea Haussk. . . 148
uniflora S. et S. 139
Urvillei Bois. 158
variegata Lam. 139
v. albida Ces. 140
Zuccariniana DC. . . 151
Centaurium minus Sieb. 299
Cephalorrhynchus Bois. 215

pag.
glandulosus Bois. . . . 215
Cerinthe L. 318
aspera Roth 318
v. pallida Guss. 318
cleiostoma B. et Spr. . . . 319
echioides L. 334
macrophylla B. et H. . . . 319
maculata L. 319
maculata Raul. 319
major L. 318
v. pallida (Guss.) 318
minor L. 319
orientalis L. 336
purpurea Vis. 318
retorta S. et S. 318
strigosa Rchb. 318
Cestrum Parqui L'Her. 369
Chaenorrhinum DC. . 417
litorale Fritsch. 418
minus Willk. et L. . . 418
rubrifolium Willk. et L. 417
Chaiturus marrubiastrum Rchb. . 534
Chamaemelum Vis. . 66
rosellum B. et O. . . . 67
Tempskyanum Fr. et Sint. 67
tenuifolium (Veit) . . . 66
trichophyllum Bois. . . . 66
v. discoideum Form. . . . 67
uniglandulosum Vis. . . . 66
Chamaepeuce Alp. . . 126
afra DC. 117
Alpini Jaub. et Sp. . . . 126
v. camptolepis Bois. . . . 127
v. mutica Bois. 127
v. polycephala DC. . . . 127
atropurpurea B. et H. . . 117
cynaroides DC. 113
fruticosa DC. 127
gnaphalodes Gelmi . . . 126
mutica DC. 126
polycephala DC. 127
stellata DC. 118
stricta DC. 117
v. graeca B. et H. . . . 117
Chamomilla meridionalis C. Koch 66
officinalis C. Koch . 66
Chartolepis lancifolia Fenzl . . 165
Chironia centaurium Curt. . . . 297
maritima W. 299
spicata W. 299
Chlora L. 296
imperfoliata Ch. et B. . . 297
intermedia Ten. 296

pag.
perfoliata L. 296
serotina Koch 297
Chondrilla L. 204
acantholepis Bois. 204
capitata Sieb. 205
juncea L. 204
v. acantholepis Bois. . . 204
nudicaulis L. 215
ramosissima S. S. . . 204
Chrysanthemum L. . 69
coronarium L. 71
v. discolor Urv. 71
corymbosum L. 69
hybridum Guss. 70
leucanthemum Dall. . . . 70
leucanthemum L. 67
montanum DC. 68
myconis L. 69
v. discolor Hal. 70
pallens Gay 68
paludosum Sieb. 70
parthenium Pers. 68
segetum L. 70
tenuifolium v. discoideum Hal. 67
trichophyllum Bois. . . . 66
viscidum Friedr. 70
Cicendia Ad. 300
filiformis Delarb. . . . 300
Cichorum L. 177
divaricatum Schousb. . . . 178
endivia L. 178
v pumilum Vis. 178
glabratum Presl 178
hybridum Heldr. 179
intybus L. 177
v. divaricatum DC. . . . 178
v. glabratum (Presl) . . . 178
minimum Port. 178
pumilum Jacq. 178
pumilum × spinosum Hal. 179
spinosum L. 178
Cineraria arachnoidea Rchb. . . 80
Aucheri Nym. 78
bicolor Willd. 77
gnaphalodes Nym. . . 78
maritima Ch. et B. . . 77
taygetea Nym. 78
thapsoides Nym. . . . 80
Cionura Grisb. 292
erecta Grisb. 292
Cirsium Tourn. 108
acarna Moench 119
afrum DC. 117

pag.
appendiculatum Grisb. . 114
v. thessalum Form. . . . 114
armatum Vel. 110
arvense Scop. 118
bracteatum Link 119
candelabrum Grisb. . . 115
chelmeum Orph. 115
creticum Urv. 116
cynaroides Spreng. . . 113
decussatum Bois. 111
dolopicum Form. 111
eriophorum Form. 109
Heldreichii Hal. . . . 111
hellenicum B. et O. . . . 111
Hippolyti Link 116
horridum Form. 110
hypopsilum B. et H. . . . 111
hypopsilum Raul. 111
italicum DC. 112
lanceolatum Scop. . . . 112
v. brunneum Form. . . . 112
v. opacum Form. 112
leucocephalum Spreng. . . 112
ligulare Bois. 109
Lobelii Ten. 111
v. hypopsilum Bois. . . . 111
maculatum Lam. 125
microcephalum Schultz. . . 116
montanum W. et K. . . . 114
morinaefolium B. et H. 111
nemorale Rchb. 112
odontolepis Bois. 110
odontolepis Heldr. 109
oleraceum Scop. . . . 115
palustre Form. 116
pauciflorum Spreng. . . 115
Pelii Form. 109
pindicolum Haussk. . . . 116
polyanthos Urv. 116
rivulare Link 118
siculum DC. 116
v. longispinum Bois. . . 116
spathulatum Gaud. . . 109
v. Parnassi Hal. . . . 109
spinosissimum Scop. . . . 119
stellatum All. 118
strictum Spreng. . . . 117
tymphaeum Haussk. . . 114
validum Form. 110
v. macrocephalum Form. . 110
v. montanum Form. . . . 110
Clinopodium Plumieri Fraas . . 540
plumosum Sieb. . . 540

pag.
vulgare L. 540
Cnicus Vaill. 171
acarna L. 119
afer Sieb. 113
afer Willd. 117
benedictus L. 171
cynaroides Willd. 113
eriophorus S. et S. . . . 111
ferox S. et S. 109
galactites Lois. 126
leucocephalus Willd. . . . 112
oleraceus L. 115
palustris S. et S. 116
pauciflorus W. et K. . . . 115
rivularis Willd. 118
siculus Guss. 116
spathulatus Mor. 109
spinosissimus L. 119
strictus Ten. 117
syriacus Willd. 119
Coleostephus Myconis Cass. . . 70
Compositae Vaill. . . . 1
Convolvulaceae Vent. . 301
Convolvulus L. 301
alceaefolius Ch. et B. . . 308
althaeoides L. 308
v. argyreus Chois. . . . 309
v. italicus (R. et Sch.) . . 308
v. pedatus Chois. 309
argyreus DC. 309
arvensis L. 307
v. Cherleri (Ag.) 307
v. linearifolius Chois. . . 307
cantabricus L. 305
Cherleri Ag. 307
cneorum L. 304
cochlearis Griseb. . . . 306
dorycnium L. 303
farinosus L. 309
hirsutus Stev. 307
v. Sibthorpii (R. et Sch.) . 308
italicus R. et Sch. 308
lanatus Vahl 303
lineatus L. 304
lineatus S. et S. 304
litoralis L. 302
oleaefolius DC. 304
parnassicus B. et O. . . . 306
pentapetaloides L. . . 306
radicosus Heldr. et Sart. 305
sagittifolius S. et S. . . . 308
sepium L. 302
sericeus Chois. 303

pag.
Sibthorpii R. et Sch. . . . 308
siculus L. 306
silvaticus W. et K. . . 302
silvestris Willd. 302
soldanella L. 303
suffruticosus Desf. 309
tenuissimus S. et S. . . 309
v virescens Hal. 309
Tournefortii Sieb. 304
tricolor L. 306
Conyza ambigua DC. 14
candida Clem. 22
candida L. 23
candida S. et S. . . . 22
gnaphalodes Sieb. . . . 78
limonifolia S. et S. . . 23
pumila S et S. 28
pygmaea Sieb. 28
rupestris Sieb. 28
saxatilis S. et S. . . . 28
squarrosa L. 20
verbascifolia W. . . . 21
vulgaris Lam. 20
Cordia myxa L. 362
Cota altissima Gay. 54
Brachmanni B. et H. . . . 55
brachycentros Gay. . . . 54
parnassica B. et H. . . . 53
tinctoria Gay. 53
Cotula abrotanifolia W. 59
complanata S. et S. . . 58
Crepis L. 216
albida Ch. et B. 220
altissima Bald. 182
auriculaefolia Sieb. . . 219
v. minor Raul. 219
Baldaccii Hal. 218
barbata L. 181
biennis L. 223
bithynica Bois. 222
blattarioides Vill. 218
bulbosa L. 225
bursifolia L. 229
cernua Ten. 224
Columnae Froel. . . . 222
corcyrensis Hal. 227
cretica Bois. 224
croatica Horn. 230
crocifolia B. et R. . . 222
Dioscoridis L. 230
divaricata B. et H. . . 219
foetida L. 227
v. glandulosa (Guss.) . . 227
v. interrupta (S. et S.) . . 227
v. lagoserioides Bald. . . 227
v. maritima Bois. 227
v. radicata S. et S. . . . 227
v. rhoeadifolia (M. B.) . . 227
Fraasii Schultz. . . . 220
v. Mungierii (B. et H.) . . 221
fuliginosa S. et S. 225
geracioides Haussk. . . 217
glandulosa Guss. 227
v. acutiloba Haussk. . . . 227
v. eglandulosa Haussk. . . 227
v. interrupta Haussk. . . 227
v. maritima Haussk. . . 227
v. simplex Haussk. . . . 227
globifera Hall. 230
hispida W. et K. 228
incana S. et S. 220
incarnata Vis. 226
interrupta Sieb. 220
interrupta S. et S. 227
lacera Ten. 229
leontodontoides Gelmi . . 225
multiflora S. et S. . . 229
Mungierii B. et H. 221
neglecta L. 224
v fuliginosa Ch. et B. . . 225
nemausensis Sieb. 177
nudicaulis Sieb. 225
polymorpha Wallr. . . . 224
v. humilis DC. 224
pulchra L. 223
v. adenoclada Haussk. . . 223
radicata S. et S. 227
Raulini Bois. 218
Reuteriana Bois. . . . 221
rhagadioloides L. 189
rigida W. et K. 219
rubra L. 226
v. integrifolia Hal. . . . 226
Sartoriana B. et H. . . . 222
setosa Hall. 228
scariosa W. 229
sibirica L. 218
Sibthorpiana B. et H. . 220
Sieberi Bois. 220
smyrnaea DC. 221
Sprengeriana W. 189
stricta Scop. 224
tectorum L. 224
tubaeformis Hal. . . . 230
turcica Deg. et Bald. . 219
vesicaria L. 228
v. scariosa (W.) 214
virens L. 224

pag.
v. arvensis Ch. et B. . . 224
v. bellidifolia Ch. et B. . 224
v. silvatica Ch. et B. . . 224
virgata Desf. 182
Cressa L. 309
cretica L. 309
Crupina Cass. 166
crupinastrum (Mor.) . . 166
Morisii Bor. 166
vulgaris Cass. 166
v. serrata Forn. 166
Cupularia graveolens G. et G. . 25
viscosa G. et G. . . . 24
Cuscuta L. 310
alba Presl 311
approximata Bab. 312
atheniensis Orph. 312
breviflora Vis. 312
calliopes Heldr. et Sart. . . 311
cassiopes Heldr. et Sart. . 311
epispartos B. et O. . . . 312
epithymum L. 311
v. alba Presl 311
v. Calliopes Bois. 311
v. macranthera Heldr. et Sart. 311
v. micrantha Bois. . . . 311
europaea L. 310
globularis Bert. 312
leucosphaera B. et H. . . 312
macranthera Heldr. et Sart. 311
micrantha Tin. 312
minor Ch. et B. 311
monogyna Vahl. . . . 310
obtusiflora Humb. et Bonpl. 312
v breviflora Engelm. . . 312
palaestina Bois. 312
planiflora Ten. 311
v. approximata Engelm. . 312
v. Tenorii Engelm. . . . 312
Cuscutaceae Endl. . . . 310
Cynanchum L. 292
acutum L 292
acutum Sieb. 293
erectum L. 292
luteum Spreng. 290
monspeliacum L 292
nigrum R. Br. 291
Cynara L. 120
cardunculus L. 120
horrida Ait. 120
humilis Heldr. 120
humilis S. et S. 120
scolymus L. 120
scolymus L. 120
Sibthorpiana B. et H. . 120
Cynoglossum L. . . . 359
albanicum Deg. et Bald. 360
apenninum S. et S. . . . 359
cheirifolium Mazz. 360
Columnae Ten. 360
v dolopicum Haussk. . . 360
creticum Vill. 359
linifolium L. 357
montanum Ch. et B. . . . 360
myosotoides Lab. 358
nebrodense Guss. . . . 359
officinale S. et S. 360
omphalodes L. 357
pictum Ait. 359
silvaticum Mazz. 360
sphacioticum B. et H. . 359
Cyrtandraceae Jack. . 300

D.

Datura L. 362
metel L. 362
stramonium L. 362
Deloderium taraxacifolium Cass. . 188
Digitalis L. 420
ambigua Murr. 421
v. obtusiloba Neilr. . . . 421
ferruginea L. 420
grandiflora Lam. 421
v obtusiflora Koch . . . 421
laevigata W. et K. . . 421
lanata Ehrh. 421
v. abbreviata Haussk. . . 421
lutea L. 422
ochroleuca Jacq. 421
viridiflora Lindl. . . . 422
Diosphaera Bus. . . . 278
asperuloides Bus. . . . 278
dubia Bus. 278
Jacquini Bus. 278
Diospyros L. 286
lotus L. 286
Diotis Desf. 39
candidissima Desf 39
maritima Sm. 39
Disandra africana Camb. . . . 422
Doronicum L. 75
bellidiastrum S. et S. . . 13
caucasicum M. B. . . 75

pag.
Columnae Link 75
Columnae Ten. 76
cordatum Schultz . . . 76
cordifolium Sternb. . . . 76
eriorrhizon Guss. 75
orientale Willd. 75
Orphanidis Bois. . . . 75
v. oligocephalum Form. . . 75
pardalianches S. et S. . . 75
Dracocephalum lamiifolium Desf. 512

E.

Ebenaceae Vent. . . . 286
Echinops L. 88
albidus B. et Spr. . . 90
v. longifolius Form. . . . 90
banaticus Roch. 90
corymbosus L. 92
creticus B. et H. 88
glandulosus Weiss. . . . 88
graecus Mill. 88
lanuginosus Lam. 88
microcephalus S. et S. 91
ritro L. 91
v. elegans Bert. 91
v. ruthenicus (M. B.) . . 91
v. Sartorianus Bois. . . . 91
v. tenuifolius DC. 91
Rochelianus Grisb. 90
ruthenicus M. B. 91
ruthenicus Roch. 90
Sartorianus Bois. 91
sphaerocephalus Ch. et B. . 90
sphaerocephalus L. . . 89
v. albidus Bois. 90
v. taygeteus Bois. . . . 89
sphaerocephalus S. et S . 88
spinosus Lieb. 88
taygeteus B. et H. . . 89
tenuissimus S. et S. . . . 89
viscosus DC. 88
Echinospermum lappula Lehm. . 361
Echium L. 337
altissimum Raul. 337
arenarium Guss. 340
calycinum Viv. 340
capitatum L. 338
creticum S. et S. 340
diffusum Ch. et B 338
diffusum S. et S. . . . 340
elegans Lehm. 338
v. incanum B. et H. . . . 339
hispidum S. et S. 338

pag.
italicum L. 337
maritimum Sieb. 339
niveum Mazz. 340
parviflorum Moench . 340
v. erectum DC. 341
petraeum Tratt. 341
plantagineum L. . . . 339
v. albiflorum Raul. . . . 340
v. versicolor Haussk. . . 339
prostratum Ten. 340
pustulatum S. et S. . . 338
pyramidatum DC. 337
rubrum Fraas 338
sericeum Vahl 339
v. diffusum Bois. 340
v. hispidum Bois. 338
setosum Ch. et B. 338
Sibthorpii R. et Sch. . . . 338
violaceum Urv. 339
vulgare L. 338
Edrajanthus DC. . . . 279
graminifolius DC. . . 279
v. australis Wettst. . . . 280
Kitaibelii DC. 280
parnassicus (B. et Spr.) 279
tennifolius Bois. 279
Elephas Columnae Guss. . . . 443
Endoptera aspera DC. 231
dichotoma B. et Bal. . 229
Dioscoridis DC. . . . 230
Erica L. 282
arborea L. 282
carnea L. 283
cinerea Pieri 282
cretica Tausch 282
herbacea L. 283
manipuliflora Salisb. . . . 282
mediterranea Sieb. 282
multiflora S. et S. 282
scoparia Pieri 282
vagans Benth. 282
verticillata Forsk. . . . 282
Ericaceae DC. 282
Erigeron L. 14
acre L. 15
alpinum L. 15
v. majus Bois. 15
canadense L. 14
crispum Pourr. 14
graveolens L. 25
linifolium W. 14
olympicum Schott 15
siculum L. 27

pag
uniflorum Bald. 15
viscosum L. 24
Erythraea Ren. . . . 297
centaurium Pers. . . . 297
v. grandiflora Pers. . . . 298
grandiflora Biv. 298
latifolia Bois. 298
v. tenuiflora Grisb. . . . 298
maritima Pers. 299
pulchella Fr. 298
ramosissima Pers. 298
spicata Pers. 299
tenuiflora Hoffm. et L. 298
virescens Willd. 299
Eufragia latifolia Grisb. . . . 437
viscosa Benth. 417
Eupatorium L. . . . 10
cannabinum L. 10
v. indivisum Bois. . , . 11
v. simplicifolium Hal. . . 11
v. syriacum Bois. 11
syriacum Jacq. 11
v. indivisum DC. 11
Euphrasia L. 439
frutescens Sieb. 438
glutinosa M B. 439
latifolia L. 436
odontites L. 439
officinalis S. et S. 439
olympica Hal. et Sint. . . 440
pectinata Ten. 439
puberula Hal. 439
salisburgensis Funk . 440
v. olympica Hal. 440
serotina Lam. 439
Sintenisii Hal. 440
tatarica Form. 439
viscosa Sieb. 437
Evax Gaertn 37
exigua DC. 38
perpusilla B. et H. . . 38
pygmaea Pers. 37
Exacum filiforme S. et S. . . . 300

F.

Fabera cretensis Schultz . . . 175
hispida Schultz 175
Filago L. 39
arvensis L. 36
v. lagopus DC. 36
canescens Jord. 34
pag.
canescens × spathulata Hssk. 36
congesta Guss. 38
eriocephala Guss. 34
eriocephala×spathulata Hssk. 36
exigua L. 38
gallica L. 37
germanica L. 37
v. canescens (Jord.) . . . 34
v. eriocephala Guss. . . . 34
v. lanuginosa DC. 34
v. lutescens (Jord.) . . . 34
intermedia Haussk. . . . 36
lagopus Parl. 36
v alpina Raul. 36
lutescens Jord. 34
maritima L. 39
minima Form. 37
prostrata Parl. 35
pygmaea L. 37
pyramidata L. 34
v canescens Form. . . . 34
similata Haussk. 36
spathulata Presl. . . . 35
v. affinis Haussk. 35
v. decalvans Hal. 35
v. prostrata Bois. 35
v. siria Heldr. 35
tenuifolia Presl 37
Fraxinus L. 289
excelsior L. 289
ornus L. 289
rotundifolia Lam. 289

G.

Galactites Moench. . . 125
tomentosa Moench. 125
v. alata DC. 125
v. elegans DC. 126
Galeobdolon luteum Huds. . . . 509
Galeopsis galeobdolon L. . . . 509
Gatyona Dioscoridis Rchb. . . . 230
globulifera Cass. 230
Gentiana L. 295
aestiva R. et Sch. 296
asclepiadea L. 296
centaurium L. 297
filiformis L. 300
lutea L. 295
maritima L. 299
perfoliata L. 296
pulchella Sw. 298
spicata L. 299
symphyandra Murb. . . 295

pag.
tergestina Beck 296
verna L. 296
v. aestiva Bald. 296
Gentianaceae Juss. . . 295
Geropogon L. 191
glaber L. 191
hirsutus Urv. 191
Glechoma L. 539
arvensis L. 531
hederacea L. 534
Gnaphalium L. 32
arvense L. 36
conglobatum Friedr. . . . 31
Fontanesii Camb. 31
gallicum L. 37
germanicum L. 34
Hoppeanum Bois. 33
italicum Roth 32
lagopus Steph. 36
luteoalbum L. 32
microphyllum W. 32
orientale L. 29
pusillum Haenke 34
Roeseri B. et H. . . . 33
scandens Sieb. 33
siculum Spreng. 30
silvaticum L. 33
Sophiae Heldr. 33
stoechas Ch. et B. 30
stoechas S. et S. 32
supinum L. 33
v. subacaule (Wahlenb.) . 34
supracanum S. et S. . . . 34
uliginosum L. 33
Gomphocarpus R. Br. . 239
fruticosus R. Br. . . . 293
Gratiola L. 407
officinalis L. 407

H.

Haberlea Heldreichii Bois. . . . 301
Hedraeanthus graminifolius Wettst. 280
Kitaibelii Wettst. 280
Hedypnois Tourn. . . 180
cretica Bois. 180
cretica Willd. 181
v. gracilior Bois. 181
v. monspeliensis (W.) . . 181
monspeliensis W. 181
paludosa Scop. 202
pendula W. 181
polymorpha DC. 181
rhagadioloides W. . . . 180
tubaeformis Ten. 180
Helianthus tuberosus L. 39
Heliotropium L. . . 314
Bocconi Guss. 316
dolosum Not. 379
Eichwaldi Bois. 317
ellipticum Led. 317
ellipticum Nym. 317
europaeum L. 316
v. hirsutum Sieb. 315
v. tenuiflorum Guss. . . . 316
macrocarpum Guss. . . . 317
odorum Stev. 316
suaveolens M. B. . . . 316
turcicum Friv 316
villosum W. 314
Helichrysum DC. . . 29
amorginum B. et O. . 29
anatolicum Bois. 30
angustifolium DC. 32
conglobatum Ung. 31
decumbens Raul. 31
Heldreichii Bois. . . . 30
italicum Guss. 32
v. microphyllum Bois. . . 32
microphyllum Camb. . . . 32
nebrodense Heldr. 30
orientale DC. 29
panormitanum Bois. . . . 30
v. angustifolium Raul. . . 30
panormitanum Tin. 30
plicatum DC. 30
v. anatolicum Haussk. . . 30
rupestre DC. 30
v Cambessedesii DC. . . 31
scandens Raul. 31
serotinum v orientale Bois. 32
siculum Bois. 30
v. brachyphyllum Spreitz. . 31
v. laureoticum Haussk. . . 31
stoechas M. et R. 31
v. incanum Bald. 31
Helminthia Juss. . . 190
asplenioides DC. 191
echioides Gaertn. . . . 190
Hieracium L. 231
abietinum (B. et H.) . 243
aetolicum Hal. 238
alpestre S. et S. 222
alpinum S. et S. 237
atticum Nym. 236

pag.
auricula L. 234
Bauhini Schult. 236
bracteolatum S. et S. . 244
v. laxum Bois. 244
v. pelium Hal. 244
bracteolatum × racemosum 245
bulbosum W. 225
chalcidicum Hal. 239
ciliatum W. 189
Columnae Ten. 222
crinitum S. et S. . . . 245
cymosum L. 234
v. Heldreichianum N. P. . 234
v. sabinum Seb. 234
dolopicum Fr et Sint. . . 239
eriobasis Fr. et Sint. . . . 242
eriopus B. et H. 245
euboeum Hal. 239
florentinum All. . . . 235
v. hololeion N. P. . . . 236
v. praealtum (Vill.) . . . 236
foetidum S. et S. 222
Friwaldii Rchb. 240
Gaudryi B et O. . . . 240
Gaudryi Heldr. 238
graecum B. et H. . . . 237
Halácsyi Heldr 235
Heldreichii Bois. 238
Heldreichianum Haussk. . . 234
Heldreichianum × praealtum Haussk. 235
incisum Hoppe 242
Koracis Bois. 244
Koracis Haussk. 244
Leithneri Heldr. et Sart. . . 242
macranthum Ten. . . . 233
macranthum × Heldreichianum Hal. 235
magyaricum N. P. 236
v. graecum N. P. 236
v. laconicum N. P. . . . 236
murorum L. 243
v. pilosissimum Bois. . . 242
murorum Sieb. 242
Naegelianum Panc. . . . 237
nudicaule Sieb. 245
oligadenum (Belli) . . . 241
oligocephalum Hal. 242
panachaicum Hal. 235
pannonicum N. P. 236
v. leiocaulon N. P. . . . 236
pannosum Bois. 239
v. Friwaldii (Rchb.) . . . 240
v. scapiferum Bois. . . . 240
v. taygeteum Bois. . . . 239
Parnassi Fr. 237
pilosella L. 233
v. anatolicum N. P. . . . 233
v. eglandulosum N. P. . . 233
v. glandulosum N. P. . . 233
v. macranthum (Ten.) . . 233
v. obscurius N. P. . . . 233
piloselloides Vill. 235
plumulosum Kern. 241
praealtum Ch. et B. . . . 236
praealtum Vill. 236
v. Bauhini Grsb. 236
v. hispidissimum Bois. . . 236
racemosum W. et K. . 245
Reinholdi Heldr. et Sart. . 244
sabinum Seb. 234
v. Heldreichianum Hal. . . 234
sabinum × Bauhini Hal. 232
Sartorianum B. et H. . 242
v Leithneri Bois. 242
scapigerum B. O. et H. 241
sermenikense Fr. et Sint. . 245
Sprengerianum L. 189
taygeteum B. et H. . . . 239
thessalum Form. . . . 241
undulatum Bois. 237
versutum Friv. 240
vulgatum Hal. 243
v abietinum B. et H. . . 243
Waldsteinii Tausch . . . 241
v. oligadenum Belli . . . 241
Hippion aestivum Schm. . . . 296
Hirtellina lanceolata Cass. . . . 100
Hyalea Olivieri Jaub. et Sp. . . 152
Hymenonema Cass. . . 172
graecum DC. 172
laconicum B. et H. . . 173
v minus Rouy 173
Tournefortii Cass. 172
Hyosciamus L. 363
albus L. 364
v. major Heldr. 364
v. minor Haussk. 364
aureus L. 365
graecus Heldr. 364
major Mill. 364
minor Mill. 364
niger L. 363
v. pallidus Koch 363
pallidus W. et K. 363
reticulatus L. 364
varians Vis. 364

pag.
Hyoseris L. 179
cretica L. 181
lucida Sieb. 179
microcephala Cass. 180
radiata L. 179
v. graeca Hal. 179
rhagadioloides L. 180
scabra L. 180
Hypochaeris L. . . . 173
acaulis Sieb. 175
cretensis Ch. et B. . . 174
dimorpha Brot. 173
dimorpha Ten. 174
glabra L. 173
v. minima DC. 173
hedypnois L. 181
minima Cyr. 173
neapolitana DC. 174
pinnatifida Hal. 175
radicata L. 174
v. heterocarpa Nov. . . . 174
v. neapolitana (DC.) . . . 174
setosa Form. 186
tenuiflora Bois. 175
Hypopitys multiflora Scop. . . . 284
Hyssopus officinalis Dall. . . . 546

I.

Jankaea Heldreichii Bois. . . . 301
Jasione L. 280
dentata (DC.) 280
Heldreichii B. et O. . . . 280
montana Form. 280
v. dentata DC. 280
orbiculata Grisb. 281
supina Sieb. 281
Jasminaceae R. Br. . . 286
Jasminum Tourn. . . . 286
fruticans L. 286
grandiflorum L. 286
humile L. 286
odoratissimum L. 286
officinale L. 286
sambac L. 286
Jasonia sicula DC. 27
Ifloga spicata Schultz 34
Inula L. 19
arabica Sieb. 23
Aschersoniana Hal. . . . 23
aspera Poir. 19

pag.
attica Hal. 23
v. limonella Heldr. . . . 23
v. macrophylla Heldr. . . 22
britannica L. 21
v. rupestris Gr. et Sch. . 21
bubonium Pieri 21
candida Bois. 22
candida Cass. 22
candida (L.) 23
v. integrifolia Raul. . . . 22
v. limonifolia DC. 23
v. radiata Bois. 22
v. rotundifolia Hal. . . . 23
v. verbascifolia Clem. . . 22
conyza DC. 20
cordata Bois. 19
crispa Pers. 26
crithmifolia W. 24
crithmoides L. 24
dentata S. et S. 26
dysenterica L. 26
ensifolia L. 20
germanica L. 19
graveolens Desf. . . . 25
helenium L. 19
hirta L. 20
limonifolia Bois. . . . 23
methanaea Haussk. . . . 23
oculus christi L. . . . 21
odora L. 25
oxylepis Schultz 22
parnassica B. et H. . . 22
v. euboea Hal. 22
pulegium Friedr. 26
pulicaria L. 26
salicina v. aspera Beck . . 19
undulata Friedr. 26
verbascifolia Form. . . . 23
verbascifolia (W.) . . . 21
v. limonifolia DC. . . . 22
viscosa Ait. 24
vulgaris Trev. 20
Ipomaea L. 301
sagittata Poir. 301
Jurinea Cass. 127
anatolica Bois. 128
chia B. et O. 128
glycacantha DC. . . . 128
mollis Rchb. 127

K.

pag.
Kentrophyllum creticum Bois. . 168
dentatum DC. . 169
dentatum Weiss . 169
lanatum Duby . 167
leucocaulon DC. . 168
rubrum Link . . 169
tauricum Raul. . 168
Koelpinia linearis Pall. 183

L.

Labiatae Juss. 463
Lactuca L. 208
acanthifolia Bois. . . . 211
albicaulis Bois. 210
amorgina Heldr. et Orph. 212
angustana All. 210
Chaixi Vill. 209
cretica Desf. 208
graeca Bois. 209
leucophaea S. et S. . . . 208
muralis L. 213
sagittata W. et K. 209
saligna L. 211
sativa L. 210
scariola L. 210
v. angustana All. . . . 210
v. coriacea Schultz . . . 210
sonchifolia Sieb. 208
tenerrima Bald. 209
Tommasiniana Schultz . . 211
viminea Presl 212
virosa L. 209
Lagoseris M. B. . . . 177
bifida Koch 177
nemausensis Koch 177
Lamium L. 509
album Friedr. 510
album L. 510
v. maculatum L. 511
amplexicaule L. . . . 513
v. clandestinum Rchb. . . 514
v. microphyllum Heldr. . 514
bifidum Cyr. 513
bithynicum Benth. 512
calycinum Urv. 510
Columnae Friedr. 512
cylleneum B. et H. . . . 512
galeobdolon L. 519
garganicum L. 511
v. glabratum Grisb. . . . 512
v. longiflorum (Ten.) . . 512
v. nivale Bald. 512
v. nivale B. et H. . . . 511
v. striatum S. et S. . . . 512
grandiflorum Sieb. 512
hybridum Vill. 513
incisum W. 513
longiflorum Ten. 512
maculatum L. 511
mesogaeum Heldr. 514
moschatum Mill. . . . 510
ochroleucum Link 510
pictum B. et H. . . . 512
pictum Hal. 512
purpureum L. 511
rugosum Ch. et B. 512
striatum S. et S. 512
v. cylleneum B. et H. . . 511
v. glabrescens Heldr. . . 512
v. hymettium Heldr. . . . 512
v. minus Hal. 512
v. pictum Bois. 512
tomentosum W. 512
Lappa Tourn. 101
major Gaertn. 101
minor DC. 102
Lappula Gilib. 361
echinata Gilib. 361
myosotis Moench 361
vulgaris Mazz. 361
Lapsana L. 182
communis L. 182
v. glandulosa Form. . . . 182
rhagadiolus L. 183
stellata L. 183
tubiflora Friedr. 183
zacintha L. 213
Laurentia Mich. . . . 281
tenella DC. 281
Lavandula L. 567
dentata L. 568
spica L. 567
stocchas L. 567
Leontodon L. 185
abietum Heldr. 187
alpinus Hoppe 202
alpinus Jacq. 188
asper Poir. 186
v. Haussknechtii (Uechtr.) . 187
v. Huetii Form. 187
v. saxatilis (Ten.) . . . 187
v. setulosus Hal. 187

pag.
asperrimus Bois. . . . 186
cichoraceum Sang. 185
creticum Bois. 188
crispus Ung. 187
daedaleum Sieb. 188
fasciculatus Nym. . . . 185
graecus B. et H. . . . 187
v. alpinus Bois. 188
v. Heldreichianus Bois. . . 188
gymnanthum Link 203
hastilis L. 186
v. banaticus Heuff. . . . 185
v. glabratus Koch . . . 186
v. hispidus Bois. 185
Heldreichianus Hal. . . . 188
hirtus L. 184
hispidus L. 185
v. hastilis (L.) 186
incanus Ch. et B. 187
laevigatus W. 202
lanatus L. 198
leptocephalum Hal. . . . 201
lividus W. et K. 202
minimus Brig. 203
muricatum L'Her. 198
palustre Hudis 202
saxatilis Rchb. 187
serotinus S. et S. 201
squamosum Pieri 187
taraxacifolius (Cass.) . 188
taraxacoides Hoppe . . . 202
taraxacum L. 201
v. laevigatum Bois. . . . 201
tuberosus L. 184
Leonurus L. 534
cardiaca L. 534
marrubiastrum L. . . 534
Lesquereuxia syriaca B. et R. . 442
Leucanthemum Tourn. 67
montanum DC. 68
pallens DC. 68
setabense Raul. 70
vulgare Lam. 67
Ligustrum L. 288
vulgare L. 288
sempervirens Mazz. . . . 288
Limnanthemum Gm. 300
nymphoides H. et L. . 300
[illegible]maria Tourn. 407
[illegible] Desf. 413
[illegible]ra Bois. 413
[illegible] B. et Bal. . . . 415
[illegible] Mill. . . . 410
[illegible]m. 410
pag.
commutata Bernh. . . . 415
v. polygonoides (Rev.) . . 416
cymbalaria Friedr. 416
cymbalaria Mill. . . . 416
v. Sieberi Chav. 416
dalmatica Mill. 408
v. macedonica Fenzl . . 409
v. ramosissima Haussk. . 409
v. thessala Haussk. . . . 409
elatine Friedr. 415
v. lasiopoda Vis. 415
v. polygonoides Rev. . . 416
v. villosa Bois. 415
genistifolia Mill. . . . 409
v. angustifolia Hal. . . . 409
v. brevicalcarata Hal. . . 409
graeca Chav. 415
v. gracilior Heldr. . . . 415
lanigera Link 414
litoralis W. 418
longipes B. et H. . . . 416
macedonica Grisb. 409
micrantha Spreng. . . 413
microcalyx Bois. . . . 417
minor Desf. 418
monspeliensis Friedr. . . . 417
olympica Form. 410
parnassica B. et H. . . . 410
parviflora Desf. 413
parviflora (Jacq.) . . . 413
Pelisseriana Mill. . . . 412
peloponnesiaca B. et H. 409
v. parnassica (B. et H.) . 410
pilosa Spreitz. 417
Prestandreae Tin. 415
purpurea Mill. 411
reflexa Desf. 411
rubrifolia Rob. et Cast. . . 417
Sibthorpiana Bois. 410
v. parnassica Bois. . . . 410
v. peloponnesiaca Bois. . . 409
Sieberi Rchb. 414
v. bombycina (B. et Bal.) . 415
simplex DC. 413
Smithii B. et O. 408
spuria Mill. 414
stricta Guss. 410
triphylla Mill. 410
vulgaris Mill. 412
Lippia L. 463
nodiflora Rich. 463
repens Spreng. 463
Lithosper[illegible]n L. . . 347

pag.
apulum L. 350
arvense L. 348
v. Sibthorpianum (Grisb.) . 349
dispermum L. 361
fruticosum L. 347
hispidulum S. et S. . . 347
incrassatum Guss. . . 349
Leithneri Heldr. et Sart. . 348
officinale L. 348
orientale L. 342
petraeum DC. 341
purpureocoeruleum L. . 348
rosmarinifolium Bois. . . . 347
Sibthorpianum Grisb. . . . 349
tenuiflorum L. 349
tenuiflorum S. et S. . . . 349
tinctorium L. 346
Zahnii Heldr. 347
Lobelia Bivonae Ten. 281
cretica Juss 281
laurentia W 281
setacea S. et S. 281
tenella Biv. 281
Lobeliaceae Juss. . . . 281
Lobostemon capitatus Buek . . 338
hispidus DC. . . . 338
Lomatolepis glomerata Cass. . . 205
Lycium L. 368
barbarum L. 368
europaeum L. 368
mediterraneum Dun. . . . 368
vulgare Dun. 368
Lycopersicum esculentum Mill. . 370
Lycopsis aegyptiaca L. 327
arvensis L. 329
bullata Cyr. 329
glomerata Urv. . . . 328
obtusifolia W. 321
Sibthorpiana R. et Sch. 321
variegata L. 329
vesicaria Friedr. . . . 321
Lycopus L. 568
albus Mazz. 568
europaeus L. 568
v. pubescens Benth. . . . 568
europaeus × exaltatus Haussk. 568
exaltatus Fraas 568
intermedius Haussk. . . . 568
mollis Kern. 568
Lyonnetia abrotanifolia Less. . . 59
pectinata DC.
pusilla Cass. 59

pag.
rigida DC. 59
tenuiloba DC. 56

M.

Macrotomia DC. . . . 391
cephalotes Bois. 341
Majorana crassifolia Benth. . . 557
microphylla Benth. . . 556
onites Benth . . 556
Mandragora Tourn. . . 366
autumnalis Spreng. . . 367
Haussknechtii Heldr. . 367
Haussknechtii × vernalis Haussk. et Heldr. . 367
hybrida Haussk. et Heldr. . 367
microcarpa Bert. 367
officinarum Bert. 367
officinarum L. 366
officinarum × Haussknechtii 367
vernalis Bert. 366
Marrubium L. 503
acetabulosum L. 532
acetabulosum S. et S. . . 533
apulum Ten. 503
candidissimum Form. . . . 503
canescens Sieb. 503
creticum Mill. 503
cylleneum B. et H. . . 504
Haussknechtii Uechtr. . . 505
hirsutum W. 532
hispanicum L. 532
pannonicum Rchb. 503
peregrinum L. 503
v. platyphyllum Haussk. . 503
peregrinum × vulgare Reichb. 503
pseudodictamnus L. . . . 533
pseudodictamnus S. et S. . 533
remotum Kit. 503
rupestre Biv. 532
thessalum B. et H. . . 504
velutinum S. et S. . . 505
v. cylleneum Bois. . . . 504
v. Frivaldszkyanum Bois. . 505
vulgare L. 503
v. apulum (Ten.) 503
v. lanatum Benth. . . . 503
vulgare × peregrinum 503
Marsdenia erecta R. Br. 293
Maruta cotula DC. 64
v. psorosperma Raul. . . 64
Matricaria L. 66
chamomilla L. 66

pag.
v. coronata Bois. 66
v. eradiata Rupr. 66
v. pappulosa M. et R. . . 66
chamaemelon Heldr. . . . 66
coronaria Desr. 71
coronata Gay. 66
montana Pieri 66
parthenium L. 68
suaveolens L. 66
trichophylla Bois. 66
v. discoidea Form. . . . 67
Mattia Schult. 358
graeca B. et H. . . . 358
Schmidtii Heldr. 358
Me ampyrum L. . . . 444
barbatum Bois. 444
ciliatum B. et H. . . . 444
Melissa L. 539
altissima S. et S. 539
calamintha Dall. 542
calamintha L. 541
cordifolia Pers. 539
cretica L. 543
grandiflora L. 540
nepeta L. 542
officinalis L. 539
Melittis L. 505
albida Gun. 406
cordifolia B. et H. 506
grandiflora Sm. 506
melissophyllum L. . . 505
Mentha L. 569
aquatica L. 571
v. crispa Benth. 572
v. hirsuta Benth. 572
canescens Sieb. 570
cephalonica Br. 572
cretica Port. 571
crispa L. 572
erinoides Heldr. 572
gibraltarica W. 573
hirsuta L. 572
ilissia Heldr. 571
incana Sm. 570
litoralis Borb. 572
longifolia Huds. . . . 569
v. candicans (Cr.) 570
v. mollissima (Borkh.) . . 570
v. minutiflora (Borb.) . . 570
v. Sieberi (Koch) 570
v. Wierzbickiana (Op.) . . 570
minutiflora Bobb. 570
mollissima Borkh. 570
mollissima × aquatica . . . 571
niliaca Link 570

pag.
piperita L. 571
pubescens W. 571
pulegioides Sieb. 572
pulegium L. 572
v. pulegioides (Sieb.) . . . 572
v. subtomentella Br. . . . 572
v. tomentella (H. et L.) . 573
v. tomentosa Vis. 573
v. villicaulis Borb. . . . 572
v. villosa Benth. 573
Reverchoni Hal. 571
Rocheliana Borb. 570
rotundifolia L. 569
rugosa Lam. 569
Sieberi C. Koch 570
v rotundifolia Rev. . . . 571
silvestris L. 569
v. candicans (Cr.) 570
v. stenotricha Borb. . . . 569
v. stenostachya Bois. . . 570
spicata L. 569
v. longifolia 569
v. rotundifolia L. 569
v. viridis L. 571
tomentella H. et L. . . . 571
tomentosa Urv. 570
v viridula Heldr. 571
undulata W. 570
viridescens Borb. 570
viridis L. 571
v graciliflora Borb. . . . 571
Wierzbickiana Op. 570
Menyanthes nymphoides S. . . . 300
Metabasis aethnensis DC. . . . 170
cretensis DC. 174
hymettia DC. 174
Microlonchus L. . . . 165
salmanticus DC. . . . 165
Micromeria Benth. . . 546
canescens M. et B. 549
cremnophila B. et H. . 547
v thessala Form. 547
graeca Benth. 548
v. villicaulis Borb. . . . 549
hispida B. et H. . . . 549
juliana Benth. 546
v. myrtifolia Bois. . . . 547
juliana × cremnophila . . . 547
marifolia Raul. 543
meteorica Haussk. 547
microphylla Benth. . . 549
myrtifolia B. et Hoh. . 547
nervosa Benth. 548
plumosa Heldr. 548

pag.
sphaciotica B. et H. 549
Micropus L. 38
bombycinus Lag. . . . 39
erectus L. 38
exiguus Urv 48
pygmaeus Desf. 37
Microrrhynchus Less. 215
nudicaulis Less. . . . 215
Millina leontodontoides Cass. . . 185
hyoserioides DC. 188
Moltkia Lehm. . . . .
petraea Grisb. 341
Molluccella L. 506
spinosa L. 506
Monotropa L. 284
hypophegea Wallr. 285
hypopitys L. 284
v. glabra Roth . . . . . , 285
v. hirsuta Roth 285
Monotropaceae Nutt. . 284
Munbya cephalotes Bois. 341
jasminiflora Orph. . . . 341
Myosotis L. 350
aenesia Heldr. 353
alpestris Hal. 354
alpestris Ung. 354
arvensis Roth 355
v versicolor Pers. . . . 356
cadmea Hal. 352
caespitosa Schultz 351
collina Hoffm. 355
v. gracillima (Losc.) . . . 356
v. grandiflora B. et H. . . 356
commutata R. et Sch. . . . 351
cretica B. et H. 353
cretica Ung. 352
cyanea B. et H. 354
echinosperma Mazz. . . . 361
gracillima Losc. 356
hispida Schlecht. 355
v pygmaea Hal. 357
idaea B. et H. 352
v. boeotica Reut. 352
v. cadmea Bois. 352
v. grandiflora Bois. . . . 352
v. Kiesenwetteri Heldr. . . 353
intermedia Link 355
lappula L. 361
lingulata Lehm. 351
litoralis Stev 353
v aenesia Heldr 353
litoralis Ung. 353
Mathildae Haussk. 355
micrantha Guss. 352
nana S. et S. 357
olympica Bois. 354
palustris With. 351
v. strigulosa G. et G. . . 351
pusilla Clem. 353
pusilla Lois. 353
refracta Bois. 357
scorpioides L. 351
v. arvensis L. 335
v. palustris L. 351
sicula Guss. 352
silvatica Hoffm. 354
v Mathildae Form. . . 355
stricta Friedr. 352
stricta Link. 356
stricta Weiss. 352
strigulosa Rchb. 351
versicolor Schlecht. . . 356

N.

Nemauchenes aculeata Cass. . . 231
Nepeta L. 534
argolica Ch. et B. 537
camphorata B. et H. . . . 538
camphorata Heldr. 539
cataria L. 534
dirphya Heldr. 538
Heldreichii Hal. . . . 539
incana Sieb. 535
italica S. et S. 536
v. Spruneri Bald. 538
laconica Orph. 537
melissaefolia Lam. . . 535
nuda L. 536
v. epirota Hal. 536
pannonica L. 536
parnassica Heldr. et Sart. 537
v Orphanidea Bois. . . . 537
scordotis L. 535
Sibthorpii Benth. . . . 536
v. dirphya Bois. 538
Sibthorpii Form. 537
Spruneri Bois. 537
tomentosa Sieb. 535
tymphrestea Heldr. et Sart. 538
violacea Vill. 536
Nerium L. 293
oleander L. 293
Nicotiana glauca Grah. 370
rustica L. 370
tabacum L. 370

pag.
Nonnea Med. 321
auriculata Bois. 331
lamprocarpa Griesb. . . . 321
lutea Ch. et B. 344
obtusifolia R. et Sch. . . 321
Sibthorpiana Don. 321
ventricosa Grisb. . . . 321
Notobasis Cass. 119
syriaca Cass. 119

O.

Ocimum basilicum L. 568
Odontites Hall. 438
Bocconi (Guss.) 438
cretica Bois. 438
divergens Jord. 439
frutescens (Sieb.) . . . 438
glutinosa Benth. . . . 439
ixodes B. et H. 439
Linkii Heldr. et Sart. . 438
serotina Rchb. 439
Omalotheca supina Cass. . . . 33
Olea L. 387
europaea L. 287
v. oleaster H. et S. . . . 287
sativa H. et L. 287
Oleaceae Lindl. . . . 287
Omphalodes Tourn. . . 357
linifolia Moench. 357
Luciliae Bois. 357
myosotoides Schrank . . . 358
verna Moench 357
Onobroma corymbosum Spreng. . 92
Onopordon L. 121
acanthium L. 121
v. parnassicum B. et R. . 121
acaulon S. et S. 129
alexandrinum Heldr. . . . 122
argolicum Bois. 121
arabicum S. et S. 122
caulescens Urv. . . . 123
elatum S. et S. 121
v. argolicum Bois. . . . 121
elongatum Lam. 123
euboeum Heldr. 123
graecum Ch. et B. 122
graecum Gou. 123
horridum Haussk. 122
illex Janka 123
illyricum L. 123

pag.
laconicum Heldr. et Sart. 124
macracanthum S. et S. . . 122
messeniacum Hal. . . 122
myriacanthum Bois. . . . 123
myriacanthum Hal. . . . 123
orientale Mill. 123
parnassicum B. et H. . . . 121
Sibthorpianum B. et H. 122
tauricum W. 121
v. elatum Bois. 121
v. elatum Spreitz. 124
Weissianum Asch. 123
v. brevicaule Rouy . . . 123
Onosma L. 332
angustifolium Form. . . . 333
angustifolium Lehm. . . . 332
arenarium W. et K. . . . 335
arenarium Ung. 336
calycinum Stev. 336
echioides Jacq. 336
echioides L. 334
v. brachycalyx Hal. . . . 334
v. brachyphyllum Hal . . 334
v. helveticum DC. . . . 335
v. sparsiflorum Friedr. . . 334
echioides S. et S. 336
echioides Sieb. 336
erectum Clem. 333
erectum S. et S. . . . 333
v. pubiflorum Hal. . . . 333
frutescens Lam. . . . 336
graecum Bois. 336
helveticum Bois. . . . 335
laconicum B. et O. . . . 332
leptanthum Heldr. . . 335
orientale Lehm. 336
Orphanidis B. et H. . . . 334
pallidum Bois. 334
rhodopeum Vel. 336
simplicissimum L. 337
Spruneri Bois. 332
stellulatum Ch. et B. . . . 334
stellulatum W. et K. . . . 333
v. angustifolium Bois. . . 332
v. erectum Bois. 333
v. pallidum Bois. 334
tauricum Pall. 332
taygeteum B. et H. . . 336
tinctorium Ch. et B. . . . 336
Visianii Clem. 336
Origanum L. 552
creticum L. 555
crassifolium (Benth.) . . . 557
dictamnus Ch. et B. . . . 553

pag.
dictamnus L. 552
dubium Bois. 556
heracleoticum L. . . . 555
v. albiflorum (Haussk.) . . 555
v. creticum Hal. 555
v. rubriflorum (Haussk.) . 555
v. trichocalycinum (Haussk.) 555
hirtum Link 555
v. albiflorum Haussk. . . 555
v. prismaticum Haussk. . 555
v. rubriflorum Haussk. . . 555
v. trichocalycinum Haussk. 555
hybridum Heldr. 554
lirium Heldr. 554
majorana L. 557
maru L. 556
microphyllum Sieb. . . . 556
neglectum Vog. 555
onites L. 556
parviflorum Urv. 555
pseudodictamnus Sieb. . . 552
pulchrum B. et H. . . 552
scabrum B. et H. . . . 553
sipyleum S. et S. 553
smyrnaeum L. 556
smyrnaeum S. et S. . . . 555
Tournefortii Sibth. . . 553
viride (Bois.) 554
viridulum Heldr 554
vulgare L. 554
v. album Fraas 554
v. elongatum Form. . . . 555
v. latebracteatum Form. . 555
v. prismaticum Raul. . . 555
v. viride Bois. 554
vulgare S. et S. 554
Ormenis Cass. 64
mixta DC. 64
Ornus europaea Pers. 289
Orobanche Tourn. . . . 444
aegyptiaca Pers. . . . 446
alba Steph. 451
v. bidentata Beck 452
v. communis Beck . . . 452
v. glabrata (C. A. Mey) . 452
v. leptocalamus Beck . . 452
v. microsepala Beck . . . 452
v. minutiflora Beck . . . 452
v. subalpina Beck . . . 452
ambigua Mor. 457
amethystea Thuill 453
v. attica Beck 453
angustisepala Schultz . . . 451
apiculata Wallr. 458

pag.
arachnoidea Schultz . . . 455
arcuata Schultz 455
arenaria M. B. 448
artemisiae campestris Vauch. 456
attica Reut. 453
australis Mor. 454
breviflora Schultz 450
bicolor C. A. May 448
canescens Presl 454
canescens Sprun. 451
carotae Desm. 457
caryophyllacea Sm. . . 449
cernua Loefl. 448
v. cumana Beck 448
chassia Form. 459
cirsii Fr. 453
coerulea Vill. 447
condensata Grisb. 455
crassistyla Schultz 451
crenata Forsk. 451
cruenta Bert. 450
cumana Wallr. 448
diaphana Schultz 451
elatior Rchb. 457
elatior Sutt. 449
epithymum DC. 451
Fraasii Schultz 446
fragrans Grisb. 445
Freynii Nym. 457
fuliginosa Reut. 457
v. oxycheilos Beck . . . 457
galii Duby 449
glabrata C. A. Mey . . . 452
gracilis Hal. 451
gracilis Sm. 450
v. Spruneri Beck 450
grandiflora Ch. et B. . . . 451
grandisepala Schultz . . . 454
Grisebachii Reut. . . . 455
hederae Duby 458
v. stenantha Lojac. . . . 459
hyalina Sprun. 458
Kochii Schultz 449
laurina Rchb. 458
lavandulacea Hal. 446
lavandulacea Rchb. . . 446
leucantha Grisb. 453
lilacina Schultz 457
livida Hal. 455
livida Sendt. 458
v. adenostyla Vis. . . . 458
loricata Rchb. 456
lycica Schultz 549
major Hal. 453
major L. 449

pag.
medicaginis Rchb. 458
minor Sutt. 457
v. adenostyla Hal. . . . 455
v. minima Beck 458
v. pumila Koch 458
Muteli Hal. 445
Muteli Schultz 445
v spissa Beck 446
nana Noe 445
nudiflora Wallr. 458
palaestina Bois. 455
parviloba Schultz 459
picridis Schultz 456
v. carotae Beck 457
platystigma Rchb. 453
prosgolica Form. 459
pruinosa Lap. 451
pubescens Urv. 454
punctata Schultz 451
purpurea Jacq 447
ramosa L. 444
Reichardiae Freyn 456
reticulata Wallr. . . . 452
v. pallidiflora W. et Gr. . 453
Reuteri Schultz 458
scabiosae Koch 453
Schultzii Mut. 447
v. pyramidalis Beck . . . 447
segetum Sprun. 451
speciosa DC. 451
Spruneri Schultz 450
stigmatodes Wim. 449
striata Reut. 455
stricta Mor. 447
subverticillata Schultz . . 459
Tommasinii Rchb. 457
tridentata Schultz 454
versicolor Schultz . . . 454
v. campuliflora Loj. . . . 455
v. lanifera Beck 455
v. macrophyllon Beck . . 455
v. monochroa Beck . . . 455
v. pallida Reut. 455
v. pseudobarbata Beck . . 455
villiflora Koch 454
villosa Schultz 454
villosiflora Schultz 454
Orobanchaceae Lindl. . 444
Orvala garganica L 511
Otanthus maritimus Link . . . 39

P.

Pallenis L. 18
spinosa Cass. 18
pag.
v. microcephala Hal. . . 18
Paracaryum Bois. . . 358
myosotoides Bois. . . . 358
Sibthorpianum Bois. . . . 358
Parentucellia Viv. . . 436
latifolia Car. 436
v. major Heldr 439
v. pycnantha Heldr. . . . 437
viscosa Car 437
Pedicularis L. 442
brachyodonta Schl. et V. 443
graeca Bunge 442
v. inermis Bald. 443
rupestris B. et O 443
Petasites Tourn. . . . 11
officinalis Moench. . . 11
Periploca L. 290
graeca L 290
Petromarula DC. . . 275
pinnata DC. 275
Phaenixopus acanthifolius Cass. . 211
Phaenopus acanthifolius DC. . . 211
vimineus DC. . . . 212
Phaeopappus Bois. . . 128
saxatilis Bois. 128
Phagnalon Cass. . . . 27
graecum B. et H. . . . 27
methanaeum Haussk. . 27
pumilum DC. 28
v. glabrum Bois. 29
v. tomentosum Bois. . . . 29
rupestre Clem. 28
rupestre (L.) 28
saxatile DC. 28
saxatile Heldr. 27
saxatile (L.) 27
Tenorii Bald. 28
Phelipaea aegyptiaca Walp. . . 446
caesia Grisb. 446
coerulea C A. Mey . 447
Fraasii Walp. 446
lavandulacea Reut. . . 446
Muteli Schultz . . . 445
v nana Reut. . . . 445
nana Rchb. . . . 445
purpurea Asch. . . . 447
pyramidalis Reut. . . 447
ramosa C. A. Mey . . 445
v. Muteli Bois. . . 445
v. nana Bois. . . . 445
Schultzii Schultz . . 447
stricta Mor. 447
Phillyrea Tourn. . . . 287

pag.
angustifolia L. 288
angustifolia Mass. 288
latifolia L. 288
latifolia S. et S. 288
media L. 287
obliqua W. 288
pedunculata Ch. et B. . . 288
spinosa Tausch 288
Phlomis L. 506
clandestina Ch. et B. . . . 497
cretica Presl 507
ferruginea Ten. 507
v. cretica Benth. 507
fruticosa L. 506
herba venti S. et S. . . . 509
v tomentosa Bois. . . . 509
lanata W 507
v biflora Hal. 508
lunarifolia S. et S. . . 509
microphylla Sieb. 507
parvifolia Presl 507
pungens W. 509
v. hispidula Hal. 509
samia L. 508
v. achaia Born. 508
v. graeca Born. 508
virens DC. 507
viscosa Poir. 507
Physalis L. 366
alkekengi L. 366
somnifera L. 365
Phyteuma canescens W. et K. . 276
dubia Friv. 278
Jacquini Sieb. 278
limonifolium S. et S. . . . 277
pinnatum L. 275
repandum S. et S. 277
rumelicum Grisb. 278
strictum Sims. 277
virgatum Ch. et B. . . . 277
Picnomon Lob. 119
acarna Cass. 119
Picridium Desf. 219
hispanicum Poir 214
intermedium Schultz . . . 214
ligulatum DC. 215
perenne Fraas 215
picroides Hal. 215
tingitanum Desf. . . . 214
vulgare Desf. 215
Picris L. 189
asplenioides L. 191
echioides L. 190
hieracioides S. et S. . . . 189

pag.
pauciflora W. 190
rhagadioloides Pers. . . . 189
spinulosa Bert. 189
Sprengeriana Lam. . . 189
stricta Jord. 189
Pinardia coronaria Less. 71
Plagius ageratifolius L'Her. . . 67
Podanthum Don. . . . 276
canescens Bois. 276
limonifolium Bois. . . 277
v. alpinum Bois. . . . 277
v. ramosum Haussk. . . 277
v. repandum (S. et S.) . . 277
Psaridis Heldr. 277
trichocalycinum Bois. . 276
v densiflorum Haussk. . . 276
Podospermum C. A. Mey 199
calcitrapifolium DC. . . . 200
canum C. A. Mey . . . 199
v. alpinum (Bois.) . . . 200
Jacquinianum Koch . . . 199
v subintegrum Bois. . . 199
laciniatum DC. 200
v calcitrapifolium Bois. . 200
octangulare DC. 199
pindicolum Haussk. . . . 199
villosum Stev 197
Polemoniaceae Vent. . . 301
Polemonium L. 301
coeruleum L. 301
Prasium L. 478
creticum Rchb. 478
majus L. 478
v. creticum (Rchb.) . . . 478
Prenanthes L. 208
acanthifolia W. 211
aconitifolia Sieb. 211
attica Spreng. 204
purpurea L. 208
ramosissima All. 212
viminea L. 212
Prismatocarpus falcatus Ten. . . 275
hybridus L'Her. . 275
pentagonus L'Her. 273
speculum L'Her. . 274
Prunella L. 496
hybrida Knaf 497
intermedia Link 497
laciniata L. 496
vulgaris L. 496
v. cretica Bois. 496
v. grandiflora Haussk. . . 496

pag.
v. Herculis Borb. 496
vulgaris × laciniata . . 497
Psilostemon creticum DC. . . . 320
Ptarmica abrotanoides Vis. . . 44
ageratifolia Nym. . . 41
aizoon Grisb. 41
ambrosiaca B. et H. . 43
Clavennae DC. 42
Fraasii Nym. 43
Pteronia chamaepeuce Spreng. . 126
Pterotheca bifida F. et M. . . . 177
nemausensis Rchb. . . 177
Ptilostemon appendiculatum Cass. 127
muticum Cass. . . 126
Pulicaria Gaertn. . . . 25
crispa Link 26
dentata DC. 26
dysenterica Gaertn. . . 26
v. dentata (S. et S.) . . . 26
v. microcephala Bois. . . 26
odora Rchb. 25
viscosa Cass. 24
vulgaris Gaertn. . . . 26
Pulmonaria L. 331
auriculata (Bois). . . . 331
mollis Bois. 331
officinalis S. et S. 331
Pyrethrum Gaertn. . . 68
cinereum Grisb. 69
corymbosum W. . . . 69
fuscatum W. 69
hybridum Guss. 70
v. discolor Guss. 70
Myconis Moench 70
parthenium Sm. . . . 68
tanacetum DC. 69
trichophyllum Grisb. . . . 66
Pyrola L. 284
clorantha Sw. 284
secunda L. 284
Pyrolaceae Lindl. . . . 284

R.

Ramonda Pers. 300
Heldreichii Benth. et Hok. 301
Nathaliae Panc. et Petr. 300
pyrenaica Pers. 301
Rhagadiolus Tourn. . 183
edulis Gaertn. 183
intermedius Ten. 183
stellatus W. 183
v. edulis DC. 183
v. hebaelenus DC. . . . 183
v. intermedius (Ten.) . . 182
v. leiocarpus DC. 183
v. leiolaenus DC. 183
Rhinanthus L. 443
crista galli L. 442
elephas L. 443
major v. parnassicus Bois. . 441
maximus W. 440
pubescens B. et H. . . . 441
trixago L. 440
viscosus Lam. 437
Rhynchocorys elephas Griesb. . 443
Rindera graeca B. et H. . . . 358
Robertia tenuiflora Bois. . . . 175
Rochelia Rchb. 361
disperma Stapf 361
stellulata Rchb. 361
Rodigia Spreng. 176
commutata Spreng. . . 176
Rosmarinus L. 491
officinalis L. 491
prostratus Mazz. 491

S.

Salvia L. 479
adulterina Haussk. 489
aethiopis L. 486
alpestris Haussk. 486
amplexicaulis Lam. . . 487
amplexicaulis × virgata Haussk. 489
argentea L. 486
v. alpina Heldr. 486
v. macedonica Grisb. . . 486
v. thessala Haussk. . . . 486
baccifera Ettl. 482
Barrelieri Ten. 488
calycina S. et S. . . . 479
candidissima Vahl . . 487
v. odorata Bald. 487
clandestina Ch. et B. . . 490
clandestina L. 490
Eichleriana Heldr. . . 481
Forskahlei L. 491
fragifera Ettl. 481

pag.
glutinosa L. 483
grandiflora Ettl. 481
horminum L. 485
v. comata Heldr. 485
v. comosula Heldr. . . . 484
v. viridis Heldr 484
inamoena Vahl 488
macedonica Hausskn. . . . 486
nemorosa L. 488
nilotica Vahl 491
odorata W. 487
officinalis L. 482
peloponnesiaca B. et H. 484
pinnata L. 491
pomifera L. 481
praecox Savi 490
pratensis S. et S. 488
ringens S. et S. . . . 483
sclarea L. 485
Sibthorpii Ch. et B. . . . 490
Sibthorpii S. et S. 489
silvestris L. 488
silvestris Mazz. 487
similata Haussk. . . . 490
Tenorii Spreng. 488
triloba L. 482
v. integrifolia Rev. . . . 482
verbenaca L. 490
v. clandestina (L.) 490
v. multifida Raul. 490
v. serotina Bois. 490
v. vernalis Bois. 490
verticillata L. 483
verticillata S. et S. . . . 484
villicaulis Borb. 487
virgata Ait. 489
viridis L. 484
viscosa Mazz. 489
Santolina alpina S. et S. . . . 56
anthemoides L. . . . 59
anthemoides Sieb. . . 56
chamaecyparissus L. . 51
incana Ch. et B. . . 56
maritima Sm. . . . 39
rigida S. et S. . . . 59
Satureia L. 550
approximata Friv. 551
canescens Link 549
capitata L. 557
cuneifolia Ten. 550
v. Wiedemanniana Bois. . 551
fasciculata Tausch. . . . 550
filiformis Sieb. 548
graeca Ch. et B. 548

pag.
graeca L. 548
juliana L. 546
montana L. 551
v. hellenica Heldr. . . . 551
v. pisidica (Wettst.) . . . 551
nervosa Desf. 547
olympica Hal. 551
parnassica Bois. , ... 551
parnassica Heldr. et Sart. 551
v. macrophylla Hal. . . . 552
v. stygia (Orph.) 552
plumosa Hampe 548
pygmaea Sieb. 551
spicata Sieb. 557
spinosa L. 552
v. parnassica Heldr. et Sart. 551
v. stygia Orph. 552
thymbra L. 550
virgata Vis. 550
Schizocalyx smyrnaeus Schule . 556
Scolymus L. 171
hispanicus L. 171
maculatus Fraas 171
maculatus L. 171
Scorodonia heteromalla Moench . 471
Scorzonera L. 195
ambigua DC. 197
angustifolia Friedr. . . . 196
araneosa Urv. 197
asperrima W. 186
austriaca W. 199
calcitrapifolia Vahl . . . 200
cretica W. 198
crocifolia S. et S. . . 196
Doriae Deg. et Bald. . 199
elongata W. 172
graminifolia Ch. et B. . . 196
hirsuta L. 199
v. Doriae Deg. et Bald. . 199
humilis L. 199
Jacquiniana Bois. 199
v. alpina Bois. 200
v. messeniaca Hal. . . . 199
laciniata Jacq. 199
lacinata L. 200
lanata M. B. 198
messeniaca Ch. et B. . . . 199
mollis M. B. 197
v. euboea Bois. 197
octangularis W 200
orientalis Dall. 215
parviflora Jacq. 196
picroides L. 215
purpurea Bois. 197

pag.
resedifolia Retz 200
rhodantha Haussk. . . 197
rosea W. et K. 196
tingitana L. 214
tomentosa L. 199
tomentosa Pall. 197
undulata Vahl. 197
villosa Friedr. 197
Scrofularia L. 348
aestivalis Grisb. . . . 399
alata Gilib. 401
aquatica L. 401
auriculata Raul. 400
auriculata Scop. 400
bicolor S. et S. 406
caesia S. et S. 403
canina L. 406
v. dolopica Haussk. . . . 406
v. pallida Haussk. . . . 406
v. ramosissima Haussk. . 406
chrysanthemifolia Ch. et B. 404
chrysanthemifolia W. . . . 406
cretica B. et H. . . . 400
Ehrharti Stev. 401
v Samaritanii B. et H. . 402
filicifolia Mill. 403
frutescens S. et S. 403
glandulosa W. et K. . . . 400
glauca S. et S. 403
grandidentata Ten. 400
heterophylla W. . . . 403
v. caesia Weiss 404
v. pinnatisecta Haussk. . 404
v. Urvilleana Weiss . . . 404
laciniata W. et K. . . 406
v. multifida Bois. 405
v. obtusiloba M. et R. . . 404
v. pindicola Form. . . . 405
v. puberula Hal. 405
laciniata Spreitz. 404
laxa B. et H. 402
lucida L. 402
v. filicifolia Benth. . . . 403
v. glauca (S. et S.) . . . 403
methanaea Haussk. . . . 403
micrantha Urv. 404
multifida W 405
Neesii Wirtg. 401
nodosa L. 401
oligantha B. et H. 400
Oliveriana Wydl. 404
peregrina L. 399
v. trachytica Haussk. . . 399
pindicola Haussk. 406
pyramidalis Wydl. 406
ramosissima Urv. 406
salicifolia Sieb. 401
Samaritanii B. et H. . 402
scoparia M. et R. 406
Scopolii Hoppe 400
v. grandidentata (Ten.) . . 400
v. oligantha Bois. . . . 400
scorodonia Pieri 399
Sibthorpiana Spreng. . . . 404
silvatica B. et H. 399
taygetea Bois. 405
tenuis Haussk. 404
Urvilleana Wydl. 404
Urvillei Wydl. 404
Scrofulariaceae Lindl. 370
Scutellaria L. 492
adenotricha B. et H. . . . 493
albida Weiss 495
alpina L. 492
Columnae All. 493
v. Sibthorpii Bald. . . . 493
v. Sibthorpii Benth. . . . 494
cretica L. 471
cretica Sieb 495
decumbens Sieb. 495
galericulata L. 495
hirta S. et S. 495
v. brachystegia Bois. . . 494
hirta Sieb. 495
Linnaeana Car. 493
orientalis L. 492
v. pinnatifida Rchb. . . . 492
peregrina L. 493
v. adenotricha B. et H. . 493
v. parnassica Bois. . . . 494
v. Sibthorpii B. et R. . . 494
v. Sibthorpii Hal. 494
rupestris B. et H. . . 494
Sibthorpii (Benth.) . . 494
Sedum aizoon S. et S. 278
Senecio L. 76
aquaticus S. et S. 84
arachnoideus Sieb. . . 80
Aucheri DC. 78
barbaraefolius Krock. . 84
barkhausiaefolius B. et H. 82
bicolor Bois. 77
cacaliaster v Gmelini Vis. . 81
coronopifolius Desf. . . 84
delubrorum Heldr. 86
erraticus Bert. 84
erucifolius L. 83
v. macrophyllus Form. . . 83

pag.
euboeus B. et H. . . . 79
foeniculaceus Ten. 86
fruticulosus S. et S. . . 82
graphalodes Sieb. . . . 78
Heldreichii Bois. . . . 79
Jacquinianus Rchb. . . 80
v. expansus B. et H. . . 81
Jacobaea L. 83
lanatus Scop. 80
lividus L. 86
macedonicus Grisb. . . 80
v. brachyphyllus Bois. . . 80
v. bracteatus (B. et O.) . 80
nebrodensis Raul. 82
nemorensis L. 81
v. dalmaticus Grisb. . . . 81
v. expansus (B. et H.) . . 81
v. Gmelini Bald. 81
Parnassi B. et H. . . 79
rupestris W. et K. . . 82
Scopolii Hoppe 80
squalidus Urv. 84
taygeteus B. et H. . . 78
tenuifolius Jacq. 83
thapsoides DC. 80
trilobus S. et S. 86
tymphresteus Helldr. . . . 85
vernalis W. et H. . . 85
v nigropunctatus Clem. . 85
vernalis × vulgaris . 86
viscosus L. 85
vulgaris L. 86
vulgaris × vernalis Zab. . 86
f. pseudovernalis Zab. 86
Weylii Vatke 86
Seriola L. 176
aethnensis L. 176
commutata Less. 176
cretensis L. 174
urens S. et S. 175
Serratula arvensis L. 118
attica Fraas 166
chamaepeuce L. . . . 126
crupina Vill. 166
fruticosa Lam. 100
leucocephala Poir. . . 112
Sesamum indicum L. 300
v. subindivisum DC. . . . 300
orientale L. 300
Sibthorpia L. 422
africana L. 422
europaea S. et S. 422
Sideritis L. 497

pag.
Boissieri Magn. 500
candicans Ait. 500
condensata B. et H. . . . 494
cretica Bois. 500
v condensata Bois. . . . 499
cretica S. et S. 499
elegans Murr. 500
euboea Heldr. 499
florida B. et H. 498
lanata L. 500
montana L. 500
v. comata Bois. 501
montana S. et S. 501
nigricans Pers. 500
peloponnesiaca B. et H. . . 498
perfoliata L. 497
pindicola Heldr. 497
purpurea Talb. 502
remota Urv. 501
Roeseri B. et H. . . . 499
v. lanceolata Hal. . . . 499
romana L. 501
v. purpurea Bald. 502
scardica Grisb. 498
v. pelia Hal. 498
scardica Heldr. 498
syriaca Ch. et B. 497
syriaca Fraas 499
syriaca L 500
v condensata B. et H. . . 499
taurica Heldr. 499
theezans B. et H. . . . 497
v. cyllenea Bois. 498
v. peloponnesiaca (B. et H.) 498
tragoriganum Lag. 502
Silybum Vaill. 124
marianum Gaertn. . . . 124
v. pygmaeum Orph. . . . 124
Siphonostegia Benth. et Hook. 442
syriaca Bois. 442
Solanaceae Juss. . . . 362
Solanum L. 369
citrullifolium R. Br. . . . 369
dulcamara L. 369
flavum Kit. 370
humile Berth. 370
melongena L. 370
miniatum Berth. 370
nigrum L. 370
v. flavum (Kit.) 370
v. humile (Berth.) . . . 370
v. miniatum (Berth.) . . 370

pag.
v. villosum L. 370
pseudocapsicum L. 370
sodomacum L 369
suffruticosum Schousb. 369
tuberosum L. 370
villosum Lam. 370
Solidago L. 17
graveolens Lam. 25
virgaurea L. 17
v. vestita Hal. 17
viscosa Lam. 35
Sonchus L. 205
arvensis S. et S. 206
asper Vill. 207
chondrilloides S. et S. . . 215
ciliatus Lam 206
divaricatus Desf. 215
glaucescens Jord. 207
graecus Reut. 207
lacerus Sieb. 205
laevis Vill. 206
maritimus L. 205
Nymani Tin. et Guss. . 206
v. versicolor Haussk. . . 206
oleraceus L. 206
v. asper L. 207
v. lacerus W. 206
v. laevis L. 206
palustris L. 205
picroides S. et S. 214
tenerrimus L. 205
v. heterophyllus Hal. . . 206
v. muriculatus Clem. . . 205
v. spinulosus Friedr. . . . 205
Specularia Heist. . . . 273
falcata DC. 275
hybrida DC. 275
pentagonia DC. . . . 273
speculum DC. 274
v. calycina DC. 274
v. racemosa Bois. 274
Stachys L. 515
abrotanifolia B. et H. . . . 576
acutifolia Ch. et B. . . 522
adenocalyx C. Koch . . . 530
alpina Form. 521
alpina L. 521
v. discolor Bois. 521
v. Reinerti Bald. 521
v. turcica Form. 521
angustifolia M. B. 526
annua L. 530
v. adenocalyx Haussk. . . 530

pag.
v. parnassica Heldr. et Sart. 530
arenaria Friedr. 526
argolica Bois. 527
arvensis L. 531
betonica Benth. 514
betonicaefolia Desf. . . . 530
candida Ch. et B. . . . 529
v. chrysantha Benth. . . 529
candida Ung. 527
canescens Ch. et B. . . 524
cassia Bois. 519
chrysantha B. et H. . . 529
cicinnata Friedr. 527
cretica Hal. 518
cretica L. 518
v. albiflora Hal. 519
v. condensata Haussk. . . 518
cretica Pieri 518
cretica Sieb. 519
dasyanthes Raf. 520
v alpina Heldr. 520
decumbens Pers. . . . 526
dolopica Form. 523
Freynii Haussk. . . . 523
germanica L. 520
v. Boissieri Briq. 518
v. Heldreichii (Bois.) . . 520
v. penicillata Bald. . . . 519
v. penicillata (Heldr. et Sart.) 520
v. stricta Form. 520
glutinosa L. 530
graeca B. et H. 522
Heldreichii Bois. 520
Heldreichii Form. 518
hirta Pieri 526
iberica M. B. 525
italica Müll. 518
v. albiflora Hal. 519
v. chondrostachys Heldr. . 519
lamioides Sieb. 531
lanata S. et S. 520
lusitanica Ch. et B. . . . 522
maritima L. 526
menthaefolia Vis. 524
messeniaca Bois. . . . 524
mollissima W. 526
mucronata Sieb. 529
orientalis L. 517
palaestina L. 531
palustris L. 523
Parolinii Vis. 524
penicillata Heldr. et Sart. . 520
pubescens Ten. 526

pag.
recta L. 525
Reinerti Heldr. 521
v. abbreviata Hal. . . . 521
salviaefolia Friedr. 518
salviaefolia Ten. 518
scardica Grisb. 515
scyronica Bois. 527
serbica Panc. 531
sericophylla Hal. . . . 519
Sieberi C. Koch 519
silvatica L. 522
v. pallida Haussk. . . . 522
spinosa L. 530
spinulosa S. et S. . . . 530
Spreitzenhoferi Heldr. . 529
Spruneri Bois. 528
Swainsoni Benth. . . . 527
v. argolica Bois. 527
v. scyronica Bois. . . . 527
Swainsoni Haussk. 527
tenuifolia Pall. . . 526
tetragona B. et H. . . 528
Tournefortii Ch. et B. . . 524
Tournefortii Poir. . . . 519
tymphaea Haussk. 521
v. lacmonica Haussk. . . 521
v. petiolata Haussk. . . . 521
virgata Ch. et B. . . . 525
viridis B. et H. . . . 523
Zuccarinii Benth. 525
Staehe na L. 100
arborescens L. 100
chamaepeuce L. 126
dubia L. 101
fruticosa L. 100
uniflosculosa S. et S. . 101
Styraceae DC. 285
Styrax Tourn. 285
officinalis L. 285
Symphyandra DC. . . 248
cretica DC. 248
samothracica Deg. 249
sporadum Hal. 248
Symphytum L. 330
anatolicum Bois. . . . 330
brochum Ch. et B. 231
bulbosum Schimp. . . . 330
macrolepis Gay. 331
officinale L. 330
officinale Pieri. 331

pag.
ottomanum Frid. . . . 331
tuberosum S. et S. . . . 331

T.

Tanacetum L. 69
annuum L. 69
balsamita L. 69
vulgare L. 69
Taraxacum Hall. . . . 201
erythrospermum Andrz. . . 202
gymnanthum Hal. 201
gymnanthum Heldr. . . . 203
gymnanthum Link. . . 203
Haussknechtii Uechtr. . 201
humifusum Heldr et Sart. . 203
laevigatum DC. . . . 202
v. polycephalum Raul. . . 202
v. scolopendrinum Heldr. . 202
v. stenolobum Heldr. . . 202
minimum Tod. 203
officinale Wigg. . . . 201
v. alpestre Ung. 202
v. alpinum (Hoppe) . . . 202
v. paludosum (Scop.) . . . 202
palustre DC. 202
vulgare Lam. 201
v. alpinum Bald. 202
v. pindicolum Bald. . . . 201
v. Steveni Bald. 202
Teucrium L. 469
Achaemenis Schreb. . . . 478
alpestre S. et S. . . . 477
v. laxior Spreitz. 477
v. major Bois. 477
brevifolium Schreb. . . 470
capitatum L. 478
capitatum S. et S. 478
chamaedrys L. 472
v. canescens Form. . . . 473
v. canum Form 473
v. glanduliferum Haussk. . 473
v. oxyodon Heldr. et Hal. . 474
v. revolutum Form. . . . 473
chamaepitys L. 467
chium Poir. 468
creticum L. 470
creticum Lam. 470
creticum Scheele 471
cuneifolium S. et S. . . 476
cuspidatum Cel. . . . 471

pag.
divaricatum Sieb. . . . 474
v. graecum (Cel.) 474
flavum L. 473
v. glaucum Haussk. . . . 474
v. humilior Ch. et B. . . 474
v. leiophyllum Cel. . . . 474
v. purpureum Benth. . . 474
graecum Cel. 474
graecum Hal. 474
v. minor Heldr. 474
Halácsyanum Heldr. . . 470
v. laxum Hal. 471
hyssopifolium Schreb. . . . 470
iva L. 467
lamiifolium Urv. 471
lanuginosum H. et S. . . . 472
lucidum S. et S. 474
marum L. 475
massiliense L. 471
microphyllum Desf. . . 475
montanum L. 475
v. hirsutum Bois. 476
v. parnassicum Cel. . . . 476
montanum S. et S. . . . 476
pannonicum Kern. 476
parnassicum Cel. 476
polium L. 477
v. Achaemenis (Schreb.) . 478
v. angustifolium Benth. . 478
v. angustissimum Cel. . . 478
v. bombycinum Heldr. . . 477
v. capitatum (L.) 478
v. gnaphalodes Raul. . . 477
v. intermedium Cel. . . . 478
v. lanuginosum Cel. . . . 477
v. pseudohyssopus (Schreb.) 477
v. purpurascens Benth. . . 478
v. roseum Bois. 478
v. virescens Hal. 477
prostratum Hal. 476
pseudohyssopus Schreb. . . 477
quadratulum S. et S. . . . 475
ramosissimum Sieb. . . . 475
rosmarinifolium Lam. . . . 470
saxatile Sieb. 477
scordioides Schreb. . . 472
scordium L. 472
scordium Pieri 472
scorodonia L. 471
Sieberi Cel. 474
silvestre Lam. 471
supinum L. 475
tenthrion Schreb. 477
Thrincia Roth 184
hirta Roth 184

pag.
Oliverii DC. 184
tuberosa DC. 184
v. Olivieri Haussk. . . . 184
Thymbra L. 557
capitata Grisb. 557
v. albospinosa (Balv.) . . 558
hirsuta Pers. 550
spicata L. 557
Thymus L. 558
acinos Ch. et B. 544
acinos L. 546
alpinus L. 543
angustifolius v. Chaubardi B. et H. 564
atticus Cel. 562
Billardieri Bois. 567
Boissieri Hal. 559
v. ciliatopubescens Hal. . 559
calamintha Fraas 542
calamintha Sm. 541
capitatus H. et L. 557
v. albospinosus Bald. . . 558
Chaubardi B. et H. . . . 564
Chaubardi Hal. 563
cherlerioides Vis. 559
ciliatopubescens Hal. . 559
creticus Brot 557
dolopicus Form. 565
euboeus Hal. 560
exiguus S. et S. 546
glabratus Ch. et B. . . . 563
graveolens M. B. 546
graveolens S. et S. . . . 566
heterotrichus Grisb. . . . 565
hirsutus Bois. 559
v. humillimus Bald. . . . 557
hirsutus Hal. 559
hirsutus M. B. 559
hirsutus Sint. 559
hirsutus Spreitz 561
hirtus Sieb. 543
humillimus Cel. 559
incanus S. et S. 542
lanceolatus S. et S. . . . 563
lanuginosus Ch. et B. . . 542
lanuginosus Mill. . . . 565
v. Marschallianus (W.) . . 565
leucadicus (Rchb.) . . . 562
leucotrichus Hal. . . . 561
v. Degenii Hal. 561
longicaulis Hal. 564
Marschallianus W. . . . 565
mastichina S. et S. . . . 547
microphyllus Urv. 549
montanus W. et K. . . 566

pag.
nepeta Sm. 542
ocheus Heldr. et Sart. . 565
patavinus Jacq. 545
parnassicus Hal. . . . 561
paronychioides Haussk. . . 563
pindicolus Form. 564
pulegioides S. et S. . . . 566
revolutus Cel. 561
v. creticus Deg. 561
serpyllum Clem. 564
v. angustifolius Bois. . . 563
v. Chaubardi Bois. . . . 564
v. latifolius Bois. 566
v. Marschallianus Bois. . . 565
v. ocheus Bois. 565
Sibthorpii Benth. . . . 563
striatus Bois. 562
striatus Raul. 561
striatus Vahl. 563
v. leucadicus Rchb. . . . 562
suaveolens S. et S. . . . 544
teucrioides B. et Spr. . 566
v. tymphaeus Form. . . . 566
tragoriganum L. 550
tymphresteus Hal. . . 560
villosus S. et S. 567
vulgaris Pieri. 557
vulgaris S. et S. 561
zygis S. et S. 562
Tolpis Gaertn. 181
altissima Pers. 182
barbata S. et S. 181
cretica Sieb. 182
quadriaristata Biv. 181
quadriaristata S. et S. . . 182
umbellata Bert. 181
virgata Bert. 182
Tragopogon L. 192
asper L. 191
australis Jord. 192
balcanicus Vel. 193
crocifolius L. 193
Cupani Guss. 192
Dalechampii L. 191
floccosus W. et K. . . 195
lanatus W. 198
longifolius Heldr. et Sart. 195
majus Jacq. 195
orientalis L. 194
picroides L. 191
porrifollus L. 192
v. brachyphyllus Bois. . . 192
pratensis L. 194
v. orientalis (L.) 194

pag.
Samaritanii Heldr. et Sart. 193
Tommasinii Schultz . . . 194
Trachelium asperuloides B. et O. 278
Jacquini Bois. . . . 278
rumelianum Hampe . 278
rumelicum Bois. . . 278
Trachystemon Don. . 320
creticum Don. 320
Trichocrepis bifida Vis. 177
Tripolium vulgare Nees. 16
Trixago apula Stev. 440
latifolia Rchb. 436
viscosa Rchb. 437
Tussilago L. 11
farfara L. 11
hybrida L. 11
petasites L. 11
Tyrimnus Cass. 125
leucographus Cass. . . 125

U.

Urospermum Juss. . . 191
asperum DC. 191
Dalechampii DC. 191
echioides Fraas 191
picroides Def. 191
v. asperum DC. 191

V.

Verbascum L. 372
acutifolium Hal. . . . 385
acutifolium × blattaria . 394
Adeliae Heldr. 390
adenotrichum Hal. . . 381
agrimoniodes Deg. et Bb. 381
ambiguus Haussk. 391
ambracicum Hal. 378
arcturus L. 395
auriculatum Mazz. 386
auriculatum S. et S. . . . 386
australe Schrad.. 375
banaticum Roch. . . . 388
blattaria L. 393
v. blattariforme (Grisb.) . 394
blattaria × phlomoides Hssk. 394
blattaria × Sartorii Haussk. . 394
blattariforme Grisb. . . . 394
blattariforme × Sartorii Hssk. 394
Boerhavii L. 384

pag.
calaurium Heldr. 383
candidissimum DC. 386
Chaixi Hal. 392
Chaixi Vill. 392
coenobitarum Hssk. et Heldr. 377
delphicum B. et H. . . 355
densiflorum Bert. . . . 374
v. thapsiforme Vis. . . . 375
dervichorum Hssk. et Heldr. 389
epirotum Hal. 380
epixanthinum B. et H. 379
v. Samaritanii Bois. . . . 379
epixanthinum Hal. 380
erraticum Haussk. 383
ferrugineum Andr. 394
flagriforme Haussk. . . . 394
floccosum W. et K. . . . 387
floccosum × phlomoides Hssk. 376
foetidum B. et H. . . 379
gloeotrichum Haussk. et Heldr. 385
graecum Heldr. et Sart. 393
v. aetolicum Bois 393
v. Zuccarinii Bois. . . . 393
Guicciardii B. et H. . . 377
Guicciardii × sinuatum Hal. 378
Haussknechtii Heldr. . 389
Haussknechtii × rigidum Hk. 389
Haussknechtii × sinuatum Haussk. 391
Heldreichii Bois. . . . 388
Heldreichii × phlomoides Hk. 377
Heldreichii × pulverulentum Haussk. 387
Heldreichii × Sartorii Haussk. et H. 377
Heldreichii × sinuatum Haussk. 391
hybridum Brot. 391
hypoleucum B. et H. . 392
leucophyllum Grisb. . . 389
v. integrifolium Haussk. . 389
limnense Fraas 390
longifolium Ten. . . . 377
lychnitis L 388
macilentum Hal. 394
macrurum Ten. 375
malacotrichum B. et H. 378
mallophorum B. et H. . 390
mallophorum × thyrsoideum Hal. 392

pag.
megaphlomos B. et H. . 386
v. macrophyllum Hal. . . 387
v. microcalyx Hal. . . . 387
meteoricum Haussk. . . 391
mucronatum Lam. . . 386
nigrum L. 392
nigrum Mazz. 392
nigrum × mallophorum B. et H. 392
nigrum × pulverulentum Hk. 387
nigrum × sinuatum Bois. . . 392
paradoxum Haussk. . . . 383
parallelum Haussk 394
pelium Hal. 382
permixtum Hal. 387
perviscosum Borb. 385
petrophilum Hal. 394
phalereum Haussk. 384
phlomoides L. 375
v. Sartorii Bois. 376
phlomoides S. et S. . . . 376
phlomoides × rigidum Hssk. 376
phlomoides × sinuatum Hssk. 377
phoeniceum L. 394
pindicolum Fr. et Sint. . . 380
pinnatifidum Vahl. . . 384
v. dentatum Haussk. . . . 384
pinnatifidum × plicatum Hssk. 383
pinnatifidum × sinuatum Haussk. 384
plicatum S. et S. 382
plicatum × sinuatum Haussk. 383
pseudoflagriforme Haussk. . 394
pseudosinuatum Haussk.. . 377
pulverulentum Hal. . . . 385
pulverulentum Vill. . . 387
pulverulentum × Heldreichii Hal. 387
pulverulentum × Sartorii Haussk. 376
pulverulentum × sinuatum Hal. 391
pulverulentum × thyrsoideum Hal. 387
Reiseri Hal. 382
repandum Grisb. 394
rigidum × Haussknechtii Hal. 389
rigidum × Sartorii Haussk. . 376
Samaritanii Hal. 381
Samaritanii Haussk. . . . 380
Samaritanii Heldr. 379
Sartorii B. et H. . . . 376
Sartorii × blattaria Hal. 394

pag.
Sartorii × Heldreichii Hal. 377
Sartorii × pulverulentum Hal. 376
Sartorii × rigidum Hal. 376
Sartorii × sinuatum Hk. 377
Schraderi Mey. 374
Schraderi Raul. 375
sinuatum L. 391
semirigidum Haussk. . . . 376
speciosum Schrad. 386
v. megaphlomos B. et H. . 386
spinosum L. 395
sterile Haussk. 387
subphlomoides Haussk. . . 377
taygeteum Hal. 380
thapsiforme Bois. 376
thapsiforme Schrad. . . . 375
thapsus Dall. 375
thapsus L. 374
thessalum Haussk. 391
thyrsoideum Host. . . 392
triste S. et S. 394
tymphaeum Fr. et Sint. . . 380
undulatum Lam. . . . 382
v. rigidum (B. et H.) . . 383
undulatum Sieb. 391
undulatum × pinnatifidum Hal. 383
undulatum × sinuatum Hal. 383
v. sinuatiforme Hal. . . . 383
v. undulatiforme Hal. . . 383
viscidulum Fr. et Sint. . . 381
viscidulum Mazz. 394
Verbena L. 462
adulterina Haussk. 463
Haussknechtii Heldr. . . . 463
nodiflora L. 463
officinalis L. 462
v. grandiflora Haussk. . . 462
officinalis × supina . . 463
supina L. 462
Verbenaceae Juss. . . . 461
Veronica L. 423
acinifolia L. 431
agrestis L. 435
v. byzantina S. et S. . . 434
alba Mazz. 435
anagallis L. 429
v. anagalliformis Bor. . . 429
anagalloides Guss. . . 429
aphylla L. 428
arvensis L. 432
australis Schrad. 425
austriaca Bois. 428
austriaca L. 427
v teucrioides (B. et H.) . 427
balcanica Vel. 430
beccabunga L. 429
Buxbaumii Ten. 434
byzantina Mazz. 435
chamaedryoides Ch. et B. . 426
chamaedrys L. 426
v. pilosa Benth. 426
v. pindica Haussk. . . . 426
Chaubardi B. et R. . . 433
Chaubardi Spreitz. 433
cretica Pall 425
cymbalaria Bod. 435
v. panormitana Tin. . . . 436
dentata Schm. 427
depauperata W. et K. . . . 428
didyma Ten. 435
erinoides B. et Spr. . . 431
filiformis DC. 435
Formanekii Heldr. 434
glauca Ung. 433
glauca S. et S. 433
graeca Sprun. 433
graminea Mazz. 429
hederifolia L. 436
v triloba (Op.) 436
Jacquini Baumg. 428
latifolia Jacq. 427
latifolia L. 426
longifolia L. 424
multifida L. 428
officinalis L. 426
v. glandulosa Form. . . . 426
pectinata L. 436
peduncularis Ch. et B. . . 433
peloponnesiaca B. et O. . 433
persica Poir. 434
polita Fr. 435
praecox All. 434
prostrata v. filicaulis Hal. . 427
pseudochamaedrys Jacq. . . 427
Sartoriana B. et H. . . 432
Schmidtii R. et Sch. . . . 427
scutellata L. 428
serpyllifolia L. 430
v. glandulosa Haussk. . . 430
v. oxya Form. 430
spicata L. 425

pag.
spuria S. et S. 424
syriaca R. et Sch. 436
teucrioides B. et H. . . . 427
teucrium Jacq. 427
teucrium L. 427
v. glandulosa Form. . . . 427
thessala Form. 434
thessalica Benth. . . . 431
thymifolia S. et S. . . 425
Tournefortii Gm. 434
triloba Op.. 436
triphyllos L. 434
tymphrestea B. et Spr. . . 425
urticaefolia Jacq. 426
verna L. 431
Villarsia nymphoides Vent. . . 300
Vinca L. 294
herbacea W. et K. : . 294
major L. 294
minor L. 294
mixta Vel. 295
Vincetoxicum Moench. 290
canescens Dec. 290
fuscatum Bois 292
fuscatum Hal. 291
laxum Sieb. 292
luteum Sieb. 290
nigrum Moench. 291
nivale B. et H. 291
v. undulatum Heldr. . . . 292
speciosum B. et Spr. . 291
pag.
triste Grisb. 291
Vitex L. 461
agnus castus L. . . . 461

W.

Wahlenbergia graminifolia DC. . 279
Withania Pauq. . . . 365
somnifera Mor. 365

X.

Xanthium L. 246
antiquorum Wallr. 246
spinosum L. 247
strumarium L. 246
v. antiquorum Bois. . . . 246
Xeranthemum L. . . 93
annuum L. 93
v. inapertum L. 93
cylindraceum S. et S. . 93
erectum Presl. 93
inapertum W. 93

Z.

Zacintha Gaertn. . . . 213
verrucosa Gaertn. 213
Zapania nodiflora Lam. 463
Zizyphora L. 491
capitata L. 491
Zollikoferia chondrilloides M. et R. 215

NAT.HIS
send to q

Stanford University Libraries
3 6105 015 853 307

DATE DUE			

STANFORD UNIVERSITY LIBRARIES
STANFORD, CALIFORNIA 94305